Contents lists available at SciVerse ScienceDirect

Chemistry and Physics of Lipids

ELSEVIER



journal homepage: www.elsevier.com/locate/chemphyslip

Steric analysis of epoxyalcohol and trihydroxy derivatives of 9-hydroperoxy-linoleic acid from hematin and enzymatic synthesis

Christopher P. Thomas^{a, c}, William E. Boeglin^a, Yoel Garcia-Diaz^b, Valerie B. O'Donnell^c, Alan R. Brash^{a,*}

^a Department of Pharmacology, Vanderbilt Institute of Chemical Biology, Vanderbilt University, Nashville, TN 37232, USA

^b Department of Chemistry, Vanderbilt Institute of Chemical Biology, Vanderbilt University, Nashville, TN 37232, USA

^c Institute of Infection & Immunity, Tenovus Building, UHW, Cardiff University, Heath Park, Cardiff CF14 4XN, United Kingdom

ARTICLE INFO

Article history: Received 31 October 2012 Received in revised form 6 December 2012 Accepted 6 January 2013 Available online xxx

Keywords: 9-HPODE Trihydroxyoctadecenoic acid Linoleic acid Epoxyalcohol synthase Epoxide hydrolase Hematin

ABSTRACT

We characterize the allylic epoxyalcohols and their trihydroxy hydrolysis products generated from 9*R*and 9*S*-hydroperoxy-octadecenoic acid (HPODE) under non-enzymatic conditions, reaction with hematin and subsequent acid hydrolysis, and enzymatic conditions, incubation with *Beta vulgaris* containing a hydroperoxide isomerase and epoxide hydrolase. The products were resolved by HPLC and the regio and stereo-chemistry of the transformations were determined through a combination of ¹H NMR and GC–MS analysis of dimethoxypropane derivatives. Four trihydroxy isomers were identified upon mild acid hydrolysis of 9*S*,10*S*-*trans*-epoxy-11*E*-13*S*-hydroxyoctadecenoate: 9*S*,10*R*,13*S*, 9*S*,12*R*,13*S*, 9*S*,10*S*,13*S* and 9*S*,12*S*,13*S*-trihydroxy-octadecenoic acids, in the ratio 40:26:22:12. We also identified a prominent δ-ketol rearrangement product from the hydrolysis as mainly the 9-hydroxy-10*E*-13-oxo isomer. Short incubation (5 min) of 9*R*- and 9*S*-HPODE with *B. vulgaris* extract yielded the 9*R*- and 9*S*-hydroxy-10*E*-12*R*,13*S*-*cis*-epoxy products respectively. Longer incubation (60 min) gave one specific hydrolysis product via epoxide hydrolase, the 9*R*/*S*,12*S*,13*S*-trihydroxyoctadecenoate. These studies provide a practical approach for the isolation and characterization of allylic epoxy alcohol and trihydroxy products using a combination of HPLC, GC–MS and ¹H NMR.

© 2013 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

The oxygenation at C9 of linoleic acid is the gateway to a diverse collection of oxylipin products (Andreou et al., 2009). C9 oxygenation in plants is catalyzed by 9-lipoxygenases (9-LOX), well-represented among the LOX genes (Feussner and Wasternack, 2002). Linoleate 9-lipoxygenases are also known in prokaryotes (Andreou et al., 2008; Gao et al., 2010; Zheng et al., 2008), and are a catalytic activity of arachidonate 12*R*-LOX in higher animals (Meruvu et al., 2005; Siebert et al., 2001; Zheng et al., 2011). Among the many possible transformations of the resulting 9-hydroperoxy-octadecenoic acid (9-HPODE), in keeping with other fatty acid hydroperoxides are rearrangements to epoxy-hydroxy derivatives

E-mail address: alan.brash@vanderbilt.edu (A.R. Brash).

(fatty acid epoxyalcohols), with subsequent hydrolysis producing a collection of trihydroxy-octadecenoates.

The non-enzymatic transformation of fatty acid hydroperoxides to epoxyalcohols is catalyzed by heme or transition metals, resulting in a mixture of isomers, as analyzed earlier utilizing 13-HPODE (Dix and Marnett, 1985; Gardner et al., 1974; Hamberg, 1975). The present study aims to address the heme transformations with 9-HPODE as starting material. Enzymatic conversion results in regioand stereo-specific formation of epoxyalcohols. Lipoxygenases can also catalyze epoxyalcohol synthesis (Garssen et al., 1976; Nigam et al., 2004; Pace-Asciak et al., 1995; Yu et al., 2003), the transformation being favored under anaerobic conditions (Zheng and Brash, 2010). Heme and lipoxygenase catalysis involves formation of alkoxyl and epoxyallylic radical intermediates, whereas oxygen transfer from the hydroperoxide with epoxidation of a remote double bond is effected by other enzymes. These include peroxygenase (Blée et al., 1993; Hamberg and Hamberg, 1996), catalase-related hemoproteins (Gao et al., 2009; Niisuke et al., 2009), as well as putative P450-like enzymes that remain to be characterized, for example in the fish fungus Saprolegnia parasitica (Hamberg et al., 1986), potato leaves (Hamberg, 1999), and beetroot (Hamberg and Olsson, 2011).

Trihydroxy hydrolysis products are readily formed from fatty acid epoxyalcohols, and in the case of potential 9-HPODE-derived

Abbreviations: CD, circular dichroism; COSY, correlation spectroscopy; DCM, dichloromethane; DMP, dimethoxypropane; GC–MS, gas chromatography–mass spectrometry; H(P)ODE, hydro(pero)xyoctadecadienoic acid; IPA, isopropyl alcohol; LOX, lipoxygenase; RP-HPLC, reversed-phase high pressure liquid chromatography; SP-HPLC, straight-phase high pressure liquid chromatography; UV, ultraviolet.

^{*} Corresponding author at: Department of Pharmacology, Vanderbilt University Medical Center, Nashville, TN 37232-6602, USA. Tel.: +1 615 343 4495; fax: +1 615 322 4707

^{0009-3084/\$ –} see front matter © 2013 Elsevier Ireland Ltd. All rights reserved. http://dx.doi.org/10.1016/j.chemphyslip.2013.01.002

products, these triols are reported in such diverse settings as; an in-vivo pathogen response in potato leaves (Göbel et al., 2002; Hamberg, 1999), in brewing, where their formation contributes to the bitter flavor of beer (Garbe et al., 2005; Hamberg, 1991b), as products in mammalian blood vessels and leukocytes (Claeys et al., 1985; Funk and Powell, 1985), and in preserved meat as a marker of lipid peroxidation (Püssa et al., 2009). The adjuvant activity of 9S,12S,13S-trihydroxyoctadecenoic acid (commonly known as pinellic acid) and its isomers in the administration of flu vaccine has also been reported (Shirahata et al., 2003). In addition, our interests in the transformations of linoleate and its esters through LOX activity in mammalian epidermis (Zheng et al., 2011) have focused our attention on the preparation and characterization of 9-HPODE-derived epoxyalcohols and the resulting triols. There are effective methods for their analysis in the literature, although the separations cannot be related to typical HPLC methodology, which was one impetus for our conducting the present study. Hamberg (1991a) made assignments based on GLC and TLC mobilities coupled with mass spectrometry and chemical degradation (oxidative ozonolysis, followed by GLC analysis of the fragments). Shirahata et al. (2006) prepared all possible stereo isomers of pinellic acid by total synthesis and compared their spectral properties by NMR. The use of an acetonide derivative to capture the vicinal 12,13-diol in the 9,12,13-triol is part of this procedure and has been applied in other studies to identify single naturally occurring fatty acid triols by NMR (Bruno et al., 1992; Cardellina and Moore, 1980). We make use of this procedure in our NMR assignments herein.

Accordingly, the objectives of the current study were (i) to develop a practical method for production and purification of allylic epoxyalcohols via hematin treatment of 9-hydroperoxy-linoleic acid, (ii) to characterize the HPLC separation of the trihydroxyoctadecenoates resulting from hydrolysis of the allylic alcohols, (iii) to utilize a GC–MS and NMR approach for the assignment of the trihydroxy-octadecenoate regio- and stereo-chemistry, (iv) to compare the enzymatic synthesis, isomeric distribution and chirality of epoxyalcohols and triols from 9*R*- and 9*S*-hydroperoxy-linoleic acid in *Beta vulgaris*, a source of epoxyalcohol synthase (Hamberg and Olsson, 2011), and (v) to characterize specific delta-ketols as products from the mild acid treatment of allylic *trans*-epoxyalcohols.

2. Experimental

2.1. Generation of 9R-HPODE and 9S-HPODE substrates

2.1.1. 9R-HPODE

Linoleic acid (10 mg) in 0.5 ml ethanol was stirred into 200 ml 50 mM Tris pH 7.5, 150 mM NaCl, 20 mM CHAPS with a strong oxygen stream on the surface of the solution. *Anabaena* LOX enzyme (a linoleate 9*R*-LOX, (Zheng et al., 2008)) was added as 150 μ l of a 20 mg/ml solution; reaction was complete in 2 min as evidenced by UV analysis of the conjugated diene chromophore of the product. The solution was then acidified to pH 4.0 and extracted with ethyl acetate, washed twice with half volume of water, taken to dryness and redissolved in a small volume of methanol for storage prior to HPLC. The 9*R*-HPODE was purified by semi-preparative SP-HPLC using a Beckman Ultrasphere silica 10 μ m column, with isocratic hexane/IPA/acetic acid (100:2:0.1), a flow rate of 4 ml/min, with UV monitoring at 235 nm.

2.1.2. 9S-HPODE

9S-HPODE was prepared using the linoleate 9S-LOX in potato (Galliard and Phillips, 1971). Yukon Gold potatoes were diced and blended with 255 ml water. To eliminate starch, the solution was then stirred with 250 µl Tween 80, 320 µl Viscozyme L (Sigma) and 160 µl Amyloglucosidase (from *Aspergillus niger* 300 units/ml,

Sigma) for 1 h at room temperature. After a low speed centrifugation (500 rpm, 10 min) the supernatant was poured through 2 layers of cheesecloth into a graduated cylinder. Sodium acetate (1 M) pH 5.5 was then added to 10% volume. The resulting enzyme solution was then suspended in a water bath at 15 °C. Linoleic acid in ethanol (1 g) was added drop wise with a strong stream of oxygen blowing on the surface of the solution and the reaction stirred for 90 min. The resulting solution was acidified to pH 4.0 and extracted with dichloromethane (DCM). The DCM was then washed twice with half volume of H₂O and taken to dryness. The 9S-HPODE was purified using SP-HPLC with a Beckman Ultrasphere silica 10 μ m column with an isocratic mobile phase of hexane/IPA/acetic acid (100:2:0.1).

2.2. Reaction of 9S-HPODE with hematin

A hematin solution (5 mg/ml) was prepared by grinding the crystals to a fine powder and stirring with 0.1 M K₂HPO₄ adjusted to pH 11.0 with NaOH. 9S-HPODE (20 mg in 0.5 ml ethanol) was mixed with 0.1 M K₂HPO₄ (20 ml, pH 8.5), 1 ml of the hematin solution was added and the reaction left for 15 min at room temperature. The sample was cooled on ice, two volumes of ice-cold DCM were added, then a predetermined mix of 1 M KH₂PO₄ (5 ml) and 1 M HCl solution, sufficient to bring the pH to ~4.5, followed immediately by vigorous mixing of the phases. The aqueous layer was removed and the DCM washed twice with ice-cold water. The DCM was transferred to a fresh vial, dried under N₂, and the sample stored at -20 °C in DCM:MeOH (2:1, v/v) prior to initial purification on an open-bed silica Bond-Elut cartridge.

2.2.1. Silica cartridge purification of extract

The extract from the hematin reaction was pre-dissolved in ethyl acetate (2 ml) then hexane (8 ml) was added and the sample immediately applied to a 1 g Bond-Elut silica cartridge preconditioned with 25% ethyl acetate in hexane. A further 10 ml rinse of the sample vial with ethyl acetate (25%) in hexane was added and followed by a further 10 ml 25% ethyl acetate. The products were then eluted with 100% ethyl acetate (10 ml), and the sample was dried under N₂ and stored for analysis in DCM:MeOH, 2:1.

2.2.2. HPLC purification of hematin products

The range of products generated from the hematin reaction was analyzed on an Agilent 1100 series HPLC using a Thomson Advantage Silica 150 Å 5 μ column (250 mm × 4.6 mm) an isocratic solvent of hexane:IPA:acetic acid (100:2:0.02), a flow rate of 1 ml/min, with diode array detection at wavelengths of 205, 220, 235 and 270 nm.

2.3. HPLC separation of trihydroxy-octadecenoate products from allylic epoxyalcohol hydrolysis

2.3.1. Acid hydrolysis of

9S,10S-trans-epoxy-11E-13S-hydroxyoctadecenoate

To the allylic epoxide dissolved in acetonitrile (5 ml), 1% aqueous acetic acid (20 ml) was added and mixed thoroughly. After standing at room temperature for 30 min, the hydrolysis products were extracted in a C18 Bond Elute cartridge, washed with two volumes of H_2O and eluted with MeOH.

2.3.2. Purification of trihydroxy-octadecenoate products

Initial RP-HPLC analysis employed a Waters Symmetry C18 column (250 mm \times 4.6 mm) eluted isocratically with methanol/water/acetic acid (80/20/0.01) at 1 ml/min. The 205 nm absorbing peaks in the triol region of the chromatogram at \sim 4 min retention time were pooled and subsequently resolved using a Chiralpak AD column (250 mm \times 4.6 mm) with an isocratic mobile

phase of hexane/methanol/ethanol/acetic acid (86.5/4.5/9/0.1) run at 1 ml/min. The peak designated as II was further separated using a Chiralpak OD-H column (250 mm × 4.6 mm) with a mobile phase of hexane/ethanol/acetic acid, (97.5/2.5/0.1) run at 1 ml/min.

2.4. Assignment of regio- and stereo-chemistry of trihydroxy-octadecenoates via ¹H NMR and GC-MS analysis

Methyl esters were prepared by dissolving the products in MeOH (typically 10 µl), followed by addition of several drops of ethereal diazomethane ($\sim 100 \,\mu$ l); after mixing and standing for 30 s on ice the sample is taken to dryness under a stream of nitrogen. Methyl esters of the triols derived from hydrolysis of methyl 9S,10S-trans-epoxy-11E-13S-hydroxyoctadecenoate were purified as a group by SP-HPLC using a Thomson Advantage Silica column ($25 \text{ cm} \times 0.46 \text{ cm}$), a solvent of hexane/IPA (90:10, by vol.) run at 1 ml/min, with UV detection at 205 nm and a retention time of 9-10 min. Acetonide derivatives were prepared by addition of DMP:acetone (1:1) containing 1 mM p-toluene sulfonic acid to the dry methyl ester derivatives (reaction is complete within 5 min at room temperature); the acid was then neutralized using an equal volume of triethylamine (1.5 mM) in DCM and the sample taken to dryness. Individual methyl ester DMP derivatives were purified by HPLC using a Thomson Scientific silica column $(25 \times 0.46 \text{ cm})$, a solvent of hexane/IPA (97:3, v/v), and a flow rate of 1 ml/min with UV detection at 205 nm; the methyl ester DMP derivatives eluted with retention times close to 5 min. Resolution of mixtures of triol methyl ester DMP derivatives was achieved using a solvent of hexane/IPA (100:1, by vol.).

GC–MS analyses were run using an Agilent/J&W DB-5MS column ($25 \text{ m} \times 0.2 \text{ mm}$, $0.33 \mu \text{m}$ film) in a ThermoFinnigan DSQ instrument operated in El mode (ion source temperature $250 \degree$ C, electron energy 70 eV) set to full scans from m/z 50 to 650.

 1H NMR spectra of the DMP derivatized trihydroxy methyl esters were recorded in C₆D₆ (150 μ l solvent in a 3 mm tube) at 25 °C using a Bruker 600 MHz instrument equipped with a 5 mm Z-gradient TCI Cryo-probe and using pulse program zg with parameter settings including a pulse length 8.50 μ s, time domain 32768, and a relaxation time of 2 s.

2.5. Enzymatic synthesis of epoxyalcohols and triols from 9S/R-HPODE with B. vulgaris extract

A single beet was peeled and diced, 100 g was added to ice cold 0.1 M potassium phosphate pH 6.7 (600 ml) and blended using a Polytron homogenizer for 3×30 s or until a smooth consistency was obtained. The solution was filtered through 2 layers of gauze and used directly for incubation with HPODE. Metabolizing activity was monitored by disappearance of the UV absorbance at 235 nm using 130 µl of homogenate in 870 µl buffer and 50 µM HPODE substrate as described (Hamberg and Olsson, 2011). For larger scale incubations, 4 mg of either 9R- or 9S-HPODE in 200 µl methanol was added to the beetroot homogenate (26 ml) with stirring at room temperature for 1 h. The incubations were then acidified to pH 3.0 with 1N HCl and extracted with 2 volumes of ethyl acetate. The ethyl acetate was washed twice with half a volume of water, taken to dryness, and the recovered products resuspended in isopropanol/chloroform (1:2). The sample was then loaded onto a Waters Sep-Pak Plus NH₂ cartridge. The reaction products were eluted in 15 mL ethyl acetate/acetic acid (98:2). HPLC purification was then performed using a Waters Symmetry C18 column with an isocratic 1 ml/min flow of methanol/water/acetic acid (80:20:0.01). Generation of the epoxyalcohol product was carried out in a similar way with the following deviations: (1) reaction time was reduced to 5 min to prevent conversion by epoxide hydrolase and (2) the unstable epoxyalcohol products were extracted with diethyl ether eliminating the requirement for the Sep-Pak chromatography.

2.5.1. Assignment of regio- and stereo-chemistry

¹H NMR and GC–MS analysis was carried out as Section 2.4 for triol analysis. Deuterated benzene was used as solvent (locked on the 7.15 ppm signal of residual C_6H_6), except for underivatized triols and their methyl esters, recorded in deuterated methanol (locked on 3.34 ppm). Epoxyalcohol methyl esters were converted to the trimethylsilyl derivative by treatment with BSTFA for 1 h (or overnight) at room temperature.

For CD analysis, approximately 50 μ g of epoxyalcohol methyl ester was converted to the 9-O-benzoate ester (1 μ l benzoyl chloride, ~1 mg dimethylaminopyridine in 30 μ l pyridine, 15 min at room temperature) and repurified by RP-HPLC (MeOH/H₂O, 95:5 solvent, flow rate 1 ml/min, retention time approximately 7 min) as described (Schneider et al., 1997). The samples were re-dissolved in acetonitrile and the concentration adjusted to give 1 AU at the 226 nm UV chromophore. CD spectra (200–300 nm) were recorded using a Jasco J-700 spectropolarimeter (3 scans of each sample averaged, scan rate 30 nm/min).

3. Results

3.1. Generation and isolation of epoxyalcohols from hematin treatment of 9-HPODE

Fig. 1A illustrates the RP-HPLC separation of the hematin products from 9S-HPODE. These include four 9,10-epoxy-13-hydroxy isomers appearing late in the chromatogram, of which the more prominent are the two trans epoxide isomers, identified earlier as 9S,10S-trans-epoxy-13S-hydroxy followed by the 9S,10S,13R diastereomer (Zheng et al., 2011). On either side of the first of these are the two less abundant 9,10-cis-epoxide isomers labeled as peaks 6 and 8, the stereochemistry of which can be deduced from earlier studies as the 9S,10R-cis-epoxy-13R-hydroxy diastereomer followed by the 9S,10R,13S (Gao et al., 2009). We also identified two 9S,10S-trans-epoxy-11-hydroxy diastereomers (Fig. 1A, peaks 4 and 5): the 11S-erythro isomer elutes earlier at \sim 9.3 min, and the 11*R*-threo-hydroxy isomer co-elutes with the first *cis*-epoxide at 11.0 min (see Supplementary Tables S9 and S10 for ¹H NMR assignments); this is consistent with previous findings that the erythro diastereomers are the less polar on SP-HPLC or TLC (Corey and Mehrotra, 1983; Corey et al., 1983; Dix and Marnett, 1985; Gardner and Kleiman, 1981; Vasiljeva et al., 1993).

3.2. Separation and characterization of trihydroxyoctadecenoate isomers from the acid hydrolysis of 9S,10S-trans-epoxy-11E, 13S-hydroxyoctadecenoate

To prepare a series of trihydroxy standards for analytical work, we resolved the products of mild acid hydrolysis of 95,10*S*-*trans*-epoxy-11*E*-13*S*-hydroxyoctadecenoate. After epoxide ring opening, attack of water on either face of the molecule at C10 or C12 is expected, giving rise to two 95,10(R/S),13*S*- and two 95,12(R/S),13*S*-triols. Using an RP-HPLC system in which the epoxyalcohol starting material elutes at 7 min, a δ -ketol by-product (Section 3.4) elutes at ~5.5 min, the four trihydroxy isomers co-elute around 4 min (Fig. 1B). Although the four triols are diastereomers, not enantiomers, use of a Chiralpak AD chiral column proved most helpful for their separation, and using a combination of successive chromatographic steps the four were resolved: Chiralpak AD separation of the collected trihydroxy isomers allows separation of 3 of the 4 trihydroxy products (labeled I, II and III) (Fig. 1C). Peak II is further separated to peaks IIa and IIb



Fig. 1. Isolation of epoxyalcohols and trihydroxyoctadecenoic acids. (A) SP-HPLC analysis of products from reaction of 9S-HPODE with hematin. The four allylic epoxy alcohols are the smaller 9,10-cis-epoxy-11E-13-hydroxy products (peak 6, 9S, 10R, 13R, and peak 8, 9S, 10R, 13S), and the more prominent trans epoxides (peak 7, 9S,10S,13S, and peak 9, 9S,10S,13R). Other products appearing on the chromatogram include the following: (1) 9-keto-octadec-10E,12Z-dienoic acid, (2) 9S,10S-epoxy-13-oxo-octadec-11E-enoic acid, (3) 9S-HODE, (4) 9S,10Strans-epoxy-11S-erythro-hydroxy-octadec-12Z-enoic acid, and (5) 9S,10S-transepoxy-11R-threo-hydroxy-octadec-12Z-enoic acid. Column: Thomson Advantage Silica 150 Å 5 μ column, 25 cm \times 0.46 cm; solvent: hexane/IPA/HAc 100/2/0.02 (by vol.), flow rate 1 ml/min. (B) RP-HPLC analysis after mild acid hydrolysis of the 9S,10S-trans-epoxy-13S-hydroxy isomer with unresolved triols (4 min), δ -ketol (5.5 min), and unreacted epoxyalcohol (6.7 min). Column: Waters Symmetry C18 column 250 mm \times 4.6 mm; solvent: methanol/water/acetic acid (80/20/0.01) with UV detection at 205 nm. (C) Further separation of the collected triols using a Chiralpak AD column produced three peaks, designated I, II and III. Solvent: hexane/methanol/ethanol/acetic acid (86.5/4.5/9/0.1), 1 ml/min, UV detection at 205 nm. (D) Resolution of II was achieved using a Chiralpak OD-H column yielding IIa and IIb. Solvent: hexane/ethanol/acetic acid, (97.5/2.5/01), 1 ml/min, UV detection at 205 nm.

using the Chiralpak OD-H column (Fig. 1D). Each peak was collected and analyzed individually.

3.2.1. GC-MS analysis of triol isomers

The four trihydroxyoctadecanoate methyl esters from acid hydrolysis of 9S,10S-*trans*-epoxy-13S-hydroxyoctadecenoate were treated with dimethoxypropane/toluene sulphonic acid, thus coupling the vicinal diol as the dimethoxypropane (DMP) acetonide derivative, then silylated on the free hydroxyl with BSTFA and analyzed by GC–MS. This clearly identified triols I and IIb as having a 13-OTMS group (prominent m/z 173), and IIa and III as 9-OTMS (m/z 259):

TriOH Peak I and IIb

The mass spectrum of the methyl ester, Me₃Si, DMP derivative of peaks I and IIb showed ions at m/z 441 (M⁺ – 15; loss of \cdot CH₃), 295 (C1–C13 – 90; loss of Me₃SiOH), 199 (C11–C18), 173 (C13–C18), 103 (Me₃SiO⁺=CH₂), 99 (C13–C18–Me₃Si⁺) and 73 (Me₃Si⁺). These fragment ions indicate a 9,10,13 trihydroxyoctadecenoate (Fig. 2A and C).

TriOH IIa and III.

Analyzed as the methyl ester, DMP, Me₃Si ether derivatives the following ions were detected; m/z 441 (M⁺ – 15; loss of CH₃), 356 (M⁺ – 100, rearrangement followed by loss of OHC-(CH₂)₄-CH₃), 298 (C1–C13–Me₃Si⁺–CH₃), 259 (C1–C9), 209 (C9–C18–Me₃SiOH), 103 (Me₃SiO⁺=CH₂), 73 (Me₃Si⁺). The m/z 259 ion representing α -cleavage next to the 9-OTMS group is diagnostic for the 9,12,13 trihydroxyoctadecenoate (Fig. 2B and D).

3.2.2. ¹ H NMR of DMP derivative of products from acid hydrolysis of 9S,10S-trans-epoxy-11E, 13S-hydroxyoctadecenoate

Table 1 details the diagnostic chemical shift, multiplicity and coupling constants for all 4 trihydroxy isomers formed (See also Supplementary Tables S1–S4). From the GC–MS analysis, I and IIb are 9,10,13-trihydroxy, and IIa and III are the 9,12,13-trihydroxy epimers. As these products are generated from 95,105-epoxy-135hydroxy, the difference in the NMR data is due to the chirality of either the 10-hydroxy or 12-hydroxy. The results are consistent with the unknown hydroxy groups in I and III having the same relative configuration (I, $J_{9,10}$ = 7.5 Hz and III, $J_{12,13}$ = 7.62 Hz with both the H10 and H12 chemical shifts at 4.06 ppm). The chemical shifts of the acetonide methyl groups (I and III - 1.47 and 1.45 ppm) indicate the syn-diol configuration (Bruno et al., 1992; Cardellina and Moore, 1980). Trihydroxy IIa and IIb show the opposite chirality of the unconfirmed hydroxyl (IIa, $J_{12,13} = 6.84$ Hz and IIb, $J_{9,10} = 6.72$ Hz with both the H10 and H12 chemical shifts at 4.41 ppm). The acetonide methyl groups in this case indicate an anti-diol (IIa and IIb – 1.56 and 1.375 ppm). All products have a trans double bond $(J_{10,11/11,12} = 15.5 \text{ Hz}).$

Peak III co-eluted with pinellic acid, (the beetroot product from 9S-HPODE (Hamberg and Olsson, 2011), see the following Section 3.3) in the Chiralpak AD separation and also shares the same ¹H NMR and GC–MS data, confirming its structure as pinellic acid, 9S,12S,13S-trihydroxyoctadec-10E-enoic acid. Triol I shares the same ¹H NMR profile as III (Table 1) and can therefore be assigned as 9S,10S,13S-trihydroxyoctadec-11E-enoic acid. The stereochemistry of IIa, the second 9,12,13-triol, has an ¹H NMR spectrum consistent with an *anti*-diol and is therefore 9S,12*R*,13Strihydroxyoctadec-10E-enoic acid. The second 9,10,13-triol, IIb, also shares this *anti*-diol and can be assigned as 9S,10*R*,13Strihydroxyoctadec-11E-enoic acid.



Fig. 2. Regio isomer assignment of 9S-HPODE-derived triols. EI mass spectra of the triols as the methyl ester, dimethoxypropane, trimethylsilyl ether derivative (M⁺ 456). A prominent m/z 173 ion indicates a 13-OTMS structure and is diagnostic of a 9,10,13-triol (I and IIb, panels A and C) with other characteristic ions at m/z 327, 295, 270, 228, 227 and 212. A prominent ion at m/z 259 indicates a 9-OTMS and is diagnostic of a 9,12,13-triol (IIa and III, panels B and D) with other characteristic ions at m/z 356, 298, 241 and 209.

3.2.3. SP-HPLC resolution of the triols as the methyl ester DMP derivative

As suggested by the results above, chromatographic separation of the four triols from the 9S,10S,13S epoxyalcohol is quite challenging as they are poorly resolved in typical RP-HPLC and SP-HPLC systems. In a late development in this work, we found that the triols are widely resolved by SP-HPLC of the methyl ester DMP derivative (Fig. 3A). To further test the capabilities of this method, we prepared the four diastereomeric triols from acid hydrolysis of the 9*S*,10*S*,13*R*-epoxyalcohol (the second of the two major epoxyalcohols from the hematin reaction with 9*S*-HPODE). These are separated into an additional three new peaks in this system (Fig. 3B). As indicated in the figure legend, the first of these new peaks (marked with an asterisk) contains two triols which can be resolved by SP-HPLC of the methyl ester derivative.

26 Table 1

Diagnostic MS and 'H NMR characteristics of triol hydrolysis products of 98 108-epoxy-138-hydroxy-octadec-116-epoic a	Dia	iagnostic MS	and	¹ H NMR	characteristics	of triol	hvdrolvsi	products	of 95 105	-enoxy-	-13S-hv	droxy-	octadec-	11E	-enoic	ac	id
---	-----	--------------	-----	--------------------	-----------------	----------	-----------	----------	-----------	---------	---------	--------	----------	-----	--------	----	----

Hydrolysis Products	Epoxyalcohol precursor: 95,105,125 R O E OH	Product stereochemistry	GC–MS (DMP)-MeTMS diagnostic ions	¹ H NMR DMP geminal (OH) proton	¹ H NMR DMP-methyl (ppm)
Ι		95,105,135 9,10-syn (<i>threo</i>)	m/z 173	H10 – 4.06 ppm $J_{9,10}$ = 7.7 Hz $J_{10,11}$ = 7.5 Hz	1.45 1.47
IIa		95,12R,13S 12,13-anti (<i>erythro</i>)	m/z 209 and 259	H12 – 4.42 ppm $J_{12,13} = 6.8$ Hz $J_{11,12} = 6.8$ Hz	1.38 1.56
IIb		95,10R,13S 9,10-anti (<i>erythro</i>)	m/z 173	H10 – 4.42 ppm $J_{9,10}$ = 7.0 Hz $J_{10,11}$ = 6.7 Hz	1.38 1.56
III		95,125,135 12,13-syn (threo)	m/z 209 and 259	H12 – 4.06 ppm $J_{12,13} = 7.6$ Hz $J_{11,12} = 7.6$ Hz	1.45 1.47
δ-ketol		9-hydroxy, 13-keto	m/z 241 and 99	Diagnostic signal H12 – doublet at 2.73 ppm J _{11,12} = 7.0 Hz	

 $R = HO_2C(CH_2)_{7-}$.

3.3. Enzymatic conversion of 9S- and 9R-HPODE by beetroot extract to epoxyalcohols and triols

Hamberg and Olsson reported the efficient enzymatic transformation of 9S-HPODE by a filtered homogenate of B. vulgaris (beetroot) to 9S-hydroxy-12R,13S-cis-epoxy-octadec-10E-enoic acid and upon longer incubation its enzymatic hydrolysis to 9S,12S,13S-trihydroxy-octadec-10E-enoate (pinellic acid); 13S-HPODE was not metabolized (Hamberg and Olsson, 2011). In extending these experiments we found that 13R-HPODE is also not metabolized, while 9R-HPODE was converted with only a slightly lower rate of reaction compared to 9S-HPODE. The 9S-HPODE enantiomer gave products as reported (Hamberg and Olsson, 2011), a single epoxyalcohol product after 5 min incubation (data not shown), and one prominent triol peak following its hydrolysis during 60 min incubation (RP-HPLC, peak at 4 min retention time, Fig. 4A with GC-MS, Fig. 4B and ¹H NMR, Table 2 and Supplementary Table S5, confirming the structure). The transformations of 9*R*-HPODE produced one main epoxyalcohol with two minor isomers upon short-term incubation (5 min). Fig. 4C. Incubation for 60 min yielded their enzymatic hydrolysis products with the polarity of triols (Fig. 4E), resolved on a Chiralpak AD column into one predominant trihydroxy isomer with two minor products (Fig. 4E, inset).

3.3.1. Identification of the main 9R-HPODE-derived epoxyalcohol formed in beetroot

The epoxyalcohol produced by 9*R*-HPODE incubation with beetroot was isolated as per methods as a single main product eluting at 6 min on RP-HPLC (Fig. 4C). The ¹H NMR spectrum clearly established the presence of a *trans* double bond with allylic *cis*-epoxide (J=4 Hz) and carbon-bearing hydroxyl on either side (Table 2, Supplementary Table S6). However, due to the symmetry of structure in this type of linoleate derivative, the NMR does not establish whether the epoxyalcohol is 9,10-epoxy-11*E*-13-hydroxy or the reversed 9-hydroxy-10*E*-12,13-epoxy.

The mass spectrum of the methyl ester TMS ether derivative (Fig. 4D) showed ions at m/z 398 (M⁺), 327 (C1–C13), 297, 259 (C1–C9), 237 (327-Me₃SiOH), 199 (C1–C10), 173 (C13–C18), 103

(Me₃SiO⁺=CH₂), 73 (Me₃Si⁺). The presence of ions at both m/z 173 ion (C13–C18, typical of 13-OTMS) and m/z 259 (C1–C9, typical of 9-OTMS) also fails to resolve the issue. Indeed, Gardner and Kleiman (1981) have shown that both m/z 173 and 259 are prominent ions



Fig. 3. SP-HPLC resolution of triols as the methyl ester DMP derivative. (A) The four triol hydrolysis products of methyl 9*S*,10*S*-epoxy-13*S*-hydroxyoctadec-11*E*-enoate were purified as a group by SP-HPLC, derivatized to the acetonide, and analyzed using a Thomson Advantage Silica 150 Å 5 μ column (25 cm × 0.46 cm) with a solvent of hexane/IPA 100/1 (by vol.) at a flow rate 1 ml/min, with UV detection at 205 nm. The relative proportions of the four triols are indicated as percentages. (B) The same four products were chromatographed with the triols (methyl ester DMP derivative) from the second major epoxyalcohol of the hematin reaction, the 13*R* diastereomer, 9*S*,10*S*-epoxy-13*R*-hydroxyoctadec-11*E*-enoate (the shaded peaks). The four 13*R* triols resolve as two pairs by SP-HPLC of the methyl ester derivative using the same Thomson silica column with a solvent of hexane/IPA 90/10 (by vol.), retention times 8.8 min (9*S*,10*R*,13*R*+9*S*,12*S*,13*R*) and 11.5 min (9*S*,10*S*, 13*R*+9*S*,12*R*,13*R*). As the Me DMP derivative, 9*S*,10*R*,13*R*+9*S*,12*R*,13*R*, isomers.



Fig. 4. Isolation and structural characterization of products from the incubation of 9*R*- and 9*S*-HPODE with *B. vulgaris* extract. (A) RP-HPLC (as per Section 2.3.2) of products from 60 min incubation of 9*S*-HPODE: trihydroxy product(s) elute at 4 min with unreacted 9-HPODE substrate and 9-HODE eluting at 15 and 17.5 min respectively. Inset: separation using a Chiralpak AD column confirms the generation of one major trihydroxy isomer. (B) EI mass spectrum of the methyl ester DMP TMS ether derivative with a diagnostic ion at m/z 259 indicating a 9-OTMS group and a 9,12,13-trihydroxyoctadecenoate structure. (C) RP-HPLC of the epoxyalcohol (retention time, 6 min) from a 5 min incubation of 9*R*-HPODE with *B. vulgaris*. (D) EI mass spectrum (methyl ester TMS ether derivative) of the 9-hydroxy-12,13-epoxyoctadecenoic acid product. (E) RP-HPLC of products from the 60 min incubation of 9*R*-HPODE with two triol peaks eluting at 4 min. Inset: the triols were resolved using a Chiralpak AD column as major (17 min) and minor (24 min) isomers. (F) EI spectrum of the major isomer from 9*R*-HPODE, a 9,12,13-trihydroxyoctadecenoate, analyzed as in (B).

in the mass spectra of the methyl ester TMS ether derivative of 9-hydroxy-12,13-epoxyoctadecenoates.

Based on its biosynthesis in beetroot and by analogy to the metabolism of 9S-HPODE (Hamberg and Olsson, 2011), an oxygen transfer mechanism converting 9*R*-HPODE to 9*R*-hydroxy-12,13*cis*-epoxyoctadecenoate is anticipated. This was confirmed by comparison of the beetroot 9*R*-HPODE-derived epoxyalcohol with the two 9,10-*cis*-epoxy-13-hydroxy diastereomers obtained from hematin treatment of 9-HPODE (Section 3.1), and the corresponding two from 13S-HPODE (9-hydroxy-10*E*-12,13-*cis*-epoxy). By RP-HPLC, three resolved from the beetroot product and one of 13S-HPODE *cis*-epoxyalcohols co-chromatographed (data not shown). This one must have the same regio- and stereochemistry as beetroot 9*R*-HPODE product or be the enantiomer. Finally, we assigned this 13*S*-HPODE-derived *cis*-epoxyalcohol with the 9*R*-hydroxy configuration by CD analysis (Gonnella et al., 1982; Schneider et al., 1997) (Fig. 5), establishing it as identical to the beetroot product which therefore has the structure, 9*R*-hydroxy-12*R*,13*S*-*cis*-epoxyoctadec-10*E*-enoic acid (Scheme 1).

3.3.2. Identification of the main 9R-HPODE-derived triols formed in beetroot

The major triols from the 9S- and 9R-HPODE incubations with beetroot show indistinguishable mass spectra (Fig. 4B, 4F), indicating they are the same regio isomers. Analyzed as the methyl ester, DMP, TMS ether derivatives the following ions were detected; m/z

Table 2

Diagnostic MS and ¹H NMR characteristics of 95/R-hydroperoxy-10E, 12Z-octadecadienoic acid and Beta vulgaris incubation products.

Incubation products	Hydroperoxide precursor: 9R R	Product stereochemistry	GC–MS (DMP)-MeTMS diagnostic ions	¹ H NMR DMP geminal (OH) proton	¹ H NMR DMP-methyl (ppm)
5 min incubation	R O OH	9R-hydroxy,12R,13S- epoxy	m/z 259	<i>cis</i> -epoxide H12 – 3.25 ppm J _{12,13} = 4.4 Hz	
60 min incubation	OH R OH OH OH OH OH OH	9R,12S,13S 12,13-syn (threo)	m/z 209 and 259	H12 – 4.06 ppm J _{12,13} = 7.3 Hz J _{11,12} = 7.3 Hz	1.46 1.48
5 min incubation	R OH	9S-hydroxy,12R,13S- epoxy	m/z 259	<i>cis</i> -epoxide H12 – 3.25 ppm J _{12,13} = 4.4 Hz	
60 min incubation	R OH OH OH	95,125,135 12,13-syn (threo)	m/z 209 and 259	H12 – 4.06 ppm $J_{12,13} = 7.1$ Hz $J_{11,12} = 7.3$ Hz	1.46 1.47

 $R = HO_2C(CH_2)_7 - .$

441 (M^+ – 15; loss of CH₃), 356 (M^+ – 100, rearrangement followed by loss of OHC-(CH₂)₄-CH₃), 298 (C1–C13–Me₃Si⁺–CH₃), 259 (C1–C9), 209 (C9–C18–Me₃SiOH), 103 (Me₃SiO⁺=CH₂), 73 (Me₃Si⁺). The presence of the ion at m/z 259, diagnostic of the 9-OTMS group, identifies these products as 9,12,13 trihydroxy (Table 2).

Table 2 includes a summary of the diagnostic ¹H NMR data for the major 9S- and 9*R*-HPODE-derived triols as the methyl ester DMP derivative (with full details given in Supplementary Tables S5 and S7). The original 9*R*/13S or 9*S*/13S configurations in the parent epoxyalcohols should be retained, and therefore the unresolved issue is the chirality at C12. The chemical shifts of the C12 proton and coupling constant for H12,13 are almost identical (4.06 ppm, $J_{12,13}$ = 7.1 or 7.3 Hz), indicating the same relative configuration in the vicinal diol: the substituents are *trans* to each other on the acetonide ring and so the hydroxyls must be in the *syn* configuration and are 12S,13S. The chemical shifts of the acetonide methyl groups also support this conclusion (Bruno et al., 1992; Cardellina and Moore, 1980). This reversal of configuration at C12 upon hydrolysis of the epoxyalcohols is expected through the actions of epoxide hydrolase (Arand et al., 2005). Finally, the coupling constants for both these products show the double bond is $trans(J_{10,11} = 15.5 \text{ Hz})$. Therefore, the 9S-HPODE-derived triol, pinellic acid, is 9S,12S,13S as already reported (Hamberg and Olsson, 2011), and the main triol from 9*R*-HPODE is 9*R*,12*S*,13*S*-trihydroxy-10*E*-octadecenoic acid (Scheme 1).

3.4. Identification of a specific δ-ketol product from mild acid hydrolysis of 9S,10S-trans-epoxy-11E,13S-hydroxyoctadecenoate

Upon RP-HPLC analysis of acid-treated 95,10S-*trans*-epoxy-11E-13S-hydroxy-octadecenoate, we consistently observed a peak, more polar than the parent epoxyalcohol yet considerably less polar than the triol hydrolysis products, and exhibiting only end



Scheme 1.



Fig. 5. CD analysis of the hydroxyl configuration in 9-hydroxy-12,13-*cis*-epoxyalcohols CD spectra were recorded using a Jasco-700 spectropolarimeter on the 9-O-benzoate esters of the 9-hydroxy-10*E*-12,13-*cis*-epoxy-octadecenoate methyl esters in acetonitrile solution with the concentration adjusted to give 1.0 AU at 227 nm in the UV (methods in Section 2.5.1). Interaction of the chromophores of the 9-O-benzoate ester and the 10*E* bond produces either a positive first Cotton effect (top) associated with a positive interaction (+) in the Newman projection, and allowing designation of the 9S configuration, or the negative Cotton effect, negative Newman projection, designating 9*R*.

absorbance in the UV (Fig. 1B). After further purification as the methyl ester (Chiralpak AD column), the main component was identified by NMR as a δ -ketol. The significant chemical shifts, multiplicity and coupling constants for the δ -ketol product are detailed in Table 1 (Supplementary Table S8). Diagnostic features of the spectrum include the doublet signal for the CH₂ at 2.7 ppm ($J_{11,12}$ = 7.02 Hz), with coupling only to the one *trans* double bond (J = 15.5 Hz), with no additional coupling due to the adjacent ketone and a second CH₂ triplet representing H14 near H2 (Supplementary Table S8). Due to the symmetry typical of many linoleate-derived products in the NMR, it was not possible to distinguish the 9-oxo-11E-13-hydroxyoctadecenoate from the 9-hydroxy-10E-13-oxo isomer. In this case, however, GC-MS analyses were definitive. The methyl ester, methoxime, TMS ether derivative gave a pair of closely eluting peaks for the syn and anti isomers on GC with slightly differing ion abundances on MS (Fig. 6A and B); both peaks gave ions at $m/z 429 (M^{+\bullet}) 396 (M^{+}-CH_{3}O)$, 356 (C1-C13), 285 (C1-C11), 270 (C9-C18), 168 (C10-C18), 73 (Me₃Si⁺). GC-MS of the methyl ester, TMS derivative yielded one main peak showing two main ions at m/z 241 (C9-C18) and 99 (C13-C18) (Table 1, Fig. 6C and D) (Gardner and Kleiman, 1979). These results, together with the NMR data, identify the δ -ketol as 9-hydroxy-13oxo-octadec-10E-enoic acid.

4. Discussion

Utility of hematin method. In the hematin treatment of fatty acid hydroperoxides under the conditions we describe, the allylic epoxyalcohols are recovered as major products when suitable precautions are taken to avoid their facile hydrolysis. To avoid hydrolysis of these highly acid-labile allylic epoxyalcohols, the samples are kept on ice during acid extraction, the organic solvent is added prior to acidification (and thus is ready for mixing the instant after acid is added), and the acidification itself uses a predetermined quantity of 1 M KH₂PO₄ and HCl to produce the desired pH, immediately followed by organic extraction; subsequent washing of the organic phase with water prior to evaporation is another essential precaution. Although the products chromatograph

satisfactorily in RP-HPLC solvent containing 0.01% glacial acetic acid, prior formation of the methyl esters (diazomethane) and elimination of the acetic acid improves the subsequent recovery of intact allylic epoxyalcohol products from the solvent.

Identification of hydrolysis products. In linoleate-derived triols arising from intermediate allylic epoxyalcohols, the structural symmetry of the 9,10,13- and 9,12,13-trihydroxy products confounds their definitive assignment solely by NMR. On the other hand, these regioisomers are readily distinguished by GC-MS of the DMP derivative (Fig. 2, Table 1 and Supplementary Tables S1-S4). Epoxide ring opening under mild acid hydrolysis induces formation of an intermediate allylic carbocation with the hydroxyl configurations at 9 and 13 fixed with respect to the substrate. When the structure of the epoxyalcohol is known, therefore, the remaining issue of stereochemistry in the resulting triols is the configuration of the hydroxy at C10 or C12. In ¹H NMR of the DMP derivative, the chemical shift of the allylic hydroxyl in the vicinal diol is diagnostic of the syn or anti configuration. Thus, the chemical shift of the allylic hydroxyl in the syn vicinal diols (products I and III, Table 1) is 0.4 ppm upfield in comparison to anti (products IIa, IIb), which is sufficient to define the complete stereochemistry.

The chemical shifts of the geminal methyl groups on the acetonide ring in the DMP derivatives also provide structural information, confirming the chirality of the unknown hydroxyl (cf. (Bruno et al., 1992; Cardellina and Moore, 1980; Shirahata et al., 2006)). When the 9,10 or 12,13 hydroxyls are *syn*, the chemical shifts of the acetonide methyl groups are almost superimposable, only ~0.02 ppm apart. When the vicinal diol hydroxyls are *anti*, the methyl groups are in a *cis* configuration and the chemical shifts differ by ~0.2 ppm. Thus, for products I and III (Table 1), the acetonide methyls have chemical shifts of 1.45 and 1.47 ppm, whereas they are separated in IIa and IIb with values of 1.38 and 1.56 ppm.

Four additional triols with the 9S,10(R or S),13R and 9S,12(R or S),13R configurations are available by acid hydrolysis of the second *trans*-epoxyalcohol from 9S-HPODE (9S,10S-epoxy-11E-13R-hydroxy). We show that these can be resolved by SP-HPLC of the methyl ester DMP derivative (Fig. 3B). Although not addressed in this study, the two *trans*-epoxyalcohols derived from the enantiomeric hydroperoxide, 9R-HPODE, will give rise to eight triol enantiomers to the products resolved here. As an alternative to the reported degradative methods of chiral analysis via ozonolysis (Hamberg, 1991a, 1991b; Garbe et al., 2005), chiral HPLC column methods could be explored as a means to their assignment.

Proportions of the hydrolysis products. Hamberg has reported the relative proportions of the four products of mild acid hydrolysis from a linoleate-derived 9,10-cis-epoxy-13-hydroxy epoxyalcohol: 61% and 15% attack on the allylic epoxy carbon (favoring reversal of configuration), 15% and 8% on the carbon α to the hydroxyl (favoring formation of the erythro vicinal diol) (Hamberg, 1991a). He found similar proportions, 61:18:15:6, in hydrolysis of the 12,13-cis-epoxy-9-hydroxy precursor of pinellic acid (Hamberg and Olsson, 2011). We found different proportions for mild acid hydrolysis of an analogous trans-epoxy product, methyl 95,10Strans-epoxy-11E-13S-hydroxy-octadecenoate. Our best estimate comes from resolution of the four diastereomers as the methyl ester DMP derivative showing a ratio of 22:26:40:12 for products I, IIa, IIb, and III respectively (Fig. 3A). A point of agreement with hydrolysis of the cis-epoxyalcohols is that the highest abundance triol is formed via attack on the allylic epoxy carbon with reversal of configuration (product IIb in our case), and the lowest is the three vicinal diol δ to the original epoxide (product III in our case).

Stereospecificity of the beetroot epoxyalcohol synthase. Here we showed that 9*R*-HPODE, enantiomeric to the 9*S*-HPODE studied by Hamberg and Olsson is also a good substrate for the beetroot enzymes (Hamberg and Olsson, 2011) (whereas neither of the 13-HPODE enantiomers react). The initial major product from



Fig. 6. Identification of a δ -ketol. (A) Total ion current chromatogram of the methyl ester, methoxime, TMS ether derivative of the δ -ketol formed during acid hydrolysis of 9S,10*S*-trans-epoxy,13*S*-hydroxy-octadec-11*E*-enoic acid showing the closely eluting *syn* and *anti* isomers at ~9.8 min. (B) El mass spectrum of the first-eluting isomer with diagnostic ions at m/z 168 and 285. (C) Total ion current chromatogram of the methyl ester, TMS ether derivative. (D) The mass spectrum of the main peak has diagnostic ions at m/z 99 and 241 (Gardner and Kleiman, 1979).

the 9*R*-hydroperoxide is the analogous 9*R*-hydroxy-12*R*,13*S*-cisepoxide, similarly formed by epoxidation at 12,13. Subsequent hydrolysis gave a major product formed via epoxide hydrolase. Recently it was shown that *cis* epoxides present in epoxyeicosatrienoic acids (EETs) are converted to the *syn*-diol by soluble epoxide hydrolase present in rat red blood cells; conversely the *trans* epoxides became *anti*-diols (Jiang et al., 2008). These results are in line with our findings. It is logical, therefore, that the main product from the enzymatic hydrolysis of 9*R*-hydroxy-12*R*,13*Strans*-epoxide is indeed the 9*R*,12*S*,13*S*-triol.

To assign the stereoconfigurations in the primary beetroot epoxyalcohol product from 9*R*-HPODE, NMR was used to establish the hydroxy-*trans* double bond-*cis*-epoxy partial structure. Again, because of the symmetry in linoleate-derived structures, NMR did not reveal the orientation of this unit in the fatty acid. Similarly, GC–MS is not definitive because the usual



Scheme 2.

"diagnostic" ions for 9-OTMS (m/z 259) or 13-OTMS (m/z 173) were both present in the EI mass spectrum of the methyl ester TMS derivative (cf. (Gardner and Kleiman, 1981)). We assigned the configuration by (i) comparison of its HPLC mobility to each of the potential *cis*-epoxy diastereomers prepared by hematin treatment of 9-HPODE (two 9,10-cis-epoxy-11E-13-hydroxy diastereomers) and 13S-HPODE (two 9-hydroxy-10E-12R,13S-cis-epoxides). The 9*R*-HPODE-derived beetroot product co-chromatographed on RP-HPLC only with one of the two 12*R*,13S-epoxides; (ii) their 9-hydroxy configuration was then established by CD analysis (Gonnella et al., 1982), proving that the beetroot product is 9*R*-hydroxy-10*E*-12*R*,13S-cis-epoxy-octadec-10*E*-enoic acid. Although we did not pursue the matter, another potential application of the CD method is assignment of the stereochemistry of the free hydroxyl in the DMP derivative of the triols.

Mechanism of transformation to delta-ketol. We observed that mild acid hydrolysis of the trans-epoxyalcohol, 9S,10S-epoxy-11E-13S-hydroxy-octadecenoate, formed not only the distinctly more polar triols, but also a peak of intermediate polarity on RP-HPLC (Fig. 1B) that we identified by NMR and GC-MS as mainly one δ -ketol. In the 1970s, Gardner reported formation of two isomeric δ -ketols from treatment of either 9-HPODE or 13-HPODE with FeCl₃/cysteine (Gardner and Kleiman, 1979), but there has been little attention to such products since. Gardner described that a single epoxyalcohol intermediate treated with a Lewis acid (BF3-etherate) gave rise to similar quantities of two isomeric ketols, 9-hydroxy-10E-13-oxo and 9-oxo-11E-13-hydroxy. We found a 10-fold relative abundance of the 9-hydroxy-13oxo isomer over the 9-oxo-13-hydroxy. Whereas Gardner and Kleiman implicated a symmetrical carbocation intermediate to account for the appearance of apparently similar amounts of two diastereomers, our results suggest asymmetry in the reaction intermediate, giving the bias toward predominantly one outcome (Scheme 2). Although delta-ketols have not appeared in the recent literature, given the similarity in the EI mass spectra of the methyl ester TMS ether derivatives of certain epoxyalcohols and delta-ketols (Gardner and Kleiman, 1979), they are a candidate that might provide an alternative explanation for the appearance of unlikely epoxyalcohol isomers in the analysis of human skin fatty acids (Zheng et al., 2011). Even more probable, we suspect a delta-ketol(s) accounts for the peak of intermediate polarity illustrated in the RP-HPLC profile in our acid hydrolysis of the arachidonate-derived epoxyalcohol, 8R-hydroxy-11R,12Rtrans-epoxy-eicosatrienoic acid (Yu et al., 2003); in that case LC-MS analysis confirmed that the intermediate polarity product(s) has the same molecular weight as the starting epoxyalcohol and potential $\delta\mbox{-}ketol$ product.

Acknowledgements

Financial support was provided by European Union FP7, Marie Curie Outgoing Fellowship (C.P.T.), by NIAMS grant AR-51968 (to A.R.B.) and a voucher for use of the mass spectrometry resource through the Vanderbilt-Ingram Cancer Center Support grant (P30 CA068485). We would also like to thank Drs. Wade Calcutt and Donald Stec for their assistance with GC–MS and ¹H NMR respectively, and Nicholas M. Adams and Dr. David Wright for the CD analysis.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at http://dx.doi.org/10.1016/j.chemphyslip. 2013.01.002.

References

- Andreou, A., Brodhun, F., Feussner, I., 2009. Biosynthesis of oxylipins in nonmammals. Progress in Lipid Research 48, 148–170.
- Andreou, A.Z., Vanko, M., Bezakova, L., Feussner, I., 2008. Properties of a mini 9Rlipoxygenase from Nostoc sp. PCC 7120 and its mutant forms. Phytochemistry 69, 1832–1837.
- Arand, M., Cronin, A., Adamska, M., Oesch, F., 2005. Epoxide hydrolases: structure, function, mechanism, and assay. Methods in Enzymology 400, 569–588.
- Blée, E., Wilcox, A.L., Marnett, L.J., Schuber, F., 1993. Mechanism of reaction of fatty acid hydroperoxides with soybean peroxygenase. Journal of Biological Chemistry 268, 1708–1715.
- Bruno, I., Dauria, M.V., Iorizzi, M., Minale, L., Riccio, R., 1992. Marine eicosanoids: occurrence of 8,11,12-trihydroxylated eicosanoic acids in starfishes. Experientia 48, 114–115.
- Cardellina, J.H., Moore, R.E., 1980. Malyngic acid, a new fatty acid from *Lyngbya* majuscula. Tetrahedron 36, 993–996.
- Claeys, M., Kivits, G.A., Christ-Hazelhof, E., Nugteren, D.H., 1985. Metabolic profile of linoleic acid in porcine leukocytes through the lipoxygenase pathway. Biochimica et Biophysica Acta 837, 35–51.
- Corey, E.J., Mehrotra, M.M., 1983. Stereochemistry of the lipoxygenase-catalyzed allylic hydroperoxide to oxiranylcarbinol rearrangement. Tetrahedron Letters 45, 4921–4922.
- Corey, E.J., Mehrotra, M.M., Cashman, J.R., 1983. New synthetic routes to leukotrienes and other arachidonate derived epoxy eicosatetraenoic acids (EPETE's) – exclusion of the hydroxy epoxide pathway for leukotriene biosynthesis. Tetrahedron Letters 24, 4917–4920.
- Dix, T.A., Marnett, L.J., 1985. Conversion of linoleic acid hydroperoxide to hydroxy, keto, epoxyhydroxy, and trihydroxy fatty acids by hematin. Journal of Biological Chemistry 260, 5351–5357.
- Feussner, I., Wasternack, C., 2002. The lipoxygenase pathway. Annual Review of Plant Biology 53, 275–297.

- Funk, C.D., Powell, W.S., 1985. Release of prostaglandins and monohydroxy and trihydroxy metabolites of linoleic and arachidonic acids by adult and fetal aortae and ductus arteriosus. Journal of Biological Chemistry 260, 7481–7488.
- Galliard, T., Phillips, D.R., 1971. Lipoxygenase from potato tubers, partial purification and properties of an enzyme that specifically oxygenates the 9-position of linoleic acid. Biochemical Journal 124, 431–438.
- Gao, B., Boeglin, W.E., Brash, A.R., 2010. Omega-3 fatty acids are oxygenated at the n-7 carbon by the lipoxygenase domain of a fusion protein in the cyanobacterium *Acaryochloris marina*. Biochimica et Biophysica Acta 1801, 58–63.
- Gao, B., Boeglin, W.E., Zheng, Y., Schneider, C., Brash, A.R., 2009. Evidence for an ionic intermediate in the transformation of fatty acid hydroperoxide by a catalaserelated allene oxide synthase from the cyanobacterium *Acaryochloris marina*. Journal of Biological Chemistry 284, 22087–22098.
- Garbe, L.A., Hübke, H., Tressl, R., 2005. Enantioselective formation pathway of a trihydroxy fatty acid during mashing. Journal of the American Society of Brewing Chemists 63, 157–162.
- Gardner, H.W., Kleiman, R., 1979. Lack of regioselectivity in formation of oxohydroxyoctadecenoic acids from the 9-hydroperoxide or 13-hydroperoxide of linoleic acid. Lipids 14, 848–851.
- Gardner, H.W., Kleiman, R., 1981. Degradation of linoleic acid hydroperoxides by a cysteine.FeCl₃ catalyst as a model for similar biochemical reactions, 2. Specificity in formation of fatty acid epoxides. Biochimica et Biophysica Acta 665, 113–125.
- Gardner, H.W., Kleiman, R., Weisleder, D., 1974. Homolytic decomposition of linoleic acid hydroperoxide – identification of fatty acid products. Lipids 9, 696–706.
- Garssen, G.J., Veldink, G.A., Vliegenthart, J.F.G., Boldingh, J., 1976. The formation of *threo*-11-hydroxy-*trans*-12: 13-epoxy-9-*cis*-octadecenoic acid by enzymic isomerisation of 13-L-hydroperoxy-9-*cis*, 11-*trans*-octadecadienoic acid by soybean lipoxygenase-1. European Journal of Biochemistry 62, 33–36.
- Göbel, C., Feussner, I., Hamberg, M., Rosahl, S., 2002. Oxylipin profiling in pathogeninfected potato leaves. Biochimica et Biophysica Acta 1584, 55–64.
- Gonnella, N.C., Nakanishi, K., Martin, V.S., Sharpless, B.K., 1982. General method for determining absolute configurations of acyclic allylic alcohols. Journal of the American Chemical Society 104, 3775–3776.
- Hamberg, M., 1975. Decomposition of unsaturated fatty acid hydroperoxides by hemoglobin – structures of major products of 13L-hydroperoxy-9,11octadecadienoic acid. Lipids 10, 87–92.
- Hamberg, M., 1991a. Regiochemical and stereochemical analysis of trihydroxyoctadecenoic acids derived from linoleic acid 9-hydroperoxide and 13-hydroperoxide. Lipids 26, 407–415.
- Hamberg, M., 1991b. Trihydroxyoctadecenoic acids in beer: qualitative and quantitative analysis. Journal of Agricultural and Food Chemistry 39, 1568–1572.
- Hamberg, M., 1999. An epoxy alcohol synthase pathway in higher plants: biosynthesis of antifungal trihydroxy oxylipins in leaves of potato. Lipids 34, 1131–1142.
- Hamberg, M., Hamberg, G., 1996. Peroxygenase-catalyzed fatty acid epoxidation in cereal seeds (sequential oxidation of linoleic acid into 9(S),12(S),13(S)trihydroxy-10(E)-octadecenoic acid). Plant Physiology 110, 807–815.
- Hamberg, M., Herman, R.P., Jacobsson, U., 1986. Stereochemistry of two epoxy alcohols from Saprolegnia parasitica. Biochimica et Biophysica Acta 879, 410–418.
- Hamberg, M., Olsson, U., 2011. Efficient and specific conversion of 9-lipoxygenase hydroperoxides in the beetroot. Formation of pinellic acid. Lipids 46, 873–878.

- Jiang, H., Zhu, A.G., Mamczur, M., Morisseau, C., Hammock, B.D., Falck, J.R., McGiff, J.C., 2008. Hydrolysis of *cis*- and *trans*-epoxyeicosatrienoic acids by rat red blood cells. Journal of Pharmacology and Experimental Therapeutics 326, 330–337.
- Meruvu, S., Walther, M., Ivanov, I., Hammarstrom, S., Furstenberger, G., Krieg, P., Reddanna, P., Kuhn, H., 2005. Sequence determinants for the reaction specificity of murine (12R)-lipoxygenase: targeted substrate modification and site-directed mutagenesis. Journal of Biological Chemistry 280, 36633–36641.
- Nigam, S., Patabhiraman, S., Ciccoli, R., Ishdorj, G., Schwarz, K., Petrucev, B., Kuhn, H., Haeggstrom, J.Z., 2004. The rat leukocyte-type 12-lipoxygenase exhibits an intrinsic hepoxilin A3 synthase activity. Journal of Biological Chemistry 279, 29023–29030.
- Niisuke, K., Boeglin, W.E., Murray, J.J., Schneider, C., Brash, A.R., 2009. Biosynthesis of a linoleic acid allylic epoxide: mechanistic comparison with its chemical synthesis and leukotriene A biosynthesis. Journal of Lipid Research 50, 1448–1455.

Pace-Asciak, C.R., Reynaud, D., Demin, P.M., 1995. Hepoxilins: a review on their enzymatic formation, metabolism and chemical synthesis. Lipids 30, 107–114.

- Püssa, T., Raudsepp, P., Toomik, P., Pallin, R., Maeorg, U., Kuusik, S., Soidla, R., Rei, M., 2009. A study of oxidation products of free polyunsaturated fatty acids in mechanically deboned meat. Journal of Food Composition and Analysis 22, 307–314.
- Schneider, C., Schreier, P., Humpf, H.-U., 1997. Exciton-coupled circular dichroism (ECCD) in acyclic hydroxylated dienes: a sensitive method for the direct stereochemical assignment of lipoxygenase products. Chirality 9, 563–567.
- Shirahata, T., Sunazuka, T., Yoshida, K., Yamamoto, D., Harigaya, Y., Kuwajima, I., Nagai, T., Kiyohara, H., Yamada, H., Omura, S., 2006. Total synthesis, elucidation of absolute stereochemistry, and adjuvant activity of trihydroxy fatty acids. Tetrahedron 62, 9483–9496.
- Shirahata, T., Sunazuka, T., Yoshida, K., Yamamoto, D., Harigaya, Y., Nagai, T., Kiyohara, H., Yamada, H., Kuwajima, I., Omura, S., 2003. Total synthesis and adjuvant activity of all stereoisomers of pinellic acid. Bioorganic and Medicinal Chemistry Letters 13, 937–941.
- Siebert, M., Krieg, P., Lehmann, W.D., Marks, F., Fürstenberger, G., 2001. Enzymic characterization of epidermis-derived 12-lipoxygenase isoenzymes. Biochemical Journal 355, 97–104.
- Vasiljeva, L.L., Manukina, T.A., Demin, P.M., Lapitskaja, M.A., Pivnitsky, K.K., 1993. Synthetic study of hepoxilins, 3. Synthesis, properties, and identification of epimeric hepoxilins (-)-(10R)-B₃ and (+)-(10S)-B₃. Tetrahedron 49, 4099–4106.
- Yu, Z., Schneider, C., Boeglin, W.E., Marnett, LJ., Brash, A.R., 2003. The lipoxygenase gene ALOXE3 implicated in skin differentiation encodes a hydroperoxide isomerase. Proceedings of the National Academy of Sciences of the United States of America 100, 9162–9167.
- Zheng, Y., Boeglin, W.E., Schneider, C., Brash, A.R., 2008. A 49- kDa mini-lipoxygenase from Anabaena sp, PCC 7120 retains catalytically complete functionality. Journal of Biological Chemistry 283, 5138–5147.
- Zheng, Y., Brash, A.R., 2010. On the role of molecular oxygen in lipoxygenase activation: comparison and contrast of epidermal lipoxygenase-3 with soybean lipoxygenase-1. Journal of Biological Chemistry 285, 39876–39887.
- Zheng, Y., Yin, H., Boeglin, W.E., Elias, P.M., Crumrine, D., Beier, D.R., Brash, A.R., 2011. Lipoxygenases mediate the effect of essential fatty acid in skin barrier formation: a proposed role in releasing omega-hydroxyceramide for construction of the corneocyte lipid envelope. Journal of Biological Chemistry 286, 24046–24056.