Stereocontrol in organic synthesis using silicon-containing compounds. A synthesis of (-)-tetrahydrolipstatin using the alkylation of a β -silyl ester and the hydroboration of an allylsilane

Ian Fleming * and Nicholas J. Lawrence

Department of Chemistry, Lensfield Road, Cambridge, UK CB2 1EW

Conjugate addition of bis(Z-tridec-1-enyl)cuprate Z-10 to (5S)-1-[(Z)-3'-dimethyl(phenyl)silylprop-2enoyl]-5-(trityloxymethyl)pyrrolidin-2-one Z-6 gave the 3*R*-imide Z-12. Subsequent enolate *n*-hexylation of the benzyl ester Z-13a derived from this imide gave the 2*R*,3*S*-ester Z-14a. Reduction of the ester group and protection of the alcohol as its TBDMS group gave the allylsilane (Z)(7*R*,8*S*)-7-(*tert*-butyldimethylsilyloxymethyl)-8-dimethyl(phenyl)silylhenicos-9-ene Z-15. Hydroboration–oxidation gave the 7*R*,8*S*,10*S*alcohol 16. Protection of the C-10 hydroxy as its benzyl ether, removal of the silyl protecting group and oxidation gave (2*R*,3*S*,5*S*)-5-benzyloxy-3-dimethyl(phenyl)silyl-2-hexylhexadecanoic acid 19. Silyl-tohydroxy conversion, β-lactone formation, and hydrogenolysis gave the known alcohol (3*S*,4*S*)-3-hexyl-4-[(*S*)-2'-hydroxytridecyl]oxetan-2-one 22, from which tetrahydrolipstatin 1 was prepared by a conventional esterification. Each of the stereochemistry determining steps, $4 \longrightarrow Z-6$, $7 \longrightarrow E-8$, $E-8 \longrightarrow Z-9$, $Z-6 + Z-10 \longrightarrow Z-12$, $Z-13a \longrightarrow Z-14a$ and $Z-15 \longrightarrow 16$, took place with a remarkably high level of open-chain stereocontrol.

Introduction

The esterase inhibitor, tetrahydrolipstatin 1,¹ supports on its carbon skeleton a 1,3-relationship and a 1,2-relationship of exactly the kind that our synthetic methods controlling stereochemistry are able to solve (Scheme 1). The 1,3-relation-



ship between C-3 and C-5 is that of a 1,3-diol **2**, where we must differentiate the two hydroxy groups in order to make the β -lactone without risk of δ -lactone formation. Hydroboration of a *cis* allylsilane **3**, can be expected to set up the masked diol with the correct *syn* relationship,² and with the two hydroxy groups about as well differentiated as they could possibly be. The 1,2-relationship between C-2 and C-3 is that which can be achieved by alkylation of an enolate carrying a β -silyl group **3**,³ and conversion of the phenyldimethylsilyl group into a hydroxy

with retention of configuration⁴ can be carried out at an appropriate stage, after the C-5 hydroxy has been protected. We therefore need to make the β -silyl ester **3**, or a derivative of it, in an enantiomerically enriched state, for which we also have methods already developed.⁵ We published a preliminary account⁶ of how we were able to put these methods together to synthesise (-)-tetrahydrolipstatin, and we now report our work in full.

Results and discussion

Our method for the control of absolute stereochemistry in the synthesis of β -silyl carbonyl compounds uses either the conjugate addition of a silylcuprate reagent to an α,β -unsaturated imide derived from Koga's chiral auxiliary **5**, or the conjugate addition of a carbon-based cuprate to a β -silylated α,β -unsaturated imide derived from the same auxiliary. In this case only the latter method will work, since we need the double bond between C-4 and C-5, and if it were present in the imide the silylcuprate would attack C-5 instead of C-3. We therefore made the β -silyl α,β -unsaturated imide Z-**6** from the acetylenic acid **4** (Scheme 2), choosing the *cis* double bond so that the



Scheme 2 Reagents: i, H2, Lindlar; ii, (COCl)2, CH2Cl2; iii, 5

product of the conjugate addition will be the enantiomer with the natural absolute configuration. We also prepared the *trans* isomer E-6, which we have made before,⁵ in order to have

available for analytical purposes the diastereoisomer of the conjugate addition product.

We needed the double bond in the ester **3** to be *cis*, and were particularly happy, given how much of the stereochemistry was to be controlled by silicon to use it to control the double bond geometry too. A completely regioselective and *syn* stereospecific silyl-cupration⁷ of the terminal acetylene **7** gave only the *trans* vinylsilane *E*-**8**, and bromodesilylation then gave the *cis* vinyl bromide *Z*-**9**, cleanly with inversion of configuration, as expected (Scheme 3).⁸ We then prepared the cuprate *Z*-**10** from



Scheme 3 Reagents: i, $(PhMe_2Si)_2CuLi\cdotLiCN$; ii, NH_4Cl , H_2O ; iii, Br_2 ; iv, NaOMe; v, Li, Et_2O ; vi, 0.5 equiv. $CuBr\cdotSMe_2$; vii, BuLi, $PhMe_2$ -SiCl; viii, $(c-C_6H_{11})_2BH$; ix, AcOH

the bromide by halogen–lithium exchange, and coordination of two of the vinyllithium groups to one copper(I) ion. For comparisons later on, we also wanted the *trans* isomer *E*-10, which we made by the complementary route using *syn* reduction of the silylated acetylene to give the *cis* vinylsilane *Z*-8, and bromination, again with inversion, to give the *trans* vinyl bromide *E*-9.

For reference, we carried out the conjugate addition of the vinylcuprate Z-10 to the trans β -silylenone E-6 that we have used before. With magnesium bromide to coordinate the two oxygen atoms of the imide, the nucleophile attacks anti to the trityloxymethyl group to give as the only detectable product (>95:5, ¹H NMR) the diastereoisomer 11 (Scheme 4). For the synthesis itself, we carried out the conjugate addition of the cis vinylcuprate Z-10 to the cis β -silylenone Z-6. Attack anti to the trityloxymethyl group now gave largely the diastereoisomer Z-12, and, since we had both diastereoisomers, we were able to measure (¹H NMR) the degree of selectivity reliably (95:5). Again for reference purposes, we also prepared the trans imide E-12 from the trans vinylcuprate E-10. This compound appeared (¹H NMR) to have the usual high degree of diastereoisomeric purity, but since we did not need to know the degree of enantiomeric purity of this compound reliably, we did not prepare the missing member of this quartet by treating the trans imide E-6 with the trans vinylcuprate E-10.

We removed the chiral auxiliary by treating the imide Z-12 with lithium benzyl oxide to give the benzyl ester Z-13a (Scheme 5), and presumably with the ratio of enantiomers the same as the ratio of diastereoisomers. We were now ready to control the relative stereochemistry between C-2 and C-3 by alkylation of the enolate derived from the ester Z-13a, which took place, as far as we could tell, with complete selectivity in favour of the *anti* isomer Z-14a. A vinyl or aryl group attached to the stereogenic centre has frequently been particularly effective in this respect, which is why we chose to deal with this





relationship before the hydroboration. We were also hopeful that, with a branched α -carbon, the ester might not be reduced by the hydroborating reagent, but this was not the case—we were unable to avoid reducing the ester group, even after making the *tert*-butyl ester Z-14b in place of the benzyl. We were able to isolate the diol, but all attempts to interrupt the reduction before it had gone to completion gave us only the alcohol derived from reduction of the ester group, but with the double bond intact. For reference later on, we also prepared the *tert*-butyl ester *E*-14b, starting with the *trans* imide *E*-12.

Unable to avoid the over-reduction, we reduced the ester Z-14a deliberately, protected the primary alcohol group as its *tert*-butyldimethylsilyl ether Z-15, and then carried out the hydroboration–oxidation reaction, which took place with high (>95:5) stereocontrol to give, as far as we could tell, only the *syn* isomer 16 (Scheme 6). Although this was very much the

CO₂Bn

OTBS

OTBS

C₆H₁₃-n

C₆H₁₃-n

Z-15 91%

SiMe₂Ph

C₆H₁₃-n

PhMe₂S

E-14b

PhMe₂Si

OH

i, ii

E-15 100%

iii. iv

SiMe₂Ph

C₆H₁₃-n

n-C11H23

n-C11H23

n-C11H29

CO₂Bu^t

OTBS

OTBS

Č₆H₁₃-n

C₆H₁₃-n

PhMe₂S

i, ii

PhMe₂S

iii, iv

OH

Z-14a

n-C11H23

n-C11H23

n-C11H23



Scheme 6 Reagents: i, LiAlH₄; ii, Bu'Me₂SiCl, imidazole; iii, 9-BBN; iv, NaOH, H₂O₂; v, BnOC(=NH)CCl₃, TfOH; vi, TBAF; vii, PDC; viii, Jones

result we expected from our work with similar systems,² we checked that we were not being deceived by the diastereoisomer 16 having a ¹H NMR spectrum unresolved from its C-5 diastereoisomer. To prepare the 3,5-anti isomer 17, we merely had to reduce the ester E-14b with a trans double bond in place of the cis, and subject the silvl ether E-15 to hydroborationoxidation. Although the stereoselectivity in the hydroboration was, as usual with a trans double bond, less impressive, the diastereoisomers proved to be easily distinguished, and there was no trace of any of the distinctive ¹H NMR signals of the anti isomer 17 in the spectrum of the syn isomer 16. To avoid the formation of a δ -lactone later on, we protected the hydroxy group as its benzyl ether 18, necessarily using the acid-catalysed method of ether formation,9 since the standard base-catalysed method would have given a silyl ether by displacement of the phenyl group.¹⁰ We then restored the correct oxidation level to the acid 19.

The silyl-to-hydroxy conversion gave the β -hydroxy acid **20**, for which we were able to use, for the first time in a synthesis, one of our newly developed one-pot procedures.⁴ We did not purify this intermediate but subjected it immediately to benzenesulfonyl chloride¹¹ to make the β -lactone **21**, and removed the protecting group. We obtained the known lactone **22**,¹² our first crystalline intermediate, in good overall yield for the three steps (Scheme 7). One recrystallisation served to give the enantiomerically and diastereomerically pure material, as judged by its sharp and correct melting point and optical



Scheme 7 Reagents: i, Hg(OAc)₂, AcOOH, AcOH; ii, PhSO₂Cl, Py; iii, H₂, Pd/C; iv, DCC, **23**, DMAP, DMF; v, AcOCHO

rotation. We knew that this recrystallisation was removing the 5% of enantiomer, which must have been present throughout. That one recrystallisation was enough, confirmed that both of the other steps controlling relative stereochemistry, $Z-13a \longrightarrow Z-14a$ and $Z-15 \longrightarrow 16$, must have taken place with a very high level of stereoselectivity, remarkable for the fact that both of them were genuinely open-chain control, without any rings or cyclic transition structures. Curiously, although the alcohol 22 was known, none of the earlier syntheses of tetrahydrolipstatin^{13,14,15,16} had used it as an intermediate, preferring instead to use its diastereoisomer at C-5 and a Mitsunobu inversion to attach the leucine residue. Two more recent syntheses have also adopted this tactic.17 Nevertheless, we, and others since,^{18,19} easily esterified the alcohol 22 with the leucine derivative 23 using dicyclohexylcarbodiimide. We then removed the benzyloxycarbonyl protecting group and replaced it with the formyl to give tetrahydrolipstatin 1, identical (mp, mixed mp, rotation, ¹H NMR and mixed ¹H NMR) with an authentic sample.

Experimental

General

The standard aqueous work-up referred to below involved the addition of aqueous ammonium chloride (adjusted to pH 8 by the addition of aqueous ammonia) to the reaction mixture, extraction with diethyl ether $(2-3 \times)$, and separation of the ether layers, which were dried (Na_2SO_4) and evaporated under reduced pressure. Ether refers to diethyl ether.

3-Dimethyl(phenyl)silylprop-1-ynoic acid 4

Ethynyldimethyl(phenyl)silane²⁰ (10 g) in THF (20 cm³) was added to methylmagnesium chloride (3 mol dm⁻³ in THF, 40 cm³) at 0 °C. The mixture was stirred at room temperature for 2 h. Carbon dioxide was bubbled through the solution at -23 °C for 30 min and at room temperature for 2 h. Hydrochloric acid (1 mol dm⁻³, 100 cm³) was added and the mixture extracted with hexane. The extracts were dried (Na₂SO₄) and evaporated under reduced pressure to give the *acid* 4 (12.6 g, 99%); *R*_f(hexane–EtOAc, 1:1) 0.17 (streak); *v*_{max}(film)/cm⁻¹ 3500– 2400 (OH), 2180 (C=C), 1680 (CO), 1250 (SiMe) and 1110 (SiPh); $\delta_{\rm H}$ (CDCl₃) 9.0–8.0 (1 H, br, OH), 7.60–7.32 (5 H, m, Ph) and 0.51 (6 H, s, SiMe₂); $\delta_{\rm C}$ (CDCl₃) 157.4, 134.0, 133.6, 130.1, 128.1, 95.4, 95.0 and -2.0; *m*/*z* 204 (0.4%, M⁺), 189 (0.5, M - Me), 161 (25, M - CO₂ + H), 160 (20, M - CO₂), 145 (100, M - CO₂ - Me) and 137 (20, MePhSiOH) (Found: M⁺, 204.0618. C₁₁H₁₂O₂ requires *M*, 204.0606).

(Z)-3-Dimethyl(phenyl)silylprop-1-enoic acid

The acid **4** (12.8 g, 62.7 mmol) was hydrogenated in methanol (100 cm³) over palladium (10% on BaSO₄, 1 g) in the presence of quinoline (20 cm³) for 3 h. The mixture was filtered over Celite after 1 equivalent of hydrogen (1405 cm³) had been consumed. Evaporation under reduced pressure gave the *acid* (8.2 g, 64%) as prisms, mp 101–103 °C; $R_{\rm f}$ (hexane–EtOAc, 1:1) 0.62; $\nu_{\rm max}$ (CH₂Cl₂) 3500–2000 (OH), 1700 (CO), 1600 (C=C), 1248 (SiMe) and 1118 (SiPh); $\delta_{\rm H}$ (CDCl₃) 11.0 (1 H, br, OH), 7.57–7.32 (5 H, m, Ph), 6.85 (1 H, d, *J* 14.3, SiCH=CH), 6.58 (1 H, d, *J* 14.3, SiCH=CH) and 0.47 (6 H, s, SiMe₂); $\delta_{\rm c}$ (CDCl₃) 172.0, 153.8, 138.7, 135.4, 133.6, 128.9, 127.7 and -2.2; *m/z* 206 (0.8%, M⁺), 205 (1.5, M – H), 191 (100, M – Me), 137 (20, MePhSiOH), 135 (100, PhMe₂Si) and 129 (60, M – Ph) (Found: C, 64.05; H, 6.75; M⁺, 206.0747. C₁₁H₁₄O₂Si requires C, 64.05; H, 6.85%; *M*, 206.0763).

(5*S*)-1-[(*Z*)-3'-Dimethyl(phenyl)silylprop-2-enoyl]-5-(trityloxy-methyl)pyrrolidin-2-one *Z*-6

Oxalyl chloride (1.53 cm³) was added to the alkenoic acid (1.67 g) in dichloromethane (15 cm³). One drop of DMF was added, the mixture warmed to room temperature, stirred for 1 h and the solvent evaporated under reduced pressure. n-Butyllithium (1.6 mol dm^{-3} in hexane, 4.6 cm³) was added to a solution of the lactam²¹ (2.68 g) in THF (5 cm³) at -20 °C to give the lithium salt 5. After 20 min the mixture was cooled to -78 °C and a solution of the crude acid chloride in THF (10 cm³) added. The mixture was warmed to room temperature and stirred for 30 min. Standard aqueous work-up and chromatography (SiO₂, hexane–EtOAc, 3:1) gave the amide (3.68 g, 94%) as prisms, mp 104–106 °C (from hexane); $[a]_{D}^{20}$ –109.7 (c 20.5 in CHCl₃); R_f(hexane-EtOAc, 3:1) 0.60; v_{max}(CDCl₃) 1733 (CO), 1670 (CO), 1248 (SiMe) and 1112 (SiPh); $\delta_{\rm H}$ (CDCl₃) 7.88 (1 H, d, J 14.1, SiCH=CH), 7.61-7.58 (2 H, m, m-SiPh), 7.43-7.20 (18 H, m, o- and p-SiPh and 3 × Ph), 6.78 (1 H, d, J 14.1, SiCH=CH), 4.44 (1 H, m, CHN), 3.58 (1 H, dd, J 3.9 and 9.7, CH_ACH_BOCPh₃), 3.15 (1 H, dd, J 2.5 and 9.7, CH_AH_BOCPh₃), 2.98 (1 H, dt, J 17.9 and 10.3, NCOCH_ACH_B), 2.50 (1 H, ddd, J 17.9, 9.2 and 2.4, NCOC_ACH_B), 2.00 (2 H, m, COCH₂CH₂), 0.47 (3 H, s, $SiMe_AMe_B$) and 0.44 (3 H, s, $SiMe_AMe_B$); δ_c(CDCl₃) 176.3, 165.9, 149.5, 143.7, 139.5, 138.0, 133.6, 128.7, 128.6, 128.0, 127.6, 127.2, 87.1, 64.0, 56.6, 33.1, 21.2, -1.7 and -2.3; m/z 545 (3%, M⁺), 530 (30, M – Me), 302 (15, M - CPh₃) and 243 (100, CPh₃) (Found: C, 76.5; H, 6.50; N, 2.50; M⁺, 545.2382. C₃₅H₃₅NO₃Si requires C, 77.0; H, 6.45; N, 2.55%; M, 545.2386).

Tridec-1-yne 7

Bromoundecane (5.13 cm³, 22.9 mmol) was added to a slurry of lithium acetylide–ethylenediamine complex (3.06 g, 33 mmol) in DMSO (15 cm³) at 0 °C. The mixture was warmed to room temperature and stirred for 2 h. Standard aqueous work-up and chromatography (SiO₂, hexane) gave tridecyne²² (4.13 g, 100%); $R_{\rm f}$ (hexane) 0.47; $v_{\rm max}$ (film)/cm⁻¹ 3310 (=CH) and 2120 (C=C); $\delta_{\rm H}$ (CDCl₃) 2.17 (2 H, td, *J* 7.0 and 2.6, CH₂C=CH), 1.93 (1 H, t, *J* 2.6, C=CH), 1.50–1.20 (18 H, m, CH₂) and 0.87 (3 H, t, *J* 6.6, Me); $\delta_{\rm C}$ (CDCl₃) 84.7, 68.0, 31.9, 29.6 (2), 29.5, 29.4, 29.1, 28.8, 28.5, 22.7, 18.4 and 14.1; *m*/*z* 165 (0.2%, M – Me), 151 (0.5, M – Et), 109 (15, C₈H₁₃), 95 (50, C₇H₁₁), 82 (55, C₆H₁₀), 81 (100, C₆H₉), 67 (70, C₅H₇) and 55 (50, C₄H₇) (Found: M⁺ – Me, 165.1648. C₁₂H₂₁ requires *M* – Me, 165.1644).

(E)-1-Dimethyl(phenyl)silyltridec-1-ene E-8

The silvllithium reagent²³ (1.14 mol dm⁻³ in THF, 73.3 cm³,

83.6 mmol) was added over 2 min to a slurry of copper(I) cyanide (3.74 g, 41.9 mmol) in THF (50 cm³) at 0 °C. The mixture was stirred at 0 °C for 20 min and cooled to -78 °C. Tridec-1yne (5.7 g, 31.7 mmol) was added in ether (15 cm³) and the mixture stirred 2 h, and for a further 30 min at 0 °C. The reaction was quenched with aqueous ammonium chloride and filtered over Celite. The aqueous layer was extracted with ether, and the organic layers were dried (Na₂SO₄) and evaporated under reduced pressure. Chromatography (SiO₂, hexane) gave the vinylsilane (10.5 g, 95%); $R_{\rm f}$ (hexane) 0.51; $v_{\rm max}$ (film)/cm⁻¹ 1610 (C=C), 1245 (SiMe) and 1110 (SiPh); $\delta_{\rm H}$ (CDCl₃) 7.54–7.32 (5 H, m, Ph), 6.12 (1 H, dt, J 18.5 and 6.2, SiCH=CH), 5.74 (1 H, dt, J 18.5 and 1.4, SiCH=CH), 2.13 (2 H, m, CH= CHCH₂), 1.40–1.20 (18 H, m, CH₂), 0.88 (3 H, t, J 6.6, Me) and 0.31 (6 H, s, SiMe₂); *m*/*z* 316 (6%, M⁺), 301 (40, M – Me), 162 $(30, M - C_{11}H_{22}), 161 (30, M - C_{11}H_{23}), 135 (50, PhMe_2Si)$ and 121 (100, MePhSiH) (Found: M⁺, 316.2572. C₂₁H₃₆Si requires M, 316.2586).

(Z)-1-Bromotridec-1-ene Z-9

Bromine (0.45 cm³, 8.75 mmol) was added to the vinylsilane E-8 (2.77 g, 8.75 mmol) in dichloromethane (20 cm³) at -78 °C and stirred for 10 min. Sodium sulfite (1 g) and methanol (10 cm³) were added at -78 °C and the mixture stirred for 10 min. The still cold mixture was poured into saturated aqueous sodium sulfite and extracted with pentane. The extracts were dried (Na_2SO_4) and evaporated under reduced pressure. Freshly prepared sodium methoxide (1 mol dm⁻³ in MeOH, 11 cm³) was added to the crude dibromide in methanol (10 cm³) at 0 °C. The solution was stirred at 0 °C for 1 h and at room temperature for 2 h. Standard aqueous work-up and chromatography (SiO₂, hexane) gave the vinyl bromide (2.19 g, 96%); $R_{\rm f}$ (hexane) 0.64; v_{max} (film)/cm⁻¹ 1623 (C=C); δ_{H} (CDCl₃) 6.12 (1 H, dt, J 6.9 and 1.2, CH=CHBr), 6.07 (1 H, q, J 6.9, CH=CHBr), 2.17 (2 H, m, CH2C=CHBr), 1.40-1.20 (18 H, m, CH2) and 0.87 (3 H, t, J 6.4, Me); $m/z \ 260 \ (5\%, M^+)$, $162 \ (5, M - C_8H_{16})$, $97 \ (65, C_7H_{13})$, 83(80, C₆H₁₁), 69 (75, C₅H₉), 57 (100, C₄H₉) and 55 (70, C₄H₇) (Found: M⁺ 260.1149. C₁₃H₂₅Br requires *M*, 260.1139).

1-Dimethyl(phenyl)silyltridec-1-yne

Tridec-1-yne (2.0 g) was stirred with butyllithium (1.6 mol dm⁻³, 7.5 cm³) in THF (10 cm³) for 30 min, and then refluxed with chlorodimethyl(phenyl)silane (1.8 g) for 18 h. Standard work-up gave the *ethynylsilane* (3.04 g, 86%); $R_{\rm f}$ (hexane) 0.32; $v_{\rm max}$ (film)/cm⁻¹ 2170 (C=C), 1250 (SiMe) and 1115 (SiPh); $\delta_{\rm H}$ (CDCl₃) 7.63–7.34 (5 H, m, Ph), 2.26 (2 H, t, *J* 7.0, CH₂C=C), 1.50 (2 H, m, CH₂CH₂C=C), 1.40–1.20 (16 H, m, CH₂), 0.88 (3 H, t, *J* 6.6, Me) and 0.38 (6 H, s, SiMe₂); $\delta_{\rm C}$ (CDCl₃) 137.8, 133.7, 129.3, 127.8, 109.7, 82.3, 32.0, 29.7 (×2), 29.6, 29.4, 29.2, 28.9, 28.7, 22.8, 20.0, 14.2 and -0.5; *m/z* 314 (0.1%, M⁺), 299 (60, M – Me), 135 (100, PhMe₂Si) and 121 (50, MePhSiH) (Found: M⁺, 314.2454. C₂₁H₃₄Si requires *M*, 314.2430).

(Z)-1-Dimethyl(phenyl)silyltridec-1-ene Z-8

The ethynylsilane (2.34 g) and dicyclohexylborane (8 mmol) were stirred in THF (7 cm³) at room temperature for 2 h. Acetic acid (1.2 cm³) was added and the mixture stirred for a further 1 h. Water was added, and the organic layer washed with aqueous sodium hydrogen carbonate (saturated, 100 cm³), dried (Na₂SO₄) and the solvent removed under reduced pressure. Chromatography of the residue [SiO₂, light petroleum (bp 30-40 °C)] gave the vinylsilane (2.04 g, 87%); $R_{\rm f}$ (hexane) 0.49; v_{max} (film)/cm⁻¹ 1605 (C=C), 1250 (SiMe) and 1110 (SiPh); $\delta_{\rm H}$ (CDCl₃) 7.56–7.31 (5 H, m, Ph), 6.42 (1 H, dt, J 14 and 7.4, SiCH=CH), 5.61 (1 H, dt, 14 and 1.1, SiCH=CH), 2.00 (2 H, m, CH₂CH=CH), 1.25 (18 H, m, CH₂), 0.88 (3 H, t, J 6.6, Me) and 0.37 (6 H, s, SiMe₂); $\delta_{\rm C}$ (CDCl₃) 151.1, 139.8, 133.7, 128.8, 127.7, 126.4, 33.8, 32.0, 29.7 (×2), 29.6 (×2), 29.5, 29.4, 29.3, 22.7, 14.1 and -0.8; m/z 316 (21%, M⁺), 301 (50, M - Me), 162 $(50, M - C_{11}H_{22})$, 161 (50, $M - C_{11}H_{23}$), 135 (85, PhMe₂Si) and

(E)-1-Bromotridec-1-ene E-9

This was prepared in the same way as the vinyl bromide Z-9 from the vinylsilane Z-8 (1.93 g) to give the vinyl bromide (1.62 g, 95%); $R_{\rm f}({\rm hexane})$ 0.61; $v_{\rm max}({\rm film})/{\rm cm}^{-1}$ 1620 (C=C) and 930 (trans-CH=CH); $\delta_{\rm H}({\rm CDCl}_3)$ 6.16 (1 H, dt, J 13.5 and 7.0, CH=CHBr), 5.99 (1 H, dt, J 13.5 and 1.2, CH=CHBr), 2.00 (2 H, m, CH₂CH=CH), 1.37–1.20 (18 H, m, CH₂) and 0.87 (3 H, t, J 6.5, Me); $\delta_{\rm C}({\rm CDCl}_3)$ 138.3, 104.0, 32.9, 31.9, 29.63 (×2), 27.56, 29.4, 29.3, 29.0, 28.6, 22.7 and 14.1; m/z 260 (3%, M⁺), 162 (3, M - C₇H₁₄), 148 (5, M - C₈H₁₆), 97 (60, C₇H₁₃), 83 (70, C₆H₁₁), 69 (75, C₅H₉), 57 (100, C₄H₉) and 55 (70, C₄H₇) (Found: M⁺, 260.1122. C₁₃H₂₅Br requires M, 260.1140).

(5*S*)-1-[(*Z*)-(3'*R*)-3'-Dimethyl(phenyl)silylhexadec-4'-enoyl]-5-(trityloxymethyl)pyrrolidin-2-one *Z*-12

The vinyl bromide Z-9 (2.95 g, 11.3 mmol) in ether (20 cm³) was added to lithium wire (containing 1% Na, 300 mg, 13 mmol) at -20 °C and stirred for 3 h. The resulting vinyllithium was added to copper bromide-dimethyl sulfide complex²⁴ (1.16 g, 5.65 mmol) in THF (32 cm³) and dimethyl sulfide (16 cm³) at -40 °C, stirred for 30 min and cooled to -78 °C. A slurry of the lactam Z-6 (2.43 g, 4.45 mmol) and anhydrous magnesium bromide²⁵ (4.36 g) in THF (30 cm³) was added very slowly to the cuprate. After complete addition, the mixture was stirred at -78 °C for 1 h and warmed to 0 °C over 90 min. Standard aqueous work-up and chromatography (SiO₂, hexane-EtOAc, 20:1 and then 10:1) gave an inseparable mixture (5:95) of 11 and the *allylsilane* (3.06 g, 94%); $[a]_{D}^{20}$ -26.3 (c 1.63 in CHCl₃); $R_{\rm f}$ (hexane-EtOAc, 5:1) 0.44; $v_{\rm max}$ (film)/cm⁻¹ 1740 (CO), 1695 (CO) and 1600 (C=C); $\delta_{\rm H}$ (CDCl₃) 7.55–7.50 (2 H, m, *o*-SiPh), 7.37-7.14 (18 H, m, CPh₃, m- and p-SiPh), 5.23 (1 H, dt, J 10.9 and 6.8, SiCHCH=CH), 5.05 (1 H, t, J 10.9, SiCHCH=CH), 4.39 (1 H, m, CHN), 3.34 (1 H, dd, J 9.6 and 4.4, CH_AH_B-OCPh₃), 3.17 (1 H, dd, J 16.4 and 11.4, SiCHCH_AH_B), 3.13 (1 H, m, CH_ACH_BOCPh₃), 2.81 (1 H, dt, J 17.8 and 10.2, $NCOCH_AH_B$), 2.73 (1 H, dd, J 16.4 and 3.0, SiCHCH_ACH_B), 2.61 (1 H, td, J 11.4 and 3.0, SiCH), 2.10-1.80 (4 H, m, NCO-CH₂CH₂ and CH₂), 1.24 (18 H, m, CH₂), 0.87 (3 H, t, J 6.6, Me), 0.33 (3 H, s, $SiMe_AMe_B$) and 0.30 (3 H, s, $SiMe_AMe_B$); $\delta_{\rm C}({\rm CDCl_3})$ 176.1, 173.4, 143.6, 137.2, 134.1, 129.8, 129.5, 129.1, 128.9, 128.5, 127.8, 127.1, 127.1, 86.9, 63.9, 56.6, 37.2, 33.0, 31.9, 29.7, 29.6, 29.5, 29.3, 27.6, 24.0, 22.7, 21.0, 14.1, -4.4 and $-5.3;\,m/z$ 727 (5%, M⁺), 484 (4, M - CPh₃), 243 (100, CPh₃) and 135 (35, PhMe₂Si) (Found: M⁺, 727.4492. C₄₈H₆₁NO₃Si requires M, 727.4420). The ratio of isomers was determined by integration of the $SiMe_2$ signals in the $^1\!H$ NMR spectrum.

(5*S*)-1-[(*Z*)(3'*S*)-3'-Dimethyl(phenyl)silylhexadec-4'-enoyl]-5-(trityloxymethyl)pyrrolidin-2-one 11

This was made in the same way as the allylsilane Z-12 from the vinylcuprate Z-10 (1.02 mmol) and the lactam $E-6^5$ (187 mg, 0.34 mmol) to give the *allylsilane* (188 mg, 75%); $[a]_{D}^{20}$ -23.9 (c 2.98 in CHCl₃); R_f (hexane-EtOAc, 5:1) 0.44; v_{max} (film)/cm⁻¹ 1740 (CO), 1690 (CO) and 1600 (SiPh); $\delta_{\rm H}$ (CDCl₃) 7.57–7.50 (2 H, m, o-SiPh), 7.44-7.10 (18 H, m, CPh₃, m- and p-SiPh), 5.25 (1 H, dt, J 10.8 and 6.8, SiCHCH=CH), 5.12 (1 H, t, J 10.8, SiCHCH=CH), 4.30 (1 H, m, CHN), 3.43 (1 H, dd, J 9.7 and 4.3, CH_ACH_BOCPh₃), 3.20-3.11 (2 H, m, CH_AH_BOCPh₃ and SiCHCH_ACH_B), 2.84 (1 H, dt, J 18.3 and 10.5, NCOCH_A-CH_B), 2.64 (1 H, m, SiCHCH_ACH_B), 2.41 (1 H, m, NCOCH_A-CH_B), 2.28 (1 H, m, SiCH), 2.00–1.80 (4 H, m, NCOCH₂CH₂ and CH₂), 1.25 (18 H, m, CH₂), 0.88 (3 H, t, J 6.5, Me), 0.324 (3 H, s, Si Me_AMe_B) and 0.321 (3 H, s, Si Me_AMe_B); $\delta_C(CDCl_3)$ 175.8, 173.3, 143.6, 137.0, 134.1, 129.2, 129.0, 128.7, 128.5, 127.8, 127.5, 127.0, 86.9, 63.7, 56.7, 37.9, 33.0, 31.0, 29.7, 29.64, 29.60, 29.5, 29.3, 27.6, 23.9, 22.6, 21.1, 14.1, -4.6 and -5.1; m/z 727 (2%, M⁺), 485 (6, M - CPh₃ + H), 243 (100, CPh₃)

and 135 (20, PhMe₂Si) (Found: M^+ , 727.4463. C₄₈H₆₁NO₃Si requires *M*, 727.4420). The diastereoisomer *Z*-12 was not detectable (¹H NMR).

(5*S*)-1-[(*E*)(3'*R*)-3'-Dimethyl(phenyl)silylhexadec-4'-enoyl]-5-(trityloxymethyl)pyrrolidin-2-one *E*-12

This was prepared in the same way as the allylsilane Z-12, from the vinyl bromide E-9(1.4 g) and the lactam Z-6(1.15 g) to give the allylsilane (1.12 g, 73%); $[a]_{D}^{21}$ -35.0 (c 1.3 in CHCl₃); $R_{\rm f}$ (hexane–EtOAc, 5:1) 0.43; $v_{\rm max}$ (film)/cm⁻¹ 1740 (CO), 1690 (CO) and 1600 (C=C); δ_H(CDCl₃) 7.53–7.47 (2 H, m, *o*-SiPh), 7.36-7.14 (18 H, m, CPh₃, m- and p-SiPh), 5.27-5.11 (2 H, m, CH=CH), 4.38 (1 H, m, CHN), 3.36 (1 H, dd, J 9.7 and 4.5, CH_AH_BOCPh₃), 3.21 (1 H, dd, J 16.4 and 11.7, SiCHCH_AH_B), 3.14 (1 H, dd, J 9.7 and 2.9, CH_AH_BOCPh₃), 2.83 (1 H, dt, J 17.7 and 10.5, NCOCH_ACH_B), 2.71 (1 H, dd, J 16.4 and 3.1, SiCHCH_ACH_B), 2.25 (1 H, ddd, J 11.7, 7.6 and 3.1, SiCH), 2.10-1.80 (4 H, m, CH₂ and COCH₂CH₂), 1.25 (18 H, m, CH₂), 0.87 (3 H, t, J 6.6, Me), 0.31 (3 H, s, $SiMe_AMe_B$) and 0.29 (3 H, s, SiMe_AMe_B); δ_C(CDCl₃) 176.1, 173.4, 143.6, 137.3, 134.1, 129.3, 129.2, 129.1, 128.5, 127.8, 127.7, 86.9, 63.8, 56.6, 36.3, 33.0, 32.7, 31.9, 29.8, 29.7 (×2), 29.5, 29.3, 29.1, 27.9, 22.7, 21.0, 14.1, -4.3 and -5.3; m/z 727 (5%, M⁺), 712 (1, M - Me), 649 (5, M - C₆H₆), 484 (5, M - CPh₃), 243 (100, CPh₃) and 135 (50, PhMe₂Si) (Found: M^+ , 727.4440. $C_{48}H_{61}NO_3Si$ requires M, 727.4220).

Benzyl (Z)(3R)-3-dimethyl(phenyl)silylhexadec-4-enoate Z-13a

The lactam Z-12 (1.61 g) in THF (7 cm³) was added to lithium benzyl oxide [prepared from *n*-butyllithium (1.6 mol dm⁻³ in hexane, 7.7 cm³) and benzyl alcohol in THF (10 cm³)] and stirred at room temperature for 24 h. Standard aqueous workup and chromatography (SiO₂, hexane-EtOAc, 10:1) gave the *ester* (0.95 g, 93%); $[a]_{D}^{20} - 12.7$ (*c* 1.90 in CHCl₃); R_{f} (hexane–EtOAc, 5:1) 0.60; v_{max} (film)/cm⁻¹ 1730 (CO), 1250 (SiMe) and 1110 (SiPh); δ_H(CDCl₃) 7.57–7.44 (2 H, m, *o*-SiPh), 7.38–7.27 (8 H, m, CH₂Ph, m- and p-SiPh), 5.31 (1 H, dt, J 10.8 and 7.0, SiCHCH=CH), 5.08 (1 H, m, SiCHCH=CH), 5.02 (1 H, d, J 12.1, CH_AH_BPh), 4.97 (1 H, d, J 12.1, CH_AH_BPh), 2.55 (1 H, td, J 11.3 and 3.5, SiCH), 2.42 (1 H, dd, J 14.6 and 3.5, CH_AH_B-CO), 2.22 (1 H, dd, J 14.6 and 11.7, CH_AH_BCO), 2.00-1.80 (2 H, m, CH₂), 1.25 (18 H, m, CH₂), 0.88 (3 H, t, J 6.5, Me), 0.28 (3 H, s, $SiMe_AMe_B$) and 0.27 (3 H, s, $SiMe_AMe_B$); $\delta_{\rm C}({\rm CDCl}_3)$ 173.3, 136.7, 134.0, 130.0, 129.2, 128.4, 128.2, 128.0, 127.7, 66.1, 35.5, 31.9, 29.7, 29.6, 29.5, 29.4, 27.6, 25.1, 22.7, 14.1, -4.6 and -5.4; m/z 478 (0.04%, M⁺), 463 (0.5, M - Me), $400 (2, M - C_6H_6), 387 (4, M - C_7H_7) \text{ and } 135 (100, PhMe_2Si)$ (Found: M⁺, 478.3236. C₃₁H₄₆O₂Si requires *M*, 478.3267).

tert-Butyl (*Z*)(3*R*)-3-dimethyl(phenyl)silylhexadec-4-enoate *Z*-13b

This was prepared in the same way as the benzyl ester Z-13a, from the lactam Z-12 (1.98 g, 2.72 mmol) and lithium *tert*-butyl oxide (30 mmol), stirring for 48 h to give the *allylsilane* (1.03 g, 85%); $R_{\rm f}$ (hexane–EtOAc, 5:1) 0.64; $v_{\rm max}$ (film)/cm⁻¹ 1730 (CO) and 1250 (SiMe); $\delta_{\rm H}$ (CDCl₃) 7.50–7.47 (2 H, m, *o*-SiPh), 7.37– 7.30 (3 H, m, *m*- and *p*-SiPh), 5.28 (1 H, dt, *J* 10.9 and 7.0, SiCHCH=CH), 5.08 (1 H, m, SiCHCH=CH), 2.50 (1 H, td, *J* 11.5 and 3.4, SiCH), 2.28 (1 H, dd, *J* 14.4 and 3.4, SiCHCH_A-CH_B), 2.04 (1 H, dd, *J* 14.4 and 11.5, SiCHCH_ACH_B), 2.00–1.80 (2 H, m, CH₂), 1.37 (9 H, s, Bu'), 1.25 (18 H, m, CH₂), 0.88 (3 H, t, *J* 6.6, Me), 0.28 (3 H, s, SiMe_AMe_B) and 0.27 (3 H, s, SiMe_AMe_B); *mlz* 444 (2.4%, M⁺), 443 (6, M – H), 309 (95, M – PhMe₂Si) and 135 (100, PhMe₂Si) (Found: M⁺, 444.3404. C₂₈H₄₈O₂Si requires *M*, 444.3423).

Benzyl (Z)(2R,3S)-3-dimethyl(phenyl)silyl-2-hexylhexadec-4enoate Z-14a

This was prepared in the same way as the ester Z-13b, from the ester Z-13a (150 mg), to give the *allylsilane* (150 mg, 85%); $[a]_{D}^{20}$

-21.4 (*c* 2.33 in CHCl₃); $R_{\rm f}$ (hexane–EtOAc, 5:1) 0.65; $v_{\rm max}$ (film)/cm⁻¹ 1730 (CO), 1250 (SiMe) and 1110 (SiPh); $\delta_{\rm H}$ (CDCl₃) 7.52–7.46 (2 H, m, *o*-SiPh), 7.40–7.26 (8 H, m, CH₂-*Ph*, *m*- and *p*-SiPh), 7.35 (1 H, dt, *J* 10.8 and 7.2, SiCHCH=CH), 4.82 (1 H, d, *J* 12.5, OCH_ACH_BPh), 4.75 (1 H, d, *J* 12.5, CH_AH_BPh), 2.43 (2 H, m, SiCHCH), 1.90–1.80 (2 H, m, CH₂), 1.25–1.10 (28 H, m, CH₂), 0.85 (6 H, m, 2 × Me), 0.29 (3 H, s, SiMe_AMe_B) and 0.24 (3 H, s, SiMe_AMe_B); $\delta_{\rm C}$ (CDCl₃) 175.5, 137.5, 136.1, 134.2, 130.5, 128.4, 128.2, 128.0, 127.7, 127.5, 127.0, 65.7, 46.1, 31.9, 31.6, 30.6, 29.7–29.1 (m), 22.7, 22.6, 14.1, 14.0, -3.4 and -4.3; *m*/*z* 471 (1.2%, M – CH₂Ph) and 135 (100, PhMe₂Si) (Found: M⁺ – CH₂Ph, 471.3693. C₃₇H₅₈O₂Si requires *M* – CH₂Ph, 471.3658).

tert-Butyl (*Z*)(2*R*,3*S*)-3-dimethyl(phenyl)silyl-2-hexylhexadec-4-enoate *Z*-14b

The ester Z-13b (365 mg) was added to LDA (0.24 mol dm⁻³ in THF, 3.4 cm³) at -78 °C and the mixture stirred for 30 min. Iodohexane (0.74 cm³) was added and the mixture stirred at -78 °C for 1 h and warmed to 0 °C over 1 h. Standard aqueous work-up gave the *allylsilane* (324 mg, 75%); $R_{\rm f}$ (hexane–EtOAc, 5:1) 0.72; $v_{\rm max}$ (film)/cm⁻¹ 1725 (CO), 1250 (SiMe) and 1110 (SiPh); $\delta_{\rm H}$ (CDCl₃) 7.52–7.47 (2 H, m, *o*-SiPh), 7.36–7.29 (3 H, m, *m*- and *p*-SiPh), 5.30 (1 H, dt, *J* 11.8 and 6.8, SiCHCH=*CH*), 5.14 (1 H, m, SiCHC*H*=CH), 2.46 (1 H, dd, *J* 11.8 and 7.5, SiCH), 2.34 (1 H, m, COCH), 1.90–1.80 (2 H, m, CH₂), 1.39 (9 H, s, Bu'), 1.25–1.20 (28 H, m, CH₂), 0.86 (6 H, m, 2 × Me) and 0.29 (6 H, s, SiMe₂); *m/z* 528 (1.6%, M⁺), 457 (4, M - C₃H₁₁), 443 (5, M - C₆H₁₅), 309 (90, M - C₆H₁₄ – PhMe₂Si) and 135 (100, PhMe₂Si) (Found: M⁺, 528.4334. C₃₄H₆₀O₂Si requires *M*, 528.4362).

tert-Butyl (*E*)(3*R*)-3-dimethyl(phenyl)silylhexadec-4-enoate *E*-13b

This was prepared in the same way as the benzyl ester Z-13a, from the lactam *E*-12 (2.98 g), to give the *allylsilane* (0.78 g, 43%); $R_{\rm f}$ (hexane–EtOAc, 5:1) 0.64; $v_{\rm max}$ (film)/cm⁻¹ 1730 (CO) and 1250 (SiMe); $\delta_{\rm H}$ (CDCl₃) 7.49–7.45 (2 H, m, *o*-SiPh), 7.36–7.31 (3 H, m, *m*- and *p*-SiPh), 5.24 (2 H, m, SiCHCH=CH), 2.30–2.10 (3 H, m, SiCHCH₂), 1.94 (2 H, m, CH₂), 1.37 (9 H, s, Bu'), 1.24 (18 H, m, CH₂), 0.87 (3 H, t, *J* 6.4, Me) and 0.26 (6 H, s, SiMe₂); *m/z* 444 (4%, M⁺), 373 (10, M – C₅H₁₁), 310 (25, M – PhMe₂Si + H), 155 (40, C₁₁H₂₃) and 135 (100, PhMe₂Si) (Found: M⁺, 444.3428. C₂₈H₄₈O₂Si requires *M*, 444.3423).

tert-Butyl (*E*)(2*R*,3*S*)-3-dimethyl(phenyl)silyl-2-hexylhexadec-4-enoate *E*-14b

This was prepared in the same way as the ester Z-14b, from the ester *E*-13b (220 mg), to give the *allylsilane* (132 mg, 51%); $R_{\rm f}$ (hexane–EtOAc, 5:1) 0.72; $v_{\rm max}$ (film)/cm⁻¹ 1725 (CO), 1250 (SiMe) and 970 (*trans*-CH=CH); $\delta_{\rm H}$ (CDCl₃) 7.52–7.46 (2 H, m, *o*-SiPh), 7.33–7.29 (3 H, m, *m*- and *p*-SiPh), 5.14 (2 H, m, SiCHC*H*=C*H*), 2.29 (1 H, m, SiCH), 2.08 (1 H, dt, *J* 3.2 and 6.7, COCH), 1.94 (2 H, m, CH₂), 1.45 (2 H, m, CH₂), 1.38 (9 H, s, Bu'), 1.25–1.10 (26 H, m, CH₂), 0.88 (6 H, m, 2 × Me), 0.29 (3 H, s, SiMe_AMe_B) and 0.28 (3 H, s, SiMe_AMe_B); *m*/*z* 528 (1, M⁺), 472 (1, M – C₄H₈), 457 (2, M – C₅H₁₁), 444 (5, M – C₆H₁₄), 309 (70, M – C₆H₁₄ – PhMe₂Si) and 135 (100, PhMe₂Si) (Found: M⁺, 528.4360. C₃₄H₆₀O₂Si requires *M*, 528.4362).

(Z)(3R)-3-Dimethyl(phenyl)silylhexadec-4-en-1-ol

Hydroboration (9-BBN, 1.5 equiv., 4 h, room temperature)² of the allylsilane Z-13a (93 mg) gave a mixture of the starting material (52 mg, 56%) and the *alcohol* (23 mg, 32%); *R*_f(hexane–EtOAc, 3:1) 0.47; *ν*_{max}(film)/cm⁻¹ 3350 (OH), 1250 (SiMe) and 1110 (SiPh); $\delta_{\rm H}$ (CDCl₃) 7.51–7.26 (5 H, m, Ph), 5.34 (1 H, dt, *J* 11.0 and 7.0, SiCHCH=CH), 5.14 (1 H, t, *J* 11.0, Si-CHCH=CH), 3.63–3.44 (2 H, m, CH₂OH), 2.10 (1 H, td, *J* 11.6 and 2.6, SiCH), 2.00–1.10 (23 H, CH₂, SiCHCH₂ and OH), 0.88

(3 H, t, J 6.1, Me), 0.28 (3 H, s, $SiMe_AMe_B$) and 0.27 (3 H, s, $SiMe_AMe_B$); m/z 259 (0.1%, M – Me) and 135 (100, PhMe₂Si) (Found: M⁺ – Me, C₂₄H₄₂OSi requires M – Me, 359.2770).

(3R,5S)-3-Dimethyl(phenyl)silylhexadecane-1,5-diol

Hydroboration (BH₃·THF, 2 equiv., 0 °C, 2 h)² of the allylsilane Z-13a (117 mg) gave the *diol* (20.1 mg, 18%); $R_{\rm f}$ (hexane– EtOAc, 3:1) 0.21; $v_{\rm max}$ (film)/cm⁻¹ 3300 (OH), 1250 (SiMe) and 1110 (SiPh); $\delta_{\rm H}$ (CDCl₃) 7.55–7.28 (5 H, m, Ph), 3.76–3.40 (3 H, m, CHOH and CH₂OH), 2.30 (2 H, s, OH), 1.80–1.10 (25 H, m, CH₂ and CH₂SiCHCH₂), 0.88 (3 H, t, *J* 6.0, Me), 0.36 (3 H, s, Si $Me_{\rm A}Me_{\rm B}$) and 0.28 (3 H, s, SiMe_A $Me_{\rm B}$); *m*/*z* 374 (0.2%, M – H₂O), 359 (0.3, M – Me – H₂O), 205 (50), 137 (50, MePhSiOH) and 135 (100, PhMe₂Si) (Found: M⁺ – H₂O, 374.2991. C₂₄H₄₂OSi requires M – H₂O, 374.3004).

(Z)(2R,3S)-3-Dimethyl(phenyl)silyl-2-hexylhexadec-4-en-1-ol

Hydroboration (9-BBN, 1.3 equiv., reflux, 3 h)² and oxidation of the allylsilane Z-14b (80 mg) gave, after chromatography (SiO₂, hexane–EtOAc, 5:1), the starting material (30 mg, 38%) and the *alcohol* (24 mg, 29%); $R_{\rm f}$ (hexane–EtOAc, 5:1) 0.50; $v_{\rm max}$ (film)/cm⁻¹ 3400 (OH), 1250 (SiMe) and 1110 (SiPh); $\delta_{\rm H}$ (CDCl₃) 7.52–7.33 (5 H, m, Ph), 5.35 (2 H, m, CH=CH), 3.50–3.30 (3 H, m, CH₂OH and CHOH), 2.00–1.00 (32 H, m, CH₂ and SiCHCH), 0.87 (6 H, m, 2 × Me), 0.32 (3 H, s, SiMe_A-Me_B) and 0.29 (3 H, s, SiMe_AMe_B); $\delta_{\rm C}$ (CDCl₃) 138.6, 133.9, 130.6, 128.9, 127.7, 127.3, 65.7, 41.6, 41.9, 31.8, 30.3, 29.8, 29.6, 29.5, 29.4, 28.6, 28.0, 27.6, 22.7, 22.6, 14.1, 14.0, -3.1 and -3.4; *m*/*z* 457 (0.1%, M – Me), 306 (6, M – PhMe₂SiOH), 152 (5, PhMe₂SiOH) and 135 (100, PhMe₂Si) (Found: M⁺ – H, 457.3847. C₃₀H₅₄OSi requires *M* – H, 457.3866).

(Z)(7R,8S)-7-(tert-Butyldimethylsilyloxymethyl)-8-dimethyl-(phenyl)silylhenicos-9-ene Z-15

The ester Z-14a (1.87 g) in ether (20 cm³) was added to a slurry of lithium aluminium hydride (0.7 g) in ether (20 cm³) and stirred at 0 °C for 3 h. Standard aqueous work-up gave crude alcohol [R_f (hexane-EtOAc, 5:1) 0.47] which was stirred with imidazole (0.91 g) and tert-butylchlorodimethylsilane in DMF (20 cm³) at room temperature for 1 h. Standard aqueous work-up and chromatography (SiO₂, hexane–EtOAc, 10:1) gave the *allylsilane* (1.75 g, 91%); $[a]_D^{20}$ –21.4 (*c* 2.33 in CHCl₃); $R_{\rm f}$ (hexane-EtOAc, 5:1) 0.74; $v_{\rm max}$ (film)/cm⁻¹ 1250 (SiMe); δ_H(CDCl₃) 7.52–7.47 (2 H, m, o-SiPh), 7.35–7.28 (3 H, m, mand p-SiPh), 5.35 (1 H, dt, J 11.4 and 7.0, SiCHCH=CH), 5.21 (1 H, t, J 11.4, SiCHCH=CH), 3.36 (1 H, dd, J 9.5 and 4.9, CH_ACH_BO), 3.20 (1 H, dd, J 9.5 and 8.4, CH_ACH_BO), 2.53 (1 H, dd, J 11.7 and 3.3, SiCH), 2.00-1.50 (3 H, m, SiCHCH and CH₂), 1.30–1.15 (28 H, m, CH₂), 0.86 (15 H, m, 2 × Me and Bu'), 0.29 (3 H, s, Si Me_AMe_BPh), 0.26 (3 H, s, Si Me_AMe_BPh), -0.04 (3 H, s, SiMe_AMe_BBu^t) and -0.06 (3 H, s, SiMe_AMe_B-Bu'); m/z 515 (0.5%, M - Bu'), 440 (0.5, M - TBDMSOH), 427 (1, M - TBDMSOCH₂), 379 (10, M - PhMe₂Si - Bu'), 365 (12, M - PhMe₂Si - Bu' - Me), 209 (50, PhMe₂SiOSi-Me₂) and 135 (100, PhMe₂Si) (Found: M⁺ - Bu^t, 515.4113. $C_{36}H_{68}OSi_2$ requires M - Bu', 515.4104).

(*E*)(7*R*,8*S*)-7-(*tert*-Butyldimethylsilyloxymethyl)-8-dimethyl-(phenyl)silylhenicos-9-ene *E*-15

This was prepared in the same way as the allylsilane Z-15, from the ester *E*-14b (119 mg), to give the *allylsilane* (130 mg, 100%); $R_{\rm f}$ (hexane–EtOAc, 5:1) 0.74; $\nu_{\rm max}$ (film)/cm⁻¹ 1250 (SiMe); $\delta_{\rm H}$ (CDCl₃) 7.50–7.46 (2 H, m, *o*-SiPh), 7.34–7.28 (3 H, m, *m*- and *p*-SiPh), 5.21 (2 H, m, SiCHC*H*=C*H*), 3.37 (1 H, dd, *J* 9.6 and 4.8, C*H*_AH_BO), 3.25 (1 H, t, *J* 9.6, CH_AC*H*_B), 2.10 (1 H, m, SiCH), 1.96 (2 H, m, CH₂), 1.30–1.10 (28 H, m, CH₂), 0.86 (15 H, m, 2 × Me and Bu'), 0.28 (3 H, s, SiMe_AMe_BPh), 0.25 (3 H, s, SiMe_AMe_BPh), -0.03 (3 H, s, SiMe_AMe_BBu') and -0.04 (3 H, s, SiMe_AMe_BBu'); *m*/*z* 515 (0.2%, M – Bu'), 440 (1, M – TBDMSOH), 427 (0.5, M – TBDMSOCH₂), 379 (12, $M - PhMe_2Si - Bu'$), 365 (12, $M - PhMe_2Si - Bu' - Me$), 209 (20, $PhMe_2SiOSiMe_2$) and 135 (100, $PhMe_2Si$) (Found: $M^+ - Bu'$, 515.4086. $C_{36}H_{68}OSi_2$ requires M - Bu', 515.4104).

(7*R*,8*S*,10*S*)-7-(*tert*-Butyldimethylsilyloxymethyl)-8-dimethyl-(phenyl)silylhenicosan-10-ol 16

Hydroboration (9-BBN, 10 equiv., 24 h, reflux)² and oxidation, of the allylsilane Z-15 (250 mg) gave the alcohol (177 mg, 68%); $[a]_{\rm D}^{20}$ -4.3; $R_{\rm f}$ (hexane-EtOAc, 5:1) 0.59; $v_{\rm max}$ (film)/cm⁻¹ 3450 (OH) and 1250 (SiMe); $\delta_{\rm H}({\rm CDCl_3})$ 7.52–7.47 (2 H, m, o-SiPh), 7.35-7.30 (3 H, m, m- and p-SiPh), 3.58 (2 H, dd overlying br m, J 10.4 and 6.1, CH_AH_BOSi and CHOH), 3.40 (1 H, t, J 10.4, CH_ACH_BOSi), 3.70 (2 H, m, CH₂), 1.30–1.10 (32 H, m, CH₂ and SiCHCH), 0.88 (15 H, m, 2 × Me and Bu^t), 0.31 (3 H, s, $SiMe_AMe_BPh$) and 0.29 (3 H, s, $SiMe_AMe_BPh$), 0.01 (3 H, s, $SiMe_AMe_BBu'$) and -0.01 (3 H, s, $SiMe_AMe_BBu'$); $\delta_C(CDCl_3)$ 139.2, 133.9, 128.8, 127.7, 70.9, 66.2, 40.8, 37.9, 34.4, 31.9, 31.8, 30-29 (m), 26.1, 25.9, 22.7, 22.6, 20.2, 18.2, 14.12, 14.07, -2.7, -3.1, -5.3 and -5.4; m/z 575 (0.2%, M - Me), 533 (1, M - Bu'), 209 (30, PhMe₂SiOSiMe₂) and 135 (100, PhMe₂Si) (Found: $M^+ - Me$, 575.4712. $C_{36}H_{70}O_2Si_2$ requires M - Me, 575.4680). The diastereoisomer 17 was not detectable (¹H NMR).

(7R,8S,10R)-7-(*tert*-Butyldimethylsilyloxymethyl)-8-dimethyl-(phenyl)silylhenicosan-10-ol 17

This was prepared in the same way as the alcohol 16, from the allylsilane E-15 (127 mg), to give a mixture of the alcohols 16 and 17 (1:4, 70 mg, 59%); [a]_D -15.8 (c 1.00 in CHCl₃); $R_{\rm f}$ (hexane-EtOAc, 5:1) 0.59; $v_{\rm max}$ (film)/cm⁻¹ 3400 (OH) and 1250 (SiMe); $\delta_{\rm H}$ (CDCl₃) 7.51–7.47 (2 H, m, *o*-SiPh), 7.34–7.30 (3 H, m, m- and p-SiPh), 3.57 (1 H, dd, J 10.4 and 5.1, CH_A-CH_BO), 3.44 (1 H, t, *J* 10.4, CH_ACH_BO), 3.20 (1 H, m, CHOH), 2.94 (1 H, s, OH), 1.70-1.15 (31 H, m, SiCH and CH₂), 0.88 (15 H, m, CH₂ and Bu'), 0.32 (6 H, s, SiMe₂Ph), 0.02 (3 H, s, $SiMe_AMe_BBu'$) and 0.00 (3 H, s, $SiMe_AMe_BBu'$); $\delta_C(CDCl_3)$ 139.0, 133.8, 128.9, 127.7, 72.8, 64.4, 41.0, 37.6, 32.2, 31.9, 31.7, 30-29.4 (m), 28.3, 25.9, 25.7, 22.8, 22.7, 22.6, 18.2, 14.1, 14.0, -3.0, -3.2 and -5.3; m/z 533 (0.4%, M - Bu') and 135 (100, PhMe₂Si) (Found: $M^+ - Bu^t$, 533.4247. $C_{36}H_{70}O_2Si_2$ requires M - Bu', 533.4210). The isomer ratio was determined from the ¹³C NMR spectrum.

(7*R*,8*S*,10*S*)-10-Benzyloxy-7-(*tert*-butyldimethylsilyloxymethyl)-8-dimethyl(phenyl)silylhenicosane 18

The alcohol **16** (1.9 g) was stirred in cyclohexane (30 cm³) and dichloromethane (30 cm³) with benzyl trichlorobenzylacetimidate⁹ (1.5 cm³) and trifluoromethansulfonic acid (0.15 cm³) at room temperature for 30 min. Standard aqueous work-up and chromatography (SiO₂, hexane–EtOAc, 10:1) gave the *benzyl ether* (1.99 g, 91%); $R_{\rm f}$ (hexane–EtOAc, 5:1) 0.57; $v_{\rm max}$ (film)/cm⁻¹ 1250 (SiMe); $\delta_{\rm H}$ (CDCl₃) 7.51–7.47 (2 H, m, *o*-SiPh), 7.35–7.26 (8 H, m, CH₂Ph, *m*- and *p*-SiPh), 4.34 (2 H, s, CH₂Ph), 3.34 (2 H, m, CH₂OTBDMS), 3.22 (1 H, m, CHOBn), 1.60–1.13 (34 H, m, CH₂ and SiCHCH), 0.87 (15 H, m, Bu' and 2 × Me), 0.31 (3 H, s, SiMe_AMe_BPh), 0.29 (3 H, s, SiMe_AMe_BPh) and -0.01 (6 H, s, SiMe₂Bu'); *mlz* 665 (0.2%, M – Me), 623 (0.5, M – Bu'), 209 (25, PhMe₂SiOSiMe₂), 135 (100, PhMe₂Si) and 91 (60, PhCH₂) (Found: M⁺ – Me, 665.5203. C₄₂H₇₃O₂Si₂ requires M – Me, 665.5149).

(2R,3S,5S)-5-Benzyloxy-3-dimethyl(phenyl)silyl-2-hexylhexadecan-1-ol

Tetrabutylammonium fluoride (1 mol dm⁻³ in THF, 30 cm³) was added to the silyl ether **18** (1.99 g) and the mixture stirred at room temperature for 1 h. Standard aqueous work-up and chromatography (SiO₂, hexane–EtOAc, 10:1) gave the *alcohol* (1.62 g, 98%); $[a]_D^{20} + 2.3$ (*c* 1.8 in CHCl₃); R_r (hexane–EtOAc, 5:1) 0.40; v_{max} (film)/cm⁻¹ 3400 (OH), 1250 (SiMe) and 1110 (SiPh); δ_H (CDCl₃) 7.53–7.38 (2 H, m, *o*-SiPh), 7.36–7.26 (8 H,

m, CH₂Ph, m- and p-SiPh), 4.34 (2 H, s, CH₂Bn), 3.52–3.43 (3 H, m, CH₂OH and CHOBn), 1.60–1.17 (34 H, m, CH₂ and SiCHCH), 0.87 (3 H, t, J 6.5, Me_A), 0.85 (3 H, t, J 6.7, Me_B), 0.33 (3 H, s, SiMe_AMe_B) and 0.32 (3 H, s, SiMe_AMe_B); m/z 551 (0.003%, M – Me), 165 (12, PhCH₂OSiMe₂), 135 (100, PhMe₂-Si) and 91 (100, CH₂Ph) (Found: M⁺ – Me, 551.4280. C₃₇H₆₂-O₂Si requires M – Me, 551.4284).

(2R,3S,5S)-5-Benzyloxy-3-dimethyl(phenyl)silyl-2-hexylhexadecanoic acid 19

The alcohol (1.62 g, 2.86 mmol) and PDC²⁶ (3.3 g, 8.60 mmol) were stirred in DMF (25 cm³) at room temperature for 2 h. Chromatography of the solution (SiO₂, hexane-EtOAc, 5:1) gave the aldehyde (1.41 g, 88%). Jones' reagent²⁷ (2.5 cm³, 6.7 mmol) was added to the aldehyde in acetone (50 cm³). The mixture was stirred at 0 °C for 45 min. The solvent was removed under reduced pressure and chromatography (SiO₂, hexane-EtOAc, 5:1) gave the *acid* **19** (0.96 g, 62%); $[a]_{D}^{20}$ +19.2 (*c* 2.7 in CHCl₃); R_{f} (hexane-EtOAc, 5:1) 0.40; v_{max} (film)/cm⁻¹ 3500-2300 (OH), 1700 (CO), 1250 (SiMe) and 1110 (SiPh); δ_H(CDCl₃) 7.52–7.49 (2 H, m, *o*-SiPh), 7.35–7.26 (8 H, m, C₂Ph, m- and p-SiPh), 4.44 (1 H, d, J 11.5, CH_AH_BPh), 4.32 (1 H, d, J 11.5, CH_ACH_BPh), 3.40 (1 H, m, CHOBn), 2.52 (1 H, m, CHCO₂H), 1.70–1.10 (33 H, m, CH₂ and SiCH), 0.87 (6 H, m, $2 \times \text{Me}$), 0.35 (3 H, s, Si Me_AMe_B) and 0.32 (3 H, s, Si Me_AMe_B); m/z 580 (0.1%, M⁺), 565 (0.7, M – Me), 503 (4, M – Ph), 472 $(5, M - BnOH), 135 (60, PhMe_2Si) and 91 (100 CH_2Ph)$ (Found: M⁺, 580.4296. C₃₇H₆₀O₃Si requires *M*, 580.4314).

(2S,3S,5S)-5-Benzyloxy-3-hydroxy-2-hexylhexadecanoic acid 20 The silvl acid 19 (43 mg) and mercuric acetate (30 mg) were stirred in peracetic acid (15% in AcOH, 1 cm³) at room temperature for 3 h. Chromatography (SiO₂, Et₂O), gave the hydroxy acid (13 mg, 38%); R_f(hexane-EtOAc, 1:1) 0.20; $v_{\text{max}}(\text{film})/\text{cm}^{-1}$ 3400–2400 (OH); $\delta_{\text{H}}(\text{CDCl}_3)$ 7.36–7.29 (5 H, m, Ph), 4.67 (1 H, d, J 11.2, CH_ACH_BOBn), 4.40 (1 H, d, J 11.2, CH_ACH_BOBn), 3.94 (1 H, dt, J 9.8 and 3.0, CHOH), 3.75 (1 H, ddd, J 13.7, 7.0 and 3.2, CHOBn), 2.35 (1 H, ddd, J 9.2, 6.0 and 3.0, CHCO₂H), 1.75 (33 H, CH₂ and OH) and 0.87 (6 H, m, $2 \times \text{Me}$; $\delta_{c}(\text{CDCl}_{3})$ 176.3, 137.4, 128.7, 128.0, 127.9, 80.3, 72.3, 70.6, 51.7, 38.5, 33.0, 31.9, 31.6, 29.8, 29.6, 29.5, 29.3, 29.1, 27.2, 24.3, 22.7, 22.5, 14.1 and 14.0; m/z 460 (0.5%, $M - 2 \times H$), 338 (1, $M - C_7 H_8 O_2$), 123 (1, $C_7 H_7 O_2$), 105 (40, PhCO) and 91 (100, PhCH₂) (Found: $M^+ - 2 \times H$, 460.3527. $C_{29}H_{50}O_4$ requires $M - 2 \times H$, 460.3553).

(3S,4S)-4-[(S)-2'-Benzyloxytridecyl]-3-hexyloxetan-2-one 21

The silyl acid 19 (0.8 g) and mercuric acetate (1 g) were stirred in peracetic acid (30 cm³) at room temperature for 18 h. The solvent was evaporated under reduced pressure. Standard aqueous work-up and chromatography (SiO2, Et2O-Me2CO, 1:1) gave crude hydroxy acid, which was dissolved in pyridine (50 cm³) at 0 °C. Benzenesulfonyl chloride (6 cm³) was added and the mixture stirred at 0 °C for 15 h. Standard aqueous work-up and chromatography (SiO₂, hexane-EtOAc, 3:1) gave the *lactone* (0.46 g, 76%); $R_{\rm f}$ (hexane-EtOAc, 3:1) 0.6; $v_{\rm max}$ - (CH_2Cl_2) 1825 (lactone CO); $\delta_H(CDCl_3)$ 7.35–7.25 (5 H, m, Ph), 4.54 (1 H, d, J 11.5, CH_ACH_BPh), 4.41 (1 H, d, J 11.5, CH_A-CH_BPh), 4.40 (1 H, m, CHOCO), 3.50 (1 H, m, C₁₁CHO), 3.24 (1 H, dt, J 4 and 7.5, CHCO₂), 2.15 (1 H, dt, J 14.4 and 6.5, OCHCH_AH_BCHO), 1.90 (1 H, m, OCHCH_AH_BCHO), 1.50-1.20 (30 H, m, $15 \times CH_2$) and 0.86 (6 H, m, $2 \times Me$); m/z 338 $(0.1\%, M - PhCHO), 275 (0.8, M - C_{10}H_{17}O_2), 107 (20,$ PhCH₂O) and 91 (100, PhCH₂) (Found: M⁺ – PhCHO, 338.3180. $C_{29}H_{48}O_3$ requires M - PhCHO, 338.3184).

(3S,4S)-3-Hexyl-4-[(S)-2'-hydroxytridecyl]oxetan-2-one 22

The benzyl ether **21** (0.46 g) and palladium (10% on charcoal, 1 g) were stirred in THF (20 cm³) under hydrogen for 15 h. Filtration and chromatography (SiO₂, hexane–EtOAc, 3:1)

Published on 01 January 1998. Downloaded on 05/11/2013 10:12:24

gave the alcohol¹² (0.34 g, 92%) as needles mp 64–65 °C (lit.¹² mp 64.5–65.5 °C); $[a]_D^{20}$ –15.3 (*c* 1.2 in CH₂Cl₂) (lit.¹² $[a]_D$ –15); R_f (hexane–EtOAc, 3:1) 0.40; v_{max} (film)/cm⁻¹ 3600 (OH), 1820 (lactone C=O); δ_H (CDCl₃) 4.46 (1 H, dt, *J* 4.1 and 6.5, CHOCO), 3.76 (1 H, m, CHOH), 3.30 (1 H, ddd, *J* 8.5, 6.8 and 4.0, CHCO), 2.00–1.15 (33 H, m) and 0.87 (6 H, m, 2 × Me); δ_C (CDCl₃) 171.4, 76.2, 69.3, 56.7, 41.1, 37.6, 31.9, 31.5, 29.6, 29.54, 29.5, 29.3, 28.9, 27.8, 26.7, 25.4, 22.6, 22.5, 14.1 and 14.0.

(S)-1'-[(2"S,3"S)-3"-Hexyl-4"-oxooxetan-2"-yl]methyldodecyl (S)-N-formylleucinate (tetrahydrolipstatin) 1

DCC (167 mg) and (S)-N-(benzyloxycarbonyl)leucine 23 (418 mg) were stirred in dichloromethane (6 cm³) at 0 °C for 15 min. The solvent was evaporated under reduced pressure and the residue dissolved in DMF (5 cm³) and added to a solution of the alcohol (56 mg) and DMAP (15 mg) in DMF (1 cm³). The mixture was stirred at room temperature for 1 h. Standard aqueous work-up and chromatography (SiO₂, hexane-EtOAc, 3:1) gave N-CBz-protected ester, $R_{\rm f}$ (hexane-EtOAc, 3:1) 0.50, which was stirred with palladium (10% on charcoal, 100 mg) in THF (10 cm³) under hydrogen at room temperature for 4 h. The solution was filtered over Celite and evaporated under reduced pressure. The crude residue was stirred with formic acetic anhydride (0.2 cm³) in dichloromethane (0.5 cm³) at room temperature for 15 min. Standard aqueous work-up and chromatography (SiO₂, hexane-EtOAc, 3:1) gave tetrahydrolipstatin (69 mg, 89%) as an amorphous solid mp 40–41 °C (lit. mp 43 °C, ¹ 41–42.5 °C, ¹³ 40–42 °C, ^{14,15} 40–41 °C, ^{16,17,18} 42–43 °C ¹⁹); $[a]_{D}^{25} - 33.4 (c \ 1.33 \text{ in CHCl}_3) [lit. <math>[a]_{D}^{20} - 32 (c \ 1 \text{ in CHCl}_3), {}^1[a]_{D}^{20} - 34.45 (c \ 1 \text{ in CHCl}_3), {}^{13}[a]_{D}^{20} - 33 (c \ 0.36 \text{ in CHCl}_3), {}^{14.15}[a]_{D}^{20} - 31.2 (c \ 0.5 \text{ in CHCl}_3), {}^{16}[a]_{D} - 33 (CHCl}_3), {}^{17}[a]_{D}^{20} - 31.8 (c \ 0.37 \text{ in CHCl}_3), {}^{18}[a]_{D}^{25} - 34.58 (c \ 0.96 \text{ in CHCl}_3), {}^{19}]; v_{max}(CHCl}_3) 1825$ (lactone CO), 1740 (NHCO₂) and 1695 (NHCHO); $\delta_{\rm H}$ (CDCl₃) 8.20 and 8.05 (1 H, s and d, J 7, NHCHO), 5.93 (1 H, d, J 9, NH), 5.02 (1 H, m, CHOleucine), 4.67 (1 H, dt, J 4.5 and 8.5, CHNH), 4.28 (1 H, 5-line m, lactone CHOCO), 3.20 (1 H, dt, J 4.0 and 7.5, lactone CHCO), 2.15 (1 H, dt, J 14.7 and 7.8, OCHCH_AH_BCHO), 1.98 (1 H, dt, J 14.7 and 4.5, OCHCH_A- $H_{\rm B}$ CHO), 1.90–1.20 (33 H, m, 15 × CH₂ and Me₂CHCH₂), 0.95 (6 H, d, J 5.1, CHMe₂) and 0.87 (6 H, m, $2 \times Me$); $\delta_{C}(CDCl_{3})$ 171.9, 170.8, 160.9, 74.7, 72.6, 56.9, 49.6, 41.3, 38.6, 34.0, 31.8, 31.4, 29.6 (2C), 29.5, 29.4, 29.3, 28.9, 27.6, 26.6, 25.0, 24.8, 22.8, 22.5, 21.7, 14.1 and 14.0.

Acknowledgements

We thank the SERC (now EPSRC) for a studentship (N. J. L.), and Dr P. Barbier of Hoffmann-La Roche for a gift of tetrahydrolipstatin.

References

 E. Hochuli, E. Kupfer, R. Maurer, W. Meister, Y. Mercadel and K. Schmidt, J. Antibiotics, 1987, 40, 1086.

- 2 I. Fleming and N. J. Lawrence, J. Chem. Soc., Perkin Trans. 1, 1992, 3309.
- R. A. N. C. Crump, I. Fleming, J. H. M. Hill, D. Parker, N. L. Reddy and D. Waterson, *J. Chem. Soc., Perkin Trans.* 1, 1992, 3277.
 I. Fleming, R. Henning, D. C. Parker, H. E. Plaut and P. E. J.
- Sanderson, J. Chem. Soc., Perkin Trans. 1, 1995, 317.
- 5 I. Fleming and N. D. Kindon, J. Chem. Soc., Perkin Trans. 1, 1995, 303.
- 6 I. Fleming and N. J. Lawrence, *Tetrahedron Lett.*, 1990, **31**, 3645. 7 I. Fleming, T. W. Newton and F. Roessler, *J. Chem. Soc.*, *Perkin*
- Trans. 1, 1981, 2527. 8 A. W. P. Jarvie, A. Holt and J. Thompson, J. Chem. Soc. (B), 1969,
- 852; R. B. Miller and G. McGarvey, J. Org. Chem., 1979, 44, 4623.
 9 T. Iversen and D. R. Bundle, J. Chem. Soc., Chem Commun., 1981, 1240.
- 10 C. C. Price and J. R. Sowa, J. Org. Chem., 1967, 32, 4126; K. Tamao, T. Yamauchi and Y. Ito, Chem. Lett., 1987, 171; I. Fleming, S. K. Patel and C. J. Urch, J. Chem. Soc., Perkin Trans. 1, 1989, 115; P. F. Hudrlik, Y. M. Abdallah and A. M. Hudrlik, Tetrahedron Lett., 1992, 33, 6747; S. C. Archibald and I. Fleming, Tetrahedron Lett., 1993, 34, 2387; A. Barbero, D. C. Blakemore, I. Fleming and R. N. Wesley, J. Chem. Soc., Perkin Trans. 1, 1997, 1329.
- 11 W. Adam, J. Baeza and J.-C. Liu, J. Am. Chem. Soc., 1972, 94, 2000; S. Mageswaran and M. U. S. Sultanbawa, J. Chem. Soc., Perkin Trans. 1, 1976, 884; J. Mulzer, A. Pointer, A. Chucholowski and G. Brüntrup, J. Chem. Soc., Chem Commun., 1979, 52.
- 12 S. Kondo, K. Uotani, M. Miyasmoto, T. Hazato, H. Naganawa, T. Aoyagi and H. Umezawa, J. Antibiotics, 1978, 31, 797.
- 13 P. Barbier and F. Schneider, Helv. Chim. Acta, 1987, 70, 196.
- 14 P. Barbier, F. Schneider and U. Widmer, *Helv. Chim. Acta*, 1987, **70**, 1412.
- 15 P. Barbier and F. Schneider, J. Org. Chem., 1988, 53, 1218.
- 16 A. Pommier, J.-M. Pons, P. Kocienski and L. Wong, *Synthesis*, 1994, 1294.
- S. Hanessian, A. Tehim and P. Chen, J. Org. Chem., 1993, 58, 7768;
 B. Giese and M. Roth, J. Braz. Chem. Soc., 1996, 7, 243.
- 18 S. C. Case-Green, S. G. Davies and C. J. R. Hedgecock, *Synlett*, 1991, 781.
- 19 N. K. Chadha, A. D. Batcho, P. C. Tang, L. F. Courtney, C. M. Cook, P. M. Wovkulich and M. R. Uskokovic, *J. Org. Chem.*, 1991, 56, 4714.
- 20 I. Fleming, K. Takaki and A. P. Thomas, J. Chem. Soc., Perkin Trans. 1, 1987, 2269.
- 21 K. Tomioka, T. Suenaga and K. Koga, *Tetrahedron Lett.*, 1986, 27, 369, and ref. 5.
- 22 J. Klein and E. Gurfinkel, Tetrahedron, 1970, 26, 2127.
- 23 M. V. George, D. J. Peterson and H. Gilman, J. Am. Chem. Soc., 1960, 82, 403; H. Gilman, R. A. Klein and H. J. S. Winkler, J. Org. Chem., 1961, 26, 2474.
- 24 H. O. House, C.-Y. Chu, J. M. Wilkins and M. J. Umen, J. Org. Chem., 1975, 40, 1460.
- 25 C. L. Stevens and S. J. Dykstra, J. Am. Chem. Soc., 1954, 76, 4402.
- 26 E. J. Corey and G. Schmidt, *Tetrahedron Lett.*, 1979, 399.
- 27 C. Djerassi, R. R. Engle and A. Bowers, J. Org. Chem., 1956, 21, 1547.

Paper 8/04275F Received 5th June 1998 Accepted 3rd July 1998