

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

SYNTHETIC COMMUNICATIONS<sup>®</sup> Vol. 33, No. 19, pp. 3317–3325, 2003

# Coupling of 3,8-Dibromo-1,10-phenanthroline with 3,5-Diethynylheptyloxybenzene: A Suzuki/Miyaura Versus a Sonogashira Perspective

Jinhua Yang,<sup>1</sup> Woon Su Oh,<sup>1</sup> Ian A. Elder,<sup>1</sup> Nicholas Leventis,<sup>2</sup> and Chariklia Sotiriou-Leventis<sup>1,\*</sup>

> <sup>1</sup>Department of Chemistry, University of Missouri-Rolla, Rolla, Missouri, USA <sup>2</sup>NASA Glenn Research Center, Materials Division, Cleveland, Ohio, USA

# ABSTRACT

We report a new application of the Suzuki-Miyaura reaction whereas two bifunctional reactants, 3,8-dibromo-1,10-phenanthroline and 3,5diethynylheptyloxylbenzene (9), yield 3,8-*bis*(3-ethynyl-5-heptyloxyphenylethynyl)-1,10-phenanthroline (2) efficiently (74% yield) without polymerization. This was achieved by reacting a stoichiometric amount of 9 and (Me<sub>3</sub>Si)<sub>2</sub>NLi to obtain quantitatively the

3317

DOI: 10.1081/SCC-120023989 Copyright © 2003 by Marcel Dekker, Inc. 0039-7911 (Print); 1532-2432 (Online) www.dekker.com

<sup>\*</sup>Correspondence: Chariklia Sotiriou-Leventis, Department of Chemistry, University of Missouri-Rolla, 21000 Brookpark Road, Rolla, MO 65409, USA; Fax: (573)341-6033; E-mail: cslevent@umr.edu.

XY

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

Yang et al.

monoacetylide anion of 9 (10). The latter was activated with B-methoxy-9-BBN and reacted in analogy to the alkynyl copper complex of a Sonogashira route. However, in the Sonogashira reaction, the alkynyl copper complex is present in small equilibrium concentrations and polymerization takes place even when reagents are mixed slowly. Actually the Sonogashira route gave no desired product 2, as the latter polymerizes easily via homo-coupling in the presence of air and Cu(I).

*Key Words:* Suzuki; Miyaura; Sonogashira; Coupling; 3,8-Dibromo-1,10-phenanthroline.

Sonogashira coupling involves the palladium(0) catalyzed reaction of terminal alkynes with vinyl or aryl halides in the presence of CuI or CuBr and an amine such as triethyl or diisopropyl amine.<sup>[1]</sup> Mechanistically, oxidative addition of the organic halide gives a Pd(II) intermediate, which undergoes transmetallation with the alkynyl copper (generated in small equilibrium quantities from the terminal alkyne, base, and copper iodide). Reductive elimination with coupling of the two organic ligands gives the product and regenerates the Pd(0) catalyst. For convenience, oftentimes the palladium catalyst is introduced as a stable, soluble Pd(II) derivative such as Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>. This complex is rapidly reduced by the terminal alkyne in situ to give the catalytic Pd(0) species. Due to the mild conditions usually employed (e.g., reaction at room temperature) and the use of inexpensive reagents, the Sonogashira coupling reaction has been a very convenient synthetic method for the preparation of aromatic alkynes.

Despite its convenience, the Sonogashira reaction cannot be used for coupling of dialkynes with dibromoaromatic compounds. First, bifunctionality on both reactants generally leads to polymerization. Furthermore, even though it might be considered that by using slow syringe-pump addition of the dialkyne might lead to the desirable product, unfortunately during work-up, the bisacetylenic product itself tends to undergo oxidative coupling in the presence of air and Cu(I), known as homocoupling,<sup>[2]</sup> leading again to polymerization.

In brief, the main causes of polymerization during the Sonogashira coupling reaction are: (a) the generation of only a small equilibrium amount of the alkynyl anion, and (b) the presence of catalytic amounts of copper (related to homocoupling). It seems that if the alkynyl anion could be generated quantitatively and activated by means other than copper, polymerization would be avoided.

An alternative procedure that also leads to aryl alkynes is the Suzuki/Miyaura coupling reaction.<sup>[3]</sup> According to that procedure,

# 3318

- 474

MARCEL DEKKER, INC. • 270 MADISON AVENUE • NEW YORK, NY 10016

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

# Suzuki/Miyaura Vs. Sonogashira Perspective

# 3319

the alkynyl anion is formed quantitatively by the reaction of a terminal alkyne with a stoichiometric amount of a strong base (e.g., (Me<sub>3</sub>Si)<sub>2</sub>NLi). Subsequently, the alkynyl anion is activated via reaction with B-methoxy-9-borabicyclo[3.3.1]nonane (B-methoxy-9-BBN) to form a stable methoxy (alkynyl)borate complex, which subsequently reacts with an aryl bromide in the presence of Pd(0) to yield identical products with the corresponding Sonogashira coupling reaction that involves the same alkyne and aryl bromide starting materials. The Suzuki/Miyaura route uses harsher conditions (strong base) and a more expensive reagent (B-methoxy-9-BBN). However, it involves quantitative formation of the alkynyl anion and it does not use copper salts, so polymerization with bifunctional terminal halides and dibromoaromatic compounds should not be an issue.

These inferences were confirmed by comparing the synthesis of 3,8bis(4-isonicotynoylphenylethynyl)-1,10-phenanthroline (1) with the synthesis of 3,8-bis(3-ethynyl-5-heptyloxyphenylethynyl)-1,10-phenanthroline (2). Compound 1 was designed as a ligand for intramolecular electron transfer quenching, while metal complexes of compound 2 were designed as building blocks of larger conjugated systems.



Sonogashira coupling of 4-(p-bromobenzoyl)pyridine (**3**) with 3,8bisethynyl-1,10-phenanthroline (**4**, prepared also via Sonogashira coupling of TMS-protected acetylene and 3,8-dibromo-1,10-phenanthroline) proceeds uneventfully giving **1** in 32% yield (Sch. 1).<sup>[4]</sup> On the other hand, direct Sonogashira coupling of 3,8-dibromo-1,10-phenanthroline with 3,5-diethynylheptyloxybenzene (**9**) under the same conditions led to an intractable polymer.

Compound 9 was synthesized according to Sch. 2. Pentabromophenol (5) was dehalogenated in benzene by  $AlCl_3$  to give 3,5-dibromophenol (6),<sup>[6]</sup> which then reacted with 1-bromoheptane to yield 3,5-dibromoheptyloxylbenzene (7). A direct Sonogashira coupling of 7 with



Scheme 1.

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.





trimethylsilylacetylene catalyzed by Pd(PhCN)<sub>2</sub>Cl<sub>2</sub> with P(*t*-Bu)<sub>3</sub> in the presence of CuI and diisopropylamine at room temperature gave compound **8**, which was subsequently deprotected in the presence of NaOH in CH<sub>2</sub>Cl<sub>2</sub>/MeOH to give compound **9** in 65% overall yield from **5**.

Of course, protection of one of the acetylenic groups of 9 could be a viable route for the synthesis of 2 via the Sonogashira reaction, however protection would introduce two more steps and would most certainly compromise the overall yield. Interestingly, both the first step of a possible protection route for the synthesis of 2, and the first step of the Suzuki/Miyaura route (Sch. 3) involve quantitative formation of the monoacetylide 10 by using a stoichiometrric amount of a strong base.

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

# Suzuki/Miyaura Vs. Sonogashira Perspective

### 3321

In the Suzuki/Miyaura route, **10** is activated with B-methoxy-9-BBN to form a stable methoxy(alkynyl)borate complex **11**, which is then added to a solution of 3,8-dibromo-1,10-phenanthroline and a catalytic amount of  $Pd(PPh_3)_4$  in THF. Intermediate **11** reacts in analogy to the alkynyl copper complex of the Sonogashira route yielding **2** in very good yield (74%).

In conclusion, we have shown that coupling via the Suzuki/Miyaura route of two bifunctional reagents, namely 3,8-dibromophenanthroline and 3,5-diethynylheptyloxybenzene (9) gives compound 2 efficiently without polymerization.

# EXPERIMENTAL

**General.** <sup>1</sup>H and <sup>13</sup>CNMR spectra were recorded on a Varian INOVA 400 NMR spectrometer and are reported as parts per million (ppm) from TMS ( $\delta$ ). Melting points were uncorrected. The Electrospray Mass Spectometry (ESI) was conducted using a TSQ7000 (ThermoFinnigan) with an API2 source and the "performance pack" (wider aperture on the skimmer and extra turbo pump on the source). The electrospray source is equipped with a stainless steel needle; the voltage was 4.5 kV. Sample solutions were injected using a 500 µL Gastight syringe (Hamilton) in a Pump 11 syringe pump (Harvard Apparatus).

**Materials.** All starting materials, reagents and solvents were from Aldrich or Acros and used as received, unless noted otherwise.  $P(t-Bu)_3$  was purchased from Strem Chemicals. For coupling reactions, all the solvents were anhydrous and were obtained directly from Aldrich or Acros.

**3,8-bis(4-Isonicotinoylphenylethynyl)-1,10-phenanthroline (1).** 4-bromobenzoylpyridine<sup>[7]</sup> (**3**, 0.68 g, 2.6 mmol), 3,8-diethynyl-1,10-phenanthroline<sup>[8]</sup> (**4**, 0.2 g, 0.88 mmol), Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> (60 mg, 0.085 mmol), and CuI (64 mg, 0.34 mmol) were added to anhydrous diisopropylamine (8 mL) and anhydrous DMF (40 mL) under nitrogen and the reaction mixture was refluxed for 24 h. After evaporation of the solvent under vacuum, the residue was diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed with 5% aqueous KCN solution (3 × 20 mL), water (5 × 100 mL), and saturated aq. NaCl (100 mL). The organic phase was dried over anhydrous sodium sulfate, concentrated in vacuo followed by the addition of diethyl ether. The solid thus obtained was recrystallized from CH<sub>2</sub>Cl<sub>2</sub>/THF to give the product (0.17 g, 32%). M.p. 266°C (dec.). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 9.32 (d, J = 1.83 Hz, 2H), 8.85 (d, J = 5.68 Hz, 4H), 8.45 (d, J = 1.83 Hz, 2H), 7.89



NI<del>I</del>

MARCEL DEKKER, INC. • 270 MADISON AVENUE • NEW YORK, NY 10016

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

#### 3322

#### Yang et al.

(s, 2H), 7.86 (d, J = 5.49 Hz, 4H), 7.76 (d, J = 8.06 Hz, 4H), 7.60 (d, J = 5.68 Hz, 4H). MS (ESI+) 591.1816 (M + H)<sup>+</sup>, (theoretical ionic mass: 590.1743).

**3.5-Dibromophenol** (6).<sup>[6]</sup> Aluminum chloride (2.67 g, 0.02 mol) was added to a solution of pentabromophenol (5, 5.0 g, 0.01 mol) in anhydrous benzene (20 mL) and the reaction mixture was heated to reflux for 3h under nitrogen. After cooling to room temperature, the reaction mixture was poured into a mixture of HCl and ice water. The organic layer was separated and the water layer was extracted with diethyl ether  $(3 \times 20 \text{ mL})$ . The combined organic layers were extracted with 10% aqueous NaOH ( $3 \times 20 \text{ mL}$ ). The combined aqueous layers were made strongly acidic with HCl. The product was then extracted with methylene chloride  $(3 \times 20 \text{ mL})$  and the combined organic layers were washed with water until the aq. layer was neutral and then dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent in vacuo, gave the crude product which was recrystallized from hexane to give off-white needles (2.3 g, yield: 88%). M.p. 81-82°C (Lit.<sup>[6a]</sup> m.p. 80°C, Lit.<sup>[6b]</sup> m.p. 76–79°C). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 4.85 (s, 1H), 6.95 (d, J = 1.65 Hz, 2H, 7.25 (t, J = 1.65 Hz, 1H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) 156.4, 126.8, 123.1, 117.8.

**3,5-Dibromoheptyloxylbenzene (7).** A mixture of 3,5-dibromophenol (6, 5.0 g, 0.02 mol), potassium carbonate (2.76 g, 0.02 mol), and 1-bromoheptane (3.1 mL, 0.02 mol) in acetone (50 mL) was stirred at reflux overnight. After cooling to room temperature, water (50 mL) was added and the reaction mixture was extracted with diethyl ether ( $3 \times 20$  mL). The combined organic layers were washed first with aqueous 10% NaOH, followed by water, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness under vacuum. The oil thus obtained was distilled under vacuum to give a colorless liquid (5.4 g, 84%). B.p. 145°C/0.5 torr. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 0.89 (t, J = 6.96 Hz, 3H), 1.29–1.46 (m, 8H), 1.72–1.79 (m, 2H), 3.90 (t, J = 6.50 Hz, 2H), 6.97 (d, J = 1.65 Hz, 2H), 7.22 (t, J = 1.65 Hz, 1H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) 160.6, 126.3, 123.3, 117.1, 68.8, 32.1, 29.3, 26.2, 22.9, 14.4.

**3,5-bis**(Trimethylsilylethynyl)heptyloxylbenzene (8).  $Pd(PhCN)_2Cl_2$  (46 mg, 0.12 mmol) and CuI (15.2 mg, 0.08 mmol) were added to a dry, septum-capped round bottom flask, purged with nitrogen. Anhydrous dioxane (2 mL),  $P(t-Bu)_3$  (0.5 mL, 0.25 M solution in dioxane),  $HN(i-Pr)_2$  (0.68 mL, 4.8 mmol), 7 (0.7 g, 2 mmol) in 0.5 mL dioxane were then added via syringe to the stirred reaction mixture. Finally, trimethylsilylacetylene (0.66 mL, 4.6 mmol) was added to the flask. In a few minutes, a precipitate was formed. After stirring the reaction mixture at room temperature for 2 h, the solvent was evaporated to dryness and

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

# Suzuki/Miyaura Vs. Sonogashira Perspective

#### 3323

the crude product was purified on a silica gel column using 2% CH<sub>2</sub>Cl<sub>2</sub> in hexane as eluent. The product was further purified by recrystallization from methanol to give a yellow solid (0.68 g, yield 89%). M.p. 47°C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 7.16 (t, J = 1.47 Hz, 1H), 6.94 (d, J = 1.28 Hz, 2H), 3.91 (t, J = 6.50 Hz, 2H), 1.71–1.79 (m, 2H), 1.28–1.46 (m, 8H), 0.89 (t, J = 5.78 Hz, 3H), 0.22 (s, 18H).

**3,5-bis(Ethynyl)heptyloxylbenzene (9). 8** (3.35 g, 8.7 mmol), acetone (30 mL), methanol (60 mL), and NaOH (0.12 g, 3 mmol) were mixed together and stir at room temperature for 2 h. Water (100 mL) was then added, followed by extraction with diethyl ether ( $3 \times 50$  mL). The combined ether extracts were washed with water until the aqueous layer was neutral, dried with anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness in vacuo to give a yellow solid (2.05 g, 98%). M.p. 32°C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 7.20 (t, J = 1.28 Hz, 1H), 6.99 (d, J = 1.28 Hz, 2H), 3.93 (t, J = 6.59 Hz, 2H), 3.05 (s, 2H), 1.72–1.80 (m, 2H), 1.28–1.46 (m, 8H), 0.89 (t, J = 6.95 Hz, 3H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) 158.6, 128.0, 123.2, 118.8, 82.6, 77.5, 68.2, 31.7, 29.1, 29.0, 25.9, 22.6, 14.1. MS (ESI+) 241.1587 (M + H),<sup>+</sup> (theoretical ionic mass: 240.1514).

**3,8**-*bis*(**3**-Ethynyl-**5**-heptyloxyphenylethynyl)-**1,10**-phenanthroline (2). Lithium bis(trimethylsilyl)amide (2.25 mmol, 2.25 mL of 1 M solution in THF) was added under nitrogen to a well stirred solution of 9 (0.86 g, 2.25 mmol) in anhydrous THF (10 mL) at -78°C. After 30 min, B-methoxy-9-BBN (2.25 mmol, 2.25 mL of 1 M solution in hexane) was added to the reaction mixture via a septum with a syringe. After stirring for 2 h at  $-78^{\circ}$ C, the solution was transferred via a cannula to a second flask containing Pd(PPh<sub>3</sub>)<sub>4</sub> (60 mg, 0.052 mmol) and 3,8-dibromo-1,10-phenanthroline (0.25 g, 0.75 mmol) in dry THF (30 mL) under nitrogen. The reaction mixture was then heated to reflux for 24h and cooled to room temperature. Methylene chloride (50 mL) was added and the organic layer was washed with water  $(20 \times 100 \text{ mL})$  dried with anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness. The crude product thus obtained was purified on silica gel column. Initial elution with CH<sub>2</sub>Cl<sub>2</sub> gave unreacted starting material. Subsequent elution with diethyl ether gave the product. After removal of the solvent in vacuo a light yellow solid was obtained (0.36 g, 74%). M.p. 85–86°C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 9.25 (d, J = 2.01 Hz, 2H), 8.34 (d, J = 2.01 Hz, 2H), 7.78 (s, 2H), 7.33 (t, J = 1.28 Hz, 2H), 7.12 (t, J = 1.46 Hz, 2H), 7.05 (t, J = 1.28 Hz, 2H), 3.98 (t, J = 6.5 Hz, 4H), 3.10 (s, 2H), 1.78–1.82 (m, 4H), 1.48 (m, 8H), 1.28–1.46 (m, 8H), 0.91 (t, J = 6.87 Hz, 6H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) 158.6, 152.1, 144.2, 137.9, 127.8, 127.5, 126.7, 123.4, 123.3, 119.2, 118.9, 118.0, 92.9, 86.4, 82.5, 77.6, 68.2, 31.7, 29.0, 28.9, 25.8, 22.5, 14.0. MS (ESI+) 657.3476 (M + H),<sup>+</sup> (theoretical ionic mass: 656.3403).

MA.

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

3324

#### Yang et al.

# ACKNOWLEDGMENTS

We gratefully acknowledge support by the Petroleum Research Fund, administered by the ACS (Grant No. 35154-ACS), and from the National Cancer Institute (Grant No. 1 R15 CA82141-01A2 to C. S.-L.). We also thank Dr. Nathan D. Leigh at the University of Missouri-Columbia for obtaining the MS spectra.

# REFERENCES

- 1. (a) Sonogashira, K.; Tohda, Y.; Hagihara, N. Convenient synthesis of acetylenes. Catalytic substitutions of acetylenic hydrogen with bromo alkenes, iodo arenes, and bromopyridines. Tetrahedron Lett. 1975, 50, 4467–4470; (b) Cassar, L. Synthesis of aryl- and vinylsubstituted acetylene derivatives by the use of nickel and palladium complexes. J. Organomet. Chem. 1975, 93, 253-257; (c) Dieck, H.A.; Heck, R.F. Palladium catalyzed synthesis of aryl, heterocyclic, and vinylic acetylene derivatives. J. Organomet. Chem. 1975, 93, 259-263; (d) Hundertmark, T.; Littke, A.F.; Buchwald, S.L.; Fu, G.C. Pd(PhCN)<sub>2</sub>Cl<sub>2</sub>/P(t-Bu)<sub>3</sub>: A versatile catalyst for sonogashira reactions of aryl bromides at room temperature. Org. Lett. 2000, 2 (12), 1729–1731; (e) Thorand, S.; Krause, N. Improved procedures for the palladium-catalyzed coupling of terminal alkynes with aryl bromides (Sonogashira coupling). J. Org. Chem. 1998, 63, 8551-8553; (f) Nakamura, K.; Okubo, H.; Yamaguchi, M. Low-temperature Sonogashira coupling reaction. Synlett 1999, 5, 549–550; (g) Bohm, V.P.W.; Herrmann, W.A. Coordination chemistry and mechanisms of metal-catalyzed C-C coupling reactions, 13: a copper-free procedure for the palladium-catalyzed Sonogashira reaction of aryl bromides with terminal alkynes at room temperature. Eur. J. Org. Chem. 2000, 3679-3681; (h) Pearson, D.L.; Schuman, J.S.; Tour, J.M. Iterative divergent/convergent approach to conjugated oligomers by a doubling of molecular length at each iteration. A rapid route to potential molecular wires. Macromolecules 1994, 27, 2348-2350.
- (a) Rossi, R.; Carpita, A.; Bigelli, C. Palladium-promoted route to 3alkyl-4-(1-alkynyl)hexa-1,5-dyn-3-enes and/or 1,3-diynes. Tetrahedron Lett. 1985, 26 (4), 523–526; (b) Ikegashira, K.; Nishihara, Y.; Hirabayashi, K.; Mori, A.; Hiyama, T. Copper (I) salt promoted homo-coupling reaction of organosilanes. Chem. Commun. 1997, 1039–1040; (c) Nishihara, Y.; Ikegashira, K.; Hirabayashi, K.;

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

# Suzuki/Miyaura Vs. Sonogashira Perspective

# 3325

Ando, J.; Mori, A.; Hiyama, T. Coupling reactions of alkynylsilanes mediated by a Cu(I) salt: Novel syntheses of conjugate diynes and disubstituted ethynes. J. Org. Chem. **2000**, *65*, 1780–1787; (d) Hay, A.S. Oxidative coupling of acetylenes. J. Org. Chem. **1960**, *25*, 1275–1276; (e) Hay, A.S. Oxidative coupling of acetylenes II. J. Org. Chem. **1962**, *27*, 3320–3321.

- (a) Soderquist, J.A.; Matos, K.; Rane, A.; Ramos, J. Alkynylboranes in the Suzuki-Miyaura coupling. Tetrahedron Lett. **1995**, *36* (14), 2401–2402; (b) Soderquist, J.A.; Rane, A.M.; Matos, K.; Ramos, J. Alkynylborinates in organoborane conversions. Tetrahedron Lett. **1995**, *36* (38), 6847–6850; (c) Castanet, A.S.; Colobert, F.; Schlama, T. Suzuki-Miyaura coupling of alkynylboronic esters generated in situ from acetylenic derivatives. Org. Lett. **2000**, *2* (23), 3559–3561.
- 4. According to the literature, direct Sonogashira couplings on 3,8dibromo-1,10-phenanthroline are complicated by complexation of the Cu(I) catalyst with the phenanthroline's nitrogens.<sup>[5a]</sup> In that respect, Sonogashira coupling of Ru(II)-coordinated 3,8-dibromo-1,10-phenanthroline works exceptionally well.<sup>[5b]</sup>
- (a) Suffert, J.; Ziessel, R. Towards molecular electronics: a new family of aromatic polyimine chelates substituted with alkyne groups. Tetrahedron Lett. **1991**, *32*, 757–760; (b) Tzalis, D.; Tor, Y. Coordination compounds as building blocks: single-step synthesis of multi-ruthenium(II) complexes. Chem. Commun. **1996**, 1043– 1044.
- (a) Ishida, Y.; Jikei, M.; Kanimoto, M. Unusual thermal behavior of the aromatic polyamide dendrons. Polym. Adv. Technol. 2000, 11, 698–704; (b) Kraus, R.; Spiteller, G. Unusual fragmentation of trimethylsilylated enols derived from *m*- and *p*-hydroxyacetophenones. Organic Mass Spectrometry 1989, 24, 861–865.
- Leventis, N.; Elder, I.A.; Gao, X.; Bohannan, E.W.; Sotiriou-Leventis, C.; Rawashdeh, A.-M.M.; Overschmidt, T.J.; Gaston, K.R. The redox chemistry of 4-benzoyl-*N*-methylkpyridinium cations in acetonitrile with and without proton donors: The role of hydrogen bonding. J. Phys. Chem. B 2001, *105*, 3663–3674.
- Joshi, H.S.; Jamshidi, R.; Tor, Y. Conjugated 1,10-phenanthrolines as tunable fluorophores. Angew. Chem. Int. Ed. 1999, 38 (18), 2722–2725.

Received in the USA April 3, 2003



) TM MARCEL DEKKER, INC. • 270 MADISON AVENUE • NEW YORK, NY 10016

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

Copyright © 2003 EBSCO Publishing

Copyright of Synthetic Communications is the property of Taylor & Francis Ltd and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.