Journal of Molecular Liquids 337 (2021) 116435



Contents lists available at ScienceDirect

Journal of Molecular Liquids

journal homepage: www.elsevier.com/locate/molliq

Tuning the extraction mechanism of uranyl ion in bicyclooctanium, propylpyridinium, piperidinium and imidazolium based ionic liquids: First ever evidence of 'cation exchange', 'anion exchange' and 'solvation' mechanism



Amit Pandey^{a,1}, S. Hashmi^{b,1}, G. Salunkhe^b, Velavan Kathirvelu^a, Keisham S. Singh^c, Rohit Singh Chauhan^b, Arijit Sengupta^{d,e,*}

^a Department of Applied Sciences, National Institute of Technology Goa, Ponda, Goa 403401, India

^b Department of Chemistry, K.J.Somaiya College of Science and Commerce, Vidya-vihar, Mumbai 400077, India

^c Bioorganic Chemistry Laboratory, CSIR-National Institute of Oceanography, Dona Paula, Goa 403004, India

^d Radiochemistry Division, Bhabha Atomic Research Centre, Trombay, Mumbai 400085, India

^e Homi Bhabha National Institute, Anushaktinagar, Mumbai 400094, India

ARTICLE INFO

Article history: Received 4 April 2021 Revised 2 May 2021 Accepted 5 May 2021 Available online 8 May 2021

Keywords: Cation exchange Anion exchange Solvation Speciation Ionic liquid

ABSTRACT

A novel diamide ligand, 5-bromo-N1, N3-diisopropylisophthalamide has been synthesized and characterized by FTIR, ¹H NMR, and ¹³C NMR. The extraction mechanism of uranyl ion with the diamide ligand has been studied in unexplored ionic liquid families. For the same metal–ligand system; cation exchange, anion exchange, and solvation mechanism can be predominately achieved by changing the ionic liquid system. The experimental results suggest that the change in extraction mechanism is induced by the cations of the ionic liquids. In bicyclooctanium ionic liquid, the extracted species was $[UO_2(NO_3)_3L_2]^{-}$; In piperidinium and pyridinium-based ionic liquid, the species was $[UO_2(NO_3)_2L$ and $UO_2(NO_3)_2L_2$ respectively. In imidazolium-based ionic liquid, the species was $[UO_2(NO_3)_2L]^{+}$. Though the extraction efficiency in methylpyridinium and bicyclooctanium based ionic liquids are better, the pipiredinium and imidazolium-based ionic liquid a high degree of radiological stability. The aqueous solution of sodium carbonate is found to be better in the back extraction of uranyl ions from the ionic liquids relative to EDTA and oxalic acid. However, the efficacy of back extraction differs largely on the ionic liquid system.

© 2021 Elsevier B.V. All rights reserved.

1. Introduction

Though ionic liquid based research evolved mainly as a potential 'green' alternatives to the volatile organic diluents due to some of the advantageous properties like lower vapor pressure, high flash point, high degree of chemical, thermal and electrochemical stability; the high 'anti microbial' characteristic of some of the ionic liquids put doubts on the 'greenness' [1–6]. Some classes of ionic liquids are neither biodegradable nor completely incinerable. However, the most interesting property of ionic liquid is its tunability due to which it is widely accepted as a 'designer solvent'. It has been reported that for N3 alkyl-substituted methylimida-

¹ These authors contributed equally.

zolium cation, the extraction mechanism depends on the length of the alkyl group. If the butyl group at the N3 position is replaced with an octyl or decyl, the extraction mechanism changes from cation exchange to solvation [7–9]. This was attributed to the ease of solubilizing ability of N3 alkyl-substituted methyl imidazolium cation in water, which reduces as the alkyl chain length increases. This also indicates that the species involved can also be modified from cationic to neutral. It was also reported in the literature that the extraction kinetics are highly dependent on the length of the alkyl groups of N3 alkyl-substituted methylimidazolium ion and even the nature of the anionic moieties of the ionic liquid [10-14]. The longer the alkyl chain length, viscous the ionic liquids are, and hence slower is the mass transfer. The ionic liquid with PF_6^- anion is more viscous compared to NTf_2^- anion and hence resulting in slower extraction kinetics. The more branched ionic liquids are more prone to radiation-induced fragmentation [15,16]. Hence depending upon the requirement, the extraction

^{*} Corresponding author at: Radiochemistry Division, Bhabha Atomic Research Centre, Trombay, Mumbai 400085, India.

E-mail address: arijita@barc.gov.in (A. Sengupta).

process can be modified by modification in ionic liquids [17]. The extraction efficiency also depends on the nature of alkyl substitution in the imidazolium cations of ionic liquid. There are several attempts to design the functionalized ionic liquid for task-specific applications, where functionalities are covalently attached to the ionic liquid moieties. Carbomyl methyl phosphine oxide (CMPO), diglycolamide (DGA), trialkyl phosphine (TRPO) functionalized ionic liquids have been extensively used for the extraction of actinides [18,19]. The extraction behavior of neodymium and other rare earth were investigated by trioctylmethylammonium dioctyl diglycolamate functionalized ionic liquid extraction of metals by functionalized ionic liquid is another new approach reported in the literature [22].

Application of ionic liquid not only improves the efficiency/ selectivity of metal extraction: but also the speciation may differ [23,24]. Vast research on the application of ionic liquids in the separation of metal ions, mainly f-block elements is centered at imidazolium-based ionic liquids only. Recently, it has been demonstrated for the first time that a sulphoxide ligand in bicyclooctanium ionic liquid can be used for separation of f-block elements [25]. However, different families of ionic liquids (pyridinium, piperidinium, phosphonium, octanium, etc.) are still unexplored and they may show some unique features. One such example is the extraction of Sr²⁺ by Calix crown ether using pyridiniumbased ionic liquid. The extraction of Sr²⁺ predominantly follows the least common 'anion exchange' mechanism through a monoanionic complex [26]. In view of that, pyridinium, piperidinium, and bicyclooctanium families of ionic liquids are considered in the present case and are compared with the most commonly used methylimidazolium-based ionic liquid.

The success of nuclear establishment highly depends on the safe management of radiotoxic material; efficient and selective separation scheme for actinides, lanthanides, and other fission products [27–29]. The tri-n-butyl phosphate is the conventional workhorse for efficient separation schemes in PUREX and THOREX processes in nuclear establishment [30,31]. Phosphine oxides, phosphonates, Cyanex 923, phosphinic acids, carbamoyl methyl phosphine oxide are some of the ligands explored for uranyl extraction [18,32–35]. However, phosphate-based ligands are not completely incinerable, hence there is always a quest for new nitrogen-based ligand functionalities for the separation of uranyl ions. Secondary amides like di-n-hexyl octanamide (DHOA), diamide, glycolamides are some of the functionalities explored for efficient uranyl extraction [36–40].

In the present case, diisopropylisophthalamide in pyridinium, piperidinium, imidazolium, and bicyclooctanium based ionic liquids have been used for the extraction of UO_2^{2+} from the aqueous nitric acid feed. The investigation deals with the understanding of the extraction mechanism, species involved, etc. The radiation and chemical stability of organic phases, stripping of metal ions from organic phases have also been investigated.

2. Experimental

2.1. Materials and methods

5-Bromoisophthalic acid, isopropylamine, triethylamine and the ionic liquids (Fig. 1); 1-Hexyl-1,4-diaza[2,2,2]bicyclooctanium bis(trifluoromethylsulfonyl)imide (IL1), 1-Propylpyridinium bis(tri fluoromethylsulfonyl)imide (IL2), 1-Butyl-1methylpiperidinium bis(trifluoromethylsulfonyl)imide (IL3), 1-Hexyl-3methylimidazolium bis(trifluoromethylsulfonyl)imide (IL4) were procured from TCI Chemical Pvt Ltd., India. Dichloromethane and thionyl chloride were purchased from SRL Chemicals. The solvents used for the synthesis of ligand were freshly distilled before use. Nitric acid (Suprapure grade) was purchased from Sigma Aldrich. The reagents like NaNO₃, N₂CO₃, n-dodecane, and oxalic acid, were of analytical grade purchased from Sigma Aldrich. Deionized water was used throughout the experiment. The U stock solutions were prepared from U₃O₈ powder by dissolving in Conc. HNO₃ followed by repeated evaporation. The stock solutions were made in 1 M HNO₃.

NMR spectra were recorded on a Bruker Avance 400 MHz spectrometer at 400 MHz (¹H) and 100 MHz (¹³C) with SiMe₄ as internal references. The IR spectrum was recorded on IR Affinity-1, Fourier Transform Infrared Spectrophotometer (Shimadzu). CHNS analysis of the ligand was carried out in Elementar vario MICRO cube CHNS Analyzer. The powder sample (3 mg) wrapped in a small tin foil was directly dropped into the combustion tube where it was fully combusted in the presence of oxygen. The MICRO cube analyzed the CHN content (weight %) of the ligand in one single run. Energy Dispersive X-ray Fluorescence (EDXRF) Spectroscopy (Make: Xenemetrix, Model: EX3600SDD) was used for the estimation of U [41–43].

2.2. Synthesis and characterization of ligands

2.2.1. 5-bromo-N1, N3-diisopropylisophthalamide (L)

Thionyl chloride (40 mmol, 2.90 mL) and 5-bromoisophthalic acid (2 mmol, 0.490 g) were taken in a round bottom flask and refluxed under nitrogen atmosphere for 16 hr (see Scheme 1). A clear transparent solution was formed. Excess thionyl chloride was removed in a vacuum to give a colorless viscous liquid, 5bromoisophthaloyl dichloride [9,38]. The acid chloride was then dissolved in dry dichloromethane (10 mL) and cooled to 0 °C. A solution of isopropylamine (8.0 mmol, 0.655 mL) and triethylamine (4.0 mmol, 0.56 mL) in dry dichloromethane (5 mL) was added in the acid chloride solution drop by drop. The resulting mixture was kept stirring at room temperature under a nitrogen atmosphere for 16 hr [44]. The reaction mixture was then dried over a vacuum. The residue was dissolved in dichloromethane and washed with aqueous ammonium chloride solution followed by water. The organic layer was dried with anhydrous Na₂SO₄ and concentrated to dryness yielded ligand (L). Purification by silica gel column chromatography with petroleum ether and ethyl acetate (0 to 30%) mixture as eluent afforded 0.536 g (82%) of white solid (L), Rf = 0.28, TLC (eluent: petroleum ether/ethyl acetate 30:70 v/v). The ¹H NMR, ¹³C NMR, and IR spectra of the ligand are shown in Figs. 2-4 respectively.

2.2.2. Analytical and spectroscopic data

CHNS analysis: Anal. Calcd: C 51.39; H 5.85; N 8.56. Found: C 51.62; H 5.85; N 8.50; FTIR (KBr, cm⁻¹): 3251 (s, N—H str. amide), 3078 (C—H str. aromatic), 2973–2874 (s, C—H str. alkyl), 1632 (s, C=O str. amide), 1551 (s, N—H bend amide) 1591–1462 (C=C str. aromatic), 894–714 (C—H bend aromatic); ¹H NMR (400 MHz, CDCl₃, δ): 8.027 (s, 1H, ArH), 7.964 (s, 2H, 2 × ArH), 7.26 (residual solvent peak), 6.206–6.190 (d, 2H, 2 × NH amide), 4.300–4.249 (m, 2H, 2 × CH isopropyl), 1.285–1.269 (d, 12H, 4 × CH₃ isopropyl). ¹³C NMR (100 MHz. CDCl₃, δ): 164.52, 136.96, 132.63, 123.84, 122.97, 42.40, 22.73.

2.3. Extraction

The extraction profile of uranyl is obtained by measuring D_U as a function of aqueous feed acidity in various ionic liquid systems at a constant ligand concentration (1 mM) and temperature (300 K). An equivalent volume of the organic (IL) and aqueous solutions were used in the extraction process. The equilibration time and



Fig. 1. The structures of the ionic liquids used in the present investigation.



Scheme 1. (a) Synthesis of 5-bromoisophthloyl dichloride; (b) Synthesis of 5-bromo-N1, N3-diisopropylisophthalamide (L).

the time of centrifugation were 180 min and 5 min, respectively. The Distribution ratio was calculated as follows [45]

$$D_U = \frac{[M]_{IL}}{[M]_{aq}} \tag{1}$$

Where $[M]_{lL}$, and $[M]_{aq}$ are the analytical concentrations of uranyl in ionic liquid and aqueous phase. The dependence of the D_U as a function of ligand concentration was investigated by keeping the acidity of the feed solution as a constant (1 M HNO₃). For nitrate ion variation, aqueous phase NaNO₃ concentration was varied keeping other parameters constant. LiNTf₂ salt and bromide salts of the corresponding ionic liquid cations are water-soluble and hence their concentration was varied in water to investigate their involvement in the extraction of uranyl ion by the ligand. Stripping studies were carried out in two steps. In the first step, the extraction process was done. while in the subsequent step, the stripping was carried out using sodium carbonate, oxalic acid, and ethylenediaminetetraacetic acid (EDTA). Several consecutive contacts of the stripping agent were made for achieving cumulative back extraction of locked up uranyl from the ionic liquid phase. The % stripping was calculated as follows [46]

$$\% Stripping = \frac{[M]_{aq}}{[M]_{IL}} \times 100\%$$
⁽²⁾



Fig. 2. ¹H NMR Spectra of 5-bromo-N1, N3-diisopropylisophthalamide.







Fig. 4. FTIR Spectra of 5-bromo-N1, N3-diisopropylisophthalamide.

For the radiation stability study, the solvent systems were irradiated with various gamma radiation doses using ⁶⁰Co as a source. With the irradiated solvent systems, the D values were calculated and the deviation was monitored.

3. Results and discussion

3.1. Extraction profile

Extraction of uranyl ion has been carried out with a 1 mM concentration of ligand. The extraction profile as a function of aqueous feed acidity is shown in Fig. 5. In IL1, the D_U increases up to 2 M HNO₃ reaching a maximum value of ~12, and remains almost constant between 2 M and 4 M HNO₃ followed by a marginal decrease. The decrease in D_U value at higher acidity is not very significant. In the case of IL2, the D_U value increases rapidly up to 1 M HNO₃ followed by a marginal increase from 1 M to 2 M HNO₃ and remained unchanged beyond 2 M HNO₃. In the plateau region, the D_U value was ~20. In IL3 based system, the trend was almost similar. The enhancement in D_U value was gradual up to 4 M HNO₃, followed by a plateau. The maximum D_U value obtained in this case was \sim 13. Interestingly, using IL4 as a solvent system, the trend in D_U vs aqueous feed acidity is unique compared to other ionic liquid systems, i.e. D_U value was maximum (~8) at 1 M HNO₃ and decreases with an increase in feed acidity. This diluent-induced difference in extraction profiles is not very common in molecular diluents. However, in ionic liquid, structural modification can bring changes in extraction properties. The above fact was verified and well-reported for imidazolium-based ionic liquids [47,48]. In the case of methylimidazolium-based ionic liquid, it was demonstrated that if the alkyl group at the 3rd position is butyl, then there is a chance of cation exchange mechanism, where, the butyl imidazolium cation is getting exchanged with the cationic metalligand complex. However, for higher homologue, i.e. C₈mim⁺ or C₁₀mim⁺ ion, water solubility is very less and hence, the ion exchange mechanism is prohibited and the extraction proceeds through a solvation mechanism involving neutral metal-ligand



Fig. 5. The variation in D_U value as a function of aqueous feed acid concentration using 1 mM ligand in different ionic liquid.

complex. In the present case, it is quite interesting to study ionic liquids that have diverse cationic motifs. There are only a few reports that describe the extraction of metal ions in ionic liquids through the anion-exchange mechanism. One such example is the extraction of uranyl ion by tri-*n*-butyl phosphate in ionic liquid from higher aqueous feed acidity [49]. Recently anion exchange mechanism was also reported for Sr^{2+} extraction by Calix crown ether using pyridinium-based ionic liquid [26]. The increasing trend in D_U with an increase in aqueous feed acidity is an indication of either the solvation mechanism or anion exchange mechanism. The initial enhancement in D_U with aqueous feed acidity is ascribed to the participation of nitrate ions. However, at higher feed acidity, plenty of H⁺ ions are available and they compete with uranyl ion resulting in either plateau or marginal reduction in D_U

values. In the case of ionic liquid IL4, the extraction profile indicates the extraction might proceed through the cation exchange mechanism.

Solvation mechanism:

$$UO_{2 aq}^{2+} + mL_{IL} + 2NO_{3 aq}^{-} \to UO_{2}(NO_{3})_{2}.mL_{IL}$$
(3)

Cation exchange mechanism:

$$UO_{2}^{2+}{}_{aq} + mL_{IL} + nNO_{3}^{-}{}_{aq} + (2-n)IL_{IL}^{+}$$

$$\rightarrow [UO_{2}(NO_{3})_{n}.mL]_{IL}^{2-n} + (2-n)IL_{aq}^{+}$$
(4)

Anion exchange mechanism:

$$UO_{2}^{2+}{}_{aq} + mL_{IL} + nNO_{3}^{-}{}_{aq} + (n-2)IL_{IL}^{-}$$

$$\rightarrow [UO_{2}(NO_{3})_{n}.mL]_{IL}^{n-2} + (n-2)IL_{IL}^{-}$$
(5)

Where, m and n are the number of ligand molecules and nitrate ions associated with each uranyl ion in the formation of the complex. IL⁺ represents the cationic part of the ionic liquid, while IL⁻ is the anionic part of the ionic liquid.

3.2. Speciation and extraction mechanism

The equilibrium constant for the uranyl extraction can be defined as follows for solvation mechanism

$$K_{ex} = \frac{[UO_2(NO_3)_2.mL_{IL}]}{\left[UO_2^{+}_{aq}\right][L_{IL}]^m[NO_3^{-}_{aq}]^2}$$
(6)

$$K_{ex} = \frac{D_U}{[L_{IL}]^m [NO_{3 \ ag}]^2}$$
(7)

Taking logarithm on both sides and rearranging, the equation takes the form

$$Log D_{U} = Log K_{ex} + mLog[L_{IL}] + 2Log[NO_{3 aq}^{-}]$$
(8)

Similarly, for the cation exchange mechanism, the equilibrium constant can be written as

$$K_{ex} = \frac{\left[\left[UO_{2}(NO_{3})_{n} \cdot mL\right]_{lL}^{2-n}\right]}{\left[UO_{2}^{2+}{}_{aq}\right]\left[L_{L}\right]^{m}\left[NO_{3}^{-}{}_{aq}\right]^{n}} \frac{\left[L_{aq}^{+}\right]^{2-n}}{\left[L_{L}^{+}\right]^{2-n}}$$
(9)

The ratio of the concentration of cationic part of ionic liquid in both the phase is known as the partition coefficient for that species. The partition coefficient is a constant for a given temperature and also considering K_{ex} as the conditional extraction, the equation can be simplified as follows

$$K_{ex} = \frac{\left[\left[UO_2(NO_3)_n . mL \right]_{IL}^{2-n} \right]}{\left[UO_2^{2+}_{aq} \right] \left[L_{IL} \right]^m \left[NO_3^{-}_{aq} \right]^n}$$
(10)

$$Log D_{U} = Log K_{ex} + mLog [L_{IL}] + nLog [NO_{3 aq}^{-}]$$
(11)

For anion exchange mechanism the Kex can be written as follows

$$K_{ex} = \frac{\left[\left[UO_2(NO_3)_n \cdot mL \right]_{lL}^{(n-2)} \right]}{\left[UO_2^{2+}{}_{aq} \right] \left[L_{LL} \right]^m \left[NO_3^{-}{}_{aq} \right]^n} \frac{\left[IL_{LL} \right]^{n-2}}{\left[IL_{LL} \right]^{n-2}}$$
(12)

Similarly, the partition coefficient for the anionic part of the ionic liquid is a constant for a given temperature, and hence equation can be written as

$$Log D_{U} = Log K_{ex} + mLog[L_{IL}] + nLog[NO_{3aa}]$$
⁽¹³⁾

Therefore, a plot of $Log(D_U)$ as a function of Log([L]) will be a straight line with a slope, which determines the metal-ligand stoichiometry, keeping nitrate ion in the aqueous phase constant. Sim-



Fig. 6. The variation of D_U values as a function of (a) ligand concentration in ionic liquid phase; (b) nitrate ion concentration in the aqueous phase.

ilarly, a plot of $Log(D_{II})$ value as a function of $Log([NO_3^-])$ would be a straight line with a slope value, which determines the number of nitrate ions associated with each metal ion in the complex. Fig. 6 (a) shows the variation in D_U value as a function of ligand concentration. An increase in D_U value with ligand concentration indicates the direct participation of ligand molecules. For ionic liquids IL1 and IL2, the slope is evaluated as 2, while it is 1 for ionic liquids IL3 and IL4. This revealed that UO_2^{2+} formed 1:2 complex in ionic liquid IL1 and IL2, whereas 1:1 metal-ligand complex with IL3 and IL4. Fig. 6 (b) represented the variation in D_U value as a function of nitrate ion concentration in the aqueous phase. The slope for IL1 and IL2 were found to be 3 and 2, respectively. However, for IL3 and IL4, the slope is evaluated as 2 and 1, respectively. These results indicate that in IL1, the species is $[UO_2(NO_3)_3L_2]^-$, in IL2 the species is $UO_2(NO_3)_2L_2$, whereas in IL3 the species is UO₂(NO₃)₂L and in IL4 the species involved is $[UO_2(NO_3)L]^+$. This implies that in IL4, the cation exchange mechanism predominates following cationic species, and the cationic part of the ionic liquid will be exchanged in place of the metal-ligand complex during extraction to maintain the electrical neutrality. On the other hand, in IL1, the anion exchange mechanism predominates involving anionic metal-ligand species. To maintain the charge neutrality in both phases, an equivalent amount of ionic liquid anion would get exchanged. However, for IL2 and IL3, the solvation mechanism is predominant involving a neutral metal-ligand complex. However, this has to be confirmed



Fig. 7. The dependence of D_U values on the concentration of (a) cations of ionic liquid and (b) anions of ionic liquids in the aqueous phase.

by further investigation on the dependence of cations or anions of ionic liquids on the extraction properties of uranyl.

To further confirm the speciation and the extraction mechanism, cation and anion dependence of ionic liquid on D_U values were monitored using the water-soluble salts of the same [Fig. 7]. For example, in cation dependence experiments, the chloride salts of the corresponding ionic liquid were taken, while anion dependence was checked using Li salt of NTf₂⁻ anion. The chloride salts used in the present case are: 1-Hexyl-1,4-diaza[2,2,2]bicy clooctanium chloride, 1-Propylpyridinium chloride, 1-Butyl-1 methylpiperidinium chloride, 1-Hexyl-3methylimidazolium chloride. In cation dependence experiments, the D_U values were independent of the concentration of ionic liquid cations in the aqueous phase for IL1, IL2, and IL3, suggesting no cation exchange mechanism involved. However, the D_U value decreases with an increase in ionic liquid cation concentration in the aqueous phase for IL4. The concentration dependence of the D_U value in IL4 confirms the participation of IL cation in the extraction process. More the initial concentration of cations in water, the equilibrium of the extraction process shifted more towards the backward direction. i.e. the extent of uranyl extraction would reduce. The slope of the linear dependence of D_{II} in IL4 is evaluated as -1. This implies that only one C₈mim⁺ ion gets exchanged during each metal-ligand complex extraction. This also proves that a singly charged metalligand complex formed during extraction of uranyl with amide

functionalized ligand in C₈mimNTf₂. On a similar note, anion dependence experiments suggested that, for IL2, IL3, and IL4, the D_{II} remained unchanged with an increase in the NTf₂⁻ ion concentration in the aqueous phase. However, for IL1, i.e. for bicyclooctanium ionic liquid, the D_U decreases with an increase in NTf_2^- ion concentration in the aqueous phase with a slope value of -1. This result confirms that, for IL2 and IL3, neither cation exchange nor anion exchange mechanism was predominant. Therefore, the extraction proceeds through the neutral metal-ligand complex with a 'solvation' mechanism. However, for IL1, i.e. bicyclooctanium ionic liquid the extraction proceeded via mono negative Uligand complex. To maintain the charge neutrality, one NTf_2^- anion would be transferred from the IL phase to the aqueous phase for the transfer of each U-ligand complex. In the case of IL4, C₈mim⁺ cation got exchanged in place of mono positive U-ligand complex and the extraction proceeded via cation exchange mechanism. This particular case is very interesting to demonstrate that by changing the cations of the ionic liquids, the extraction species and mechanism can be changed from solvation to cation exchange or anion exchange.

Fig. 8 depicts the most plausible uranyl-ligand complexes in different ionic liquids. In IL2 and IL3, neutral species were formed. However, the speciation and mode of coordination are quite different. In IL2, two diamides, and two nitrate ions were found to be coordinating in a monodentate fashion, whereas in IL3, one diamide was found to be coordinating through bidentate fashion, while two nitrates are in monodentate mode. In IL1, the species was found to be anionic with two ligands and three nitrates all are in the monodentate mode of coordination. However, in IL4 the species was mono positive with one diamide and one nitrate both in the bidentate mode of coordination with uranyl to satisfy the 8 coordination number.

3.3. Radiolytic stability

For long time use of the ligands, the solvent system must be radiologically stable, since this would experience lots of radiation coming out of the isotopes processed using the solvent system [50,51]. To understand the radiolytic stability, the solvent systems were exposed to gamma rays of different dose rates, and the D_U values were evaluated. The change in D_U values in irradiated solvent compared to the unirradiated solvent would pertain to the radiological damage faced by them. From Fig. 9, it is clear that for IL3 and IL4, the D_U values of unirradiated systems were ~8 and ~4, respectively. On irradiation with gamma rays, no significant change in D_U values observed. This implies that these two solvent systems were highly radiologically stable. However, IL2 showed a maximum reduction in D_U values. In the unirradiated condition, the D_U was 17, which was drastically reduced to 2 at a dose of 500 kGy. IL1 also showed similar results, i.e. unirradiated system showed D_U of 11, which reduced to ~ 3.8 after a dose of 500 kGy. Therefore, high radiological stability along with the efficiency of the solvent should be considered for processing high radioactive isotopes. In view of that, IL4 might be the best solvent among the four ionic liquids.

3.4. Back extraction

The reusability of the solvent system is another important aspect to be looked into. The expectation is to have solvent systems having high extraction efficiency as well as can be regenerated very easily after extraction. In view of this, stripping behavior needs to be investigated. It is well-known fact that only dilute acid might not be sufficient to back extract the complexed uranium from ionic liquid. Several methodologies including electro-deposition have



Fig. 8. The U complexes with the diamides in different ionic liquids involved in the separation.



Fig. 9. The radiolytic stability of the solvent systems under the influence of gammaray exposure.

been employed for back extraction [52]. However, another wellaccepted approach is to make use of the complexing agent in water [53]. The idea is to make a stronger complex of uranyl in water so that the uranium-amide complex in ionic liquid breaks up. Based on the literature, 1 mM aqueous solution of oxalic acid, sodium carbonate, and ethylenediaminetetraacetic acid have been chosen as aqueous complexing agents for back extraction. Fig. 10(a) shows the % of stripping of uranium using the stripping solution in a single contact. The results indicate that an aqueous solution of sodium carbonate is the best out of these three stripping solutions. Using aqueous sodium carbonate as a stripping solution, the effect of a



Fig. 10. (a) The stripping behavior of uranium from the uranyl-diamide complex in ionic liquid using different aqueous complexing agents; (b) Cumulative % of stripping of locked up uranium using sodium carbonate in multiple contacts.



Fig. 11. The variation in D_{U} as a function of contact time of organic phase with 1 M $\text{HNO}_{3\text{.}}$

different number of contact cycles on % stripping was investigated. Four contacts of IL1 were required for about 99.5% back extraction, while 2 contacts of IL2 and IL4 were sufficient for almost 99% back extraction. Back extraction from IL3 was not as satisfactory as observed in other cases. Even after five contacts, the cumulative back extraction was slightly less than 99%. Fig. 10 (b) depicts the cumulative % back extraction of uranyl ion from the uranyldiamide complex in ionic liquid.

The variation of D_U values for the ionic liquids having prior contact (1 hr) with 1 M HNO₃ as a function of time is shown in Fig. 11. This investigation reveals the chemical stability of the solvent systems in the presence of 1 M HNO₃. The D values were monitored up to 72 h i.e. 3 days of contact time. D_U value is almost independent of time from 1 hr onwards, suggesting that all the ionic liquid based solvent systems are chemically stable. Table 1 summarizes the mechanism, speciation and other chanrecteristic of extraction for the four ionic liquid system used in the study.

4. Conclusion

This manuscript describes the first-ever report of tuning the extraction mechanism of uranyl ion from cation exchange to anion exchange or solvation mechanism by changing the cationic part of the ionic liquid for the same uranyl-diamide systems. The novel diamide ligand (5-Bromo-N1, N3-diisopropylisophthalamide) was synthesized and characterized by ¹H NMR, ¹³C NMR and FTIR spectroscopy. In imidazolium-based ionic liquid, the extraction proceeds via monopositive $[UO_2(NO_3)L]^+$ species. For pyridinium ionic liquid, the extraction proceeds via neutral $UO_2(NO_3)_2L_2$ species, whereas for piperidiniumionic liquid, neutral $UO_2(NO_3)_2L$ species.

Table 1

The extraction characteristics of the ionic liquid based solvent systems.

cies is predominating. However, for bicyclooctanium ionic liquid, monoanionic complex $[UO_2(NO_3)_3L_2]^-$ was predominating. The radiation stability of pyridinium and imidazolium-based ionic liquids was more than that of the other two ionic liquid systems. An aqueous solution of sodium carbonate (1 mM) back extracts the uranyl ion effectively from the ionic liquid phase compared to oxalic acid and EDTA. Four contacts of eluent are required for back extraction of uranyl from bicyclooctanium ionic liquid, whereas 2 contacts were sufficient for imidazolium and pipyredinium based ionic liquid. In pyridinium-based ionic liquid, the uranyl complex was so strong, and hence can not be stripped quantitatively even after five contacts.

CRediT authorship contribution statement

Amit Pandey: Formal analysis, Investigation **S. Hashmi:** Data curation, Investigation **G. Salunkhe:** Validation. **Velavan Kathivelu:** Supervision, Writing, Review and Editing. **K.S.Singh:** Methodology. **A. Sengupta:** Concceptualization, Supervision, Writing Original Draft, Writing, Review and Editing. **R.S.Chauhan:** Supervision.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgment

The author acknowledges Dr. P.K.Pujari, Director, RC & I Group, Head, Radiochemistry Division, and Dr. R. Acharya, Head, Actinide Spectroscopy section, Radiochemistry Division, Bhabha Atomic Research Centre, Mumbai, India. Dr. Velavan Kathirvelu acknowledges the financial support by the Science and Engineering Research Board, India (SB/FT/CS-119/2014)

References

- D. Han, K.H. Row, Recent applications of ionic liquids in separation technology, Molecules 15 (2010) 2405–2426, https://doi.org/10.3390/molecules15042405.
- [2] A. Berthod, M.J. Ruiz-Ángel, S. Carda-Broch, Recent advances on ionic liquid uses in separation techniques, J. Chromatogr. 1559 (2018) 2–16, https://doi. org/10.1016/j.chroma.2017.09.044.
- [3] G.-T. Wei, Z. Yang, C.-J. Chen, Room temperature ionic liquid as a novel medium for liquid/liquid extraction of metal ions, Anal. Chim. Acta 488 (2) (2003) 183–192, https://doi.org/10.1016/S0003-2670(03)00660-3.
- [4] I.A. Shkrob, T.W. Marin, M.P. Jensen, Ionic liquid based separations of trivalent lanthanide and actinide ions, Ind. Eng. Chem. Res. 53 (9) (2014) 3641–3653, https://doi.org/10.1021/ie4036719.
- [5] M. Kamaz, A. Sengupta, S.S. DePaz, Y.-H. Chiao, S. Ranil Wickramasinghe, Poly (ionic liquid) augmented membranes for π electron induced separation/ fractionation of aromatics, J. Membr. Sci. 579 (2019) 102–110, https://doi.org/ 10.1016/j.memsci.2019.01.044.
- [6] M. Kamaz, R.J. Vogler, M. Jebur, A. Sengupta, R. Wickramasinghe, π Electron induced separation of organic compounds using supported ionic liquid

Metal	Ligand	IL	D (1 M HNO ₃)	Mechanism	Species	No of contact for stripping
UO2 ⁺	5-bromo-N1, N3- diisopropylisophthalamide	1-Hexyl-1,4-diaza[2,2,2]bicyclooctaniumbis (trifluoromethylsulfonyl)imide	11	Anion exchange	$[UO_2(NO_3)_3L_2]^-$	4
		1-Propylpyridinium bis(trifluoromethylsulfonyl)imide	16	Solvation	$UO_2(NO_3)_2L_2$	2
		1-Butyl-1methylpiperidinium bis (trifluoromethylsulfonyl)imide	3.7	Solvation	$UO_2(NO_3)_2L$	even 5 is not sufficient
		1-Hexyl-3methylimidazolium bis (trifluoromethylsulfonyl)imide	8	Cation exchange	$\left[\mathrm{UO}_2(\mathrm{NO}_3)\mathrm{L}\right]^+$	2

membranes, Sep. Purif. Technol. 236 (2020) 116237, https://doi.org/10.1016/j. seppur.2019.116237.

- [7] M.L. Dietz, J.A. Dzielawa, I. Laszak, B.A. Young, M.P. Jensen, Influence of solvent structural variations on the mechanism of facilitated ion transfer into roomtemperature ionic liquids, Green Chem. 5 (6) (2003) 682–685, https://doi.org/ 10.1039/B310507P.
- [8] M.P. Jensen, J.A. Dzielawa, P. Rickert, M.L. Dietz, EXAFS investigations of the mechanism of facilitated ion transfer into a room-temperature ionic liquid, J. Am. Chem. Soc. 124 (36) (2002) 10664–10665, https://doi.org/10.1021/ja027476y.
- [9] A. Pandey, S. Hashmi, V. Kathirvelu, K.S. Singh, A. Sengupta, Effect of alkyl chain length on the extraction properties of U and Th using novel CnmimNTf2/ isophthalamide systems, J. Mol. Liq. 323 (2021) 114944–114948, https://doi. org/10.1016/j.molliq.2020.114944.
- [10] A. Sengupta, P.K. Mohapatra, M. Iqbal, J. Huskens, W. Verboom, A highly efficient solvent system containing functionalized diglycolamides and an ionic liquid for americium recovery from radioactive wastes, Dalton Trans. 41 (2012) 6970–6979, https://doi.org/10.1039/C2DT12364A.
- [11] A. Sengupta, P.K. Mohapatra, M. Iqbal, W. Verboom, J. Huskens, S.V. Godbole, Extraction of Am(iii) using novel solvent systems containing a tripodal diglycolamide ligand in room temperature ionic liquids: a 'green' approach for radioactive waste processing, RSC Adv. 2 (2012) 7492–7500, https://doi. org/10.1039/C2RA20577G.
- [12] P.K. Mohapatra, A. Sengupta, M. Iqbal, J. Huskens, W. Verboom, Highly efficient diglycolamide-based task-specific ionic liquids: Synthesis, unusual extraction behaviour, irradiation, and fluorescence studies, Chem. Eur. J. 19 (9) (2013) 3230–3238, https://doi.org/10.1002/chem.201203321.
- [13] P.K. Mohapatra, A. Sengupta, M. Iqbal, J. Huskens, W. Verboom, Diglycolamidefunctionalized calix[4]arenes showing unusual complexation of actinide ions in room temperature ionic liquids: Role of ligand structure, radiolytic stability, emission spectroscopy, and thermodynamic studies, Inorg. Chem. 52 (5) (2013) 2533–2541, https://doi.org/10.1021/ic302497k.
- [14] A. Sengupta, P.K. Mohapatra, M. Iqbal, J. Huskens, W. Verboom, Solvent systems containing diglycolamide-functionalised calix[4]arenes in room temperature ionic liquid for metal ion extraction: studies with simulated high-level wastes, Supramol. Chem. 26 (7-8) (2014) 612–619, https://doi.org/ 10.1080/10610278.2014.885025.
- [15] I.A. Shkrob, S.D. Chemerisov, J.F. Wishart, The initial stages of radiation damage in ionic liquids and ionic liquid-based extraction systems, J. Phys. Chem. B 111 (40) (2007) 11786–11793, https://doi.org/10.1021/jp073619x10.1021/ jp073619x.s00110.1021/jp073619x.s002.
- [16] J.F. Wishart, Ionic liquids and ionizing radiation: Reactivity of highly energetic species, J. Phys. Chem. Lett. 1 (21) (2010) 3225–3231, https://doi.org/10.1021/ jz101096b.
- [17] N.K. Gupta, Ionic liquids for transuranic extraction (TRUEX)–Recent developments in nuclear waste management: A review, J. Mol. Liq. 269 (2018) 72–91, https://doi.org/j.molliq.2018.08.036.
- [18] M. Paramanik, D.R. Raut, A. Sengupta, S.K. Ghosh, P.K. Mohapatra, A trialkyl phosphine oxide functionalized task specific ionic liquid for actinide ion complexation: extraction and spectroscopic studies, RSC Adv. 6 (24) (2016) 19763–19767, https://doi.org/10.1039/C5RA24174J.
- [19] A. Sengupta, P.K. Mohapatra, P. Pathak, T.K. Ghanty, W. Verboom, Studies on neptunium complexation with CMPO- and diglycolamide-functionalized ionic liquids: experimental and computational studies, New J. Chem. 41 (2) (2017) 836–844, https://doi.org/10.1039/C6NJ03102A.
- [20] A. Rout, K. Binnemans, Solvent Extraction of Neodymium(III) by Functionalized Ionic Liquid Trioctylmethylammonium Dioctyl Diglycolamate in Fluorine-free Ionic Liquid Diluent, Ind. Eng. Chem. Res. 53 (15) (2014) 6500–6508, https:// doi.org/10.1021/ie404340p.
- [21] G. Jing, Y. Qian, X. Zhou, B. Lv, Z. Zhou, Designing and screening of multiamino-functionalized ionic liquid solution for co2 capture by quantum chemical simulation, ACS Sustain. Chem. Eng. 6 (1) (2018) 1182–1191, https://doi.org/10.1021/acssuschemeng.7b03467.
- [22] T. Vander Hoogerstraete, B. Onghena, K. Binnemans, Homogeneous liquidliquid extraction of metal ions with a functionalized ionic liquid, J. Phys. Chem. Lett. 4 (10) (2013) 1659–1663, https://doi.org/10.1021/jz4005366.
- [23] D. Das, S. Hashmi, A. Sengupta, S. Kannan, C.P. Kaushik, Understanding the extraction behaviour of UO²⁺₂ and Th⁴⁺ using novel picolinamide/N-oxo picolinamide in ionic liquid: A comparative evaluation with molecular diluent, J. Mol. Liq. 332 (2021) 115773, https://doi.org/10.1016/j.molliq.2021.115773.
- [24] M. Singh, A. Sengupta, M.S. Murali, S.K. Thulasidas, R.M. Kadam, Selective separation of uranium from nuclear waste solution by bis(2,4,4trimethylpentyl)phosphinic acid in ionic liquid and molecular diluents: a comparative study, J. Radioanal. Nucl. Chem. 309 (3) (2016) 1199–1208, https://doi.org/10.1007/s10967-016-4691-y.
- [25] N.K. Gupta, A. Sengupta, Sulphoxide ligands inbicyclooctanium based ionic liquid: Novel solvent systems for the extraction of f-elements, Sep. Sci. Technol. 54 (8) (2019) 1312–1324, https://doi.org/10.1080/ 01496395.2018.1533869.
- [26] A. Sengupta, P.K. Mohapatra, Extraction of radiostrontium from nuclear waste solution using crown ethers in room temperature ionic liquids, Supramol. Chem. 24 (11) (2012) 771–778, https://doi.org/10.1080/ 10610278.2012.716840.
- [27] K. Raj, K.K. Prasad, N.K. Bansal, Radioactive waste management practices in India, Nucl. Eng. Des. 236 (7-8) (2006) 914–930, https://doi.org/10.1016/j. nucengdes.2005.09.036.

- [28] R.D. Ambashta, M.E.T. Sillanpää, Membrane purification in radioactive waste management: a short review, J. Environ. Radioact. 105 (2012) 76–84, https:// doi.org/10.1016/j.jenvrad.2011.12.002.
- [29] J.N. Mathur, M.S. Murali, K.L. Nash, Actinide partitioning-A review, Solvent Extr. Ion Exch. 19 (3) (2001) 357–390, https://doi.org/10.1081/SEI-100103276.
- [30] B.J. Mincher, G. Modolo, S.P. Mezyk, Review article: The effects of radiation chemistry on solvent extraction: 1. Conditions in acidic solution and a review of TBP radiolysis, Solvent Extr. Ion Exch. 27 (1) (2009) 1–25, https://doi.org/ 10.1080/07366290802544767.
- [31] J.N. Mathur, M.S. Murali, P.R. Natarajan, L.P. Badheka, A. Banerji, A. Ramanujam, P.S. Dhami, V. Gopalakrishnan, R.K. Dhumwad, M.K. Rao, Partitioning of actinides from high-level waste streams of purex process using mixtures of CMPO and TBP in dodecane, Waste Manage. 13 (4) (1993) 317–325, https://doi.org/10.1016/0956-053X(93)90060-A.
- [32] P. Tkac, G.F. Vandegrift, G.J. Lumetta, A.V. Gelis, Study of the interaction between HDEHP and CMPO and its effect on the extraction of selected lanthanides, Ind. Eng. Chem. Res. 51 (31) (2012) 10433–10444, https://doi.org/ 10.1021/ie300326d.
- [33] A. Sengupta, S.M. Ali, K.T. Shenoy, Understanding the complexation of the Eu3+ ion with TODGA, CMPO, TOPO and DMDBTDMA: Extraction, luminescence and theoretical investigation, Polyhedron 117 (2016) 612–622, https://doi.org/ 10.1016/j.poly.2016.06.037.
- [34] M.P. Jensen, A.H. Bond, Comparison of covalency in the complexes of trivalent actinide and lanthanide cations, J. Am. Chem. Soc. 124 (33) (2002) 9870–9877, https://doi.org/10.1021/ja0178620.
- [35] V.N. Romanovskiy, I.V. Smirnov, V.A. Babain, T.A. Todd, R.S. Herbst, J.D. Law, K. N. Brewer, The universal solvent extraction (unex) process. I. Development of the unex process solvent for the separation of cesium, strontium, and the actinides from acidic radioactive waste, Solvent Extr. Ion Exch. 19 (1) (2001) 1–21, https://doi.org/10.1081/SEI-100001370.
- [36] G.M. Casparini, G. Grossi, Application of n, n-dialkyl aliphatic amides in the separation of some actinides, Sep. Sci. Technol. 15 (4) (1980) 825–844, https:// doi.org/10.1080/01496398008076273.
- [37] S.D. Reilly, A.J. Gaunt, B.L. Scott, G. Modolo, M. Iqbal, W. Verboom, M.J. Sarsfield, Plutonium(iv) complexation by diglycolamide ligands—coordination chemistry insight into TODGA-based actinide separations, Chem. Commun. 48 (2012) 9732–9734, https://doi.org/10.1039/C2CC34967A.
- [38] A. Pandey, G. Salunkhe, V. Kathirvelu, K.S. Singh, A. Sengupta, Exploring novel functionality for efficient extraction of UO₂⁺ and Th⁴⁺in ionic liquid: Mechanism, speciation, selectivity, stability and stripping, J. Mol. Liq. 324 (2021) 114716, https://doi.org/10.1016/j.molliq.2020.114716.
- [39] N. Condamines, C. Musikas, The extraction by n.N-dialkylamides. II. Extraction of actinide cations, Solvent Extr. Ion Exch. 10 (1) (1992) 69–100, https://doi. org/10.1080/07366299208918093.
- [40] Y. Sasaki, Y. Sugo, K. Morita, K.L. Nash, The effect of alkyl substituents on actinide and lanthanide extraction by diglycolamide compounds, Solvent Extr. Ion Exch. 33 (7) (2015) 625–641, https://doi.org/10.1080/ 07366299.2015.1087209.
- [41] J. Sk, M.s. Sunita Pathak, A. Sengupta, A comprehensive investigation along with the statistical evaluation for the characterization of ilmenite mineral by x-ray fluorescence spectrometry and optical emission spectrometry, ChemistrySelect 6 (8) (2021) 1911–1919, https://doi.org/10.1002/slct. v6.810.1002/slct.202004205.
- [42] S.k. Jayabun, S. Pathak, A. Sengupt, Determination of major, minor and trace metallic constituents in sillimanite: Comparative evaluation of EDXRF and DC arc carrier distillation AES, At. Spectrosc. 41 (4) (2020) 162–168.
- [43] A. Sengupta, B. Rajeswari, R. Mahadeo Kadam, AES and XRF based comparative evaluation of metallic constituents at trace and minor levels in contaminated neoprene gauntlets and cellulosic materials, ChemistrySelect 5 (13) (2020) 3763–3769, https://doi.org/10.1002/slct.v5.1310.1002/slct.202000156.
- [44] P. Devi, S.M. Barry, K.M. Houlihan, M.J. Murphy, P. Turner, P. Jensen, P.J. Rutledge, Synthesis and structural characterisation of amides from picolinic acid and pyridine-2,6-dicarboxylic acid, Sci. Rep. 5 (2015) 9950-, https://doi. org/10.1038/srep09950.
- [45] A. Rout, S. Karmakar, K.A. Venkatesan, T.G. Srinivasan, P.R. Vasudeva Rao, Room temperature ionic liquid diluent for the mutual separation of europium (III) from americium(III), Sep. Purif. Technol. 81 (2) (2011) 109–115, https:// doi.org/10.1016/j.seppur.2011.04.033.
- [46] C. Marie, P. Kaufholz, V. Vanel, M.-T. Duchesne, E. Russello, F. Faroldi, L. Baldini, A. Casnati, A. Wilden, G. Modolo, M. Miguirditchian, Development of a selective americium separation process using H4TPAEN as water-soluble stripping agent, Solvent Extr. Ion Exch. 37 (5) (2019) 313–327, https://doi.org/ 10.1080/07366299.2019.1643569.
- [47] A. Sengupta, P.K. Mohapatra, M. Iqbal, J. Huskens, W. Verboom, A diglycolamide-functionalized task specific ionic liquid (TSIL) for actinide extraction: Solvent extraction, thermodynamics and radiolytic stability studies, Sep. Purif. Technol. 118 (2013) 264–270, https://doi.org/10.1016/j. seppur.2013.07.005.
- [48] A. Rout, K. Binnemans, Influence of the ionic liquid cation on the solvent extraction of trivalent rare-earth ions by mixtures of Cyanex 923 and ionic liquids, Dalton Trans. 44 (3) (2015) 1379–1387, https://doi.org/10.1039/ C4DT02766C.
- [49] P. Giridhar, K.A. Venkatesan, T.G. Srinivasan, P.R. Vasudeva Rao, Extraction of uranium(VI) from nitric acid medium by 1.1M tri-n-butylphosphate in ionic liquid diluent, J. Radioanal. Nucl. Chem. 265 (1) (2005) 31–38, https://doi.org/ 10.1007/s10967-005-0785-7.

A. Pandey, S. Hashmi, G. Salunkhe et al.

- [50] B.J. Mincher, J.F. Wishart, The radiation chemistry of ionic liquids: A review, Solvent Extr. Ion Exch. 32 (6) (2014) 563–583, https://doi.org/10.1080/ 07366299.2014.925687.
- [51] I.A. Shkrob, T.W. Marin, J.R. Bell, H. Luo, S. Dai, J.L. Hatcher, R.D. Rimmer, J.F. Wishart, Radiation-induced fragmentation of diamide extraction agents in ionic liquid diluents, J. Phys. Chem. B 116 (7) (2012) 2234–2243, https://doi. org/10.1021/jp2117483.
- [52] P.-Y. Chen, The assessment of removing strontium and cesium cations from aqueous solutions based on the combined methods of ionic liquid extraction and electrodeposition, Electrochim. Acta 52 (17) (2007) 5484–5492, https:// doi.org/10.1016/j.electacta.2007.03.010.
- [53] P.S. Kulkarni, Recovery of uranium(VI) from acidic wastes using tri-noctylphosphine oxide and sodium carbonate based liquid membranes, Chem. Eng. J. 92 (1-3) (2003) 209–214, https://doi.org/10.1016/S1385-8947(02)00255-3.