

Journal Pre-proofs

Optimisation of Estrogen Receptor Subtype-Selectivity of a 4-Aryl-4*H*-Chromene Scaffold Previously Identified by Virtual Screening

Miriam Carr, Andrew J.S. Knox, Daniel K. Nevin, Niamh O'Boyle, Shu Wang, Billy Egan, Thomas McCabe, Brendan Twamley, Daniela M. Zisterer, David G. Lloyd, Mary J. Meegan

PII: S0968-0896(19)31245-3
DOI: <https://doi.org/10.1016/j.bmc.2019.115261>
Reference: BMC 115261


To appear in: *Bioorganic & Medicinal Chemistry*

Received Date: 22 July 2019
Revised Date: 3 December 2019
Accepted Date: 9 December 2019

Please cite this article as: M. Carr, A.J.S. Knox, D.K. Nevin, N. O'Boyle, S. Wang, B. Egan, T. McCabe, B. Twamley, D.M. Zisterer, D.G. Lloyd, M.J. Meegan, Optimisation of Estrogen Receptor Subtype-Selectivity of a 4-Aryl-4*H*-Chromene Scaffold Previously Identified by Virtual Screening, *Bioorganic & Medicinal Chemistry* (2019), doi: <https://doi.org/10.1016/j.bmc.2019.115261>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier Ltd.



Optimisation of Estrogen Receptor Subtype-Selectivity of a 4-Aryl-4*H*-Chromene Scaffold Previously Identified by Virtual Screening.

Miriam Carr^{1,2#}, Andrew J.S. Knox^{1,3#*}, Daniel K. Nevin¹, Niamh O'Boyle², Shu Wang², Billy Egan², Thomas McCabe⁴, Brendan Twamley⁴, Daniela M. Zisterer¹, David G. Lloyd¹ and Mary J. Meegan²

¹School of Biochemistry and Immunology, Trinity Biomedical Sciences Institute, 152 - 160 Pearse Street Trinity College Dublin, Dublin 2, Ireland

²School of Pharmacy and Pharmaceutical Sciences, Trinity Biomedical Sciences Institute, 152 - 160 Pearse Street Trinity College Dublin, Dublin 2, Ireland

³School of Biological and Health Sciences, Technology University Dublin, Dublin City Campus, Kevin St., Dublin 8, D08 NF82, Ireland

⁴School of Chemistry, Trinity College Dublin, Dublin 2, Ireland

Abstract

4-Aryl-4*H*-Chromene derivatives have been previously shown to exhibit anti-proliferative, apoptotic and anti-angiogenic activity in a variety of tumor models in vitro and in vivo generally via activation of caspases through inhibition of tubulin polymerisation. We have previously identified by Virtual Screening (VS) a 4-aryl-4*H*-chromene scaffold, of which two examples were shown to bind Estrogen Receptor α and β with low nanomolar affinity and <20-fold selectivity for α over β and low micromolar anti-proliferative activity in the MCF-7 cell line. Thus, using the 4-aryl-4*H*-chromene scaffold as a starting point, a series of compounds with a range of basic aryloxy groups at C-4 and modifications at the C3-ester substituent of the benzopyran ring were synthesised, producing some potent ER antagonists in the MCF-7 cell line which were highly selective for ER α (compound **35**; 350-fold selectivity) or ER β (compound **42**; 170-fold selectivity).

Keywords

Estrogen receptor alpha, Estrogen receptor beta, Isoform selectivity, Subtype Selectivity, 4-Aryl-4*H*-Chromene, Benzopyran, Anticancer, Anti-proliferative, Breast cancer, Cytotoxic, Knoevenagel condensation, Molecular Modeling

Abbreviations

1. Introduction

Estrogen Receptors (ER) are hormone regulated receptors that are known to produce their long-term effects on cell structure and function *via* intracellular signalling and subsequent modulation of gene expression. In addition to genomic signalling, a rapid non-genomic signalling pathway has also been identified which is mediated by cell membrane-associated estrogen receptors. In fact a G-Protein Coupled Estrogen Receptor (GPR30/GPER) has been identified that binds estrogen, initiating a plasma membrane-initiated signaling cascade¹⁻⁴. Focusing on 'classical' ER, two subtypes exist and are expressed in a wide-range of tissues and cells throughout the body; ER alpha (ER α) is predominantly found in the female bone, uterus, ovary (thecal cells), mammarys, adipose tissue, cardiovascular system and brain. ER beta (ER β) is found also in the adipose tissue, cardiovascular system and brain but also in the lung, bladder, colon and granulosa cells of the ovary⁵⁻¹⁴. The concept of Selective Estrogen Receptor Modulation (SERM)¹⁵ was first demonstrated with the clinical drug Tamoxifen^{16,17} which was introduced in the 1970s as a treatment for advanced breast cancer in postmenopausal women. Tamoxifen was shown to have anti-estrogenic function in the breast, however partial estrogenic effects in other target tissues such as bone and the cardiovascular system where it is beneficial but conversely harmful in the endometrium¹⁸⁻²¹. However a strong association between its use and the development of endometrial carcinoma emerged fuelling research for the discovery of alternative compounds with mixed agonistic/antagonistic profiles (Figure 1).

Figure 1.

Figure 1: Structures of selected SERM structural classes including triarylethylenes tamoxifen and endoxifen, benzothiophenes raloxifene and arzoxifene, indoles pipendoxifene and bazedoxifene, benzopyran acolbifene and the tetrahydronaphthalene lasofoxifene.

It has been since hypothesised that the optimal route to the development of an ‘ideal’ SERM can only occur when delineation of the full array of the biological roles orchestrated by ER α or ER β in different tissues is revealed, and this is best achieved through the design of subtype-selective ER ligands. To date, a plethora of both ER α and ER β selective agonists and antagonists have been described (compounds **1-12**, Figure 2)²²⁻⁵⁷. The ER α agonist 16 α -LE2 (**1**) has been reported to inhibit cardiac hypertrophy⁵⁸. The synthesis and characterisation of a series of pyrazole-based compounds resulted in a >400-fold selective agonist for ER α , 1,3,5-tris(4-hydroxyphenyl)-4-propyl-1*H*-pyrazole, (**2**) (PPT)⁴⁴. Mortenson et al subsequently synthesised a library of heterocycle-based (furans, thiophenes, and pyrroles) ER agonists, the most ER α selective (65-fold) of which was compound **3** - 2,4,5-tris(4-hydroxyphenyl)-3-methylfuran³⁹. Several SERMs have also been reported with ER α selectivity, for example; benzothiophenes (**4**), tetrahydroisoquinolines (**5**)⁵⁹ and pyrazoles⁴⁵, with the most selective ER α antagonist to date being a pyrazole core incorporating a basic side-chain termed MPP (**6**)(~220-fold)⁶⁰. To facilitate the study of ER β specific mechanisms of action, the first reported full agonist discovered some time ago by Katzenellenbogen and co-workers, diarylpropionitrile(**7**) (DPN), exhibited 70-fold preferential binding for ER β over ER α and 78-fold selectivity in transcriptional assays³⁷. ERB-041 (Prinaberel)(**8**), WAY-200070 and WAY-202196(**9**) were all developed by Wyeth and reported in 2004-5⁶¹⁻⁶³, followed by a series of ER β selective benzopyrans developed by Lilly in 2006^{41,64,65}. The wide range of activities of these compounds both *in vitro* and *in vivo* has made it particularly difficult to assign clinical relevance to them and has also been additionally hampered by the existence of different isoforms of ER β . ER β 1 appears to be the isoform displaying the strongest activity, however, some variant forms may interact with ER β 1 and also with ER α presenting a challenge in understanding the full complexity of the biology of ER β ⁶⁶. To date, very few examples of ER β antagonists have been reported (Figure 2). Two 30-fold ER β selective antagonists have been described; one based on a triazine scaffold (**10**) was discovered by

GSK³² and another by researchers at Organon⁶⁷ belonging to the class of 10-aryl substituted benzo[*b*]fluorenes (**11**). The (*R,R*)-*cis*-tetrahydrochrysenes (**12**) (THC) was subsequently demonstrated by the Katzenellenbogen lab to behave as a full agonist in ER α and full antagonist in ER β ³⁶. Surprisingly, as THC lacks a basic side-chain, the term ‘passive antagonism’ was proposed for its mechanism of action supported by crystal structures of THC bound to both ER α and ER β ligand binding domains (LBDs)⁶⁸.

We have described in earlier work the identification by *in silico* screening of a novel 4-aryl-4*H*-chromene scaffold which was shown to potently bind both ER isoforms⁶⁹. The two 4-aryl-4*H*-chromene analogs (**13** and **14**) displayed modest ER α selectivity (20-fold) and also antiproliferative activity in an ER positive breast cancer cell line, MCF-7, (Figure 3). Importantly, it was noted that neither analog incorporated a typical basic arylether and this formed the premise of the current study. In the present work, a series of 4-aryl-4*H*-chromene derivatives were designed and synthesised some of which incorporated the typical basic arylethers usually required in ER antagonists (Figure 2). In addition, we have now carried out modifications at the C3 ester position of the benzopyran ring to probe potential differences in ER binding pockets thereby maximising isoform selectivity. The most active compound in the series exhibited 350-fold selectivity for ER α and interestingly one of the series showed 170-fold selectivity for ER β . The 4-aryl-4*H*-chromene scaffold has been reported in numerous studies to possess anti-proliferative, apoptotic and anti-angiogenic activity in a variety of tumor models *in vitro* and *in vivo*, generally *via* activation of caspases through inhibition of tubulin polymerisation⁷⁰. In addition, the inhibition of insulin-regulated aminopeptidase by related 4-aryl-4*H*-chromenes has been recently reported⁷¹. The 4-aryl-4*H*-chromene Crolibulin (**15**), Figure 3, (a microtubule destabilizing agent that disrupts vascular endothelial cells, and in turn, blood flow to the tumor) is currently in Phase I/II clinical trials assessing its toxicity levels when co-administered with cisplatin and progression-free survival (PFS) in adults with anaplastic thyroid cancer (ATC)⁷². We

envisage that further biochemical analysis of our 4-aryl-4*H*-chromene compound series could potentially lead to the development of novel clinically relevant ER modulating compounds featuring the 4-aryl-4*H*-chromene scaffold structure.

Figure 2. Subtype selective ER ligands

Figure 3. 4-Aryl-4*H*-chromenes **13**, **14** and Crolibulin (**15**)

2. Results

2.1 Chemistry

Many approaches to the preparation of 4-aryl-4*H*-chromenes have been described^{71,73–76}. In the present study, the synthetic route to the 4-aryl-4*H*-chromenes (**16-25**) initially identified for the study is outlined in Scheme 1. These compounds contain methyl ester, ethyl ester or nitrile substituents at C-3 of the chromene ring and also benzyl, ethyl, methyl ether or phenol substituents on the C-4 aryl ring. The appropriate arylaldehyde is reacted with malononitrile or the cyanoacrylate ester and resorcinol in the presence of base (piperidine or triethylamine) in a one-pot synthesis. The reaction proceeds through the formation of an intermediate cyanoacrylate ester or benzyldiene malonitrile by Knoevenagel condensation from the cyanoacetate ester, malonitrile and the benzaldehyde. This intermediate cyanoacrylate ester can be observed as a solid forming within ten minutes of heating the mixture. Further reaction of the resorcinol phenolic OH with the nitrile and subsequent electrophilic ring closure afforded the desired final chromene products, **16-25**. The cyanoacrylate intermediate product was isolated in the case of **26-28** and subsequently reacted with resorcinol to afford the desired products. The phenolic methyl ester **19** was also obtained from the benzyl ether **21** by hydrogenation (H₂/Pd/C) (Scheme 1).

The series of 4-aryl-4*H*-chromenes containing a basic ether substituent at C-4 (compounds

24-36) were obtained as outlined in Scheme 2. These compounds also contain methyl, ethyl, *n*-butyl or *tert*-butyl ester or nitrile substituents at C-3 of the chromene. The benzaldehydes **29-31** were obtained by alkylation of 4-hydroxybenzaldehyde with 1-(2-chloroethyl)pyrrolidine, 1-(2-chloroethyl)piperidine or 1-(2-chloroethyl)morpholine respectively as previously reported⁷⁷ to afford the pure aldehydes in 65-97% yield. Subsequent reaction of the aldehydes **29-31** with malononitrile or the appropriate cyanoacrylate ester in the presence of piperidine or triethylamine afforded desired products **32-44** in moderate yield. The cyanoacrylate intermediate **45** was also isolated on reaction of the aldehyde **30** with ethyl cyanoacrylate and subsequently reacted with resorcinol to afford the desired product **38**.

While attempting the synthesis of the methyl esters **16, 19, 25, 32, 37, 42** using ethanol as solvent, it was noted that the final product isolated in each case was in fact the corresponding ethyl ester **17, 20, 21, 33, 38, 43** respectively. It appeared that in the presence of ethanol, a facile transesterification was occurring. Several different solvents were then employed in an attempt to obtain the methyl ester products, e.g. methanol, ethyl acetate, acetone, dichloromethane and tetrahydrofuran. Dichloromethane was found to be successful for the synthesis of all other methyl ester compounds (**16, 19, 25, 32, 37** and **42**), however yields were low. Details of the reaction conditions for the synthesis of compounds **16-25** and **32-44** are presented in Table 1. Attempts to improve the yield using potassium carbonate or cetyltrimethylammonium chloride in aqueous conditions were not successful^{75,76}. The most efficient method of preparation for the methyl ester products was by replacing the initially used triethylamine with piperidine as the base of choice and by performing the reactions under microwave conditions.

The stability of representative examples of the target compounds **38** and **40** was determined in phosphate buffer and the half life was determined to be greater than 24 h for each

compound at pH values 7.4 and 9. The compounds **38** and **40** degraded at lower pH = 4, with 27% and 32% remaining after 30 min respectively. The compounds showed very high stability in human blood plasma, and the half-life was determined to be greater than 24 h.

The molecular structure of a representative example of the 4-aryl-4*H*-chromenes series, compound **22** was determined by single crystal X-ray crystallography (*S* enantiomer shown). The ORTEP diagram is displayed in Figure 4 (thermal ellipsoids at 50% probability). It can be seen that the benzopyran ring is planar, and the aryl ring at C-4 positioned at a dihedral angle of 78.403(77) ° with respect to the benzopyran ring (Figure 4).

Figure 4. ORTEP representation of the X-Ray crystal structure of **22**, ellipsoids at 50% probability

Scheme 1. Synthesis of 4-aryl-4*H*-chromenes **16-25**

Scheme 2. Synthesis of 4-aryl-4*H*-chromenes **32-44**

2.2 Biological results and discussion


All compounds were initially evaluated for antiproliferative effects in MCF-7 cells using the MTT assay (72 h) and the results are displayed in Table 1. The most potent compound was identified as compound **41** (which contains piperidine aryl ether at C-4 and a *tert*-butyl ester at C-3 of the chromene ring) which showed comparative results to that of tamoxifen in the antiproliferative assay with IC₅₀ value of 2.65 μM (IC₅₀ value determined for control drug tamoxifen = 4.12 μM). Compounds **16**, **21**, **35** and **38** also demonstrated low micromolar antiproliferative effects with IC₅₀ values of 10.82, 4.50, 7.73 and 8.75 μM respectively. Selected compounds were concurrently tested to assess the extent of their cytotoxicity using a LDH assay. Compounds displayed negligible cytotoxicity in LDH assay with 0% cell death observed for compounds **32**, **33**, **37**, **38**, **43** and **44**.

Selected compounds from the series **16-23** and **32-44** were then screened for their binding affinity to ER α and ER β in a fluorescence polarisation assay, with tamoxifen as a positive control. The results are displayed in Table 1. For compounds **16** and **17** tested, the methyl and ethyl ester compounds showed similar affinity to each other for ER α and ER β . The cyano compound **18** did not show any binding affinity for ER α at concentrations up to 10 μ M. Compound **19** with the phenolic substituent at C-4 showed a slightly greater affinity for ER α than the benzyloxy ether **21**, however it showed markedly less affinity for ER β . The ethyl ether compounds **22** and the 3,4,dimethoxyether **23** having little antiproliferative activity, were not further evaluated.








On addition of the basic ether substituent to the core phenol **20**, whether pyrrolidine, piperidine or morpholine, greater ER α binding ability was observed. For the pyrrolidine containing compounds, the methyl (**32**), *n*-butyl (**35**) and *tert*-butyl (**36**) esters all displayed similar affinity to ER α with IC₅₀ binding of less than 100 nM each (e.g. 20 nM, 20 nM and 10 nM respectively). The ethyl ester (**33**) demonstrated slightly less affinity with binding of 1.41 μ M. The C-3 cyano compound **34** as observed with the unsubstituted compounds had decreased affinity for ER α (IC₅₀ = 0.89 μ M), and little affinity for ER β (IC₅₀ > 10 μ M). All compounds showed selectivity for ER α over ER β with significant difference noted in binding affinity, this was consistent across all compounds tested. For the compounds with the piperidine containing side chain, the methyl (**37**), ethyl (**38**) and *tert*-butyl (**41**) esters all demonstrated good affinity to ER α with ethyl ester compound **38** showing greatest activity with IC₅₀ values of 20 nM and 60 nM for ER α and ER β respectively. The *tert*-butyl ester (**41**) which was the most active in the antiproliferative assay, also demonstrated good affinity for ER α (IC₅₀ = 0.090 μ M) and also sub-micromolar binding to ER β (IC₅₀ = 370 μ M). Once again the cyano compound **39** was the least effective in the series in the ER α and ER β binding assays, with IC₅₀ values of 1.47 and 10.42 μ M respectively. Of the compounds

examined containing the morpholine side-chain (**42-44**), the ethyl ester **43** showed the greatest ER α binding activity (IC₅₀ = 220 nM), while the methyl ester **42** was the most effective as a ligand for ER β (IC₅₀ = 0.63 μ M).

Table 1. ER α and ER β binding effects and antiproliferative activity in MCF-7 cells of selected 4-aryl-4*H*-chromenes.



Compound	R ₁	R ₂	ER α IC ₅₀ (μ M) ^d	ER β IC ₅₀ (μ M) ^d	β/α	Antiproliferative activity MCF-7 cells IC ₅₀ (μ M) ^a
16	H	-CO ₂ CH ₃	3.30	1.59	0.48	10.82
17	H	-CO ₂ CH ₂ CH ₃	2.36	0.61	0.26	101.17
18	H	-CN	>10	8.39	0.84	182.05
19	OH	-CO ₂ CH ₃	0.79	10.02	12.67	156.80
20	OH	-CO ₂ CH ₂ CH ₃	nd	nd	-	>100
21	OCH ₂ C ₆ H ₅	-CO ₂ CH ₂ CH ₃	40.32	>100	-	4.5
22	OCH ₂ CH ₃	-CO ₂ CH ₂ CH ₃	nd	nd	-	>100
23	3,4-(OCH ₃) ₂	-CO ₂ CH ₂ CH ₃	nd	nd	-	>100
32		-CO ₂ CH ₃	0.02	0.35	14.31	19.60
33		CO ₂ CH ₂ CH ₃	1.41	3.10	2.21	27.29
34		-CN	0.89	>10	11.23	>50
35		-CO ₂ CH ₂ CH ₂ CH ₂ CH ₃	0.02	6.55	353.34	7.73
36		-CO ₂ C(CH ₃) ₃	0.01	0.67	49.49	10.03
37		-CO ₂ CH ₃	0.05	0.22	4.77	16.2

38		-CO ₂ CH ₂ CH ₃	0.02	0.06	2.56	8.75
39		-CN	1.47	10.42	7.10	23.21
40		-CO ₂ CH ₂ CH ₂ CH ₂ CH ₃	0.03	2.35	87.29	31.82
41		-CO ₂ C(CH ₃) ₃	0.09	0.37	3.93	2.65
42		-CO ₂ CH ₃	109.10	0.63	0.005	103.22
43		-CO ₂ CH ₂ CH ₃	0.22	2.44	11.02	18.79
44		-CN	13.97	>50	3.58	35.31
Tamoxifen^{b, c}	-	-	0.070	0.170	2.43	4.12

^aIC₅₀ values are half maximal inhibitory concentrations required to block the growth stimulation of MCF-7 cells. Values represent the mean for at least two experiments performed in triplicate. ^bThe IC₅₀ value obtained for Tamoxifen is 4.12 μM is in good agreement with the reported IC₅₀ value for tamoxifen using the MTT assay in human MCF-7 cells ⁷⁸. ^cThe ER binding values obtained are in agreement with the reported ER IC₅₀ binding data for tamoxifen (ERα 60.9 nM, ERβ 188 nM, Panvera/Invitrogen); ^dValues are an average of at least nine replicate experiments, for ERα with typical standard errors below 15%, and six replicate experiments for ERβ, with typical standard errors below 15%

2.3 Computational

To rationalise the subtype selectivity observed for compounds **35** (350-fold ERα selective) and **42** (170-fold ERβ selective) we carefully selected the most appropriate X-ray structure to utilise for subsequent docking experiments, through visual analysis of the binding modes made by all co-crystallised SERMs/SERDs with their respective ER isoforms. Lasofoxifene co-crystallised with ERα (PDBID 2OUZ) was considered to be the most suitable LBP to dock compound **35**, however, the electron withdrawing oxygen of the morpholino-ring of **42** has the potential to H-bond unlike the other pyrrolidine/piperidine side-chains in the chromene series. Taking this into account, we reasoned that PDBID 1R5K whose co-crystallised ligand GW5638, uses a bridging water molecule that is perfectly positioned for substitution by the morpholino-ring oxygen of **42** to enable H-bonding with the backbone amide N-H of Leu536

(H12) would be most appropriate for docking in this case. As no suitable ER β isoforms of these structures existed, we constructed refined models of each as described in the experimental section. Briefly, the iTASSER server was used for protein structure prediction using 2OUZ and 1R5K as templates to guide the prediction and the co-ordinates of each were firstly morphed ⁷⁹ to the co-ordinates of 2OUZ and 1R5K respectively followed by refinement using 3DRefine ⁸⁰. Reconstruction of any missing residues in 2OUZ and 1R5K was also undertaken using the iTASSER server.

Conformers of all R and S enantiomers of compound **35** and **42** were docked and in the LBP our refined models of ER α and β using FRED⁸¹, and scored using Chemgauss4.

Marvin pKa prediction ⁸² for compound **35** revealed the tertiary amine of the piperidine side-chain of compound **35** is protonated at pH7.4. Interestingly, Marvin predicted that only 15% of compound **42** would have the tertiary amine of its morpholine ring protonated at pH7.4 with the remaining >80% predominantly neutral.

Only in the case of compound **35**, was there a clear binding preference to both ER subtypes for the R enantiomer (*see supplementary information*) with 1.6 and 1.9-fold differences observed in scoring for top scoring docked poses, indicating that enantiomeric separation of the R enantiomer might afford additional binding affinity and selectivity. Figure 5 highlights the key interactions made by compound **35** in both isoforms, and as is commonly observed with most Selective Estrogen Receptor Modulators (SERMs), it strongly interacts with Asp351 (303) via a salt bridge and is stabilised within the binding pocket through a series of hydrophobic interactions (e.g. with Phe404 (356), Met343 (295)). Methionine residues are relatively unique in the protein core whereby they can form hydrophobic interactions but also engage polar oxygen (e.g. carbonyl) as recently reported by Pal et al ⁸³. In the case of compound **35**, direct contact of the nucleophilic oxygen of the carbonyl group of **35** with divalent sulfur of methionine (Met343 and Met421) through a hypervalent nonbonded S---X

interaction is apparent contributing approximately 2.5 kcal/mol for this interaction ⁸⁴. This interaction is not present in ER β as the mutation of Met421 \rightarrow Ile373 allows **35** to slightly rotate in the LBP and H-bonding with Phe346, Leu339 and Arg346 occurs rather than H-bonding to Glu353 and Arg394 observed in ER α ⁸⁵. This mutation has been reported by Nilsson et al ¹¹ as being critical in achieving subtype-selective ER based therapeutics. A similar trend is also seen for the bulky *tert*-butyl moiety of **36** whereby alpha selectivity is observed for the same reasons but to a lesser degree than **35** (~50-fold). Interestingly, switching from the pyrrolidinyl (**36**) to piperidinyl basic side-chain (**41**) led to a marked reduction in ER α selectivity (>12-fold) (Table 1).

Figure 5. Top ranked poses of Compound **35** in ER α (green) and ER β (grey) indicating key hydrogen bonding interactions.

Figure 6. Side-view of top ranked poses of Compound **35** in ER α (green) and ER β (grey) indicating shift in orientation due to Leu384 \rightarrow Met336 mutation.

Compound **42** (Figure 7) is stabilised in the binding site of both ER α and ER β via hydrogen bonding interactions with Glu353 (305) and Arg394 (346) and also through additional H-bonding contacts to the morpholino oxygen of **42** (Leu536; 2.12 Å (Val487; 2.32 Å)) and the hydroxy group of Thr347 (299) which is well positioned to H-bond with the morpholino tertiary amine, irrespective of its protonation state. Interestingly, Figure 7 highlights the additional polar contact made between Asp303 and Tyr488 in ER β as a result of rotation of the carboxylate group of Asp303 which potentially would reduce the conformational entropy of the ligand bound state of **42**, translating to an increase in binding affinity.

Figure 7. Top ranked poses of Compound **42** (R Enantiomer) in ER α (green) and ER β (grey) indicating key interactions.

4. Conclusion

We have previously demonstrated the utility of the 4-aryl-4*H*-chromene scaffold as a potent modulator of ER activity. Although two 4-aryl-4*H*-chromene analogues displayed anti-proliferative activity in the ER positive MCF-7 cell line, ER α / β selectivity was modest⁶⁹. Following from these findings, the rationale that 4-aryl-4*H*-chromene analogues incorporating a typical basic side-chain could improve ER α / β selectivity was explored. Alongside this approach, modifications at the 3-position of the benzopyran ring were also investigated with a view to maximising ER isoform selectivity *via* probing of binding pocket topology. Several of these compounds possessed potent ER binding activity indicative of potential ER antagonistic effects, the most active compound (**35**) displayed 350-fold selectivity for ER α , whilst another (**42**) showed 170-fold selectivity for ER β . Our computational study suggests that compound **35** achieves its selectivity in a completely different manner to compound **42**. In the case of compound **35**, selectivity is achieved through exploitation of the differing LBP size of each isoform which forms as a result of the mutational differences (Met421 \rightarrow Ile373) and results in a hypervalent nonbonded S---X interaction with Met343 and Met421 in ER α only. Increasing the size of the carboxylate alkyl chain increases the binding affinity and selectivity for ER α . Compound **42** appears to exhibit differences in interaction potential to H12 of ER β via Asp303 (H3) H-bonding to Tyr488 (H12) which may stabilize the ligand bound state and reduce its conformational entropy in order to achieve this selectivity.

5. Experimental section

5.1 Chemistry

Uncorrected melting points were measured on a Gallenkamp apparatus. Infra-red (IR) spectra were recorded on a Perkin Elmer FT-IR Paragon 1000 spectrometer. ^1H and ^{13}C nuclear magnetic resonance (NMR) spectra were recorded at 27 °C on a Bruker DPX 400 spectrometer (400.13 MHz, ^1H ; 100.61 MHz, ^{13}C ; 376.47 MHz, ^{19}F) in either CDCl_3 (internal standard tetramethylsilane (TMS)) or CD_3OD or DMSO-d_6 . For CDCl_3 , ^1H -NMR spectra were assigned relative to the TMS peak at 0.00 δ and ^{13}C -NMR spectra were assigned relative to the middle CDCl_3 triplet at 77.00 ppm. For CD_3OD , ^1H and ^{13}C -NMR spectra were assigned relative to the center peaks of the CD_3OD multiplets at 3.30 δ and 49.00 ppm respectively. Coupling constants are reported in Hertz. For ^1H -NMR assignments, chemical shifts are reported: shift value (number of protons, description of absorption, coupling constant(s) where applicable). Low resolution mass spectra were run on a Hewlett-Packard 5973 MSD GC-MS system in an electron impact mode. Electrospray ionisation mass spectrometry (ESI-MS) was performed in the positive ion mode on a liquid chromatography time-of-flight mass spectrometer (Micromass LCT, Waters Ltd., Manchester, UK). The samples were introduced into the ion source by an LC system (Waters Alliance 2795, Waters Corporation, USA) in acetonitrile : water (60:40 %v/v) at 200 $\mu\text{L}/\text{min}$. The capillary voltage

of the mass spectrometer was at 3 kV. The sample cone (de-clustering) voltage was set at 40 V. For exact mass determination, the instrument was externally calibrated for the mass range m/z 100 to m/z 1000. A lock (reference) mass (m/z 556.2771) was used. Mass measurement accuracies of $< \pm 5$ ppm were obtained. R_f values are quoted for thin layer chromatography on silica gel Merck F-254 plates, unless otherwise stated. Flash column chromatography was carried out on Merck Kieselgel 60 (particle size 0.040-0.063 mm), Aldrich aluminium oxide, (activated, neutral, Brockmann I, 50 mesh) or Aldrich aluminium oxide, (activated, acidic, Brockmann I, 50 mesh). Chromatographic separations were also carried out on Biotage SP4 instrument. All products isolated were homogenous on TLC. Microwave experiments were carried out with in the Biotage initiator and Discover CEM microwave synthesisers.

5.1.2 General procedures for synthesis of benzopyrans.

Method A: To the appropriate aldehyde (benzaldehyde, 4-(2-pyrrolidin-1-ylethoxy)benzaldehyde or 4-(2-piperidin-1-ylethoxy)benzaldehyde) (10 mmol, 1 equiv) was added the cyanoacetate ester (10 mmol, 1 equiv) or malononitrile (10 mmol, 1 equiv) and resorcinol (10 mmol, 1 equiv). Ethanol (or dichloromethane for methyl ester) (30 mL) was added and the reaction was heated briefly to reflux. The solution was then cooled and triethylamine (0.5 mL) was added dropwise, and the reaction solution was then heated at reflux for 4 h. Solvent was removed *in vacuo* and the product was recrystallized in ethanol or methanol (for the methyl esters). **Method B:** To the appropriate aldehyde (4-hydroxybenzaldehyde, 4-benzyloxybenzaldehyde, 4-ethoxybenzaldehyde, 3,4-dimethoxybenzaldehyde, 4-(2-pyrrolidin-1-ylethoxy)benzaldehyde, 4-(2-piperidin-1-ylethoxy)benzaldehyde or 4-(2-morpholin-1-ylethoxy)benzaldehyde) (10 mmol, 1 equiv) was added the cyanoacetate ester (10 mmol, 1 equiv) or malononitrile (10 mmol, 1 equiv) and resorcinol (10 mmol, 1 equiv). Ethanol (dichloromethane for methyl ester) (30 mL) was added and the reaction was heated briefly to reflux. The solution was then cooled and

piperidine (0.5 mL) was added dropwise and the reaction solution was then heated at reflux for 4 h. The solvent was removed *in vacuo* and product was recrystallized in ethanol (or methanol for methyl esters). **Method C:** To the appropriate aldehyde (4-benzyloxybenzaldehyde, 4-(2-pyrrolidin-1-ylethoxy)benzaldehyde, 4-(2-piperidin-1-ylethoxy)benzaldehyde or 4-(2-morpholin-1-yl)ethoxybenzaldehyde, (5 mmol, 1 equiv) was added the cyanoacetate ester (5 mmol, 1 equiv) or malononitrile (5 mmol, 1 equiv) together with resorcinol (5 mmol, 1 equiv). Ethanol (10 mL), (dichloromethane for methyl esters) was added and the reaction heated in the microwave reactor for 5 min. The solution was then cooled and piperidine (0.25 mL) was added dropwise and the reaction solution was then heated in the microwave reactor for a further 30 min. The solvent was removed *in vacuo* and the product was recrystallized from ethanol (or methanol for the methyl esters).

5.1.2.1. 2-Amino-7-hydroxy-4-phenyl-4*H*-chromene-3-carboxylic acid methyl ester (16).

Preparation following method C from benzaldehyde (5 mmol, 0.51 mL), resorcinol (5 mmol, 550 mg) and methyl cyanoacetate (5 mmol, 0.50 g). Yield 5.5%, yellow powder, M.p. 231°C. IR ν_{\max} (KBr) cm^{-1} : 3417.5, 3305.5 (NH_2), 1664.7 ($\text{C}=\text{O}$) cm^{-1} . ^1H NMR (400 MHz, CDCl_3): δ 3.55 (s, 3H, O- CH_3), 4.88 (s, 1H, CH), 6.50 (d, 1H, $J=2.52$ Hz, Ar-H), 6.55 (dd, 1H, $J=2.52$ Hz, 8.4 Hz, Ar-H), 7.00 (d, 1H, $J=8.56$ Hz, Ar-H), 7.06-7.09 (m, 1H, Ar-H), 7.20-7.23 (m, 4H, Ar-H). ^{13}C NMR (100 MHz, CDCl_3): δ 39.1 (CH₃), 49.4 (O- CH_3), 78.1 (C), 101.8 (CH), 111.6 (CH), 117.3 (C), 125.3 (CH), 126.7 (CH), 127.7 (CH), 129.5 (CH), 149.2 (C), 156.3 (C), 160.8 (C), 168.7 ($\text{C}=\text{O}$), 168.8 (C- NH_2). HRMS: Found 320.0898; $\text{C}_{17}\text{H}_{15}\text{NO}_4\text{Na}$ requires 320.0899 (M^++Na).

5.1.2.2. 2-Amino-7-hydroxy-4-phenyl-4*H*-chromene-3-carboxylic acid ethyl ester (17).

Preparation following Method A from benzaldehyde (10 mmol, 1.01 mL), resorcinol (1.10 g) and ethyl cyanoacetate (10 mmol, 1.13 g). Yield 17%, white powder, M.p. 243°C. IR ν_{\max} (KBr) cm^{-1} : 3306.4, 3300.0 (NH_2), 1662.8 ($\text{C}=\text{O}$) cm^{-1} . ^1H NMR (400 MHz, CDCl_3): δ 1.11 (t, 3H, $J=7.02$ Hz, $-\text{CH}_3$), 2.06 (m, 2H, NH_2), 4.00 (q, 2H, $J=7.01$ Hz, CH_2), 4.89 (s, 1H, CH), 6.51 (d, 1H, $J=2.64$ Hz, Ar-H), 6.54 (dd, 1H, $J=2.52$ Hz, 6.00 Hz, Ar-H), 7.00 (d, 1H, $J=8.00$ Hz, Ar-H), 7.08-7.11 (m, 1H, Ar-H), 7.19-7.25 (m, 4H, Ar-H). ^{13}C NMR (100 MHz, CDCl_3): δ 13.33 (CH_3), 39.21 (CH), δ 58.18 (CH_2), 77.01 (C), 101.84 (CH), 111.59 (CH), 117.12 (C), 125.24 (CH), 126.87 (CH), 127.55 (CH), 129.58 (CH), 148.48 (C), 149.12 (C), 156.30 (C), 160.59 (C- NH_2), 168.37 (C=O). HRMS: Found 334.1042; $\text{C}_{18}\text{H}_{17}\text{NO}_4\text{Na}$ requires 334.1055 ($\text{M}^+ + \text{Na}$).

5.1.2.3. 2-Amino-7-hydroxy-4-phenyl-4H-chromene-3-carbonitrile (18).

Preparation following Method A from benzaldehyde (10 mmol, 1.01 mL), resorcinol (1.10 g, 10 mmol) and malononitrile (10 mmol, 0.66 g). Yield 56%, Yellow powder, M.p. 246°C⁷³. IR ν_{\max} (KBr) cm^{-1} : 3401.1 (OH), 3336.3, 3218.99 (NH_2), 2180.9 (CN), 1638.46, 1624.90 ($\text{C}=\text{C}$) cm^{-1} . ^1H NMR (400 MHz, CDCl_3): δ 1.54 (s(br), 2H, NH_2), 2.76 (s(br), 1H, OH), 4.68 (s, 1H, CH), 6.51 (d, 1H, $J=2.00$ Hz, Ar-H), 6.57 (dd, 1H, $J=2.34$ Hz, 8.76 Hz, Ar-H), 6.86 (d, 1H, $J=8.76$ Hz, Ar-H), 7.22-7.25 (m, 3H, Ar-H), 7.31 (d, 2H, $J=7.6$ Hz, Ar-H). ^{13}C NMR (100 MHz, CDCl_3): δ 40.21 (CH), 58.21 (C), 101.96 (CH), 111.99 (CH), 113.79 (CN), 118.99 (C), 126.26 (CH), 127.19 (CH), 128.08 (CH), 129.71 (CH), 145.88 (C), 148.89 (C), 156.83 (C-O), 159.59 (C- NH_2). HRMS: $\text{C}_{16}\text{H}_{12}\text{N}_2\text{NaO}_2$ requires 287.0796, found 287.0794 ($\text{M}^+ + \text{Na}$).

5.1.2.4. Ethyl 2-amino-7-hydroxy-4-(4-hydroxyphenyl)-4H-chromene-3-carboxylate 20.

Preparation following the general method B above from ethylcyanoacetate (10 mmol, 1.13 g), resorcinol (1.10 g, 10 mmol) and 4-hydroxybenzaldehyde (10 mmol, 1.22 g). The product

was obtained as a dark brown solid which purified by flash column chromatography over silica gel (eluent: CH₂Cl₂ – MeOH; 85:15) and then recrystallised from ethanol. Yield: 1.04 g, 32%, M.p. 228-230°C⁸⁶. IR ν_{\max} (KBr): 3614.49, 3595.36, (OH), 3485.10, 3357.31 (NH₂), 1659.71 (C=O), 1629.41 (C=C), 1630.48 (C=C), 1591.36, 1538.14 cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆) δ 0.98 (t, 3H, *J*=7.2, CH₃), 4.28 - 4.47 (m, 2H, OCH₂), 4.68 (s, 1H, CH), 6.13 - 6.23 (m, 3H, Ar-H), 6.61 - 6.68 (m, 2H, Ar-H), 6.85 - 6.99 (m, 2H, Ar-H), 8.14 (s (br), 1H, OH), 9.15 (s (br), 2H, NH₂). ¹³C NMR (101 MHz, DMSO-*d*₆): δ 14.31 (CH₃), 56.03 (CH), 58.49 (CH₂), 76.97 (C), 101.96 (CH), 102.44 (CH), 106.17 (CH), 114.75 (C), 127.81 (CH), 129.70 (CH), 132.74 (C), 152.56 (C-O), 154.93 (C-OH), 155.24 (C-OH), 158.44 (O-C-N), 160.88 (C=O). HRMS: C₁₈H₁₇NNaO₅ requires 350.1004, found 350.0997(M⁺+Na).

5.1.2.5. 2-Amino-4-(4-benzyloxyphenyl)-7-hydroxy-4*H*-chromene-3-carboxylic acid ethyl ester 21.

Preparation following the general method B above from ethyl cyanoacetate (10 mmol, 1.13 g), resorcinol (1.10 g, 10 mmol) and 4-benzyloxybenzaldehyde (10 mmol, 2.12 g). The product was obtained as yellow powder which was recrystallised from ethanol. Yield: 1.54 g, 37%. Mp 192-194°C. IR ν_{\max} (KBr): 3421.76 (OH), 3304.84, 3248.32 (NH₂), 1659.05 (C=O), 1610.67 (C=C), 1598.45 (C=C), 1503.32, 1454.01, 1311.69 cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.10 (t, 3H, *J*=7 Hz, CH₃), 2.07 (s (br), 2H, NH₂), 3.96 (q, 2H, *J*=7.04 Hz, OCH₂), 4.75 (s, 1H, CH), 5.01 (s, 2H, OCH₂), 6.43 (d, 1H, *J*=6.52 Hz, Ar-H), 6.47 (d, 1H, *J*=8.04 Hz, Ar-H), 6.94 (d, 1H, *J*=8.52 Hz, Ar-H), 6.94 (d, 2H, *J*=8.52 Hz, Ar-H), 7.23 (d, 2H, *J*=8 Hz, Ar-H), 7.31 (d, 1H, *J*=7.04 Hz, Ar-H), 7.34 - 7.39 (m, 2H, Ar-H), 7.42 (d, 1H, *J*=6.76 Hz, Ar-H), OH not observed. ¹³C NMR (100 MHz, DMSO-*d*₆): δ 14.78 (CH₃), 38.58 (CH), 58.19 (CH₂), 69.58 (CH₂), 77.56 (C), 102.46 (CH), 112.53 (CH), 114.80 (C), 117.53 (CH), 128.08 (CH), 128.35 (CH), 128.85 (CH), 130.24 (CH), 137.68 (C), 141.58 (C), 149.51

(C-O), 156.89 (C-OH), 157.09 (C-O), 161.39 (O-C-N), 168.87 (C=O). HRMS: $C_{25}H_{23}NNaO_5$ requires 440.1474, found 440.1463 ($M^+ + Na$).

5.1.2.6. 2-Amino-4-(4-ethoxyphenyl)-7-hydroxy-4*H*-chromene-3-carboxylic acid ethyl ester 22.

Preparation following the general method B above from ethyl cyanoacetate (10 mmol, 1.13 g), resorcinol (1.10 g, 10 mmol) and 4-ethoxybenzaldehyde (10 mmol, 1.50 g). The product was obtained as yellow crystals following recrystallisation from ethanol, yield 0.78 g, 22%. M.p. 258-263°C. IR ν_{max} (KBr): 3563.54 (OH), 3431.59, 3419.15 (NH₂), 1660.91 (C=O), 1634.86 (C=C), 1613.89 (C=C), 1583.43, 1568.42, 1458.83 cm^{-1} . ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.05 (t, 3H, $J=7.04$ Hz, CH₃), 1.25 (t, 3H, $J=7.02$ Hz, CH₃), 3.93 - 4.18 (m, 4H, 2xOCH₂), 4.72 (s, 1H, CH), 6.42 - 6.50 (m, 2H, Ar-H), 6.73 (d, 2H, $J=9.04$ Hz, Ar-H), 6.94 (d, 2H, $J=8.04$ Hz, Ar-H), 7.01 (d, 1H, $J=8.12$ Hz, Ar-H), OH not observed, NH₂ not observed. ¹³C NMR (100 MHz, DMSO-*d*₆): δ 14.31 (CH₃), 14.68 (CH₃), 38.09 (CH), 58.51 (CH₂), 62.79 (CH₂), 76.81 (C), 101.97 (CH), 112.04 (CH), 113.91 (CH), 117.13 (C), 127.84 (CH), 129.76 (CH), 140.71 (C), 149.02 (C-O), 156.53 (C-OH), 160.91 (O-C-N), 168.40 (C=O). HRMS: $C_{20}H_{22}NO_5$ requires 356.1498, found 356.1488($M^+ + H$).

5.1.2.7. Ethyl 2-amino-7-hydroxy-4-(3,4-dimethoxyphenyl)-4*H*-chromene-3-carboxylate 23

(*E*)-Ethyl 2-cyano-3-(3,4-dimethoxyphenyl)acrylate **28** (10 mmol, 2.62 g) and resorcinol (1.10 g, 10 mmol) were dissolved in ethanol and the reaction mixture was briefly heated to reflux. Piperidine (0.5 mL) was added dropwise and the reaction solution was heated at reflux for 4 h. following the general method B above. The solvent was removed *in vacuo* and product was recrystallized from ethanol. Yield: 0.07 g, 2%, yellow crystals, M.p. 284-288°C ⁷¹. IR ν_{max} (KBr): 3488.34 (OH), 3431.11, 3412.48 (NH₂), 1681.40 (C=C), 1642.76 (C=C), 1629.45 (C=O), 1562.75 cm^{-1} . ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.09 (t, 3H, $J=7.04$ Hz,

CH₃), 3.66 (s, 3H, OCH₃), 3.71 (s, 3H, OCH₃), 3.95 (q, 2H, $J=7.04$ Hz, OCH₂), 4.73 (s, 1H, CH), 6.42 (s, 1H, Ar-H), 6.46 (dd, 1H, $J=2.48$, $J=7.02$ Hz, Ar-H), 6.55 (d, 1H, $J=9.04$ Hz, Ar-H), 6.77 (d, 2H, $J=9.04$ Hz, Ar-H), 6.99 (d, 1H, $J=8.52$ Hz, Ar-H), 7.55 (s (br), 2H, NH₂), 9.57 (s (br), 1H, OH). ¹³C NMR (100 MHz, DMSO-d₆): δ 14.36 (CH₃), 38.45 (CH), 55.40 (CH₃), 58.53 (CH₂), 76.72 (C), 101.97 (CH), 110.23 (CH), 111.15 (CH), 111.86 (C), 117.04 (CH), 118.64 (CH), 129.73 (CH), 141.44 (C), 146.83 (C-OCH₃), 148.19 (C-OCH₃), 149.03 (C-O), 156.60 (C-O), 160.97 (O-C-N), 168.43 (C=O). HRMS: C₂₀H₂₁NNaO₆ requires 394.1267, found 394.1262(M⁺+Na).

5.1.2.8. 2-Amino-4-(4-benzyloxyphenyl)-7-hydroxy-4H-chromene-3-carbonitrile (24).

Preparation following the general method B above from malononitrile (0.66 g, 10 mmol) and resorcinol (1.1 g, 10 mmol) and 4-benzyloxybenzaldehyde (2.12 g, 10 mmol). The product was obtained as a pink solid, 2.08 g (56%) following recrystallisation from ethanol. M.p. 264-266 °C. IR ν_{\max} : 3258.49 (OH), 3032.13 (CH), 2191.16 (CN), 1637.44 (C=C) cm⁻¹. ¹H NMR (400 MHz, DMSO-d₆) δ 9.71 (s, 1H, OH), 7.24 - 7.51 (m, 5H, ArH), 7.11 (d, $J=8.54$ Hz, 2H, ArH), 6.96 (d, $J=8.54$ Hz, 2H, ArH), 6.75 - 6.89 (m, 3H, ArH), 6.41 - 6.60 (m, 2H, ArH), 5.05 (s, 2H, CH₂), 4.59 (s, 1H, CH). ¹³C NMR (101 MHz, DMSO-d₆) δ 160.2 (O-C-N), 157.2 (C-O), 157.1 (C-O), 148.8 (C-O), 138.8 (C), 137.1 (C), 130.0 (CH), 128.5 (CH), 128.4 (CH), 127.8 (CH), 127.7 (CH), 120.8 (CN), 114.8 (CH), 114.1 (CH), 112.4 (C), 102.2 (CH), 69.3 (CH₂), 56.7 (C), 39.3 (CH). HRMS: C₂₃H₁₇N₂O₃ requires 369.1239; found 369.1232 [M-H]⁺.

5.1.2.9. 2-Amino-4-(4-benzyloxyphenyl)-7-hydroxy-4H-chromene-3-carboxylic acid methyl ester (25).

Preparation as above using Method C from 4-benzyloxybenzaldehyde (10 mmol, 2.12 g), resorcinol (1.10 g, 10 mmol) and methyl cyanoacetate, (10 mmol, 0.99 g). Yield 46%, cream

powder, M.p. 216 - 218°C. IR ν_{\max} (KBr) cm^{-1} : 3415.4, 3302.9 (NH_2), 1660.7 ($\text{C}=\text{O}$) cm^{-1} . ^1H NMR (400 MHz, Acetone- d_6): δ 3.57 (s, 3H, O- CH_3), 4.84 (s, 1H, CH), 5.04 (s, 2H, CH_2), 6.50 (d, 1H, $J=2.04$ Hz, Ar-H), 6.55 (dd, 1H, $J=2.26$ Hz, 6.04 Hz, Ar-H), 6.85 (d, 2H, $J=8.52$ Hz, Ar-H), 7.00 (d, 1H, $J=8.56$ Hz, Ar-H), 7.14 (d, 2H, $J=8.56$ Hz, Ar-H), 7.31 (d, 1H, $J=7.04$ Hz, Ar-H), 7.37 (t, 2H, $J=7.78$ Hz, Ar-H), 7.44 (d, 2H, $J=7.88$ Hz, Ar-H). ^{13}C NMR (100 MHz, Acetone- d_6): δ 38.2 (CH), 49.4 (O- CH_3), 68.9 (O- CH_2), 77.4 (C), 101.8 (CH), 111.6 (CH), 113.9 (CH), 117.6 (C), 127.1 (CH), 127.6 (CH), 127.9 (CH), 129.4 (CH), 137.2 (C), 140.8 (C), 149.2 (C), 156.2 (C), 160.6 (C-O), 160.7 (C- NH_2), 168.7 (C=O). HRMS: Found 426.1297; $\text{C}_{24}\text{H}_{21}\text{NO}_5\text{Na}$ requires 426.1317($\text{M}^+ + \text{Na}$).

5.1.3. 2-Amino-7-hydroxy-4-(4-hydroxyphenyl)-4*H*-chromene-3-carboxylic acid methyl ester (19).

2-Amino-4-(4-benzyloxyphenyl)-7-hydroxy-4*H*-chromene-3-carboxylic acid methyl ester (25) (1.6 mmol, 0.77 g) was dissolved in ethyl acetate (20 mL) and Pd/C (10%, 0.8 g) was added. The mixture was stirred under an atmosphere of hydrogen for 12 h until reaction was complete as monitored by TLC. The catalyst was removed by filtrations and the residue was purified by chromatography to afford the product as a waxy solid, IR: KBr ν_{\max} : 3415.4, 3302.9 (NH_2), 1660.7 ($\text{C}=\text{O}$) cm^{-1} . ^1H NMR (400 MHz, CDCl_3): δ 3.57 (s, 3H, O- CH_3), 4.79 (s, 1H, CH), 6.49 (d, 1H, $J=2.00$ Hz, Ar-H), 6.56 (dd, 1H, $J=2.02$ Hz, 6.00 Hz, Ar-H), 6.69 (d, 2H, $J=8.52$ Hz, Ar-H), 6.99-7.04 (m, 3H, Ar-H). ^{13}C NMR (100 MHz, CDCl_3): δ 38.2 (CH), 49.4 (O- CH_3), 68.9 (O- CH_2), 77.4 (C), 101.8 (CH), 111.6 (CH), 113.9 (CH), 117.6 (C), 127.1 (CH), 127.6 (CH), 127.9 (CH), 129.4 (CH), 137.2 (C), 140.8 (C), 149.2 (C), 156.2 (C), 160.6 (C-O), 160.7 (C- NH_2), 168.7 (C=O). Found 336.0852; $\text{C}_{17}\text{H}_{15}\text{NO}_5\text{Na}$ requires 336.0848($\text{M}^+ + \text{Na}$).

5.1.4. General procedure for synthesis of cyanoacrylate esters 26-28.

A solution of methyl or ethyl cyanoacetate (10 mmol) and the appropriate arylaldehyde (10 mmol) in methanol 30 mL was heated briefly to reflux. The solution was cooled and piperidine (0.5 mL) was added dropwise and the reaction solution was then heated at reflux for 4 h. The solution was cooled and the solvent was removed *in vacuo* and product was recrystallized from ethanol.

5.1.4.1. (E)-Methyl 2-cyano-3-(4-hydroxyphenyl)acrylate 26.

A mixture of methyl cyanoacetate (10 mmol, 1.13 g) and 4-hydroxybenzaldehyde (10 mmol, 1.22 g) was melted under vacuum in a Kugelrohr short-path vacuum distillation apparatus at 170 °C for 40 min. The product was then dried at 80 °C and then recrystallized from ethanol as beige coloured crystals. Yield: 1.802 g, 90%, Mp: 208-210 °C⁶⁹. IR ν_{\max} (KBr): 3415.50 (OH), 2224.80 (CN), 1725.19 (C=O), 1589.33 (C=C) cm^{-1} . ¹H NMR (400 MHz, DMSO-*d*₆): 3.78 (s, 3H, OCH₃), 6.91 (d, *J*=8.54 Hz, 2H), 7.94 (d, *J*=9.16 Hz, 2H), 8.18 (s, 1H, CH=C). ¹³C NMR (100 MHz, CDCl₃): δ 51.99 (OCH₃), 97.62 (C), 115.53 (CN), 115.88 (CH), 123.06 (C), 133.50 (CH), 153.89 (CH), 162.08 (C-OH), 162.76 (C=O).

5.1.4.2. (E)-Ethyl 2-cyano-3-(4-benzyloxyphenyl)acrylate 27.

Following the procedure outlined above, a solution of ethyl cyanoacetate (10 mmol, 1.13 g) and 4-benzyloxybenzaldehyde (10 mmol, 2.12 g) in benzene (20 mL) was heated briefly to reflux and then cooled. Piperidine (0.99 mL) was added dropwise and the reaction solution was then heated at reflux for 4 h. The solution was washed with HCl (10 %) and then extracted with dichloromethane. The solvent was removed *in vacuo* and the product was recrystallized from ethanol as cream coloured crystals. Yield: 2.53 g, 83%, M.p. 99-100 °C⁸⁷. IR ν_{\max} (KBr): 2221.96 (CN), 1714.07 (C=O), 1688.82 (C=C), 1563.74, 1473.48 cm^{-1} . ¹H NMR (400 MHz, CDCl₃): 1.40 (t, *J*=7.02 Hz, 3 H, CH₃), 4.38 (q, *J*=7.12 Hz, 2 H, CH₂), 5.16 (s, 2 H, CH₂), 7.07 (d, *J*=9.16 Hz, 2 H, ArH), 7.32 - 7.47 (m, 5 H, ArH), 8.01 (d, *J*=8.54 Hz, 2 H, ArH), 8.18 (s, 1 H, CH=C).

5.1.4.3. (E)-Ethyl 2-cyano-3-(3,4-dimethoxyphenyl)acrylate 28.

Following the procedure outlined above, a solution of ethyl cyanoacetate (10 mmol, 1.13 g) and 3,4-dimethoxybenzaldehyde (10 mmol, 1.66 g) in ethanol 30 mL was heated briefly to reflux and then cooled. Piperidine (0.5 mL) was then added dropwise and the reaction solution heated at reflux for 4 h. The solvent was removed *in vacuo* and product was recrystallized from ethanol to afford the product as yellow crystals. Yield: 0.34 g, 13%, M.p. 160 °C⁴¹. IR ν_{\max} (KBr): 2244.68 (CN), 1673.71 (C=C), 1643.64 (C=O), 1563.74, 1473.48 cm^{-1} . ¹H NMR (400 MHz, DMSO-*d*₆): δ 0.94 (t, 3H, *J*=7.14 Hz, CH₃), 3.72 (s, 6H, 2xOCH₃), 3.86 (q, 2H, *J*=7.04, CH₂), 6.38 (d, 1H, *J*=2.28, Ar-H), 6.45 – 6.51 (m, 1H, Ar-H), 7.18 – 7.27 (m, 1H, Ar-H), 8.46 (d, 1H, *J*=6.98, CH). ¹³C NMR (100 MHz, DMSO-*d*₆): δ 14.33 (CH₃), 58.85 (OCH₃), 62.03 (CH₂), 78.01 (C), 106.63 (CH), 112.53 (CN), 120.75 (CH), 128.77 (C), 148.70 (C-OCH₃), 149.77 (C-OCH₃), 157.73 (CH), 168.72 (C=O). HRMS: C₁₄H₁₆NO₄ requires 262.1079, found 262.1086(M⁺+H).

5.1.5. General procedure for preparation of 29-31.

4-Hydroxybenzaldehyde (16 mmol (1 equiv)) was dissolved in ethyl acetate (20 mL). To this solution was added anhydrous potassium carbonate (18 mmol, 2.50 g) (1.125 equiv) and *tetra-n*-butylammonium bromide (0.054 g, 0.2 mmol). The mixture was then heated at reflux for 10 min and the appropriate basic ether was then added (32 mmol, 2 equiv). [The basic ethers 1-(2-chloroethyl)pyrrolidine, 1-(2-chloroethyl)piperidine and 1-(2-chloroethyl)morpholine were obtained as their HCl salt and were washed with water (45 mL) and sodium hydroxide (10.5 g) in toluene (6 mL) to extract the free base for use in the reaction]. The reaction mixture was heated at reflux until reaction was complete as monitored by thin layer chromatography. On completion, the solution was filtered and the solvent was removed under reduced pressure to afford product, which was used in the following reactions without further purification.

5.1.5.1. 4-(2-Pyrrolidin-1-ylethoxy)benzaldehyde (29).

The preparation was according to the general procedure above. The product was isolated as orange oil⁷⁷ yield 65%. ¹H NMR (400 MHz, CDCl₃): δ 1.59 (s, 4H, (CH₂)₂), 2.41 (s, 4H, CH₂-N-CH₂), 2.70 (2H, t, *J*=5.78 Hz, -CH₂), 3.96 (t, *J*=5.78 Hz, 2H, O-CH₂), 6.79 (d, 2H, *J*=8.56 Hz, Ar-H), 7.58 (d, 2H, *J*=8.52 Hz, Ar-H), 9.64 (s, 1H, CHO).

5.1.5.2 4-(2-Piperidin-1-ylethoxy)benzaldehyde (30).

The preparation was according to the general procedure above (32 mmol scale). The product was isolated as orange coloured oil, (5.42 g, 73%)⁷⁷. ¹H NMR (400 MHz, CDCl₃): δ 1.53-1.55 (m, 6H, 3xCH₂), 2.51-2.66 (m, 6H, CH₂), 4.29-4.36 (m, 2H, CH₂), 7.13 (d, 2H, *J*=8.0 Hz, Ar-H), 7.85 (d, 2H, *J*=8.0 Hz, Ar-H), 9.89 (s, 1H, CHO).

5.1.5.3. 4-(2-Morpholin-1-ylethoxy)benzaldehyde (31).

The preparation was according to the general procedure above and the product was isolated as an orange coloured oil (97%)⁷⁷. ¹H NMR (400 MHz, CDCl₃): δ 2.60 (m, 4H, 2xCH₂), 2.68 (s, 2H, CH₂), 3.56 (m, 4H, 2xCH₂), 4.05 (m, 2H, 2xO-CH₂), 6.86 (m, 2H, Ar-H), 7.67 (m, 2H, Ar-H), 9.71 (s, 1H, CHO).

5.1.6.1. 2-Amino-7-hydroxy-4-[4-(2-pyrrolidin-1-ylethoxy)phenyl]-4*H*-chromene-3-carboxylic methyl ester (32).

Preparation was as described above from 4-(2-pyrrolidin-1-ylethoxy)benzaldehyde (29) (10 mmol, 2.19 g), resorcinol (1.10 g, 10 mmol) and methyl cyanoacetate (10 mmol, 0.99 g), using general Method C. Yield 10%, M.p. 167°C, yellow crystals. IR ν_{\max} (KBr) cm⁻¹: 3413.53, 3296.89 (NH₂), 1701.9(C=O), 1656.4, 1610.6(C=C) cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 1.74-1.77 (m, 4H, C-(CH₂)₂-C), 2.64-2.67 (m, 4H, CH₂-N-CH₂), 2.87-2.90 (t, 2H, *J*=5.88 Hz, CH₂-N), 3.57 (s, 3H, O-CH₃), 4.05-4.08 (t, 2H, *J*=5.90 Hz, CH₂-N), 4.83 (s, 1H, CH), 6.50-6.51 (d, 2H, *J*=2.52 Hz, Ar-H), 6.54-6.57 (dd, 1H, *J*=2.36 Hz, 6.04 Hz, Ar-H),

6.77-6.79 (d, 2H, $J=8.76$ Hz, Ar-H), 6.99-7.02 (m, 1H, Ar-H), 7.11-7.13 (d, 2H, $J=8.52$ Hz, Ar-H). ^{13}C NMR (151 MHz, $\text{DMSO-}d_6$) δ 161.5 (Ar-C), 156.9 (Ar-C), 149.4 (Ar-C), 141.0 (Ar-C), 128.0 (Ar-CH), 117.6 (Ar-CH), 114.4 (Ar-CH), 112.4 (Ar-CH), 102.4 (Ar-CH), 77.0 (-C-), 66.9 (NCH₂), 54.7 (OCH₃), 54.3 (NCH₂), 40.37 (CH), 38.3 (CH₂), 23.4 (CH₂). HRMS: Found 411.1927; C₂₃H₂₇N₂O₅ requires 411.1920(M⁺+H).

5.1.6.2. 2-Amino-7-hydroxy-4-[4-(2-pyrrolidin-1-ylethoxy)phenyl]-4H-chromeme-3-carboxylic acid ethyl ester (33).

Preparation was as described above from 4-(2-pyrrolidin-1-ylethoxy)benzaldehyde (**29**) (10 mmol, 2.19 g), resorcinol (1.10 g, 10 mmol) and ethyl cyanoacetate (10 mmol, 1.13 g) using Method B. Purification was achieved by flash column chromatography over silica gel (eluent: dichloromethane:methanol; 9:1) to afford the product as a yellow gel, yield 10%. IR ν_{max} (film) 1671.15 (C=O) cm⁻¹. ^1H NMR (400 MHz, CDCl₃): δ 1.15 (t, 3H, $J=7.28$ Hz, -CH₃), 1.76-1.90 (m, 4H, C-(CH₂)₂-C), 2.03-2.07 (m, 4H, CH₂-N-CH₂), 2.93-2.94 (t, 2H, $J=5.78$ Hz, CH₂-N), 4.01-4.06 (q, 2H, $J=7.02$ Hz, CH₂-N), 4.22-4.24 (t, 2H, $J=4.76$ Hz, O-CH₂), 4.80 (d, 1H, CH), 5.12 (bs, 1H, OH), 6.47-6.49 (d, 2H, $J=3$ Hz, Ar-H), 6.85-6.90 (dd, 3H, $J=5.04$ Hz, 8.78 Hz, Ar-H), 7.14 (d, 2H, $J=9.04$ Hz, Ar-H). ^{13}C NMR (100 MHz, CDCl₃): δ 13.09 (CH₃), 22.14 (CH₂), 38.40 (CH), 53.32 (CH₂), 53.84 (CH₂), 58.53 (CH₂), 62.56 (CH₂), 77.09 (C), 101.73 (Ar-CH), 111.44 (Ar-CH), 113.62 (Ar-CH), 116.96 (ArC), 127.76 (Ar-CH), 129.22 (Ar-C), 131.38 (Ar-C), 129.88 (Ar-CH), 138.48 (Ar-C), 148.92 (Ar-C), 155.48 (Ar-C-O), 156.12 (Ar-C-O), 162.50 (Ar-C-N), 169.209 (C=O). HRMS: Found 425.2081; C₂₄H₂₉N₂O₅ requires 425.2076(M⁺+H).

5.1.6.3. 2-Amino-7-hydroxy-4-[4-(2-pyrrolidin-1-ylethoxy)phenyl]-4H-chromeme-3-carbonitrile (34).

Preparation following General Method B from 4-(2-pyrrolidin-1-ylethoxy)benzaldehyde (**29**) (1.3 mmol, 0.285 g), resorcinol (143 mg, 1.30 mmol) and malononitrile (1.3 mmol, 85.9 mg).

Yield 20.4%, M.p. 181°C, cream powder. IR ν_{\max} (KBr) 3439.6, 3360.1 (NH₂), 2185.1 (CN) cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 1.74 (s, 4H, C-(CH₂)₂-C), 2.64 (t, 4H, $J=5.10$ Hz, CH₂-N-CH₂), 2.80 (t, 2H, $J=6.02$ Hz, CH₂-N), 4.01 (t, 2H, $J=6.04$ Hz, CH₂-N), 4.55 (s, 1H, CH), 6.10 (bs, 1H, OH), 6.38 (d, 1H, $J=2.36$ Hz, Ar-H), 6.46 (dd, 1H, $J=2.56$ Hz, Ar-H), 6.78 (d, 1H, $J=8.48$ Hz, Ar-H), 6.85 (d, 2H, $J=4.96$ Hz, Ar-H), 7.13-7.15 (d, 2H, $J=8.52$ Hz). ¹³C NMR (100 MHz, CDCl₃): δ 23.12 (CH₂), 26.45 (CH₂), 38.15 (CH), 46.66 (CH₂), 53.97 (CH₂), 54.35 (CH₂), 56.52 (C), 66.63 (CH₂), 102.09 (Ar-CH), 106.15 (CH), 112.35 (CH), 113.94 (C), 114.40 (CH), 120.75 (CH), 128.42 (CH), 129.88 (CH), 138.48 (C), 148.76 (C), 157.13 (C), 157.22 (C), 158.49 (C-O), 160.09 (C-N). HRMS: Found 378.1801; C₂₂H₂₄N₃O₃ requires 378.1818(M⁺+H).

5.1.6.4. 2-Amino-7-hydroxy-4-[4-(2-pyrrolidin-1-ylethoxy)phenyl]-4H-chromene-3-carboxylic *n*-butyl ester (35).

Preparation following the general Method A from 4-(2-pyrrolidine-1-yl-ethoxy)-benzaldehyde (**29**) (2.5 mmol, 0.547 g), resorcinol (2.5 mmol, 280 mg), and *n*-butyl cyanoacetate (2.5 mmol, 353 mg). Yield 14.2%, yellow powder. IR ν_{\max} (KBr) cm⁻¹. 3407.44, 3294.68 (NH₂), 1673.91 (C=O) cm⁻¹. ¹H NMR (400 MHz, DMSO-d₆): δ 0.78 (t, 1H, $J=7.54$ Hz, CH₂), 1.06 (t, 3H, $J=3.52$ Hz, CH₃), 1.40 (m, 1H, CH₂), 1.66 (bs, 4H, (CH₂)₂), 2.51 (d, 6H, $J=11.04$ Hz, (CH₂)₃), 2.73 (t, 2H, $J=5.76$ Hz, N-CH₂), 3.71 (m, 4H, 2 x O-CH₂), 4.73 (s, 1H, CH), 6.42 (d, 1H, $J=2.17$ Hz, Ar-H), 6.48 (dd, 1H, $J=2.24$ Hz, 6.04 Hz, Ar-H), 6.78 (d, 2H, $J=8.52$ Hz, Ar-H), 6.99 (d, 1H, $J=8.28$ Hz, Ar-H), 7.03 (m, 2H, Ar-H). ¹³C NMR (100 MHz, DMSO-d₆): δ 14.32 (CH₃), 18.56 (CH₂), 23.10 (CH₂), 30.50 (CH₂), 38.10 (CH), δ 53.95 (CH₃), 54.30 (CH₂), 54.32 (CH₂), 58.52 (CH₂), 62.20 (CH₂), 66.50 (CH₂), 76.81 (C), 101.97 (CH), 112.04 (CH), 113.98 (CH), 117.05 (C), 127.74 (CH), 127.86 (CH), 129.75 (CH), 140.83 (C), 148.85 (CH), 149.02 (C), 156.51 (C), 156.61 (C), 160.91 (C), 160.99 (C), 168.49 (C=O). HRMS: Found 453.2393; C₂₆H₃₃N₂O₅ requires 453.2389(M⁺+H).

5.1.6.5. 2-Amino-7-hydroxy-4-[4-(2-pyrrolidine-1-ylethoxy)phenyl]-4*H*-chromene-3-carboxylic tert-butyl ester (36).

Preparation following general method A above from 4-(2-pyrrolidine-1-ylethoxy)benzaldehyde (**29**) (2.5 mmol, 0.547 g), resorcinol (2.5 mmol, 280 mg) and *tert*-butyl cyanoacetate (2.5 mmol, 353 mg). Yield 63%, white powder. IR ν_{\max} (KBr) cm^{-1} : 3407.03, 3291.65 (NH_2), 1672.28 ($\text{C}=\text{O}$) cm^{-1} . ^1H NMR (400 MHz, DMSO-d_6): δ 1.36 (s, 9H, $(\text{CH}_3)_3$), 1.78 (s, 4H, $(\text{CH}_2)_2$), 2.47 (bs, 4H, $(\text{CH}_2)_2$), 2.72 (t, 2H, $J=6.04$ Hz, CH_2), 3.97 (t, 2H, $J=5.78$ Hz, CH_2), 4.65 (s, 1H, CH), 6.39 (d, 1H, $J=2.48$ Hz, Ar-H), 6.44 (dd, 1H, $J=2.26$ Hz, 8.52 Hz, Ar-H), 6.77 (d, 2H, $J=8.52$ Hz, Ar-H), 6.93 (d, 1H, $J=8.56$ Hz, Ar-H), 7.01 (d, 2H, $J=8.52$ Hz, Ar-H). 7.48 (s, 2H, NH_2), 9.57 (s, 1H, OH). ^{13}C NMR (100 MHz, DMSO-d_6): δ 23.11 (CH_2), 28.10 (CH_3), 38.58 (CH), 53.99 (CH_2), 54.34 (CH_2), 56.03 (CH_2), 66.59 (CH_2), 77.86 (C), 78.01 (C), 101.90 (CH), 111.92 (CH), 113.90 (CH), 116.93 (C), 127.88 (CH), 129.82 (CH), 141.19 (C), 148.85 (C), 156.46 (C), 156.52 (C), 160.81 (C), 168.50 ($\text{C}=\text{O}$). HRMS: Found 453.2393; $\text{C}_{26}\text{H}_{33}\text{N}_2\text{O}_5$ requires 453.2389($\text{M}^+\text{+H}$).

5.1.6.6. 2-Amino-7-hydroxy-4-[4-(2-piperidin-1-ylethoxy)phenyl]-4*H*-chromene-3-carboxylic methyl ester (37).

Preparation following the general method C above from 4-(2-piperidine-1-ylethoxy)benzaldehyde (**30**) (5 mmol, 1.16 g) and methyl cyanoacetate (5 mmol, 495 mg) and resorcinol (5 mmol, 560 mg). Yield 25%, M.p. 180-181°C, yellow powder. IR ν_{\max} (KBr) cm^{-1} : 3400.00, 3299.51 (NH_2), 1660.34 ($\text{C}=\text{O}$) cm^{-1} . ^1H NMR (400 MHz, DMSO-d_6): δ 1.35 (d, 2H, $J=4.88$ Hz, CH_2), 1.46 (t, 4H, $J=5.38$ Hz, $(\text{CH}_2)_2$), 2.39 (bs, 4H, $(\text{CH}_2)_2$), 2.51 (t, 2H, $J=1.96$ Hz, CH_2), 3.51 (s, 3H, O- CH_3), 3.96 (t, 2H, $J=5.86$ Hz, CH_2), 4.75 (s, 1H, CH), 6.44 (d, 1H, $J=2.44$ Hz, Ar-H), 6.48 (dd, 1H, $J=2.20$ Hz, 6.36 Hz, Ar-H), 6.78 (d, 2H, $J=8.80$ Hz, Ar-H), 6.99 (d, 1H, $J=8.32$ Hz, Ar-H), 7.03 (d, 2H, $J=8.80$ Hz, Ar-H). ^{13}C NMR (100 MHz, DMSO-d_6): δ 24.39 (CH_2), 26.03 (CH_2), 38.47 (CH), 50.86 (CH_3), 54.85 (CH_2), 57.89

(CH₂), 65.85 (CH₂), 77.07 (C), 102.50 (CH), 112.52 (CH), 114.58 (CH), 117.79 (C), 128.12 (CH), 130.07 (CH), 141.08 (C), 149.54 (C), 157.08 (C), 161.64 (C), 169.20 (C=O). HRMS: Found 425.2082; C₂₄H₂₉N₂O₅ requires 425.2076(M⁺+H).

5.1.6.7. 2-Amino-7-hydroxy-4-[4-(2-piperidin-1-ylethoxy)phenyl]-4H-chromene-3-carboxylic ethyl ester (38).

(i) Following the method outlined above for **28**, a solution of ethyl cyanoacetate (10 mmol, 1.13 g) and 4-(2-piperidin-1-ylethoxy)benzaldehyde (**30**) (10 mmol, 2.33 g) in ethanol (30 mL) was heated briefly to reflux and then cooled. Piperidine (0.5 mL) was then added dropwise and the reaction solution heated at reflux for 4 h. The solvent was removed *in vacuo* to afford the product (*E*)-ethyl 2-cyano-3-[4-(2-piperidin-1-ylethoxy)phenyl]acrylate **45** which was recrystallized from ethanol as orange coloured crystals and used immediately in the next reaction. Yield: 334 mg, 10.2%, M.p. 88-90 °C. IR ν_{\max} (KBr): 2213(CN), 1702(C=O), 1610(C=C) cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 8.06 (s, 1H, CH=C), 7.92 (d, *J* = 9.16 Hz, 2H, ArH), 6.86 (d, *J* = 8.54 Hz, 2H, ArH), 4.34 (q, *J* = 7.12 Hz, 2H, OCH₂), 3.45 (br. s., 4H, 2xCH₂), 1.69 (br. s., 6H, 3xCH₂), 1.38 (t, *J* = 7.32 Hz, 3H, OCH₂CH₃). ¹³C NMR (100 MHz, CDCl₃): δ 14.26 (CH₃), 24.31 (CH₂), 25.32 (CH₂), 48.10 (CH₂), 61.90 (CH₂), 94.64 (C=CH), 113.24 (CH), 117.38 (CN), 120.08 (C), 134.06 (CH), 154.01 (C), 154.21 (CH=C), 164.14 (C=O). (ii) Preparation following the general method B above from (*E*)-ethyl 2-cyano-3-[4-(2-piperidin-1-ylethoxy)phenyl]acrylate **45** (10 mmol, 2.33 g) and resorcinol (10 mmol, 1.13 g). Yield 20%, M.p. 176-178 °C, yellow powder. IR ν_{\max} (KBr) cm⁻¹: 3415.04, 3293.18 (NH₂), 1672.17 (C=O) cm⁻¹. ¹H NMR (400 MHz, DMSO-d₆): δ 1.04 (m, 9H, CH₃, CH₂), 1.98 (m, 4H, CH₂), 2.78 (2H, CH₂), 3.94 (t, 4H, *J*=6.13 Hz, CH₂), 4.74 (s, 1H, CH), 6.44 (d, 1H, *J*=2.44 Hz, Ar-H), 6.48 (dd, 1H, *J*=2.44 Hz, 5.88 Hz, Ar-H), 6.69 (d, 2H, *J*=8.80 Hz, Ar-H), 6.88 (d, 1H, *J*=8.32 Hz, Ar-H), 7.02 (d, 2H, *J*=6.84 Hz, Ar-H), 7.09 (d, 1H, *J*=8.80 Hz, Ar-H). ¹³C NMR (100 MHz, DMSO-d₆): δ 14.32 (CH₃), 23.92 (CH₂),

25.55 (CH₂), 38.09 (CH), 54.39 (CH₂), 57.41 (CH₂), 58.52 (CH₂), 65.39 (CH₂), 76.81 (C), 101.99 (CH), 112.05 (CH), 114.02 (CH), 117.12 (C), 127.85 (CH), 129.75 (CH), 140.82(C), 149.02 (C), 156.53 (C), 156.60 (C), 160.91 (C), 168.39 (C=O). HRMS: Found 439.2227; C₂₅H₃₁N₂O₅, requires 439.2233(M⁺+H).

5.1.6.8. 2-Amino-7-hydroxy-4-[4-(2-piperidin-1-ylethoxy)phenyl]-4H-chromeme-3-carbonitrile (39).

Preparation as described for **38** above from 4-(2-piperidine-1-ylethoxy)benzaldehyde (**30**) (5 mmol, 1.16 g), malononitrile (5 mmol, 330 mg) and resorcinol (5 mmol, 560 mg) following method A. Yield 35%, M.p. 194-196 °C, beige powder. IR ν_{\max} (KBr) cm⁻¹: 3401.16, 3300.00 (NH₂), 2186.78 (CN) cm⁻¹. ¹H NMR (400 MHz, DMSO-d₆): δ 1.43 (m, 2H, CH₂), 1.56 (m, 4H, CH₂), 2.48 (m, 4H, CH₂), 2.68 (t, 2H, *J*=6.02 Hz, N-CH₂), 4.07 (t, 2H, *J*=6.04 Hz, O-CH₂), 4.63 (s, 1H, CH), 6.52 (d, 1H, *J*=2.48 Hz, Ar-H), 6.60 (dd, 1H, *J*=2.52 Hz, 6.00 Hz, Ar-H), 6.78 (s, 1H, Ar-H), 6.87 (dd, 2H, *J*=2.34 Hz, 4.48 Hz, Ar-H), 7.15 (m, 2H, Ar-H). ¹³C NMR (100 MHz, DMSO-d₆): δ 24.39 (CH₂), 26.03 (CH₂), 38.47 (CH), 50.86 (CH₃), 54.85 (CH₂), 57.89 (CH₂), 65.85 (CH₂), 77.07 (C), 102.50 (CH), 112.52 (CH), 114.58 (CH), 117.79 (C), 128.12 (CH), 130.07 (CH), 141.08 (C), 149.54 (C), 157.08 (C), 161.64 (C), 169.20 (C=O). HRMS: Found 392.1957; C₂₃H₂₆N₃O₃ requires 392.1974(M⁺+H).

5.1.6.9. 2-Amino-7-hydroxy-4-[4-(2-piperidin-1-ylethoxy)phenyl]-4H-chromeme-3-carboxylic *n*-butyl ester (40).

Preparation as described for **39** above from 4-(2-piperidine-1-ylethoxy)benzaldehyde (**30**) (5 mmol, 1.16 g), *n*-butylcyanoacetate (5 mmol, 705.8 mg) and resorcinol (5 mmol, 560 mg) using general method A. Yield 13%, M.p. 158 °C, yellow powder. IR ν_{\max} (KBr) cm⁻¹: 3415.93, 3294.68 (NH₂), 1672.10 (C=O) cm⁻¹. ¹H NMR (400 MHz, DMSO-d₆): δ 1.06 (t, 3H, *J*=7.02 Hz, CH₂), 1.45 (m, 8H, (CH₂)₄), 2.40 (bs, 4H, (CH₂)₂), 2.51 (s, 2H, CH₂), 2.60 (t, 2H, *J*=5.52 Hz, N-CH₂), 3.95 (m, 4H, (CH₂)₂), 4.73 (s, 1H, CH), 6.42 (d, 1H, *J*=2.48 Hz, Ar-H),

6.46 (dd, 1H, $J=2.28$ Hz, 6.00 Hz, Ar-H), 6.78 (d, 2H, $J=8.52$ Hz, Ar-H), 6.93 (d, 1H, $J=8.52$ Hz, Ar-H), 7.01 (d, 2H, $J=8.52$ Hz, Ar-H). ^{13}C NMR (100 MHz, DMSO- d_6): δ 14.32 (CH₃), 23.88 (CH₂), 25.50 (CH₂), 30.51 (CH₂), 38.10 (CH), 54.36 (CH₃), 56.03 (CH₂), 57.36 (CH₂), 58.52 (CH₂), 62.20 (CH₂), 65.35 (CH₂), 76.82 (C), 101.99 (CH), 106.17 (CH), 112.06 (CH), 114.03 (CH), 117.09 (C), 127.73 (CH), 129.74 (CH), 140.84 (C), 149.02 (C), 156.52 (C), 158.45 (C), 160.91 (C), 168.39 (C=O). HRMS: Found 467.2558; C₂₇H₃₅N₂O₅ requires 467.2546(M⁺+H).

5.1.6.10. 2-Amino-7-hydroxy-4-[4-(2-piperidin-1-ylethoxy)phenyl]-4H-chromeme-3-carboxylic tert-butyl ester (41).

Preparation as described for **39** above from 4-(2-piperidine-1-ylethoxy)benzaldehyde (**30**) (2.5 mmol, 0.583 g), *tert*-butylcyanoacetate (2.5 mmol, 360 mg) and resorcinol (5 mmol, 560 mg) using general Method A. Yield 27%, colourless oil. IR ν_{max} (KBr) cm⁻¹: 3415.98, 3289.85 (NH₂), 1672.28 (C=O) cm⁻¹. ^1H NMR (400 MHz, DMSO- d_6): δ 1.27 (s, 9H, (CH₃)₃), 1.36 (d, 2H, $J=5.28$ Hz, CH₂), 1.48 (m, 4H, (CH₂)₂), 2.51 (bs, 4H, (CH₂)₂), 2.58 (t, 2H, $J=5.84$ Hz, N-CH₂), 3.96 (t, 2H, $J=5.84$ Hz, CH₂), 4.65 (s, 1H, CH), 6.39 (d, 1H, $J=2.36$ Hz, Ar-H), 6.45 (dd, 1H, $J=2.32$ Hz, 6.16 Hz, Ar-H), 6.77 (d, 2H, $J=8.48$ Hz, Ar-H), 6.93 (d, 1H, $J=8.48$ Hz, Ar-H), 7.01 (d, 2H, $J=8.76$ Hz, Ar-H). 7.48 (s, 2H, NH₂), 9.56 (s, 1H, OH). ^{13}C NMR (100 MHz, DMSO- d_6): δ 23.93 (CH₂), 25.56 (CH₂), 28.11 (CH₃), 38.58 (CH), 54.42 (CH₂), 57.40 (CH₂), 65.46 (CH₂), 77.86 (C), 78.02 (C), 101.90 (CH), 111.92 (CH), 113.95 (CH), 116.94 (C), 127.87 (CH), 129.83 (CH), 141.18 (C), 148.85 (C), 156.47 (C), 156.52 (C), 160.52 (C), 168.50 (C=O). HRMS: Found 467.2551; C₂₇H₃₅N₂O₅ requires 467.2546(M⁺+H).

5.1.6.11. 2-Amino-7-hydroxy-4-[4-(2-morpholin-4-ylethoxy)phenyl]-4H-chromeme-3-carboxylic methyl ester (42).

Preparation as described for **39** above from 4-(2-morpholine-1-ylethoxy)benzaldehyde (**31**) (5 mmol, 1.176 g), methyl cyanoacetate methyl cyanoacetate (5 mmol, 495 mg) and

resorcinol (5 mmol, 560 mg) using general method C. Yield 226 mg, 10.6% as a brown oil. IR ν_{\max} (KBr) cm^{-1} : 3432.55, 3367.14 (NH_2), 1663.47 ($\text{C}=\text{O}$) cm^{-1} . ^1H NMR (400 MHz, DMSO-d_6): δ 1.07 (t, 3H, $J=7.02$ Hz, CH_3), 2.42 (s, 4H, $(\text{CH}_2)_2$), 2.62 (t, 2H, $J=5.52$ Hz, CH_2), 3.54 (t, 4H, $J=4.52$ Hz, CH_2), 3.96 (m, 4H, $2\times\text{CH}_2$), 4.72 (s, 1H, CH), 6.41 (d, 1H, $J=2.00$ Hz, Ar-H), 6.46 (dd, 1H, $J=2.26$ Hz, 6.04 Hz, Ar-H), 6.78 (d, 1H, $J=8.52$ Hz, Ar-H), 6.93 (d, 2H, $J=8.04$ Hz, Ar-H), 7.01 (d, 2H, $J=8.52$ Hz, Ar-H), 7.60 (s(br), 2H, NH_2), 9.65 (s, 1H, OH). ^{13}C NMR (100 MHz, DMSO-d_6): δ 14.32 (CH_3), 38.11 (CH), 53.60 (CH_2), 57.04 (CH_2), 58.53 (CH_2), 65.13 (CH_2), 66.14 (CH_2), 76.81 (C), 101.99 (CH), 112.06 (CH), 114.03 (CH), 117.11 (C), 127.87 (CH), 129.76 (CH), 140.89 (C), 149.03 (C), 156.48 (C), 156.62 (C), 160.91 (C), 168.40 ($\text{C}=\text{O}$). HRMS: Found 427.1873; $\text{C}_{23}\text{H}_{27}\text{N}_2\text{O}_6$ requires 427.1869(M^++H).

5.1.6.12. 2-Amino-7-hydroxy-4-[4-(2-morpholin-4-ylethoxy)phenyl]-4H-chromene-3-carboxylic ethyl ester (43).

Preparation as described for **39** above following the general Method B from 4-(2-morpholine-1-ylethoxy)benzaldehyde (**31**) (5 mmol, 1.176 g), ethyl cyanoacetate (5 mmol, 565 mg) and resorcinol (5 mmol, 560 mg). Yield 24%, M.p. 144-146 °C, yellow powder. IR ν_{\max} (KBr) cm^{-1} : 3425.00, 3285.02 (NH_2), 1674.22 ($\text{C}=\text{O}$) cm^{-1} . ^1H NMR (400 MHz, DMSO-d_6): δ 1.07 (t, 3H, $J=7.02$ Hz, CH_3), 2.42 (s, 4H, $(\text{CH}_2)_2$), 2.62 (t, 2H, $J=5.52$ Hz, CH_2), 3.54 (t, 4H, $J=4.52$ Hz, CH_2), 3.96 (m, 4H, CH_2), 4.72 (s, 1H, CH), 6.41 (d, 1H, $J=2.00$ Hz, Ar-H), 6.46 (dd, 1H, $J=2.26$ Hz, 6.04 Hz, Ar-H), 6.78 (d, 1H, $J=8.52$ Hz, Ar-H), 6.93 (d, 2H, $J=8.04$ Hz, Ar-H), 7.01 (d, 2H, $J=8.52$ Hz, Ar-H), 7.60 (bs, 2H, NH_2), 9.65 (s, 1H, OH). ^{13}C NMR (100 MHz, DMSO-d_6): δ 14.32 (CH_3), 38.11 (CH), 53.60 (CH_2), 57.04 (CH_2), 58.53 (CH_2), 65.13 (CH_2), 66.14 (CH_2), 76.81 (C), 101.99 (CH), 112.06 (CH), 114.03 (CH), 117.11 (C), 127.87 (CH), 129.76 (CH), 140.89 (C), 149.03 (C), 156.48 (C), 156.62 (C), 160.91 (C), 168.40 ($\text{C}=\text{O}$). HRMS: Found 441.2044; $\text{C}_{24}\text{H}_{29}\text{N}_2\text{O}_6$ requires 441.2026(M^++H).

5.1.6.13. 2-Amino-7-hydroxy-4-[4-(2-morpholin-4-ylethoxy)phenyl]-4H-chromene-3-carbonitrile (44).

Preparation was as described for **39** above from 4-(2-morpholine-1-ylethoxy)benzaldehyde (**31**) (5 mmol, 1.176 g), malononitrile (5 mmol, 330 mg) and resorcinol (5 mmol, 560 mg) following the general method B. Yield 51%, M.p. 176°C, brown powder. IR ν_{\max} (KBr) cm^{-1} : 3400.0, 3333.3 (NH_2), 2187.7 (CN) cm^{-1} . ^1H NMR (400 MHz, DMSO-d_6): δ 2.36 (m, 4H, CH_2), 2.57 (t, 2H, $J=5.76$ Hz, CH_2), 3.36 (m, 4H, CH_2), 3.95 (t, 2H, $J=5.76$ Hz, CH_2), 4.29 (bs, 2H, NH_2), 4.47 (s, 1H, CH), 6.31 (d, 1H, $J=2.04$ Hz, Ar-H), 6.39 (dd, 1H, $J=2.50$ Hz, 6.04 Hz, Ar-H), 6.69 (d, 1H, $J=8.04$ Hz, Ar-H), 6.77 (t, 3H, $J=8.28$ Hz, Ar-H), 6.97 (d, 1H, $J=8.52$ Hz, Ar-H). ^{13}C NMR (100 MHz, DMSO-d_6): δ 37.8 (CH), 53.61 (CH_2), 56.04 (CH_2), 56.53 (C), 65.21 (CH_2), 66.16 (CH_2), 102.10 (CH), 112.31 (CH), 114.05 (C), 114.45 (CH), 120.73 (C), 128.41 (CH), 129.91 (CH), 138.54 (C), 148.76 (C), 156.96 (C), 157.19 (C), 160.08 (C=O). HRMS: Found 394.1760; $\text{C}_{22}\text{H}_{24}\text{N}_3\text{O}_4$ requires 394.1767(M^+H).

5.2. Stability Study for compounds 38 and 40

Analytical high-performance liquid chromatography (HPLC) stability studies were performed using a Symmetry® column (C_{18} , 5 μm , 4.6×150 mm), a Waters 2487 Dual Wavelength Absorbance detector, a Waters 1525 binary HPLC pump and a Waters 717plus Autosampler. Samples were detected at wavelength of 254 nm. All samples were analysed using acetonitrile (80%): water (20%) as the mobile phase over 10 min and a flow rate of 1 mL/min. Stock solutions are prepared by dissolving 5 mg of compound **38** and **40** in 10 mL of mobile phase. Phosphate buffers at the desired pH values (4, 7.4, and 9) were prepared in accordance with the British Pharmacopoeia monograph 2016. 30 μL of stock solution was diluted with 1 mL of appropriate buffer, shaken and injected immediately. Samples were withdrawn and analysed at time intervals of $t=0$ min, 5 min, 30 min, 60 min, 90 min, 120 min and 21 hours.

5.3. Plasma stability study for compounds 38 and 40

Analytical high-performance liquid chromatography (HPLC) stability studies for plasma samples were performed using a Waters HPLC 2965-2487 system with a Hypersil Gold column; $l=0.15$ m, i.d.=4.6 mm, C18, 5 μ m with flow rate: 1.0 mL/min, sample temperature: 25 °C, column temperature: 25 °C; mobile phase H₂O:ACN:MeOH. *Method 1*: (10% DMSO): PBS (1.6 mL) added to blood plasma (2.0 mL) and pre-warmed to 37 °C in a water bath. To this solution, 400 μ L of 1.0 mg/mL stock of compounds **38** and **41** was added. Immediately a 100 μ L aliquot was withdrawn and added to 900 μ L acetonitrile giving a final injection concentration of 0.01 mg/mL. The samples were then centrifuged at 5000 rpm for 10 minutes and filtered through a 0.2 micron filter and injected according to the HPLC conditions listed above. Further samples were taken in the same manner every 15 minutes for the first hour and hourly thereafter up to 3 hours. A final sample was taken after 24 hours.

5.4. Biochemical evaluation of activity

5.4.1. Antiproliferative studies

All assays were performed in triplicate for the determination of mean values reported. Compounds were assayed as the free bases isolated from reaction. The human breast cancer cell line MCF-7 was cultured in Eagles minimum essential medium in air supplemented with 5% CO₂ atmosphere with 10% fetal calf serum, 2 mM L-glutamine and 100 μ g/mL penicillin/streptomycin. The medium was supplemented with 1% non-essential amino acids.

Cells were trypsinised and seeded at a density of 2.5×10^4 cells/mL into a 96-well plate and incubated at 37 °C, in air supplemented with 5% CO₂ atmosphere for 24 h. After this time they were treated with 2 μ L volumes of test compound which had been pre-prepared as stock solutions in ethanol to furnish the final concentration range of study, 1 nM-100 μ M, and re-incubated for a further 72 h. Control wells contained the equivalent volume of the vehicle ethanol (1% v/v). The culture medium was then removed and the cells washed with 100 μ L

phosphate buffer saline (PBS) and 50 μL of 1 mg/mL MTT solution was added. Cells were incubated for 2 h in darkness at 37 °C. At this point solubilisation was begun through the addition of 200 μL DMSO and the cells maintained at room temperature in darkness for 20 min to ensure thorough colour diffusion before reading the absorbance at 595 nm. The absorbance value of control cells (vehicle treated) was set to 100% cell viability and from this graphs of absorbance versus cell density per well were prepared to assess cell viability and from these, graphs of percentage cell viability versus concentration of subject compound were drawn.

5.4.2. Lactate Dehydrogenase Assay for Measurement of Cytotoxicity.

Cytotoxicity was determined using the CytoTox 96 nonradioactive cytotoxicity assay (Promega) following the manufacturer's protocol. In this assay, the release of cytoplasmic lactate dehydrogenase (LDH) is used as a measure of cell lysis. MCF-7 cells were seeded at a density of 1×10^4 cells/well in a 96-well plate and incubated for 24 hours. The cells were then dosed with 2 μL volumes of the test compounds, over the concentration range 1 nM – 50 μM . After dosing, the plates are incubated for 72 hours. Control wells contained the equivalent volume of the vehicle, ethanol (1% v/v). After incubation, 30 μL of medium is removed from each well and transferred to a fresh 96-well plate. To this, 30 μL of substrate mix from the Promega cytotoxicity assay kit was added. The plate was left in darkness for 20 minutes at room temperature. Stop solution (30 μL) was added to the wells. The plates were read at a wavelength of 490 nm using an automated *Molecular Devices* micro-plate reader. A positive control of 100% lysis was determined for a set of untreated cells, which were lysed by the addition of lysis solution to the medium 45 minutes prior to harvesting. The percentage cell lysis was calculated for the treated cells relative to the control data. Assays were repeated in three experiments performed in triplicate (unless otherwise stated) and reported results represent the mean value \pm standard error mean. The data was processed using

PRISM (Version 4. Graphpad Software, Inc., 2236 Avenida de la Playa, La Jolla, CA 92037, USA).

5.4.3. Estrogen Receptor Binding

ER α and ER β fluorescence polarization based competitor assay kits were obtained from Panvera at Invitrogen Life Technologies, Invitrogen Corporation, 5791 Van Allen Way, PO Box 6482, Carlsbad, California 92008. The recombinant ER (insect expressed, full length, untagged human ER obtained from recombinant baculovirus-infected insect cells) and the fluorescent estrogen ligand were removed from the -80°C freezer and thawed on ice for one hour prior to use. The fluorescent estrogen ligand (2 nM) was added to the ER (40 nM for ER α and 30 nM for ER β) and screening buffer (100 mM potassium phosphate (pH 7.4), 100 μ g/mL BGG, 0.02% NaN₃ was added to make up to a final volume that was dependent on the number of tubes used (number of tubes (e.g. 50) x volume of complex in each tube (50 μ L) = total volume (e.g. 2500 μ L). Test compound (1 μ L, concentration range 100 nM to 1 mM) was added to 49 μ L screening buffer in each borosilicate tube (6 mm diameter). To this 50 μ L of the fluorescent estrogen/ER complex was added to make up a final volume of 100 μ L. A vehicle control contained 1% (v/v) of ethanol; a negative control was used to determine the theoretical maximum polarization (50 μ L of screening buffer and 50 μ L of fluorescent estrogen/ER complex). The tubes were incubated in the dark at room temperature for 2 h and were mixed by shaking on a plate-shaker. Polarization values were measured on a Beacon single-tube fluorescent polarization instrument fitted with 485 nm excitation and 530 nm emission interference filters. For ER α and ER β , graphs of anisotropy (mA) versus compound concentration were obtained for the determination of IC₅₀ values.

5.5. Computational Procedures.

All compounds were drawn in Accelrys Draw v4.1⁸⁸ with absolute stereochemistry assigned for both R and S enantiomers. Enantiomers were protonated at pH7.4 with MarvinView and subsequently converted to 3D using Molconvert (version v15.6.29.0)⁸². 10 conformers of each 3D structure were subsequently enumerated using OMEGA (version 2.5.1.4)^{89,90}.

To establish a reliable model of ER β , as no X-ray crystal structure of human ER β in an antagonist conformation currently exists, the sequence from Uniprot⁹¹ entry Q92731 (ESR2_HUMAN) was downloaded and submitted to the iTASSER server^{92,93}. PDB entry 1R5K was specified as a template to guide the structure prediction. The resulting model was submitted to the Yale Protein Multichain Morphing Server⁷⁹ to transform the atomic coordinates to those of PDB entry 1R5K. Finally, 3Drefine^{80,94,95} was used to refine our model of ER β . PDB ID 2OUZ was selected for modeling of the compound series in ER α as the co-crystallised ligand (Lasofoxifene) was structurally similar to compound **35**. The iTASSER server was also used to model any missing residues with 2OUZ used to guide the structural prediction and again 3Drefine^{80,94,95} was used to refine our model of ER α . All conformers were docked using FRED (OEDocking version 3.0.1)^{81,96} into the both Ligand Binding Pockets (LBPs). Energy minimisation of the top scoring docked pose for all enantiomers was performed using the Amber12:EHT force field (gradient of 0.05kcal/mol) as implemented in MOE (version 2012.10)⁹⁷.

5.6. X-Ray Crystallography

Crystals of compound **22** were obtained by slow crystallisation from a dilute solution of methanol over a period of 4-8 weeks. The data for the crystal structures **22** were collected on a Rigaku Saturn 724 CCD Diffractometer. A suitable crystal from each crystal compound was selected and mounted using inert oil on a 0.3 mm diameter glass fiber tip or loop and placed on the goniometer head in a 150K N₂ gas stream. Each data set was collected using Crystalclear-SM 1.4.0 software. Data integration, reduction and correction for absorption and

polarization effects were all performed using Crystalclear-SM 1.4.0 software. Space group determination, structure solution and refinement were obtained using Bruker Shelxtl Ver. 6.14 software⁹⁸. Each structure was solved with Direct Methods using the SHELXTL program and refined against IF2I with the program XL from SHELX-97 using all data. Non-hydrogen atoms were refined with anisotropic thermal parameters. Hydrogen atoms were placed into geometrically calculated positions and refined using a riding model.

Crystal data for **22** are as follows: Empirical formula: C₂₀H₂₁NO₅, M = 355.38, T = 108(2) K, Wavelength 0.71075 Å, Crystal system: Monoclinic, Space group: P 2/c, Unit cell dimensions: a = 16.982(7) Å, b = 5.340(2) Å, c = 20.683(9) Å, α = 90°, β = 91.189(6)°, γ = 90°. Volume: 1875.2(13) Å³ Z: 4, Density (calculated): 1.259 Mg/m³, Absorption coefficient: 0.091 mm⁻¹; F(000): 752; Crystal size: 0.240 x 0.210 x 0.180 mm³; Theta range for data collection: 2.328 to 24.999°. Index ranges -20 ≤ h ≤ 20, -5 ≤ k ≤ 6, -23 ≤ l ≤ 24; Reflections collected: 13677, Independent reflections: 3240 [R(int) = 0.0641]; Completeness to theta = 25.242°; 95.2 %; Absorption correction: Semi-empirical from equivalents, max. and min. transmission: 1.0000 and 0.6725, Refinement method: Full-matrix least-squares on F², Data / restraints / parameters: 3240 / 0 / 238, Goodness-of-fit on F² :1.173, Final R indices: [I > 2σ(I)]: R1 = 0.0840, wR2 = 0.2055, R indices (all data): R1 = 0.1059, wR2 = 0.2203, Extinction coefficient: n/a, Largest diff. peak and hole: 0.216 and -0.253 e.Å⁻³. CCDC deposition CCDC 1014637.

Ancilliary Information

Supporting Information

Additional figures illustrating top ranked poses of Compound **42** (S Enantiomer) in ER_α and ER_β, co-ordinates of refined models of ER_α and ER_β, chemgauss4 scores for docked poses of enantiomers of compounds **35** and **42** in ER_α and ER_β.

Corresponding Author Information

*E-mail: andrew.knox@dit.ie. Phone: (+353) 1-896-4777.

Author Contributions

M.C. and A.J.S.K. contributed equally.

Notes

The authors declare no competing financial interest.

Acknowledgements

This work was supported through funding from the Trinity College IITAC research initiative (HEA PRTLTI), Enterprise Ireland (EI), Science Foundation Ireland (SFI), with additional support for computational facilities from the Wellcome Trust. Postgraduate research awards from the Irish Research Council Government of Ireland (GOIPD/2013/188; NMO'B, EPSPD/2012/360; DKN) and the Health Research Board (PD2009/33; AJSK) respectively are gratefully acknowledged. The Trinity Biomedical Sciences Institute is supported by a capital infrastructure investment from Cycle 5 of the Irish Higher Education Authority's Programme for Research in Third Level Institutions (PRTLTI). We would also like to thank Openeye Scientific Software, Chemaxon and Chemical Computing Group for their continuing support of this research.

Abbreviations Used

EI	Electron Impact
ER	Estrogen receptor
GTP	Guanidine triphosphate
HRMS	High Resolution Molecular Ion Determination
IC	Inhibitory concentration
IR	Infrared
LDB	Ligand binding domain
LBP	Ligand binding Pocket

LRMS	Low Resolution Mass Spectra
MTD	Maximum tolerated dose
MTT	3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide
NMR	Nuclear Magnetic Resonance
ORTEP	Oak Ridge Thermal Ellipsoid Plot
PDB	Protein Data Bank
PBS	Phosphate buffer saline
SAR	Structure-Activity Relationship
SERM	Selective Estrogen Receptor Modulation
TBDMS	<i>tert</i> -Butyldimethylchlorosilane
THF	Tetrahydrofuran
TLC	Thin Layer Chromatography
TMCS	Trimethylchlorosilane
TMS	Tetramethylsilane

References

- (1) Barton, M. Position Paper: The Membrane Estrogen Receptor GPER--Clues and Questions. *Steroids* **2012**, *77* (10), 935–942.
- (2) Filardo, E. J. GPER and ER: Estrogen Receptors with Distinct Biological Roles in Breast Cancer. *Immunology, Endocrine & Metabolic Agents - Medicinal Chemistry* **2011**, *11* (4), 243–254.
- (3) Filardo, E. J.; Quinn, J. A.; Bland, K. I.; Frackelton, A. R., Jr. Estrogen-Induced Activation of Erk-1 and Erk-2 Requires the G Protein-Coupled Receptor Homolog, GPR30, and Occurs via Trans-Activation of the Epidermal Growth Factor Receptor through Release of HB-EGF. *Mol. Endocrinol.* **2000**, *14* (10), 1649–1660.
- (4) Maggiolini, M.; Vivacqua, A.; Fasanella, G.; Recchia, A. G.; Sisci, D.; Pezzi, V.; Montanaro, D.; Musti, A. M.; Picard, D.; Andò, S. The G Protein-Coupled Receptor GPR30 Mediates c-Fos up-Regulation by 17beta-Estradiol and Phytoestrogens in Breast Cancer Cells. *J. Biol. Chem.* **2004**, *279* (26), 27008–27016.
- (5) Luksha, L.; Kublickiene, K. The Role of Estrogen Receptor Subtypes for Vascular Maintenance. *Gynecol. Endocrinol.* **2009**, *25* (2), 82–95.
- (6) Ellem, S. J.; Risbridger, G. P. The Dual, Opposing Roles of Estrogen in the Prostate. *Ann. N. Y. Acad. Sci.* **2009**, *1155*, 174–186.
- (7) Thomas, C.; Gustafsson, J.-Å. The Different Roles of ER Subtypes in Cancer Biology and Therapy. *Nat. Rev. Cancer* **2011**, *11* (8), 597–608.
- (8) Harris, H. A.; Katzenellenbogen, J. A.; Katzenellenbogen, B. S. Characterization of the Biological Roles of the Estrogen Receptors, ERalpha and ERbeta, in Estrogen Target Tissues in Vivo through the Use of an ERalpha-Selective Ligand. *Endocrinology* **2002**, *143* (11), 4172–4177.
- (9) Shanle, E. K.; Xu, W. Selectively Targeting Estrogen Receptors for Cancer Treatment. *Adv. Drug Deliv. Rev.* **2010**, *62* (13), 1265–1276.

- (10) Nilsson, S.; Gustafsson, J.-Å. Estrogen Receptors: Therapies Targeted to Receptor Subtypes. *Clin. Pharmacol. Ther.* **2011**, *89* (1), 44–55.
- (11) Nilsson, S.; Koehler, K. F.; Gustafsson, J.-Å. Development of Subtype-Selective Oestrogen Receptor-Based Therapeutics. *Nat Rev Drug Discov* **2011**, *10* (10), 778–792.
- (12) Heldring, N.; Pike, A.; Andersson, S.; Matthews, J.; Cheng, G.; Hartman, J.; Tujague, M.; Ström, A.; Treuter, E.; Warner, M.; Gustafsson, J.-A. Estrogen Receptors: How Do They Signal and What Are Their Targets. *Physiol. Rev.* **2007**, *87* (3), 905–931.
- (13) Minutolo, F.; Macchia, M.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. Estrogen Receptor β Ligands: Recent Advances and Biomedical Applications. *Med Res Rev* **2011**, *31* (3), 364–442.
- (14) Mohler, M. L.; Narayanan, R.; Coss, C. C.; Hu, K.; He, Y.; Wu, Z.; Hong, S.-S.; Hwang, D. J.; Miller, D. D.; Dalton, J. T. Estrogen Receptor Beta Selective Nonsteroidal Estrogens: Seeking Clinical Indications. *Expert Opin Ther Pat* **2010**, *20* (4), 507–534.
- (15) Meegan, M. J.; Lloyd, D. G. Advances in the Science of Estrogen Receptor Modulation. *Curr. Med. Chem.* **2003**, *10* (3), 181–210.
- (16) Cole, M. P.; Jones, C. T.; Todd, I. D. A New Anti-Oestrogenic Agent in Late Breast Cancer. An Early Clinical Appraisal of ICI46474. *Br. J. Cancer* **1971**, *25* (2), 270–275.
- (17) Ward, H. W. Anti-Oestrogen Therapy for Breast Cancer: A Trial of Tamoxifen at Two Dose Levels. *Br Med J* **1973**, *1* (5844), 13–14.
- (18) Jordan, V. C. Antiestrogens and Selective Estrogen Receptor Modulators as Multifunctional Medicines. 1. Receptor Interactions. *J. Med. Chem.* **2003**, *46* (6), 883–908.
- (19) Jordan, V. C. Tamoxifen: Toxicities and Drug Resistance during the Treatment and Prevention of Breast Cancer. *Annu. Rev. Pharmacol. Toxicol.* **1995**, *35*, 195–211.

- (20) Jordan, V. C. Is Tamoxifen the Rosetta Stone for Breast Cancer? *J. Natl. Cancer Inst.* **2003**, *95* (5), 338–340.
- (21) Jordan, V. C. Tamoxifen: A Most Unlikely Pioneering Medicine. *Nat Rev Drug Discov* **2003**, *2* (3), 205–213.
- (22) Alexi, X.; Kasiotis, K. M.; Fokialakis, N.; Lambrinidis, G.; Meligova, A. K.; Mikros, E.; Haroutounian, S. A.; Alexis, M. N. Differential Estrogen Receptor Subtype Modulators: Assessment of Estrogen Receptor Subtype-Binding Selectivity and Transcription-Regulating Properties of New Cycloalkyl Pyrazoles. *J. Steroid Biochem. Mol. Biol.* **2009**, *117* (4–5), 159–167.
- (23) Asim, M.; Asim, M.; El-Salfiti, M.; Qian, Y.; Choueiri, C.; Salari, S.; Cheng, J.; Shadnia, H.; Bal, M.; Christine Pratt, M. A.; Carlson, K. E.; Katzenellenbogen, J. A.; Wright, J. S.; Durst, T. Deconstructing Estradiol: Removal of B-Ring Generates Compounds Which Are Potent and Subtype-Selective Estrogen Receptor Agonists. *Bioorg. Med. Chem. Lett.* **2009**, *19* (4), 1250–1253.
- (24) Bertini, S.; De Cupertino, A.; Granchi, C.; Bargagli, B.; Tuccinardi, T.; Martinelli, A.; Macchia, M.; Gunther, J. R.; Carlson, K. E.; Katzenellenbogen, J. A.; Minutolo, F. Selective and Potent Agonists for Estrogen Receptor Beta Derived from Molecular Refinements of Salicylaldoximes. *Eur J Med Chem* **2011**, *46* (6), 2453–2462.
- (25) Carroll, V. M.; Jeyakumar, M.; Carlson, K. E.; Katzenellenbogen, J. A. Diarylpropionitrile (DPN) Enantiomers: Synthesis and Evaluation of Estrogen Receptor β -Selective Ligands. *J. Med. Chem.* **2012**, *55* (1), 528–537.
- (26) Chen, H. Y.; Dykstra, K. D.; Birzin, E. T.; Frisch, K.; Chan, W.; Yang, Y. T.; Mosley, R. T.; DiNinno, F.; Rohrer, S. P.; Schaeffer, J. M.; Hammond, M. L. Estrogen Receptor Ligands. Part 1: The Discovery of Flavanoids with Subtype Selectivity. *Bioorg. Med. Chem. Lett.* **2004**, *14* (6), 1417–1421.

- (27) Chesworth, R.; Zawistoski, M. P.; Lefker, B. A.; Cameron, K. O.; Day, R. F.; Mangano, F. M.; Rosati, R. L.; Colella, S.; Petersen, D. N.; Brault, A.; Lu, B.; Pan, L. C.; Perry, P.; Ng, O.; Castleberry, T. A.; Owen, T. A.; Brown, T. A.; Thompson, D. D.; DaSilva-Jardine, P. Tetrahydroisoquinolines as Subtype Selective Estrogen Agonists/Antagonists. *Bioorg. Med. Chem. Lett.* **2004**, *14* (11), 2729–2733.
- (28) Compton, D. R.; Sheng, S.; Carlson, K. E.; Rebacz, N. A.; Lee, I. Y.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. Pyrazolo[1,5-A]pyrimidines: Estrogen Receptor Ligands Possessing Estrogen Receptor Beta Antagonist Activity. *J. Med. Chem.* **2004**, *47* (24), 5872–5893.
- (29) De Angelis, M.; Stossi, F.; Carlson, K. A.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. Indazole Estrogens: Highly Selective Ligands for the Estrogen Receptor Beta. *J. Med. Chem.* **2005**, *48* (4), 1132–1144.
- (30) Dykstra, K. D.; Guo, L.; Birzin, E. T.; Chan, W.; Yang, Y. T.; Hayes, E. C.; DaSilva, C. A.; Pai, L.-Y.; Mosley, R. T.; Kraker, B.; Fitzgerald, P. M. D.; DiNinno, F.; Rohrer, S. P.; Schaeffer, J. M.; Hammond, M. L. Estrogen Receptor Ligands. Part 16: 2-Aryl Indoles as Highly Subtype Selective Ligands for ERalpha. *Bioorg. Med. Chem. Lett.* **2007**, *17* (8), 2322–2328.
- (31) García-Becerra, R.; Borja-Cacho, E.; Cooney, A. J.; Smith, C. L.; Lemus, A. E.; Pérez-Palacios, G.; Larrea, F. Synthetic 19-Nortestosterone Derivatives as Estrogen Receptor Alpha Subtype-Selective Ligands Induce Similar Receptor Conformational Changes and Steroid Receptor Coactivator Recruitment than Natural Estrogens. *J. Steroid Biochem. Mol. Biol.* **2006**, *99* (2–3), 108–114.
- (32) Henke, B. R.; Consler, T. G.; Go, N.; Hale, R. L.; Hohman, D. R.; Jones, S. A.; Lu, A. T.; Moore, L. B.; Moore, J. T.; Orband-Miller, L. A.; Robinett, R. G.; Shearin, J.; Spearing, P. K.; Stewart, E. L.; Turnbull, P. S.; Weaver, S. L.; Williams, S. P.; Wisely,

- G. B.; Lambert, M. H. A New Series of Estrogen Receptor Modulators That Display Selectivity for Estrogen Receptor Beta. *J. Med. Chem.* **2002**, *45* (25), 5492–5505.
- (33) Hsieh, R. W.; Rajan, S. S.; Sharma, S. K.; Guo, Y.; DeSombre, E. R.; Mrksich, M.; Greene, G. L. Identification of Ligands with Bicyclic Scaffolds Provides Insights into Mechanisms of Estrogen Receptor Subtype Selectivity. *J. Biol. Chem.* **2006**, *281* (26), 17909–17919.
- (34) Kim, S.; Wu, J. Y.; Birzin, E. T.; Frisch, K.; Chan, W.; Pai, L.-Y.; Yang, Y. T.; Mosley, R. T.; Fitzgerald, P. M. D.; Sharma, N.; Dahllund, J.; Thorsell, A.-G.; DiNinno, F.; Rohrer, S. P.; Schaeffer, J. M.; Hammond, M. L. Estrogen Receptor Ligands. II. Discovery of Benzoxathiins as Potent, Selective Estrogen Receptor Alpha Modulators. *J. Med. Chem.* **2004**, *47* (9), 2171–2175.
- (35) Larrea, F.; García-Becerra, R.; Lemus, A. E.; García, G. A.; Pérez-Palacios, G.; Jackson, K. J.; Coleman, K. M.; Dace, R.; Smith, C. L.; Cooney, A. J. A-Ring Reduced Metabolites of 19-nor Synthetic Progestins as Subtype Selective Agonists for ER Alpha. *Endocrinology* **2001**, *142* (9), 3791–3799.
- (36) Meyers, M. J.; Sun, J.; Carlson, K. E.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. Estrogen Receptor Subtype-Selective Ligands: Asymmetric Synthesis and Biological Evaluation of Cis- and Trans-5,11-Dialkyl- 5,6,11, 12-Tetrahydrochrysenes. *J. Med. Chem.* **1999**, *42* (13), 2456–2468.
- (37) Meyers, M. J.; Sun, J.; Carlson, K. E.; Marriner, G. A.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. Estrogen Receptor-Beta Potency-Selective Ligands: Structure-Activity Relationship Studies of Diarylpropionitriles and Their Acetylene and Polar Analogues. *J. Med. Chem.* **2001**, *44* (24), 4230–4251.
- (38) Moon, J. T.; Ha, S. H.; Lee, S. H.; Kwon, T. H.; Oh, C. R.; Kim, Y. D.; Kim, J.; Choo, D. J.; Lee, J. Y. Total Synthesis and Biological Evaluation of Methylgerambullone. *Bioorg. Med. Chem. Lett.* **2010**, *20* (1), 52–55.

- (39) Mortensen, D. S.; Rodriguez, A. L.; Carlson, K. E.; Sun, J.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. Synthesis and Biological Evaluation of a Novel Series of Furans: Ligands Selective for Estrogen Receptor Alpha. *J. Med. Chem.* **2001**, *44* (23), 3838–3848.
- (40) Muthyala, R. S.; Sheng, S.; Carlson, K. E.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. Bridged Bicyclic Cores Containing a 1,1-Diarylethylene Motif Are High-Affinity Subtype-Selective Ligands for the Estrogen Receptor. *J. Med. Chem.* **2003**, *46* (9), 1589–1602.
- (41) Norman, B. H.; Richardson, T. I.; Dodge, J. A.; Pfeifer, L. A.; Durst, G. L.; Wang, Y.; Durbin, J. D.; Krishnan, V.; Dinn, S. R.; Liu, S.; Reilly, J. E.; Ryter, K. T. Benzopyrans as Selective Estrogen Receptor Beta Agonists (SERBAs). Part 4: Functionalization of the Benzopyran A-Ring. *Bioorg. Med. Chem. Lett.* **2007**, *17* (18), 5082–5085.
- (42) Roelens, F.; Heldring, N.; Dhooge, W.; Bengtsson, M.; Comhaire, F.; Gustafsson, J.-A.; Treuter, E.; De Keukeleire, D. Subtle Side-Chain Modifications of the Hop Phytoestrogen 8-Prenylnaringenin Result in Distinct Agonist/Antagonist Activity Profiles for Estrogen Receptors Alpha and Beta. *J. Med. Chem.* **2006**, *49* (25), 7357–7365.
- (43) Shiau, A. K.; Barstad, D.; Loria, P. M.; Cheng, L.; Kushner, P. J.; Agard, D. A.; Greene, G. L. The Structural Basis of Estrogen Receptor/Coactivator Recognition and the Antagonism of This Interaction by Tamoxifen. *Cell* **1998**, *95* (7), 927–937.
- (44) Stauffer, S. R.; Coletta, C. J.; Tedesco, R.; Nishiguchi, G.; Carlson, K.; Sun, J.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. Pyrazole Ligands: Structure-Affinity/Activity Relationships and Estrogen Receptor-Alpha-Selective Agonists. *J. Med. Chem.* **2000**, *43* (26), 4934–4947.

- (45) Stauffer, S. R.; Huang, Y. R.; Aron, Z. D.; Coletta, C. J.; Sun, J.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. Triarylpyrazoles with Basic Side Chains: Development of Pyrazole-Based Estrogen Receptor Antagonists. *Bioorg. Med. Chem.* **2001**, *9* (1), 151–161.
- (46) Sun, J.; Baudry, J.; Katzenellenbogen, J. A.; Katzenellenbogen, B. S. Molecular Basis for the Subtype Discrimination of the Estrogen Receptor-Beta-Selective Ligand, Diarylpropionitrile. *Mol. Endocrinol.* **2003**, *17* (2), 247–258.
- (47) Sun, J.; Meyers, M. J.; Fink, B. E.; Rajendran, R.; Katzenellenbogen, J. A.; Katzenellenbogen, B. S. Novel Ligands That Function as Selective Estrogens or Antiestrogens for Estrogen Receptor-Alpha or Estrogen Receptor-Beta. *Endocrinology* **1999**, *140* (2), 800–804.
- (48) Tan, Q.; Blizzard, T. A.; Morgan, J. D., 2nd; Birzin, E. T.; Chan, W.; Yang, Y. T.; Pai, L.-Y.; Hayes, E. C.; DaSilva, C. A.; Warriar, S.; Yudkovitz, J.; Wilkinson, H. A.; Sharma, N.; Fitzgerald, P. M. D.; Li, S.; Colwell, L.; Fisher, J. E.; Adamski, S.; Reszka, A. A.; Kimmel, D.; DiNinno, F.; Rohrer, S. P.; Freedman, L. P.; Schaeffer, J. M.; Hammond, M. L. Estrogen Receptor Ligands. Part 10: Chromanes: Old Scaffolds for New SERAMs. *Bioorg. Med. Chem. Lett.* **2005**, *15* (6), 1675–1681.
- (49) Veeneman, G. H. Non-Steroidal Subtype Selective Estrogens. *Curr. Med. Chem.* **2005**, *12* (9), 1077–1136.
- (50) Waibel, M.; De Angelis, M.; Stossi, F.; Kieser, K. J.; Carlson, K. E.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. Bibenzyl- and Stilbene-Core Compounds with Non-Polar Linker Atom Substituents as Selective Ligands for Estrogen Receptor Beta. *Eur J Med Chem* **2009**, *44* (9), 3412–3424.
- (51) Waibel, M.; Kieser, K. J.; Carlson, K. E.; Stossi, F.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. Phenethyl Pyridines with Non-Polar Internal Substituents as

- Selective Ligands for Estrogen Receptor Beta. *Eur J Med Chem* **2009**, *44* (9), 3560–3570.
- (52) Wang, P.; Min, J.; Nwachukwu, J. C.; Cavett, V.; Carlson, K. E.; Guo, P.; Zhu, M.; Zheng, Y.; Dong, C.; Katzenellenbogen, J. A.; Nettles, K. W.; Zhou, H.-B. Identification and Structure-Activity Relationships of a Novel Series of Estrogen Receptor Ligands Based on 7-thiabicyclo[2.2.1]hept-2-Ene-7-Oxide. *J. Med. Chem.* **2012**, *55* (5), 2324–2341.
- (53) Wilkening, R. R.; Ratcliffe, R. W.; Fried, A. K.; Meng, D.; Sun, W.; Colwell, L.; Lambert, S.; Greenlee, M.; Nilsson, S.; Thorsell, A.; Mojena, M.; Tudela, C.; Frisch, K.; Chan, W.; Birzin, E. T.; Rohrer, S. P.; Hammond, M. L. Estrogen Receptor Beta-Subtype Selective Tetrahydrofluorenones: Use of a Fused Pyrazole as a Phenol Bioisostere. *Bioorg. Med. Chem. Lett.* **2006**, *16* (15), 3896–3901.
- (54) Wilkening, R. R.; Ratcliffe, R. W.; Tynebor, E. C.; Wildonger, K. J.; Fried, A. K.; Hammond, M. L.; Mosley, R. T.; Fitzgerald, P. M. D.; Sharma, N.; McKeever, B. M.; Nilsson, S.; Carlquist, M.; Thorsell, A.; Locco, L.; Katz, R.; Frisch, K.; Birzin, E. T.; Wilkinson, H. A.; Mitra, S.; Cai, S.; Hayes, E. C.; Schaeffer, J. M.; Rohrer, S. P. The Discovery of Tetrahydrofluorenones as a New Class of Estrogen Receptor Beta-Subtype Selective Ligands. *Bioorg. Med. Chem. Lett.* **2006**, *16* (13), 3489–3494.
- (55) Xin, D.; Wang, H.; Yang, J.; Su, Y.-F.; Fan, G.-W.; Wang, Y.-F.; Zhu, Y.; Gao, X.-M. Phytoestrogens from *Psoralea Corylifolia* Reveal Estrogen Receptor-Subtype Selectivity. *Phytomedicine* **2010**, *17* (2), 126–131.
- (56) Yang, W.; Wang, Y.; Ma, Z.; Golla, R.; Stouch, T.; Seethala, R.; Johnson, S.; Zhou, R.; Güngör, T.; Feyen, J. H. M.; Dickson, J. K., Jr. Synthesis and Structure-Activity Relationship of 3-Arylbenzoxazines as Selective Estrogen Receptor Beta Agonists. *Bioorg. Med. Chem. Lett.* **2004**, *14* (9), 2327–2330.

- (57) Yang, C.; Xu, G.; Li, J.; Wu, X.; Liu, B.; Yan, X.; Wang, M.; Xie, Y. Benzothiophenes Containing a Piperazine Side Chain as Selective Ligands for the Estrogen Receptor Alpha and Their Bioactivities in Vivo. *Bioorg. Med. Chem. Lett.* **2005**, *15* (5), 1505–1507.
- (58) Pelzer, T.; Jazbutyte, V.; Hu, K.; Segerer, S.; Nahrendorf, M.; Nordbeck, P.; Bonz, A. W.; Muck, J.; Fritzeimer, K.-H.; Hegele-Hartung, C.; Ertl, G.; Neyses, L. The Estrogen Receptor- α Agonist 16 α -LE2 Inhibits Cardiac Hypertrophy and Improves Hemodynamic Function in Estrogen-Deficient Spontaneously Hypertensive Rats. *Cardiovasc Res* **2005**, *67* (4), 604–612.
- (59) Renaud, J.; Bischoff, S. F.; Buhl, T.; Floersheim, P.; Fournier, B.; Geiser, M.; Halleux, C.; Kallen, J.; Keller, H.; Ramage, P. Selective Estrogen Receptor Modulators with Conformationally Restricted Side Chains. Synthesis and Structure-Activity Relationship of ERalpha-Selective Tetrahydroisoquinoline Ligands. *J. Med. Chem.* **2005**, *48* (2), 364–379.
- (60) Sun, J.; Huang, Y. R.; Harrington, W. R.; Sheng, S.; Katzenellenbogen, J. A.; Katzenellenbogen, B. S. Antagonists Selective for Estrogen Receptor α . *Endocrinology* **2002**, *143* (3), 941–947.
- (61) Malamas, M. S.; Manas, E. S.; McDevitt, R. E.; Gunawan, I.; Xu, Z. B.; Collini, M. D.; Miller, C. P.; Dinh, T.; Henderson, R. A.; Keith, J. C., Jr; Harris, H. A. Design and Synthesis of Aryl Diphenolic Azoles as Potent and Selective Estrogen Receptor-Beta Ligands. *J. Med. Chem.* **2004**, *47* (21), 5021–5040.
- (62) Mewshaw, R. E.; Edsall, R. J., Jr; Yang, C.; Manas, E. S.; Xu, Z. B.; Henderson, R. A.; Keith, J. C., Jr; Harris, H. A. ERbeta Ligands. 3. Exploiting Two Binding Orientations of the 2-Phenyl-naphthalene Scaffold to Achieve ERbeta Selectivity. *J. Med. Chem.* **2005**, *48* (12), 3953–3979.

- (63) Hillisch, A.; Peters, O.; Kosemund, D.; Müller, G.; Walter, A.; Schneider, B.; Reddersen, G.; Elger, W.; Fritzscheier, K.-H. Dissecting Physiological Roles of Estrogen Receptor Alpha and Beta with Potent Selective Ligands from Structure-Based Design. *Mol. Endocrinol.* **2004**, *18* (7), 1599–1609.
- (64) Richardson, T. I.; Dodge, J. A.; Durst, G. L.; Pfeifer, L. A.; Shah, J.; Wang, Y.; Durbin, J. D.; Krishnan, V.; Norman, B. H. Benzopyrans as Selective Estrogen Receptor Beta Agonists (SERBAs). Part 3: Synthesis of Cyclopentanone and Cyclohexanone Intermediates for C-Ring Modification. *Bioorg. Med. Chem. Lett.* **2007**, *17* (17), 4824–4828.
- (65) Norman, B. H.; Dodge, J. A.; Richardson, T. I.; Borromeo, P. S.; Lugar, C. W.; Jones, S. A.; Chen, K.; Wang, Y.; Durst, G. L.; Barr, R. J.; Montrose-Rafizadeh, C.; Osborne, H. E.; Amos, R. M.; Guo, S.; Boodhoo, A.; Krishnan, V. Benzopyrans Are Selective Estrogen Receptor Beta Agonists with Novel Activity in Models of Benign Prostatic Hyperplasia. *J. Med. Chem.* **2006**, *49* (21), 6155–6157.
- (66) Leung, Y.-K.; Mak, P.; Hassan, S.; Ho, S.-M. Estrogen Receptor (ER)-Beta Isoforms: A Key to Understanding ER-Beta Signaling. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103* (35), 13162–13167.
- (67) Veeneman, G.; Loozen, H.; Mestres, J.; De Zwart, E. 10-ARYL-11-HBENZO [b]FLUORENE DERIVATIVES AND ANALOGS FOR MEDICINAL USE. WO/2002/016316, March 1, 2002.
- (68) Shiau, A. K.; Barstad, D.; Radek, J. T.; Meyers, M. J.; Nettles, K. W.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A.; Agard, D. A.; Greene, G. L. Structural Characterization of a Subtype-Selective Ligand Reveals a Novel Mode of Estrogen Receptor Antagonism. *Nat. Struct. Biol.* **2002**, *9* (5), 359–364.


- (69) Knox, A. J. S.; Meegan, M. J.; Sobolev, V.; Frost, D.; Zisterer, D. M.; Williams, D. C.; Lloyd, D. G. Target Specific Virtual Screening: Optimization of an Estrogen Receptor Screening Platform. *J. Med. Chem.* **2007**, *50* (22), 5301–5310.
- (70) Cai, S. X.; Drewe, J. A.; Kasibhatla, S.; Kemnitzer, W. D.; Tseng, B. Y.; Blais, C.; Labrecque, D.; Gourdeau, H. Substituted 4-Aryl-Chromene as Activator of Caspases and Inducer of Apoptosis and as Antivascular Agent and the Use Thereof. 7968595, June 28, 2011.
- (71) Mountford, S. J.; Albiston, A. L.; Charman, W. N.; Ng, L.; Holien, J. K.; Parker, M. W.; Nicolazzo, J. A.; Thompson, P. E.; Chai, S. Y. Synthesis, Structure-Activity Relationships and Brain Uptake of a Novel Series of Benzopyran Inhibitors of Insulin-Regulated Aminopeptidase. *Journal of medicinal chemistry* **2014**, *57* (4), 1368–1377.
- (72) Cai, S. X.; Drewe, J.; Kemnitzer, W. Discovery of 4-Aryl-4H-Chromenes as Potent Apoptosis Inducers Using a Cell- and Caspase-Based Anti-Cancer Screening Apoptosis Program (ASAP): SAR Studies and the Identification of Novel Vascular Disrupting Agents. *Anti-Cancer Agent Me* **2009**, *9* (4), 437–456.
- (73) Kolla, S. R. L., Yong-Rok. Ca(OH)₂-Mediated Efficient Synthesis of 2-Amino-5-Hydroxy-4H-Chromene Derivatives with Various Substituents. *Tetrahedron* **2011**, *67* (43), 8271–8275.
- (74) Shestopalov, A. M.; Emelianova, Y. M.; Nesterov, V. N. One-Step Synthesis of Substituted 2-Amino-4H-Chromenes and 2-Amino-4H-Benzo[f]chromenes. Molecular and Crystal Structure of 2-Amino-3-Cyano-6-Hydroxy-4-Phenyl-4H-Benzo[f]chromene. *Russ Chem B+* **2002**, *51* (12), 2238–2243.
- (75) Kidwai, M.; Saxena, S.; Khan, M. K.; Thukral, S. S. Aqua Mediated Synthesis of Substituted 2-Amino-4H-Chromenes and in Vitro Study as Antibacterial Agents. *Bioorganic & medicinal chemistry letters* **2005**, *15* (19), 4295–4298.


- (76) Ballini, R.; Bosica, G.; Conforti, M. L.; Maggi, R.; Mazzacani, A.; Righi, P.; Sartori, G. Three-Component Process for the Synthesis of 2-Amino-2-Chromenes in Aqueous Media. *Tetrahedron* **2001**, *57* (7), 1395–1398.
- (77) Yadav, Y.; MacLean, E. D.; Bhattacharyya, A.; Parmar, V. S.; Balzarini, J.; Barden, C. J.; Too, C. K. L.; Jha, A. Design, Synthesis and Bioevaluation of Novel Candidate Selective Estrogen Receptor Modulators. *Eur J Med Chem* **2011**, *46* (9), 3858–3866.
- (78) Bardon, S.; Vignon, F.; Montcourrier, P.; Rochefort, H. Steroid Receptor-Mediated Cytotoxicity of an Antiestrogen and an Antiprogestin in Breast Cancer Cells. *Cancer research* **1987**, *47* (5), 1441–1448.
- (79) *Morph2 Server*; (<http://www2.molmovdb.org>).
- (80) Bhattacharya, D.; Nowotny, J.; Cao, R.; Cheng, J. 3Drefine: An Interactive Web Server for Efficient Protein Structure Refinement. *Nucleic Acids Res.* **2016**, *44* (W1), W406-409.
- (81) *FRED, v3.0.1*; OpenEye Scientific Software (www.eyesopen.com): Santa Fe, New Mexico, USA.
- (82) *MarvinView, v15.7.27*; Chemaxon (<http://www.chemaxon.com>): ChemAxon Kft. Záhony u. 7, Building HX 1031 Budapest, Hungary.
- (83) Pal, D.; Chakrabarti, P. Non-Hydrogen Bond Interactions Involving the Methionine Sulfur Atom. *J. Biomol. Struct. Dyn.* **2001**, *19* (1), 115–128.
- (84) Iwaoka, M.; Isozumi, N. Hypervalent Nonbonded Interactions of a Divalent Sulfur Atom. Implications in Protein Architecture and the Functions. *Molecules* **2012**, *17* (6), 7266–7283.
- (85) Haldosén, L.-A.; Zhao, C.; Dahlman-Wright, K. Estrogen Receptor Beta in Breast Cancer. *Molecular and Cellular Endocrinology* **2014**, *382* (1), 665–672.
- (86) Elagamey, A. G. A.; Eltaweel, F. M. A. A. Nitriles in Heterocyclic Synthesis - Synthesis of Condensed Pyrans. *Indian J Chem B* **1990**, *29* (9), 885–886.

- (87) Wiles, C. . W., P. ..Haswell, S. J. ..Pombo-Villar, E. The Preparation and Reaction of Enolates within Micro Reactors. *Tetrahedron* **2005**, *61* (45), 10757–10773.
- (88) *Accelrys Draw v4.1*; Biovia (<http://www.biovia.com>): 5005 Wateridge Vista Drive, San Diego, CA 92121 USA.
- (89) Hawkins, P. C. D.; Skillman, A. G.; Warren, G. L.; Ellingson, B. A.; Stahl, M. T. *OMEGA*, v2.5.1.4; OpenEye Scientific Software (www.eyesopen.com): Santa Fe, New Mexico, USA.
- (90) Hawkins, P. C. D.; Skillman, A. G.; Warren, G. L.; Ellingson, B. A.; Stahl, M. T. Conformer Generation with OMEGA: Algorithm and Validation Using High Quality Structures from the Protein Databank and Cambridge Structural Database. *J. Chem. Inf. Model.* **2010**, *50* (4), 572–584.
- (91) *UniProt (Universal Protein Resource)*; (<http://www.uniprot.org/>).
- (92) Roy, A.; Kucukural, A.; Zhang, Y. I-TASSER: A Unified Platform for Automated Protein Structure and Function Prediction. *Nat Protoc* **2010**, *5* (4), 725–738.
- (93) Zhang, Y. I-TASSER Server for Protein 3D Structure Prediction. *BMC Bioinformatics* **2008**, *9*, 40.
- (94) Bhattacharya, D.; Cheng, J. i3Drefine Software for Protein 3D Structure Refinement and Its Assessment in CASP10. *PLoS ONE* **2013**, *8* (7), e69648.
- (95) Bhattacharya, D.; Cheng, J. 3Drefine: Consistent Protein Structure Refinement by Optimizing Hydrogen Bonding Network and Atomic-Level Energy Minimization. *Proteins* **2013**, *81* (1), 119–131.
- (96) McGann, M. FRED Pose Prediction and Virtual Screening Accuracy. *J. Chem. Inf. Model.* **2011**, *51* (3), 578–596.
- (97) *Molecular Operating Environment (MOE)*, 2012.10; Chemical Computing Group (<http://www.chemcomp.com>): Chemical Computing Group Inc., 1010 Sherbooke St. West, Suite #910, Montreal, QC, Canada, H3A 2R7, 2015.


- (98) Sheldrick, G. M. S. An Integrated System for Data Collection, Processing, Structure Solution and Refinement. Software Reference Manual; Bruker AXS, Inc: Madison, WI. 2001.

For Table of contents only







Tamoxifen $R^1 = H$; $R^2 = CH_3$
 Endoxifen $R^1 = OH$; $R^2 = H$




Raloxifene $R = H$; $X = CO$
 Arzoxifene $R = CH_3$; $X = O$















Pipendoxifene (ERA923) $n=1$
 Bazedoxifene (TSE424) $n=2$




Lasofoxifene




Acolbifene (EM652)

ER α selective agonists**ER α selective antagonists****ER β selective agonists****ER β selective antagonists**



13 $R^1 = F$, $R^2 = H$

14 $R^1 = H$, $R^2 = CH_3$



15 Crolibulin





Figure 4: ORTEP representation of the X-Ray crystal structure of **22**, ellipsoids at 50% probability




Figure 5. Top ranked poses of Compound **35** in ER α (green) and ER β (grey) indicating key hydrogen bonding interactions.




Figure 6. Side-view of top ranked poses of Compound **35** in ER α (green) and ER β (grey) indicating shift in orientation due to Leu384 \rightarrow Met336 mutation.






Figure 7. Top ranked poses of Compound 42 (R Enantiomer) in ER α (green) and ER β (grey) indicating key interactions.



Scheme 1: Synthesis of 4-aryl-4H-chromenes **16-25** and cyanoacrylate esters **26-28**.


Reagents and conditions: Compounds **16-18**, Method A: (i) $\text{R}^3\text{-CH}_2\text{CN}$, ($\text{R}^3 = \text{CO}_2\text{CH}_3, \text{CO}_2\text{CH}_2\text{CH}_3, \text{CN}$), (solvent CH_2Cl_2 for **16**, solvent EtOH for **17, 18**), Et_3N , reflux, 4h. (6-56%). Compounds **19-24**, Method B: (i) $\text{R}^3\text{-CH}_2\text{CN}$, ($\text{R}^3 = \text{CO}_2\text{CH}_3, \text{CO}_2\text{CH}_2\text{CH}_3, \text{CN}$), solvent EtOH; (2-57%). Compound **25**, Method C: (i) $\text{CH}_3\text{CO}_2\text{CH}_2\text{CN}$, solvent CH_2Cl_2 , piperidine, microwave 5-30 min, (46%); (ii) H_2 , Pd/C, ethyl acetate, (46%); (iii) $\text{CH}_3\text{O}_2\text{CCH}_2\text{CN}$, 170°C , 40 min, (90%); Compounds **27, 28**: $\text{CH}_3\text{CH}_2\text{O}_2\text{CCH}_2\text{CN}$, MeOH, piperidine 4 h, (13-83%). (iv) Resorcinol, EtOH, piperidine, reflux, 4 h.








Scheme 1: Synthesis of 4-aryl-4H-chromenes **16-25** and cyanoacrylate esters **26-28**.

Reagents and conditions: Compounds **16-18**, Method A: (i) $\text{R}^3\text{-CH}_2\text{CN}$, ($\text{R}^3 = \text{CO}_2\text{CH}_3, \text{CO}_2\text{CH}_2\text{CH}_3, \text{CN}$), (solvent CH_2Cl_2 for **16**, solvent EtOH for **17, 18**), Et_3N , reflux, 4h. (6-56%). Compounds **19-24**, Method B: (i) $\text{R}^3\text{-CH}_2\text{CN}$, ($\text{R}^3 = \text{CO}_2\text{CH}_3, \text{CO}_2\text{CH}_2\text{CH}_3, \text{CN}$), solvent EtOH; (2-57%). Compound **25**, Method C: (i) $\text{CH}_3\text{CO}_2\text{CH}_2\text{CN}$, solvent CH_2Cl_2 , piperidine, microwave 5-30 min, (46%); (ii) H_2 , Pd/C, ethyl acetate, (46%); (iii) Compound **26**; $\text{CH}_3\text{O}_2\text{CCH}_2\text{CN}$, 170 °C, 40 min, (90%); Compounds **27, 28**: $\text{CH}_3\text{CH}_2\text{O}_2\text{CCH}_2\text{CN}$, MeOH, piperidine 4 h, (13-83%). (iv) Resorcinol, EtOH, piperidine, reflux, 4 h.

Table 1: ER α and ER β binding effects and antiproliferative activity in MCF-7 cells of selected 4-aryl-4*H*-chromenes



Compound	R ¹	R ²	ER α IC ₅₀ (μ M) ^{c,d}	ER β IC ₅₀ (μ M) ^{c,d}	β/α	Antiproliferative activity MCF-7 cells IC ₅₀ (μ M) ^{a,e}
16	H	-CO ₂ CH ₃	3.30 \pm 0.30	1.59 \pm 0.17	0.48	10.82 \pm 1.15
17	H	-CO ₂ CH ₂ CH ₃	2.36 \pm 1.62	0.61 \pm 0.05	0.26	>100
18	H	-CN	>10	8.39 \pm 3.32	0.84	>100
19	OH	-CO ₂ CH ₃	0.79 \pm 0.40	10.02 \pm 1.39	12.67	>100
20	OH	-CO ₂ CH ₂ CH ₃	nd	nd	-	>100
21	OCH ₂ C ₆ H ₅	-CO ₂ CH ₂ CH ₃	40.32	>100	-	4.5
22	OCH ₂ CH ₃	-CO ₂ CH ₂ CH ₃	nd	nd	-	>100
23	3,4-(OCH ₃) ₂	-CO ₂ CH ₂ CH ₃	nd	nd	-	>100
32		-CO ₂ CH ₃	0.02 \pm 0.02	0.35 \pm 0.25	14.31	19.60 \pm 20.78
33		CO ₂ CH ₂ CH ₃	1.41 \pm 0.40	3.10 \pm 1.57	2.21	27.29 \pm 23.78
34		-CN	0.89 \pm 0.19	>10	11.23	>50
35		-CO ₂ CH ₂ CH ₂ CH ₂ CH ₃	0.02 \pm 0.01	6.55 \pm 2.72	353.34	7.73 \pm 7.54
36		-CO ₂ C(CH ₃) ₃	0.01 \pm 0.01	0.67 \pm 0.40	49.49	10.03 \pm 0.48
37		-CO ₂ CH ₃	0.05 \pm 0.00	0.22 \pm 0.14	4.77	16.2 \pm 5.72
38		-CO ₂ CH ₂ CH ₃	0.02 \pm 0.00	0.06 \pm 0.02	2.56	8.75 \pm 4.19
39		-CN	1.47 \pm 0.44	10.42 \pm 8.57	7.10	23.21 \pm 2.81

40		-CO ₂ (CH ₂) ₃ CH ₃	0.03 ±0.03	2.35 ±0.47	87.29	31.82 ±30.54
41		-CO ₂ C(CH ₃) ₃	0.09 ±0.00	0.37 ±0.02	3.93	2.65 ± 0.15
42		-CO ₂ CH ₃	109.10 ±6.9	0.63 ±0.03	0.0054	>100
43		-CO ₂ CH ₂ CH ₃	0.22 ±0.02	2.44 ±0.73	11.02	18.79 ±21.29
44		-CN	13.97 ±1.66	>50	3.58	35.31 ± 10.30
Tamoxifen	-	-	0.070 ^c	0.170 ^c	2.43	4.12 ± 0.038 ^b

^aIC₅₀ values are half maximal inhibitory concentrations required to block the growth stimulation of MCF-7 cells. Values represent the mean ± S.E.M (error values x 10⁻⁶) for at least two experiments performed in triplicate.

^bThe IC₅₀ value obtained for Tamoxifen is 4.12 ± 0.038 μM is in good agreement with the reported IC₅₀ value for tamoxifen using the MTT assay on human MCF-7 cells⁷⁸. ^cThe ER binding values obtained are in agreement with the reported ER IC₅₀ binding data for tamoxifen (ERα 60.9 nM ERβ 188 nM, Panvera/Invitrogen). ^dValues are an average of at least nine replicate experiments, for ERα with typical standard errors below 15%, and six replicate experiments for ERβ, with typical standard errors below 15%. ^e0% cell death observed for compounds **32**, **33**, **37**, **38**, **43** and **44** in LDH assay at 10 μM concentration, 13.4% cell death for tamoxifen in LDH assay at 10 μM concentration.