

# In Vivo and Mechanistic Studies on Antitumor Lead 7-Methoxy-4-(2-methylquinazolin-4-yl)-3,4-dihydroquinoxalin-2(1H)-one and Its Modification as a Novel Class of Tubulin-Binding Tumor-Vascular Disrupting Agents

Mu-Tian Cui,<sup>†</sup> Li Jiang,<sup>†</sup> Masuo Goto,<sup>‡</sup> Pei-Ling Hsu,<sup>‡</sup> Linna Li,<sup>||</sup> Qi Zhang,<sup>||</sup> Lei Wei,<sup>†</sup> Shou-Jun Yuan,<sup>||</sup> Ernest Hamel,<sup>§</sup> Susan L. Morris-Natschke,<sup>‡</sup> Kuo-Hsiung Lee,<sup>\*,†,||,⊥</sup> and Lan Xie<sup>\*,†,‡</sup>

<sup>†</sup>Beijing Institute of Pharmacology and Toxicology, 27 Tai-Ping Road, Beijing 100850, China

<sup>‡</sup>Natural Products Research Laboratories, UNC Eshelman School of Pharmacy, University of North Carolina, Chapel Hill, North Carolina 27599, United States


<sup>||</sup>Beijing Institute of Radiation Medicine, 27 Tai-Ping Road, Beijing 100850, China

<sup>§</sup>Screening Technologies Branch, Developmental Therapeutics Program, Division of Cancer Treatment and Diagnosis, National Cancer Institute, Frederick National Laboratory for Cancer Research, National Institutes of Health, Frederick, Maryland 21702, United States

<sup>⊥</sup>Chinese Medicine Research and Development Center, China Medical University and Hospital, Taichung 40402, Taiwan


## S Supporting Information

**ABSTRACT:** 7-Methoxy-4-(2-methylquinazolin-4-yl)-3,4-dihydroquinoxalin-2(1H)-one (**2**), a promising anticancer lead previously identified by us, inhibited tumor growth by 62% in mice at 1.0 mg/kg without obvious signs of toxicity. Moreover, compound **2** exhibited extremely high antiproliferative activity in the NIH-NCI 60 human tumor cell line panel, with low to subnanomolar GI<sub>50</sub> values (10<sup>-10</sup> M level). It also showed a suitable balance between aqueous solubility and lipophilicity, as well as moderate metabolic stability in vivo. Mechanistic studies using Mayer's hematoxylin and eosin and immunohistochemistry protocols on xenograft tumor tissues showed that **2** inhibited tumor cell proliferation, induced apoptosis, and disrupted tumor vasculature. Moreover, evaluation of new synthetic analogues (**6a–6t**) of **2** indicated that appropriate 2-substitution on the quinazoline ring could enhance antitumor activity and improve druglike properties. Compound **2** and its analogues with a 4-(2-methylquinazolin-4-yl)-3,4-dihydroquinoxalin-2(1H)-one scaffold thus represent a novel class of tubulin-binding tumor-vascular disrupting agents (tumor-VDAs) that target established blood vessels in tumors.



## INTRODUCTION

During the past decade, a novel class of antitumor agents termed tumor-vascular disrupting agents (tumor-VDAs) have been investigated intensively.<sup>1,2</sup> Unlike angiogenesis inhibitors (AIs), tumor-VDAs target established tumor blood vessels and exert an almost immediate effect on these vessels, thus leading to a rapid vascular collapse and subsequent tumor cell death via extensive necrosis and apoptosis.<sup>3</sup> Due to the morphological vasculature difference between normal tissues and solid tumors, tumor-VDAs preferentially block the blood flow of solid tumors, while the blood flow in normal tissues remains relatively intact.<sup>4</sup> Currently, more than a dozen tumor-VDA drug candidates are undergoing clinical trials,<sup>5,6</sup> and the most advanced candidate, CA-4P, a phosphate prodrug of combretastatin A-4 (CA-4, Figure 1), is in phase III trials for anaplastic thyroid cancer and Phase II trials for non-small-cell lung cancer (NSCLC).<sup>7</sup> Meanwhile, many diverse small molecule tumor-VDAs with high antiproliferative activity have also been reported recently.<sup>8,9</sup> Even though the exact



**Figure 1.** Colchicine and some clinical trial VDA candidates.

mechanism(s) of action of tumor-VDAs are still not known, most of these compounds show common pharmacological characteristics, including inhibition of microtubule polymerization, competitive binding to the colchicine binding site on

**Received:** February 21, 2017

tubulin, and disruption of the tumor cell cytoskeleton, thus classifying them as tubulin-binding tumor-VDAs,<sup>4</sup> such as CA-4.

In our prior studies on novel antitumor agents,<sup>10–12</sup> we discovered a series of new *N*-aryl-1,2,3,4-tetrahydroquinoline compounds with high antiproliferative activity in cellular assays, exemplified by prior leads **1a** and **1b**<sup>11</sup> shown in Figure 2.




Figure 2. Leads and new derivatives.


Compounds **1a** and **1b** show high antiproliferative activity, with GI<sub>50</sub> values of 1.5–1.7 and 11–18 nM, respectively, and they also inhibit tubulin polymerization with IC<sub>50</sub> values of 0.93 and 1.0 μM, respectively. On the other hand, compared with **1a**, compound **1b** displays improved aqueous solubility, a better log *P* value, and greater metabolic stability. Subsequent lead optimization on the C-ring (skeleton hopping) led to the discovery of 7-methoxy-4-(2-methylquinazolin-4-yl)-3,4-dihydroquinoxalin-2(1*H*)-one (**2**) as a new lead compound.<sup>13</sup> Compared with the corresponding prior lead **1b**, compound **2**, which has a novel structural scaffold with an altered C-ring, exhibits more potent activities in cytotoxicity and tubulin assays and better drug-like properties (Figure 2), including its log *P* value (2.97, pH 7.4), aqueous solubility (8.28 μg/mL, pH 7.4), and metabolic stability in vitro (*t*<sub>1/2</sub> 55 min in human liver microsomes).<sup>13</sup> In the current study, we screened **2** in an extensive cellular panel (NCI-60 cell lines) to confirm the antitumor activity and determine its antitumor spectrum. Moreover, we also evaluated **2** in several in vivo systems, including a H-460 xenograft model and vascular disruption assays. Based on these results, the new scaffold in **2** was

confirmed to be important for enhanced antitumor activity. Apart from the scaffold hopping, our prior structure optimizations on lead **1a**<sup>12</sup> also revealed that the 2-substituent on the quinazoline ring was modifiable to improve druglike properties without loss of antitumor potency, especially with the introduction of an alkylamino side chain. Our prior modeling studies further demonstrated that both **1a** and **2** could dock within the tubulin–colchicine binding site and superimposed well with DAMA-colchicine (the bound ligand in the tubulin crystal structure, PDB code 1SA0).<sup>12,13</sup> In addition, the docking study showed that the 2-substituent on the quinazoline ring in series-1 analogues overlapped well with the 7-side chain in DAMA-colchicine (Figure 1), thus indicating that the quinazoline 2-position has sufficient chemical space for structural modification. Consequently, in our current study, a series of new 4-(2-substituted quinazolin-4-yl)-3,4-dihydroquinoxalin-2(1*H*)-one derivatives (**6a–6t**) were designed by maintaining the new scaffold in **2** and introducing various substituents on the quinazoline ring 2-position. Our aims were to simultaneously enhance antitumor activity and improve molecular drug-like properties. All synthesized new compounds were first evaluated in a human tumor cell line (HTCL) panel. Subsequently, new active compounds were further tested in tubulin assembly assays and assessed for their essential drug-like properties.

## CHEMISTRY


New compounds **6a–6p** with various 2-alkylamino or 2-alkoxy substituents on the quinazoline ring were synthesized as shown in Scheme 1. The commercially available 2,4-dichloroquinazoline was coupled with 4-methoxy-2-nitroaniline (**3**) in *i*-PrOH in the presence of a catalytic amount of HCl to obtain 2-chloro-4-(4-methoxy-2-nitrophenyl)aminoquinazoline (**4**) in 85% yield. The nitro group was then reduced to an amino group by using zinc powder in acetic acid at 0 °C. Reaction of this intermediate with 2-chloroacetyl chloride produced **5**. Next, compound **5** underwent an intramolecular cyclization in DMF in the presence of K<sub>2</sub>CO<sub>3</sub> to yield 4-(2-chloroquinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1*H*)-one (**6a**). Displacement of the 2-chloro group on the quinazoline ring of **6a** with various amines led to a series of 2-alkylamino-derivatives **6b–6n**. The reactions were performed in anhydrous *i*-PrOH

Scheme 1<sup>a</sup>



R see Table 3

<sup>a</sup>Reagents and conditions: (i) *i*-PrOH/HCl, rt, 5–6 h; (ii) Zn/CH<sub>3</sub>COOH, CH<sub>2</sub>Cl<sub>2</sub>, 0.5 h; (iii) ClCH<sub>2</sub>COCl/K<sub>2</sub>CO<sub>3</sub>, 0 °C, 1 h; (iv) K<sub>2</sub>CO<sub>3</sub>/DMF, 100 °C, 0.5 h; (v) amine, microwave (MW), *i*-PrOH, 60–120 °C, 15–30 min (for **6b–d**, **6f–g**, **6k**), or with H<sub>2</sub>SO<sub>4</sub>, 120 °C, 20 min (for **6e**, **6j**, **6l**, **6m**), or with K<sub>2</sub>CO<sub>3</sub>, 60–120 °C, 15–20 min (for **6h**, **6i**, **6n**); (vi) EtOH or cyclopropylcarbinol/NaH at 0 °C, then MW, 95 °C, 5 min (for **6o**, **6p**).

Scheme 2<sup>a</sup>

<sup>a</sup>Reagents and conditions: (i)  $\text{ClCH}_2\text{CN}/\text{HCl}$  (conc),  $90^\circ\text{C}$ , 24 h; (ii)  $\text{TsCl}$ , DMAP,  $\text{Et}_3\text{N}$ , rt, 3 h; (iii)  $i\text{-PrOH}$ ,  $45^\circ\text{C}$ , 6 h; (iv)  $\text{KOAc}$ ,  $55^\circ\text{C}$ , 6 h; (v)  $\text{Pd/C}$ ,  $\text{H}_2$ , rt, 2 h; (vi)  $\text{ClCH}_2\text{COCl}/\text{K}_2\text{CO}_3$ , acetone,  $0^\circ\text{C}$ , 1 h; (vii)  $\text{K}_2\text{CO}_3/\text{DMF}$ ,  $100^\circ\text{C}$ , 1 h; (viii)  $\text{NaOH}$  (10%), rt, 1 h; (ix)  $\text{Br}(\text{CH}_2)_3\text{COOEt}/\text{Cs}_2\text{CO}_3$ ,  $90^\circ\text{C}$ , 4 h; (x)  $\text{NH}_2\text{OH}\cdot\text{HCl}$ ,  $0^\circ\text{C}$ , 4 h.

Table 1. Antiproliferative Potencies ( $\text{GI}_{50}$ ,<sup>b</sup> nM) of 2 against the NCI-60 Panel<sup>a</sup>

Leukemia		Colon Cancer		Ovarian Cancer		CNS Cancer	
CCRF-CEM	0.551	COLO205	0.322	IGROV1	0.830	SF-268	0.788
HL-60(TB)	0.361	HCC-2998	0.371	OVCAR-3	0.266	SF-295	0.318
K562	0.359	HCT-116	0.369	OVCAR-4	2.43	SF-539	0.298
MOLT-4	0.605	HCT-15	0.413	OVCAR-5	>100	SNB-19	0.655
RPMI-8226	0.603	HT29	0.307	OVCAR-8	0.524	SNB-75	0.360
SR	0.301	KM12	0.353	NCI/ADR-RES	0.301	U251	0.437
<b>Non-Small-Cell Lung Cancer</b>		SW-620	0.487	SK-OV-3	0.409	<b>Prostate Cancer</b>	
A549/ATCC	0.588	<b>Melanoma</b>		<b>Renal Cancer</b>		PC-3	0.636
EKVX	0.632	LOX IMVI	0.526	786-0	3.46	DU-145	0.409
HOP-62	0.690	MALME-3M	>100	A498	0.376	<b>Breast Cancer</b>	
HOP-92	>100	M14	0.336	ACHN	0.926	MCF7	0.363
NCI-H226	3.03	MDA-MB-435	0.208	CAKI-1	0.497	MDA-MB-231/ATCC	0.914
NCI-H23	0.458	SK-MEL-28	>100	RXF-393	0.246	HS 578T	0.601
NCI-H322M	0.904	SK-MEL-5	0.400	SN12C	1.01	BT-549	2.10
NCI-H460	0.397	UACC-257	>100	TK-10	>100	T-47D	<sup>c</sup>
NCI-H522	0.294	UACC-62	0.432	UO-31	10.0	MDA-MB-468	0.415

<sup>a</sup>The data were provided by the NCI, USA. The TGI and  $\text{LC}_{50}$  values are  $>1.0 \times 10^{-7}$  M in most tested cell lines in the same assays (see SI). <sup>b</sup>The concentration that corresponds to 50% growth inhibition ( $\text{GI}_{50}$ ). <sup>c</sup>Not available.

under microwave (MW) heating at less than  $120^\circ\text{C}$  for 20 min in the presence of acidic or weakly basic catalysts, for example,  $\text{H}_2\text{SO}_4$  for **6e**, **6l**, and **6m** and  $\text{K}_2\text{CO}_3$  for **6h**, **6i**, **6k**, and **6n**. Compound **6f** was prepared in  $N,N$ -dimethylacetamide (DMA) at  $155^\circ\text{C}$ . The syntheses of 2-alkoxy-compounds **6o** and **6p** were performed in the presence of  $\text{NaH}$ , used to deprotonate and enhance the nucleophilicity of the alcohol reactant. Moreover, compounds **6q–6t**, which have a methylene ( $\text{CH}_2$ ) group inserted between the quinazoline ring and the remainder of the 2-side chain, were synthesized as shown in Scheme 2. Using a literature method,<sup>14</sup> 2-chloromethyl-3H-quinazolin-4-one (**8**) was prepared from the commercial


reagents methyl 2-aminobenzoate and 2-chloroacetonitrile in 85% yield. Treatment of **8** with  $p$ -toluenesulfonyl chloride in  $\text{CH}_2\text{Cl}_2$  in the presence of trimethylamine and a catalytic amount of dimethylaminopyridine (DMAP) produced the quinazoline sulfonate ester **9** in 88% yield. A substitution reaction between **9** and **3** gave intermediate **10**. The 2-chloromethyl on the quinazoline ring in **10** was converted to the 2-acetoxymethyl in intermediate **11**. Next, compound **6q** was obtained from **11** by successive catalytic hydrogenation, amidation, and intramolecular cyclization. Notably, the cyclization of intermediate **13** was performed in DMA, rather than DMF, in 99% yield. Compound **6q** was then hydrolyzed

under basic conditions to produce hydroxymethyl compound **6r** in 98% yield. Subsequently, compound **6r** was reacted with ethyl 4-bromobutanoate<sup>15</sup> in the presence of  $\text{Cs}_2\text{CO}_3$  to yield **6s** with an extended side chain on the quinazoline ring. Finally, compound **6s** was treated with hydroxylamine to produce hydroxamide compound **6t**. Compounds **6s** and **6t** were purified by recrystallization from EtOAc.

## RESULTS AND DISCUSSION

**Testing of Compound 2. NCI-60 Cell Line Screening.** To determine its antitumor spectrum, compound **2** was tested in the National Cancer Institute's cellular panel of 60 human tumor cell lines (NCI-60). As shown in Table 1, compound **2** exhibited excellent potency with low to subnanomolar  $\text{GI}_{50}$  values (at  $10^{-10}$  M level) against most tumor cell lines, including many multidrug resistant phenotypes, in the standard NCI testing protocol.<sup>16</sup> However, the inhibitory potency of **2** was much reduced against six tumor cell lines (HOP-92, MALME-3M, SK-MEL-28, UACC-257, OVCAR-5, and TK-10). Interestingly, MALME-3M and TK10 cell lines were also significantly resistant toward a tubulin polymerization inhibitory 4-substituted-5-methyl-furo[2,3-*d*]pyrimidine derivative.<sup>17</sup> These cell lines may be less sensitive to polymerization inhibitors, due to certain biological properties, such as overexpression or ligand binding site mutations on tubulin. Thus, **2** is an extremely potent antitumor agent with a broad spectrum of activity.

**Antitumor Activity in Vivo.** Next, compound **2** was evaluated in vivo due to its remarkable antiproliferative activity in vitro, as well as strong inhibitory effects on tubulin assembly, moderate in vitro metabolic stability ( $t_{1/2}$  55 min in human liver microsomes), and good aqueous solubility and lipophilicity. Pilot experiments showed that 4 mg/kg (one dose treatment) was the lethal dose of **2** in nude mice bearing MCF-7 xenograft tumors.<sup>13</sup> This result allowed us to better assess the in vivo pharmacokinetic properties of **2**. Consequently, the assessment was performed in SD rats ( $n = 3$ ) by intravenous (iv) administration at a single-dose of 1.0 mg/kg.<sup>18</sup> Compound **2** was relatively stable metabolically in rats with a terminal half-life of 52 h, providing an experimental basis for establishing administration schedules in a subsequent pharmacodynamic study. We established nude mouse subcutaneous xenograft models of NCI-H460 lung cancer and sorted the mice into five groups ( $n = 8$  animals per group) with almost equal mean tumor size ( $100 \text{ mm}^3$ ). The mice in three groups were treated with **2** by iv injection at doses of 1.0, 0.5, or 0.25 mg/kg every 5 days for 3 weeks. Control mice and paclitaxel-treated mice were treated in the same way, receiving vehicle solution or paclitaxel (15 mg/kg), respectively (Figure 3). All of the animals survived through the entire experiment, and no obvious toxic effects were observed, although there was some effect on animal weights with the two higher doses of **2** and with paclitaxel. At the end of treatment, the mice were sacrificed, and the tumors were recovered and weighed. As shown in Table 2 and Figure 3, compound **2** significantly inhibited the growth of the NCI-H460 xenograft tumors in a dose-dependent manner. The average tumor weights of **2**-treated groups were less than those of control mice (statistical significance). The tumor growth inhibition rates were 17.8%, 36.8%, and 61.9% at doses of 0.25, 0.5, and 1.0 mg/kg, respectively. Meanwhile, the reference group treated with paclitaxel at a 15 mg/kg dose showed a tumor inhibition rate of 60.4%, comparable to what was



**Figure 3.** Dose-dependent anticancer effects of compound **2** on NCI-H460 lung cancer xenografts in nu/nu mice. Treatments started when tumors reached a mean volume of around  $100 \text{ mm}^3$ . Mice were treated with **2** at an iv dose of 0.25, 0.5, or 1 mg/kg or paclitaxel at 15 mg/kg as reference or the vehicle as control ( $n = 8$ ), every 5 days for 3 weeks. (A) Growth difference of tumor volumes at different time points. (B) Images of resected NCI-H460 xenograft tumors. (C) Tumors were resected and weighed at the end of experiment, (●) indicates the weight value of each tumor; (—) indicates average value of tumor weights. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  vs vehicle controls (one-way analysis of variance with Tukey's post hoc method).

observed with **2** at the 1.0 mg/kg dose. Therefore, compound **2** had significant in vivo antitumor effects.

**Testing of New Derivatives (6a–6t) of Compound 2. Antiproliferative Activity in Cellular Assays and SAR Analysis.** The newly synthesized  $N^4$ -(2-substituted quinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one compounds (**6a**–



Table 2. Tumor Growth Inhibition of **2** on Human Lung H460 Xenograft Tumors in Nude Mice<sup>a</sup>

	dose (mg/kg)	animals start/end	body weight change			antitumor activity	
			start (g)	end (g)	%	tumor weight (g)	inhibition rate (%)
vehicle		8/8	22.50 ± 1.23	26.40 ± 1.43	17.42	2.71 ± 1.07	
<b>2</b>	0.25	8/8	22.44 ± 1.27	25.96 ± 1.54	15.71	2.23 ± 0.57	17.8
<b>2</b>	0.5	8/8	22.28 ± 1.33	23.85 ± 1.57	7.30	1.72 ± 0.78*	36.8
<b>2</b>	1.0	8/8	22.13 ± 1.24	20.50 ± 3.27	−7.23	1.03 ± 0.71***	61.9
paclitaxel	15	8/8	22.50 ± 1.29	23.59 ± 2.66	4.99	1.08 ± 0.78**	60.4


<sup>a</sup>Administration by iv. Schedule: Q5D × 4, that is, every 5 days for 3 weeks. Data are presented as the mean ± SD, \**p* < 0.05, \*\**p* < 0.01, \*\*\**p* < 0.001 vs control.

**6t**) with the same scaffold as **2** were initially evaluated in a HTCL panel, including A549 (lung carcinoma), KB (originally isolated from epidermoid carcinoma of the nasopharynx), multidrug-resistant (MDR) cell line KB-VIN (vincristine-resistant KB subline), MDA-MB-231 (triple-negative breast cancer), and MCF-7 (estrogen receptor-positive breast cancer),<sup>19,20</sup> in parallel with paclitaxel as an experimental control. The activity in the cellular assays was determined using the established sulforhodamine B (SRB) method.<sup>21</sup> The antiproliferative activities (GI<sub>50</sub>) and structures are summarized in Table 3.

As expected, 4-(2-chloroquinazo-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1*H*)-one (**6a**) exhibited significant antiproliferative activity with low nanomolar GI<sub>50</sub> values of 0.53–2.01 nM (cf, **1a**, GI<sub>50</sub> 1.5–1.7 nM, Figure 2). After the scaffold change in the C-ring from piperidine to piperazin-2-one, the potency increased 3–6-fold (compare prior leads **1a** and **1b** with new compounds **6a** and **2**, respectively), which supported our previous hypothesis that the lactam C-ring (3,4-dihydropyrazin-2(1*H*)-one) might be an important moiety to enhance antiproliferative potency.

In addition, other series-6 compounds with the same new scaffold but different substituents at the quinazoline 2-position showed significant antiproliferative activities in the tested HTCL panel with GI<sub>50</sub> values ranging from subnanomolar to submicromolar levels (Table 3). Among them, except for **6h** and **6k**, most 2-alkylamino-compounds with either a linear side chain (**6a**–**6e**) or a small cycloalkylamino substituent (**6f**, **6g**, **6i**, and **6j**) were highly potent with GI<sub>50</sub> values of 0.54–10.6 nM, with comparable or greater potency than paclitaxel in the same assays against chemosensitive cell lines. With GI<sub>50</sub> values of >10–80 nM, the 2-(*N*-pyrrolidinyl) (**6h**) and 2-(*N*-(3-methoxyazetidyl)) (**6k**) compounds exhibited lower potency than the above-mentioned compounds.

Moreover, differing results were obtained when a 2-arylamino substituent was introduced at the quinazoline 2-position. Compound **6l** with a 2-(4-cyanophenyl)amino substituent displayed high potency with low nanomolar GI<sub>50</sub> values (<10 nM), but compound **6m** with a 2-(3-cyanophenyl)-amino group displayed at least ten times decreased potency (GI<sub>50</sub> 61.5–384 nM), suggesting that the bulkiness of the 2-substituent, for example, a *m*-substituted phenyl moiety, might obstruct the small molecule from fitting into the binding pocket on its biological target. Additionally, compound **6n**, another 2-arylamino [2-(4-pyridinylmethyl)amino] substituted compound, showed high potency only against A549 and KB cell growth (GI<sub>50</sub> 6.62 and 7.50 nM, respectively) but decreased potency toward drug-resistant cell lines KB-VIN (GI<sub>50</sub> 48.8 nM) and MDA-MB-231 (GI<sub>50</sub> 73.6 nM), as well as MCF-7 (GI<sub>50</sub> 71.8 nM).

Table 3. Antiproliferative Activity of New Series Compounds (**6a**–**6t**) against Human Tumor Cell Lines


Compound	R	GI <sub>50</sub> (nM) ± SD <sup>a</sup>				
		A549	MDA-MB-231	KB	KB-VIN	MCF-7
<b>6a</b>	–Cl	0.53 ± 0.03	2.01 ± 0.73	0.59 ± 0.09	0.66 ± 0.17	0.76 ± 0.03
<b>6b</b>		1.01 ± 0.08	10.6 ± 1.71	1.68 ± 0.15	1.22 ± 0.10	0.84 ± 0.06
<b>6c</b>		0.57 ± 0.01	0.79 ± 0.01	0.54 ± 0.00	0.55 ± 0.01	0.84 ± 0.03
<b>6d</b>		5.25 ± 0.02	7.19 ± 0.01	5.25 ± 0.15	5.33 ± 0.00	7.88 ± 0.05
<b>6e</b>		5.18 ± 0.11	10.2 ± 0.68	4.92 ± 0.19	10.5 ± 0.42	7.43 ± 0.38
<b>6f</b>		1.12 ± 0.31	10.5 ± 0.94	2.11 ± 0.27	1.17 ± 0.24	1.08 ± 0.22
<b>6g</b>		7.67 ± 0.02	17.6 ± 1.77	6.92 ± 0.22	6.72 ± 0.39	7.67 ± 0.23
<b>6h</b>		45.0 ± 2.25	80.1 ± 4.62	33.2 ± 0.64	57.8 ± 0.79	NT
<b>6i</b>		5.71 ± 0.18	8.26 ± 0.36	4.66 ± 0.31	4.74 ± 0.17	8.31 ± 0.07
<b>6j</b>		5.47 ± 0.05	8.86 ± 0.52	4.76 ± 0.06	4.80 ± 0.06	8.59 ± 0.09
<b>6k</b>		17.5 ± 0.59	53.1 ± 3.73	8.26 ± 0.39	39.6 ± 0.51	8.60 ± 0.03
<b>6l</b>		6.17 ± 0.09	8.31 ± 0.07	5.39 ± 0.26	8.02 ± 0.25	9.19 ± 0.24
<b>6m</b>		75.4 ± 2.62	97.1 ± 2.77	61.5 ± 1.77	487 ± 20.0	384 ± 198
<b>6n</b>		6.62 ± 0.28	73.6 ± 4.84	7.50 ± 0.68	48.8 ± 0.86	71.8 ± 7.76
<b>6o</b>		16.7 ± 0.43	15.4 ± 0.38	11.6 ± 0.67	10.8 ± 0.9	8.59 ± 0.19
<b>6p</b>		15.8 ± 4.09	45.2 ± 1.71	14.2 ± 3.16	7.23 ± 0.15	7.88 ± 0.05
<b>6q</b>		5.52 ± 0.12	8.95 ± 0.65	5.05 ± 0.20	6.38 ± 0.37	33.0 ± 2.58
<b>6r</b>		5.35 ± 0.17	9.70 ± 0.32	5.29 ± 0.21	5.36 ± 0.16	10.8 ± 1.04
<b>6s</b>		202 ± 28.9	975 ± 51.2	280 ± 11.0	373 ± 42.9	1281 ± 63.2
<b>6t</b>		550 ± 18.0	1015 ± 27.7	307 ± 5.05	646 ± 15.7	1213 ± 59.2
<b>2</b> <sup>b</sup>	–Me	3.20 ± 0.00	NT	2.30 ± 0.00	2.20 ± 0.00	NT <sup>c</sup>
Paclitaxel <sup>d</sup>		7.03 ± 0.47	10.0 ± 0.37	5.12 ± 0.28	2716 ± 9.06	11.9 ± 0.71


<sup>a</sup>The GI<sub>50</sub> values are the concentrations corresponding to 50% cell growth inhibition and are expressed as the mean ± SD from the dose–response curves of at least three independent experiments. <sup>b</sup>Data published in ref 13. <sup>c</sup>NT: not tested. <sup>d</sup>Paclitaxel served as positive control in the same assays.

Next, we evaluated the presence of a 2-alkoxy or 2-alkoxymethyl substituent on the quinazoline ring. The corresponding 2-ethoxy (**6o**), 2-cyclopropylmethoxy (**6p**), 2-acetoxymethyl (**6q**), and 2-hydroxymethyl (**6r**) substituted compounds all exhibited high potency against tumor cell growth, generally showing single to double digit nanomolar GI<sub>50</sub> values in the HTCL panel. However, the presence of a longer linear 2-substituent (more than six atoms) reduced antiproliferative potency, see compounds **6s** and **6t** with GI<sub>50</sub> values of 0.2–1.28 and 0.31–1.21 μM, respectively.

**Table 4.** Drug-like Properties and Inhibition of Tubulin Polymerization and Colchicine Binding of Active Compounds 6a–6g, 6i–6j, and 6q–6r

	parameters (pH 7.4)		tubulin assembly <sup>a</sup> IC <sub>50</sub> (μM) ± SD	colchicine binding inhibition <sup>b</sup> (%)	
	aq. sol. (μg/mL)	log P		5 μM	1 μM
6a	13.4 ± 0.98	2.94 ± 0.15	1.60 ± 0.04	99 ± 1	88 ± 3
6b	28.6 ± 7.35	2.77 ± 0.40	2.80 ± 0.3	88 ± 0.9	53 ± 5
6c	2.74 ± 0.11	2.56 ± 0.10	0.63 ± 0.09	96 ± 1	78 ± 4
6d	1.34 ± 0.10	3.91 ± 0.03	0.42 ± 0.01	96 ± 1	83 ± 0.8
6e	80.9 ± 17.8	2.60 ± 0.27	1.90 ± 0.04	77 ± 0.5	<sup>c</sup>
6f	67.8 ± 10.3	2.26 ± 0.13	2.90 ± 0.3	87 ± 0.7	47 ± 4
6g	1.45 ± 0.12	2.33 ± 0.00	0.59 ± 0.01	70 ± 2	<sup>c</sup>
6i	0.46 ± 0.01	2.48 ± 0.17	0.97 ± 0.1	90 ± 1	43 ± 4
6j	0.58 ± 0.00	2.14 ± 0.01	0.48 ± 0.02	97 ± 2	85 ± 0.3
6q	8.07 ± 0.47	2.70 ± 0.01	0.62 ± 0.05	95 ± 0.9	67 ± 1
6r	19.1 ± 1.58	2.14 ± 0.01	0.58 ± 0.01	97 ± 0.7	75 ± 4
2 <sup>d</sup>	11.1 ± 0.30	2.63 ± 0.00	0.77 ± 0.07	99 ± 0.02	93 ± 0.8
CA-4 <sup>e</sup>			0.54 ± 0.06	98 ± 1	82 ± 2

<sup>a</sup>The tubulin assembly assay measured the extent of assembly of 10 μM tubulin after 20 min at 30 °C. <sup>b</sup>Inhibition of [<sup>3</sup>H]colchicine binding was performed by incubating with tested compounds at 5 or 1 μM for 10 min at 37 °C, with tubulin at 1.0 μM. <sup>c</sup>Not determined. <sup>d</sup>Tubulin assay data see ref 13. <sup>e</sup>Reference compound is a drug candidate in phase II/III clinical trials.

**Figure 4.** Predicted binding mode of 2 (brown stick) and 6e (green stick) with tubulin (PDB code SLYJ) and overlapping with CA-4 (pink, the bound of ligand of SLYJ). Surrounding amino acid side chains are shown in gray stick format and are labeled. Hydrogen bonds are shown by green dashed lines, and the distance between ligands and protein is less than 3 Å.

In general, most 6-series compounds showed similar potency against A549, MCF-7, KB, and resistant KB-VIN cell lines, but were less potent against the triple-negative breast cancer cell line MDA-MB-231. Because antiproliferative activity was mostly comparable against the KB and MDR KB-VIN cell lines, the new compounds are likely not substrates of P-glycoprotein (Table 3).

Overall, the current results demonstrated that the new structural scaffold, 4-(quinazolin-4-yl)-3,4-dihydroquinoxalin-2(1H)-one, is very important for high anticancer activity, while a suitable 2-substituent on the quinazoline ring can enhance or maintain pharmacological activity. The new SAR

correlations revealed that (1) linear alkyamino substituents are more favorable than alkoxy and arylamino groups, which is a consistent finding with our prior results,<sup>12</sup> (2) H-bond donor groups (i.e., NH or OH) favor high antiproliferative potency, and (3) the length and bulky volume of the 2-substituent should be limited.


**Assessments of Physicochemical Properties.** Because potential drug candidates should have a good balance between potency and druglike properties, molecular physicochemical properties of 11 (6a–6g, 6i, 6j, 6q, and 6r) of our highly potent compounds (GI<sub>50</sub> ≤ 10 nM) were further assessed in parallel with 2. Aqueous solubility and log P values were measured at

pH 7.4 according to methods described previously.<sup>22</sup> As shown in Table 4, six compounds (**6a**, **6b**, **6e**, **6f**, **6q**, and **6r**) showed moderate aqueous solubility ranging from 8 to 87  $\mu\text{g/mL}$ , which were similar to or improved compared with that of **2** (11  $\mu\text{g/mL}$ ). Among them, **6b**, **6e**, **6f**, and **6r** have at least one H-bond donor group (NH or OH) in the quinazoline 2-substituent, thus indicating that H-bond donor groups favor increased aqueous solubility. On the other hand, lipophilicity is another major determinant of many ADME/Tox properties, as well as of pharmacological activity. Log *P* values were also measured at pH 7.4 to estimate molecular lipophilicity. Except for **6d**, all compounds in Table 4 showed lower log *P* values than those of **1a** and **1b** (4.13 and 3.65, respectively) and fell into a desirable druglike range ( $1 < \log P_{7.4} < 3$ ), which might lead to increased intestinal absorption. Thus, compound **2** and its new derivatives reach an acceptable balance between solubility and passive diffusion permeability, as well as increased metabolic stability.<sup>23</sup> Therefore, our current results indicate that the new structural scaffold in **2** and the series-6 compounds makes a major contribution to reduced molecular lipophilicity, while the changeable quinazoline 2-substituents can enhance anticancer activity and improve aqueous solubility.

**Inhibition of Tubulin Polymerization and Colchicine Binding.** To measure effects of the new compounds on microtubule polymerization, the same 11 highly active compounds (**6a-g**, **6i-j**, and **6q-r**;  $\text{GI}_{50} \leq 10 \text{ nM}$ ) were further tested in tubulin assembly and colchicine binding assays<sup>24</sup> in parallel with CA-4, a clinical trial drug candidate and highly potent competitive inhibitor of the binding of colchicine to tubulin.<sup>25</sup> Consistent with our previous results, all of the compounds were highly active in the tubulin assembly assay with low to submicromolar inhibitory  $\text{IC}_{50}$  values (0.42–2.90  $\mu\text{M}$ ), comparable with those of **2** (0.77  $\mu\text{M}$ ) and CA-4 (0.54  $\mu\text{M}$ ) in the same assay. However, an obvious difference was seen in the competitive colchicine–tubulin binding assay. With more than 97% and 83% inhibition at 5 and 1  $\mu\text{M}$ , respectively, compounds **6a**, **6d**, and **6j** showed similar potency to CA-4. The eight remaining series-6 compounds were less potent than CA-4. Moreover, compound **2** was more potent (99% at 5  $\mu\text{M}$  and 93% at 1  $\mu\text{M}$ ) than CA-4 in the colchicine binding inhibition assay. These results demonstrated that active series-6 compounds, with the same scaffold as **2**, also inhibit tubulin polymerization by targeting the tubulin–colchicine binding site; however, different 2-substituents on the quinazoline affect the molecular affinity with tubulin. Next, compounds **6e** and **2** were docked at the colchicine binding site with the tubulin crystal structure (PDB code 5LYJ). As shown in Figure 4, molecules **6e** and **2** superimposed well with the original ligand CA-4 in the binding site, while the outspread C2-side chain in **6e** formed two hydrogen bonds with a.a. Tyr202 and Asp251. These findings support the postulate that, similar to CA-4, this compound series targets the tubulin–colchicine binding site.


As representatives, compounds **6d** and **6e** were further investigated for their effects on the cell cycle (Figure 5). Similarly to CA-4, both compounds induced cell cycle arrest at the G2/M phase. Therefore, this series of 4-(quinazolin-4-yl)-3,4-dihydroquinoxalin-2(1H)-one derivatives, including **2** and active series-6, have been identified to have characteristics similar to those of CA-4 and, like CA-4, have potential as new antitumor agents.

**Testing of Compound 2 and New Derivatives (6a–t) for Antivascular Effects in Vivo.** Our prior and current studies have revealed that compound **2**<sup>13</sup> and its analogues



**Figure 5.** Effects of compounds on cell cycle. A549 cells were treated for 24 h with **6d** or **6e** at 15 nM. CA-4 (20 nM) or DMSO was used as a tubulin polymerization inhibitor or control, respectively. Fixed and propidium iodide (PI)-stained cells were analyzed by flow cytometry.

series-6 derivatives not only exhibit high antiproliferative activity but also inhibit tubulin polymerization and competitive colchicine binding to tubulin (Table 4), as well as arrest cells in the G2/M phase of the cell cycle, indicating that this compound type has characteristics similar to those of CA-4. Thus, we postulated that they might act as new tubulin-binding tumor-VDAs. The antivascular activity of **2** was evaluated on a xenograft tumor tissue section by staining with hematoxylin–eosin (HE) and by immunohistochemistry (IHC) using antibodies to CD31, cleaved caspase-3, and Ki67 (Figure 6).



**Figure 6.** Compound **2** treatment resulted in apoptosis and vascular disruption of NCI-H460 xenograft tumors. Representative images of HE staining and IHC staining of CD31 (endothelial marker), cleaved caspase-3 (apoptosis marker), and Ki67 (proliferation marker) in different treatment groups. Sections were counterstained with hematoxylin. Red arrows in CD31 staining images indicate microvessels. Scale bar, 100  $\mu\text{m}$ .



HE staining showed that the nuclei of tumor cells in vehicle controls were large and hyperchromatic, while the nuclei of 2-treated tumor cells became pyknotic. IHC staining of CD31 (endothelial marker) further demonstrated that the numbers and, especially, the sizes of the vessels in the 2-treated groups were reduced (Figure 6). In addition, 2-treated tumors had stronger cleaved caspase-3 (apoptosis marker) staining and weaker Ki67 (proliferation marker) staining, compared to control xenograft tumors. All of these results confirmed that 2 can inhibit proliferation, induce apoptosis, and disrupt tumor vasculature. Therefore, our current study results validate that, like CA-4, compound 2 is a new tubulin-binding tumor-VDA. Most likely, the 6-series compounds, which are analogues of 2, should belong to the same class of new antitubulin agents.

## CONCLUSION

In this study, we have identified 7-methoxy-4-(2-methylquinazolin-4-yl)-3,4-dihydroquinoxalin-2(1H)-one (2) as a new tubulin-binding tumor-VDA. Compound 2 showed significant dose-dependent antitumor efficacy in vivo in nude mouse H460 subcutaneous xenograft models, suppressing tumor growth by 61.9% at a dose of 1 mg/kg by iv injection every 5 days for 3 weeks without obvious signs of toxicity. It also displayed high antitumor potency in the NIH-NCI60 panel in vitro with low nanomolar GI<sub>50</sub> values and reasonable druglike properties, such as an acceptable balance between aqueous solubility and lipophilicity, and moderate metabolic stability in vitro and in vivo. Therefore, compound 2 should be a promising antitumor drug candidate as a new tumor-VDA and merits further development for clinical treatment of various tumors and cancers. With the same scaffold, a new compounds series (6) also exhibited high antiproliferative potency (GI<sub>50</sub> < 10 nM) and inhibited tubulin polymerization comparably to paclitaxel and CA-4 in corresponding assays. Thus, these compounds most likely also act as a new class of tubulin-binding tumor-VDA. New SAR results indicated that (1) the 2-substituent on the quinazoline ring can be modified to enhance antitumor activity and improve druglike properties, (2) the presence of a H-bond donor in the 2-substituent is beneficial to antitumor activity and aqueous solubility, and (3) the length and bulk of 2-substituents should be restricted.

## EXPERIMENTAL SECTION

**Chemistry.** The proton and carbon nuclear magnetic resonance (<sup>1</sup>H NMR and <sup>13</sup>C NMR) spectra were measured on a JNM-ECA-400 (400 MHz) spectrometer using tetramethylsilane (TMS) as internal standard and DMSO as solvent unless otherwise indicated. Mass spectra (MS) and high resolution mass spectra (HRMS) were measured on an API-150 mass spectrometer (ABI, Inc.) and Waters Xevo G2 with an electrospray ionization source, respectively. Melting points were measured with a SGW X-4 Micro-Melting point detector without correction. The MW reactions were performed on a MW reactor from Biotage, Inc. Medium-pressure column chromatography was performed using a CombiFlash Companion system from ISCO, Inc. Thin-layer chromatography (TLC) and preparative TLC were performed on silica gel GF<sub>254</sub> plates. Silica gel GF<sub>254</sub> was from Qingdao Haiyang Chemical Company. All commercial chemical reagents were purchased from Beijing Chemical Works or InnoChem, Inc. Pooled human liver microsomes (lot no. 20150323H) were purchased from iPhase Bioscience (Beijing) Ltd. Reagents. NADPH, MgCl<sub>2</sub>, KH<sub>2</sub>PO<sub>4</sub>, K<sub>2</sub>HPO<sub>4</sub>, reference compounds, and propranolol, as well as HPLC grade acetonitrile (ACN) and methanol (MeOH), were purchased from Sigma-Aldrich. Purities of target compounds were determined by HPLC using the following instruments and conditions: Agilent HPLC-1200 with UV detector and an Agilent Eclipse XDB-C18 column (150

mm × 4.6 mm, 5 μm), flow rate 0.8 mL/min, UV detection at 254 nm, and injection volume of 15 μL. Mobile elution was conducted with a mixture of solvents A and B (condition 1, ACN/H<sub>2</sub>O; condition 2, MeOH/H<sub>2</sub>O). Target compounds were found to be at least 95% pure in the above two HPLC analytical conditions.

**2-Chloro-4-(4-methoxy-2-nitro-phenyl)aminoquinazoline (4).** A mixture of 2,4-dichloro-quinazoline (1 g, 5.1 mmol) and 4-methoxy-2-nitroaniline (3, 0.86 g, 5.1 mmol) in anhydrous *i*-PrOH (50 mL) with a catalytic amount of HCl (conc, 10 drops) was stirred at room temperature (rt) for 5–6 h and monitored by TLC until the reaction was complete. The light yellow solid was removed by filtration and washed with a small amount of *i*-PrOH. Next, the collected solid was stirred in a saturated aqueous NaHCO<sub>3</sub> solution, refiltered, washed with water until pH = 7, and dried to obtain 4 as an orange solid (1.42 g, 85% yield). Mp 214–215 °C; <sup>1</sup>H NMR δ 3.93 (3H, s, OCH<sub>3</sub>), 7.44 (1H, dd, *J* = 9.0 and 2.8 Hz, ArH-5'), 7.57 (1H, d, *J* = 2.8 Hz, ArH-3'), 7.59 (1H, d, *J* = 9.0, ArH-6'), 7.65 (1H, t, *J* = 7.6 Hz, ArH-6), 7.70 (1H, d, *J* = 7.6 Hz, ArH-5), 7.92 (1H, t, *J* = 7.6 Hz, ArH-7), 8.46 (1H, d, *J* = 7.6 Hz, ArH-8), 10.61 (1H, s, NH); MS *m/z* (%) 331 (M + 1, 100), 333 (M + 3, 33).

**4-(2-Chloroacetyl-amino-4-methoxyphenyl)amino-2-chloroquinazoline (5).** A mixture of 4 (330 mg, 1 mmol) and zinc powder (660 mg, 1 mmol) in 50 mL of CH<sub>2</sub>Cl<sub>2</sub> in the presence of 0.4 mL of HOAc was stirred at 0 °C for 0.5 h. After the solid was removed, the filtrate was concentrated to obtain the amine product. It was immediately dissolved in acetone (25 mL), anhydrous K<sub>2</sub>CO<sub>3</sub> (138 mg, 1 mmol) was added, and the mixture was cooled to 0 °C. Chloroacetyl chloride (0.1 mL, excess) was dropped slowly into the mixture, which was kept at 0 °C with stirring for another 1–2 h. Then the mixture was poured into ice–water. The precipitated solid was removed by filtration, washed successively with water to neutral and brine, and dried to obtain 5 as a pink solid (210 mg, 56% yield). Mp 199–200 °C; <sup>1</sup>H NMR δ 3.78 (3H, s, OCH<sub>3</sub>), 4.23 (2H, s, CH<sub>2</sub>), 6.83 (1H, dd, *J* = 8.4 and 2.8 Hz, PhH-5'), 7.34 (1H, d, *J* = 2.8 Hz, PhH-3'), 7.48 (1H, d, *J* = 8.4 Hz, PhH-6'), 7.59 (1H, t, *J* = 7.6 Hz, ArH-6), 7.67 (1H, d, *J* = 7.6 Hz, ArH-5), 7.83 (1H, t, *J* = 7.6 Hz, ArH-7), 8.41 (1H, d, *J* = 7.6 Hz, ArH-8), 9.57 (1H, s, NH), 10.87 (1H, br, NH); MS *m/z* (%) 377 (M + 1, 100), 379 (M + 3, 58).

**4-(2-Chloroquinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6a).** A mixture of 5 (380 mg, 1 mmol) and anhydrous K<sub>2</sub>CO<sub>3</sub> (138 mg, 1 mmol) in DMF (10 mL) was heated to 100 °C for 1 h until the reaction was complete as judged by TLC monitoring. The mixture was poured into ice–water, and the solid product 6a was removed by filtration, washed with water, and dried to give 333 mg in a 98% yield. Yellow solid; mp 250–251 °C; <sup>1</sup>H NMR δ 3.74 (3H, s, OCH<sub>3</sub>), 4.01 (2H, s, CH<sub>2</sub>), 6.42 (1H, dd, *J* = 8.8 and 2.8 Hz, ArH-6'), 6.65 (1H, d, *J* = 2.8 Hz, ArH-8'), 6.84 (1H, d, *J* = 8.8 Hz, ArH-5'), 7.35 (1H, t, *J* = 8.4 Hz, ArH-6), 7.41 (1H, d, *J* = 8.4 Hz, ArH-5), 7.77 (1H, t, *J* = 8.4 Hz, ArH-7), 7.78 (1H, d, *J* = 8.4 Hz, ArH-8), 10.87 (1H, s, NH); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 51.7, 55.9, 102.6, 108.2, 115.6, 122.2, 122.8, 126.3, 126.6, 128.0, 132.7, 134.7, 153.4, 155.8, 157.8, 161.2, 167.8; MS *m/z* (%) 341 (M + 1, 100), 343 (M + 3, 34); HRMS *m/z* calcd for C<sub>17</sub>H<sub>14</sub>ClN<sub>4</sub>O<sub>2</sub> [M + H]<sup>+</sup> 341.0805, found 341.0867, 343.0843; HPLC purity 98.2%.

**7-Methoxy-4-(2-(methylamino)quinazolin-4-yl)-3,4-dihydroquinoxalin-2(1H)-one (6b).** A mixture of 6a (100 mg, 0.29 mmol) and methylamine (1 mL) was heated via MW to 80 °C for 20 min. The mixture was poured into ice–water, and the pH was adjusted to 7 with dilute aq. HCl. The solid was collected, washed with water, and dried to give pure 6b as a yellow solid (59 mg, 61% yield). Mp 274–275 °C; <sup>1</sup>H NMR δ 2.85 (3H, d, *J* = 4.8 Hz, NCH<sub>3</sub>), 3.67 (3H, s, OCH<sub>3</sub>), 4.30 (2H, s, CH<sub>2</sub>), 6.38 (1H, dd, *J* = 8.8 and 2.8 Hz, ArH-6), 6.52 (1H, d, *J* = 8.8 Hz, ArH-5), 6.61 (1H, d, *J* = 2.8 Hz, ArH-8), 6.85 (1H, t, *J* = 8.0 Hz, ArH-6'), 7.05 (1H, q, *J* = 4.8 Hz, NH-2'), 7.18 (1H, d, *J* = 8.0 Hz, ArH-5'), 7.39 (1H, d, *J* = 8.0 Hz, ArH-8'), 7.47 (1H, t, *J* = 8.0 Hz, ArH-7'), 10.73 (1H, s, NH); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 28.5, 51.5, 55.8, 102.6, 108.0, 120.8, 124.9, 126.2(x2), 131.7, 133.4 (x2), 154.5, 156.4 (x2), 159.9, 160.5, 168.4; MS *m/z* (%) 336 (M + 1, 100); HRMS *m/z* calcd for C<sub>18</sub>H<sub>18</sub>N<sub>5</sub>O<sub>2</sub> [M + H]<sup>+</sup> 336.1460, found 336.1501; HPLC purity 97.5%.



**General Procedure for Preparing Compounds 6c–6n under MW Irradiation.** A mixture of **6a** and an amine (excess) in anhydrous *i*-PrOH was heated under MW irradiation for 15–30 min. The reaction was monitored by TLC or LC-MS until complete. The mixture was poured into ice–water and was neutralized with dilute HCl (ca.10%) to pH 7. Precipitated solid was collected, washed with water, and dried to obtain the corresponding products.

**4-(2-(Ethylamino)quinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6c).** A mixture of **6a** (50 mg, 0.14 mmol) and ethylamine (65–75%, 1 mL, excess) in anhydrous *i*-PrOH (1 mL) was heated to 80 °C under MW irradiation for 30 min to obtain pure **6c** as a yellow solid (25 mg, 48% yield). Mp 277–278 °C; <sup>1</sup>H NMR δ 1.13 (3H, t, *J* = 7.2 Hz, CH<sub>3</sub>), 3.34 (2H, q, *J* = 7.2 Hz, NCH<sub>2</sub>), 3.67 (3H, s, OCH<sub>3</sub>), 4.31 (2H, s, CH<sub>2</sub>), 6.37 (1H, dd, *J* = 8.8 and 2.8 Hz, ArH-6), 6.51 (1H, d, *J* = 8.8 Hz, ArH-5), 6.61 (1H, d, *J* = 2.8 Hz, ArH-8), 6.84 (1H, t, *J* = 8.0 Hz, ArH-6'), 7.07 (1H, bs, NH), 7.16 (1H, d, *J* = 8.0 Hz, ArH-5'), 7.36 (1H, d, *J* = 8.0, ArH-8'), 7.46 (1H, t, *J* = 8.0 Hz, ArH-7'); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 15.3, 36.0, 51.5, 55.8, 102.6, 108.0, 112.7, 120.7, 120.8, 124.8, 126.2(x2), 131.7, 133.4, 154.4, 156.4, 159.2, 160.6, 168.4; MS *m/z* (%) 350 (*M* + 1, 100); HRMS *m/z* calcd for C<sub>19</sub>H<sub>20</sub>N<sub>5</sub>O<sub>2</sub> [*M* + H]<sup>+</sup> 350.1617, found 350.1626; HPLC purity 96.2%.

**4-(2-(*n*-Butylamino)quinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6d).** A mixture of **6a** (60 mg, 0.18 mmol) and *n*-butylamine (1 mL, excess) in anhydrous *i*-PrOH (1 mL) was heated to 120 °C under MW irradiation for 20 min. The mixture was poured into ice–water and was neutralized with dilute HCl (ca.10 mL) to pH 7. The mixture was extracted with EtOAc three times. The combined organic phases were washed with water and brine successively. After solvent was removed under reduced pressure, residue was washed with a small amount of cold acetone to give pure **6d** as a yellow solid (20 mg, 29% yield). Mp 204–205 °C; <sup>1</sup>H NMR δ 0.87 (3H, t, *J* = 7.6 Hz, CH<sub>3</sub>), 1.32 and 1.52 (each 2H, m, *J* = 7.6 Hz, CH<sub>2</sub>), 3.31 (2H, m, NCH<sub>2</sub>), 3.67 (3H, s, OCH<sub>3</sub>), 4.31 (2H, s, CH<sub>2</sub>), 6.38 (1H, dd, *J* = 8.8 and 2.8 Hz, ArH-6), 6.51 (1H, d, *J* = 8.8 Hz, ArH-5), 6.61 (1H, d, *J* = 2.8 Hz, ArH-8), 6.83 (1H, t, *J* = 8.0 Hz, ArH-6'), 7.08 (1H, m, NH), 7.16 (1H, d, *J* = 8.0 Hz, ArH-5'), 7.36 (1H, d, *J* = 8.0 Hz, ArH-8'), 7.46 (1H, t, *J* = 8.0 Hz, ArH-7'), 10.73 (1H, s, NH); MS *m/z* (%) 378 (*M* + 1, 100); HRMS *m/z* calcd for C<sub>21</sub>H<sub>24</sub>N<sub>5</sub>O<sub>2</sub> [*M* + H]<sup>+</sup> 378.1930, found 378.2017; HPLC purity 97.8%.

**4-(2-(3-Hydroxypropylamino)quinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6e).** A mixture of **6a** (100 mg, 0.29 mmol) and 3-aminopropanol (1 mL, excess) in anhydrous *i*-PrOH (0.5 mL) with one drop of H<sub>2</sub>SO<sub>4</sub> (conc) was heated to 120 °C under MW irradiation for 20 min to obtain **6e** (75% yield, white solid), mp 179–180 °C; <sup>1</sup>H NMR δ 1.69 (2H, m, CH<sub>2</sub>), 3.38 and 3.45 (each 2H, m, OCH<sub>2</sub> and NCH<sub>2</sub>), 3.67 (3H, s, OCH<sub>3</sub>), 4.31 (2H, s, CH<sub>2</sub>), 4.59 (1H, br, OH), 6.38 (1H, dd, *J* = 9.2 and 2.8 Hz, ArH-6), 6.53 (1H, d, *J* = 9.2 Hz, ArH-5), 6.60 (1H, d, *J* = 2.8 Hz, ArH-8), 6.85 (1H, t, *J* = 8.0 Hz, ArH-6'), 7.06 (1H, t, *J* = 5.6 Hz, NH), 7.17 (1H, d, *J* = 8.0 Hz, ArH-5'), 7.35 (1H, d, *J* = 8.0 Hz, ArH-8'), 7.47 (1H, t, *J* = 8.0 Hz, ArH-7'); MS *m/z* (%) 380 (*M* + 1, 100); HRMS *m/z* calcd for C<sub>20</sub>H<sub>22</sub>N<sub>5</sub>O<sub>3</sub> [*M* + H]<sup>+</sup> 380.1723, found 380.1730; HPLC purity 99.1%.

**4-(2-(Cyclopropylamino)quinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6f).** A mixture of **6a** (100 mg, 0.29 mmol) and cyclopropylamine (1 mL, excess) in DMA (2 mL) was heated to 155 °C under MW irradiation for 20 min to give pure **6f** as a yellow solid (68 mg, 65% yield). Mp 269–270 °C; <sup>1</sup>H NMR δ 0.49 and 0.65 (each 2H, m, CH<sub>2</sub>), 2.81 (1H, m, CH), 3.67 (3H, s, OCH<sub>3</sub>), 4.32 (2H, s, CH<sub>2</sub>), 6.38 (1H, dd, *J* = 8.8 and 2.8 Hz, ArH-6), 6.54 (1H, d, *J* = 8.8 Hz, ArH-5), 6.61 (1H, d, *J* = 2.8 Hz, ArH-8), 6.87 (1H, t, *J* = 8.0 Hz, ArH-6'), 7.15 (1H, d, *J* = 8.0 Hz, ArH-5'), 7.29 (1H, m, NH), 7.41 (1H, d, *J* = 8.0 Hz, ArH-8'), 7.48 (1H, t, *J* = 8.0 Hz, ArH-7'), 10.72 (1H, s, NH); MS *m/z* (%) 362 (*M* + 1, 100); HRMS *m/z* calcd for C<sub>20</sub>H<sub>20</sub>N<sub>5</sub>O<sub>2</sub> [*M* + H]<sup>+</sup> 362.1617, found 362.1658; HPLC purity 95.0%.

**4-(2-(Cyclopentylamino)quinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6g).** A mixture of **6a** (100 mg, 0.29 mmol) and cyclopentylamine (1 mL, excess) in *i*-PrOH (1 mL) was heated to 120

°C under MW irradiation for 30 min to give pure **6g** as a yellow solid (70 mg, 62% yield). Mp 268–269 °C; <sup>1</sup>H NMR δ 1.51 (4H, m, CH<sub>2</sub> × 2), 1.66 (2H, m, CH<sub>2</sub>), 1.90 (2H, m, CH<sub>2</sub>), 3.67 (3H, s, OCH<sub>3</sub>), 4.27 (1H, m, NCH), 4.31 (2H, s, CH<sub>2</sub>), 6.36 (1H, dd, *J* = 8.8 and 2.8 Hz, ArH-6), 6.51 (1H, d, *J* = 8.8 Hz, ArH-5), 6.61 (1H, d, *J* = 2.8 Hz, ArH-8), 6.84 (1H, t, *J* = 8.0 Hz, ArH-6'), 7.07 (1H, br, NH), 7.16 (1H, d, *J* = 8.0 Hz, ArH-5'), 7.31 (1H, d, *J* = 8.0 Hz, ArH-8'), 7.48 (1H, t, *J* = 8.0 Hz, ArH-7'), 10.72 (1H, br, NH); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 24.0(x2), 32.9(x2), 51.5, 52.7, 55.8, 102.6, 108.0, 112.5, 120.7, 120.8, 124.9, 126.1(x2), 131.7, 133.4, 154.4, 156.4, 158.9, 160.5, 168.4; MS *m/z* (%) 390 (*M* + 1, 100); HRMS *m/z* calcd for C<sub>22</sub>H<sub>24</sub>N<sub>5</sub>O<sub>2</sub> [*M* + H]<sup>+</sup> 390.1930, found 390.1984; HPLC purity 99.1%.

**7-Methoxy-4-(2-(pyrrolidin-1-yl)quinazolin-4-yl)-3,4-dihydroquinoxalin-2(1H)-one (6h).** A mixture of **6a** (50 mg, 0.15 mmol), 4-picolyamine (1 mL, excess), and K<sub>2</sub>CO<sub>3</sub> (30 mg, 0.21 mmol) in *i*-PrOH (2 mL) was heated to 60 °C under MW irradiation for 15 min to obtain **6h** as a white solid (42 mg, 58% yield). Mp 217–218 °C; <sup>1</sup>H NMR δ 1.90 (4H, m, CH<sub>2</sub> × 2), 3.55 (4H, m, NCH<sub>2</sub> × 2), 3.67 (3H, s, OCH<sub>3</sub>), 4.36 (2H, s, CH<sub>2</sub>), 6.38 (1H, dd, *J* = 8.8 and 2.8 Hz, ArH-6), 6.51 (1H, d, *J* = 2.8 Hz, ArH-8), 6.62 (1H, d, *J* = 8.8 Hz, ArH-5), 7.17 (1H, t, *J* = 8.0 Hz, ArH-6'), 7.19 (1H, d, *J* = 8.0 Hz, ArH-5'), 7.38 (1H, d, *J* = 8.0 Hz, ArH-8'), 7.47 (1H, t, *J* = 8.0 Hz, ArH-7'), 10.72 (1H, s, NH); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 25.6(x2), 46.9(x2), 51.4, 55.9, 102.6, 107.9, 111.8, 120.7(x2), 124.8, 126.2, 126.3, 131.7, 133.4, 154.7, 157.4, 160.2, 168.5; MS *m/z* (%) 376 (*M* + 1, 100); HRMS *m/z* calcd for C<sub>21</sub>H<sub>22</sub>N<sub>5</sub>O<sub>2</sub> [*M* + H]<sup>+</sup> 376.1773, found 376.1784; HPLC purity 95.4% (MeOH/H<sub>2</sub>O).

**4-(2-(3,3-Difluoropyrrolidin-1-yl)quinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6i).** A mixture of **6a** (50 mg, 0.15 mmol) and 3,3-difluoropyrrolidine hydrochloride (30 mg, 0.29 mmol) and K<sub>2</sub>CO<sub>3</sub> (30 mg, 0.21 mmol) in *i*-PrOH (2 mL) was heated to 120 °C under MW irradiation for 20 min to obtain **6i** as a brown solid (52 mg, 85% yield). Mp 247–248 °C; <sup>1</sup>H NMR δ 2.51 (2H, m, CH<sub>2</sub>), 3.68 (3H, s, OCH<sub>3</sub>), 3.79 (2H, t, *J* = 7.2 Hz, NCH<sub>2</sub>), 3.98 (2H, t, *J*<sub>H-F</sub> = 12.8 Hz, NCH<sub>2</sub>CF<sub>2</sub>), 4.41 (2H, s, 3'-CH<sub>2</sub>), 6.38 (1H, dd, *J* = 8.8 and 2.8 Hz, ArH-6), 6.56 (1H, d, *J* = 8.8 Hz, ArH-5), 6.63 (1H, d, *J* = 2.8 Hz, ArH-8), 6.92 (1H, t, *J* = 8.0 Hz, ArH-6'), 7.23 (1H, d, *J* = 8.0 Hz, ArH-5'), 7.45 (1H, d, *J* = 8.0 Hz, ArH-8'), 7.53 (1H, t, *J* = 8.0 Hz, ArH-7'), 10.87 (1H, s, NH); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 33.8, 44.5, 51.4, 53.6, 55.8, 102.7, 108.0, 112.2, 120.9, 121.6, 124.4, 126.2, 126.5, 128.7, 132.0, 133.7, 154.2, 156.6, 157.1, 160.4, 168.4; MS *m/z* (%) 412 (*M* + 1, 100); HRMS *m/z* calcd for C<sub>21</sub>H<sub>20</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub> [*M* + H]<sup>+</sup> 412.1585, found 412.1664; HPLC purity 95.6%.

**4-(2-(3,3-Difluoroazetidin-1-yl)quinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6j).** A mixture of **6a** (100 mg, 0.29 mmol) and 3,3-difluoroazetidine hydrochloride (45 mg, 1.1 mmol) in *i*-PrOH (2 mL) in the presence of one drop H<sub>2</sub>SO<sub>4</sub> (conc) was heated to 120 °C under MW irradiation for 20 min. The crude product was dissolved in MeOH and left uncovered overnight. The precipitated solid was collected to obtain pure **6j** as a yellow solid (40 mg, 35% yield). Mp 280–281 °C; <sup>1</sup>H NMR δ 3.68 (3H, s, OCH<sub>3</sub>), 4.41 (2H, s, CH<sub>2</sub>), 4.51 (4H, t, *J*<sub>H-F</sub> = 12.8 Hz, NCH<sub>2</sub> × 2), 6.39 (1H, dd, *J* = 8.8 and 2.8 Hz, ArH-6), 6.61 (1H, d, *J* = 8.8 Hz, ArH-5), 6.63 (1H, d, *J* = 2.8 Hz, ArH-8), 6.99 (1H, t, *J* = 8.0 Hz, ArH-6'), 7.25 (1H, d, *J* = 8.0 Hz, ArH-5'), 7.51 (1H, d, *J* = 8.0 Hz, ArH-8'), 7.58 (1H, t, *J* = 8.0 Hz, ArH-7'), 10.78 (1H, s, NH); MS *m/z* (%) 398 (*M* + 1, 100); HRMS *m/z* calcd for C<sub>20</sub>H<sub>18</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub> [*M* + H]<sup>+</sup> 398.1429, found 398.1441 (*M* + 1, 100); HPLC purity 97.8%.

**7-Methoxy-4-(2-(3-methoxyazetidin-1-yl)quinazolin-4-yl)-3,4-dihydroquinoxalin-2(1H)-one (6k).** A mixture of **6a** (50 mg, 0.15 mmol), methoxyazetidine (50 mg, 0.41 mmol), and K<sub>2</sub>CO<sub>3</sub> (50 mg, 0.36 mmol) in anhydrous *i*-PrOH (2 mL) was heated to 120 °C under MW irradiation for 15 min to obtain pure **6k** as a gray solid (23 mg, 39% yield) after recrystallization from MeOH. Mp 221–222 °C; <sup>1</sup>H NMR δ 3.23 (3H, s, OCH<sub>3</sub>), 3.68 (3H, s, OCH<sub>3</sub>), 3.88 (2H, d, *J* = 7.2 Hz, NCH<sub>2</sub>), 4.25 (2H, d, *J* = 7.2 Hz, NCH<sub>2</sub>), 4.28 (1H, m, CH), 4.36 (2H, s, CH<sub>2</sub>), 6.38 (1H, dd, *J* = 8.8 and 2.8 Hz, ArH-6), 6.54 (1H, d, *J* = 8.8 Hz, ArH-5), 6.62 (1H, d, *J* = 2.8 Hz, ArH-8), 6.90 (1H, t, *J* = 8.0 Hz, ArH-6'), 7.20 (1H, d, *J* = 8.0 Hz, ArH-5'), 7.42 (1H, d, *J* = 8.0 Hz, ArH-8'), 7.51 (1H, t, *J* = 8.0 Hz, ArH-7'), 10.72 (1H, s, NH); <sup>13</sup>C

NMR (DMSO- $d_6$ )  $\delta$  51.6, 55.8, 55.9, 57.4, 69.8(x2), 102.6, 108.0, 112.5, 120.9, 121.5, 126.2, 127.4, 131.8, 133.6(x2), 154.1, 156.6, 160.0, 160.4, 168.4; MS  $m/z$  (%) 392 (M + 1, 100); HRMS  $m/z$  calcd for  $C_{21}H_{22}N_5O_2$  [M + H]<sup>+</sup> 392.1723, found 392.1729; HPLC purity 96.0%.

**4-(2-(4-Cyanophenyl)aminoquinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6l).** A mixture of **6a** (50 mg, 0.15 mmol) and 4-aminobenzonitrile (30 mg, 0.25 mmol) in *i*-PrOH (2 mL) in the presence of  $H_2SO_4$  (conc, 1 drop) was heated to 120 °C under MW irradiation for 20 min to obtain pure **6l** as a yellow solid (50 mg, 79% yield) after being recrystallized from MeOH. Mp 183–184 °C; <sup>1</sup>H NMR  $\delta$  3.71 (3H, s, OCH<sub>3</sub>), 4.57 (2H, s, CH<sub>2</sub>), 6.47 (1H, dd,  $J$  = 8.8 and 2.8 Hz, ArH-6), 6.66 (1H, d,  $J$  = 8.8 Hz, ArH-5), 6.89 (1H, d,  $J$  = 2.8 Hz, ArH-8), 7.16 (1H, t,  $J$  = 8.0 Hz, ArH-6'), 7.37 (2H, d,  $J$  = 8.2 Hz, 2'-PhH), 7.65 (1H, d,  $J$  = 8.0 Hz, ArH-5'), 7.70 (1H, t,  $J$  = 8.0 Hz, ArH-7'), 7.75 (2H, d,  $J$  = 8.2 Hz, 2'-PhH), 7.95 (1H, d,  $J$  = 8.0 Hz, ArH-8'), 10.58 (1H, brs, NH), 10.87 (1H, s, NH); MS  $m/z$  (%) 423 (M + 1, 100); HRMS  $m/z$  calcd for  $C_{24}H_{19}N_6O_2$  [M + H]<sup>+</sup> 423.1569, found 423.1649; HPLC purity 95.3%.

**4-(2-(3-Cyanophenyl)aminoquinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6m).** A mixture of **6a** (50 mg, 0.15 mmol) and 3-aminobenzonitrile (30 mg, 0.25 mmol) in *i*-PrOH (2 mL) in the presence of  $H_2SO_4$  (conc, 1 drop) was heated to 120 °C under MW irradiation for 20 min to obtain **6m** as a yellow solid (55 mg, 88% yield). Mp 262–263 °C; <sup>1</sup>H NMR  $\delta$  3.71 (3H, s, OCH<sub>3</sub>), 4.57 (2H, s, CH<sub>2</sub>), 6.50 (1H, dd,  $J$  = 8.8 and 2.8 Hz, ArH-6), 6.65 (1H, d,  $J$  = 2.8 Hz, ArH-8), 7.00 (1H, d,  $J$  = 8.8 Hz, ArH-5), 7.22 (1H, m, ArH-6'), 7.41 (1H, s, ArH-3'), 7.56 (2H, m, ArH-5' and ArH-4''), 7.65 (1H, d,  $J$  = 7.6 Hz, ArH-8'), 7.76 (1H, t,  $J$  = 7.6 Hz, ArH-7'), 7.92 (1H, m, ArH-5''), 8.13 (1H, m, ArH-6''), 10.44 (1H, br, NH), 10.84 (1H, s, 1-NH); <sup>13</sup>C NMR (DMSO- $d_6$ )  $\delta$  52.2, 56.0, 102.6, 108.6, 112.3, 119.1, 121.3, 124.1, 124.3, 124.5, 125.2, 127.0, 127.1, 127.3, 128.6, 130.9, 133.3, 135.9, 139.0, 151.9, 152.1, 158.8, 161.6, 167.1; MS  $m/z$  (%) 423 (M + 1, 100); HRMS  $m/z$  calcd for  $C_{24}H_{19}N_6O_2$  [M + H]<sup>+</sup> 423.1569, found 423.1571; HPLC purity 96.6%.

**7-Methoxy-4-(2-((pyridin-4-ylmethyl)amino)quinazolin-4-yl)-3,4-dihydroquinoxalin-2(1H)-one (6n).** A mixture of **6a** (35 mg, 0.1 mmol), pyridin-4-ylmethanamine (1 mL, excess), and  $K_2CO_3$  (30 mg, 0.21 mmol) in *i*-PrOH (2 mL) was heated to 120 °C under MW irradiation for 1 h to obtain **6n** as a yellow solid (20 mg, 47% yield). Mp 219–220 °C; <sup>1</sup>H NMR  $\delta$  3.71 (3H, s, OCH<sub>3</sub>), 4.31 (2H, s, 3-CH<sub>2</sub>), 4.55 (2H, d,  $J$  = 6.0 Hz, CH<sub>2</sub>), 6.36 (1H, dd,  $J$  = 8.8 and 2.8 Hz, ArH-6), 6.55 (1H, d,  $J$  = 8.8 Hz, ArH-5), 6.60 (1H, d,  $J$  = 2.8 Hz, ArH-8), 6.67 (1H, t,  $J$  = 7.6 Hz, ArH-6'), 7.18 (1H, d,  $J$  = 7.6 Hz, ArH-5'), 7.33 (3H, m, ArH-8' and PyH), 7.47 (1H, t,  $J$  = 7.6 Hz, ArH-7'), 7.73 (1H, m, NH), 8.43 (2H, m, PyH), 10.73 (1H, br, 1-NH); MS  $m/z$  (%) 413 (M + 1, 100); HRMS  $m/z$  calcd for  $C_{23}H_{21}N_6O_2$  [M + H]<sup>+</sup> 413.1726, found 413.1799; HPLC purity 97.8%.

**4-(2-Ethoxyquinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6o).** Ethanol (2 mL) was cooled to 0 °C, and NaH (260 mg, 11 mmol) was carefully added in portions with stirring over 1 h. Compound **6a** (100 mg, 0.29 mmol) was then added into the solution with stirring and kept at the same temperature for another 0.5 h until hydrogen was no longer formed. The mixture was then heated to 95 °C under MW irradiation for 5 min and monitored by TLC and LC-MS until the reaction was complete. The mixture was poured into ice-water, and the precipitated solid was collected, washed with water, and dried to obtain pure **6o** as a brown solid (85 mg, 85% yield). Mp 233–234 °C; <sup>1</sup>H NMR  $\delta$  1.34 (3H, t,  $J$  = 7.2 Hz, CH<sub>3</sub>), 3.68 (3H, s, 7'-OCH<sub>3</sub>), 4.41 (4H, m, OCH<sub>2</sub> and 3'-CH<sub>2</sub>), 6.39 (1H, dd,  $J$  = 8.8 and 2.8 Hz, ArH-6), 6.65 (2H, m, ArH-5 and -8), 7.12 (1H, m, ArH-6'), 7.35 (1H, d,  $J$  = 8.0 Hz, ArH-5'), 7.63 (2H, m, ArH-8', ArH-7'), 10.78 (1H, br s, NH); <sup>13</sup>C NMR (DMSO- $d_6$ )  $\delta$  15.0, 51.6, 55.9, 62.9, 102.7, 108.1, 114.2, 121.4, 123.7, 124.0, 126.2, 127.3, 132.3, 134.0, 153.5, 157.0, 161.9, 162.2, 167.9; MS  $m/z$  (%) 351 (M + 1, 100); HRMS  $m/z$  calcd for  $C_{19}H_{19}N_4O_3$  [M + H]<sup>+</sup> 351.1457, found 351.1507; HPLC purity 96.2%.

**4-(2-(Cyclopropylmethoxy)quinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6p).** The preparation was the same as described for **6o**. The mixture of cyclopropylcarbinol (2 mL), NaH

(260 mg, 11 mmol), and **6a** (100 mg, 0.29 mmol) was stirred at 0 °C for about 1.5 h and then heated to 95 °C under MW irradiation for 5 min to obtain pure **6p** as a yellow solid (90 mg, 83% yield). Mp 215–216 °C; <sup>1</sup>H NMR  $\delta$  0.34 (2H, m, CH<sub>2</sub>), 0.55 (2H, m, CH<sub>2</sub>), 0.81 (1H, m, CH), 3.71 (3H, s, OCH<sub>3</sub>), 4.17 (2H, d,  $J$  = 7.2 Hz, OCH<sub>2</sub>), 4.41 (2H, s, CH<sub>2</sub>), 6.39 (1H, dd,  $J$  = 8.8 and 2.8 Hz, ArH-6), 6.64 (2H, m, ArH-5 and -8), 7.13 (1H, m, ArH-6'), 7.35 (1H, d,  $J$  = 8.0 Hz, ArH-5'), 7.63 (2H, m, ArH-8' and -7'), 10.77 (1H, br, NH); <sup>13</sup>C NMR (DMSO- $d_6$ )  $\delta$  3.74(x2), 10.5, 51.6, 55.8, 71.8, 102.6, 108.1, 114.2, 121.4, 123.7, 123.9, 126.2, 127.3, 132.2, 134.0, 153.2, 157.0, 161.9, 162.2, 168.2; MS  $m/z$  (%) 377 (M + 1, 100); HRMS  $m/z$  calcd for  $C_{21}H_{21}N_4O_3$  [M + H]<sup>+</sup> 377.1614, found 377.1615; HPLC purity 95.1%.

**4-(2-Acetoxymethylquinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6q).** A mixture of **13** prepared as described below (390 mg, 0.94 mmol) and anhydrous  $K_2CO_3$  (260 mg, 1.88 mmol) in DMA (3 mL) was heated at 100 °C for 1 h. After the reaction was complete as judged by LC-MS monitoring, the mixture was poured into ice water and extracted with EtOAc three times. The combined organic phases were washed successively with water and brine and dried over anhydrous  $Na_2SO_4$ . After removal of solvent in vacuo, the residue was purified by flash column chromatography (gradient elution, petroleum ether/EtOAc, 0–30%) to afford **6q** as a yellow solid (355 mg, 99% yield). Mp 192–194 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.26 (3H, s, CH<sub>3</sub>), 3.80 (3H, s, OCH<sub>3</sub>), 4.66 (2H, s, NCH<sub>2</sub>), 5.35 (2H, s, OCH<sub>2</sub>), 6.40 (1H, dd,  $J$  = 8.8 and 2.8 Hz, ArH-6), 6.54 (1H, d,  $J$  = 2.8 Hz, ArH-8), 6.63 (1H, d,  $J$  = 8.8 Hz, ArH-5), 7.25 (1H, t,  $J$  = 8.0 Hz, ArH-6'), 7.48 (1H, d,  $J$  = 8.0 Hz, ArH-5'), 7.72 (1H, t,  $J$  = 8.0 Hz, ArH-7'), 7.92 (1H, d,  $J$  = 8.0 Hz, ArH-8'), 8.37 (1H, s, NH); <sup>13</sup>C NMR (400 Hz, CDCl<sub>3</sub>)  $\delta$  21.0, 51.1, 55.7, 66.4, 102.4, 108.6, 115.7, 120.8, 123.9, 125.6, 125.8, 128.8, 130.4, 133.2, 152.1, 157.2, 159.5, 160.4, 169.3, 170.9; MS  $m/z$  (%) 379 (M + 1, 100); HRMS  $m/z$  calcd for  $C_{20}H_{19}N_4O_4$  [M + H]<sup>+</sup> 379.1406, found 379.1431; HPLC purity 95.5%.

**4-(2-(Hydroxymethyl)quinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6r).** To a solution of **6q** (355 mg, 0.94 mmol) in MeOH (8 mL) was added aqueous NaOH (10%, ca. 1.0 mL, excess) at rt. The mixture was stirred for 1.5 h. Then, the pH was adjusted to 7 with aq. HCl (10%), and the mixture was extracted with EtOAc three times. The combined organic layer was washed successively with water and brine and dried over anhydrous  $Na_2SO_4$ . After removal of solvent in vacuo, **6r** was obtained as a yellow powder (310 mg, 98% yield). Mp 216–218 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  3.80 (3H, s, OCH<sub>3</sub>), 4.00 (1H, s, OH), 4.67 (1H, s, CH<sub>2</sub>), 4.85 (1H, s, OCH<sub>2</sub>), 6.41 (1H, dd,  $J$  = 8.8 and 2.8 Hz, ArH-6), 6.57 (1H, d,  $J$  = 2.8 Hz, ArH-8), 6.64 (1H, d,  $J$  = 8.8 Hz, ArH-5), 7.26 (1H, t,  $J$  = 8.0 Hz, ArH-6'), 7.51 (1H, d,  $J$  = 8.0 Hz, ArH-5'), 7.73 (1H, t,  $J$  = 8.0 Hz, ArH-7'), 7.92 (1H, d,  $J$  = 8.0 Hz, ArH-8'), 8.82 (1H, s, NH); <sup>13</sup>C NMR (400 Hz, CDCl<sub>3</sub>)  $\delta$  51.2, 55.7, 64.4, 102.5, 108.7, 115.9, 120.8, 123.9, 125.6, 125.8, 128.3, 130.4, 133.4, 151.4, 157.2, 159.6, 163.8, 169.2; MS  $m/z$  (%) 337 (M + 1, 100); HRMS  $m/z$  calcd for  $C_{18}H_{17}N_4O_3$  [M + H]<sup>+</sup> 337.1301, found 337.1290; HPLC purity 96.3%.

**4-(2-((4-Ethoxy-4-oxobutoxy)methyl)quinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6s).** A mixture of **6r** (165 mg, 0.5 mmol) and  $Cs_2CO_3$  (325 mg, 1 mmol) in  $CH_3CN$  (10 mL) was heated to reflux for 0.5 h. Then, ethyl 4-bromobutyrate (325 mg, 1.0 mmol) was added dropwise at the same temperature, and stirring was continued for 3 h. The mixture was passed through Celite, which was washed several times with  $CH_3CN$ . The combined filtrate was concentrated in vacuo to obtain **6s** as a yellow oil, which was used in the next reaction without further purification. However, pure **6s** was obtained as a yellow solid by crystallization from EtOAc. Mp 160–161 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.26 (3H, t,  $J$  = 7.2 Hz, CH<sub>3</sub>), 2.01 (2H, m, CH<sub>2</sub>), 2.45 (2H, t,  $J$  = 7.2 Hz, COCH<sub>2</sub>), 3.86 (3H, s, OCH<sub>3</sub>), 4.07 (2H, t,  $J$  = 7.2 Hz, OCH<sub>2</sub>), 4.15 (2H, q,  $J$  = 7.2, OCH<sub>2</sub>), 4.65 (2H, s, CH<sub>2</sub>), 4.83 (2H, s, ArCH<sub>2</sub>O), 6.42 (1H, dd,  $J$  = 8.8 and 2.4 Hz, ArH-6), 6.66 (1H, d,  $J$  = 8.8 Hz, ArH-5), 7.01 (1H, d,  $J$  = 2.4 Hz, ArH-8), 7.24 (1H, t,  $J$  = 8.0 Hz, ArH-6'), 7.46 (1H, d,  $J$  = 8.0 Hz, ArH-5'), 7.72 (1H, t,  $J$  = 8.0 Hz, ArH-7'), 7.92 (1H, d,  $J$  = 8.0 Hz, ArH-8'); <sup>13</sup>C NMR (400 Hz, CDCl<sub>3</sub>)  $\delta$  14.3, 22.1, 31.1, 41.7, 51.7, 55.9, 60.8, 64.5,



103.0, 108.0, 115.9, 121.0, 125.7 ( $\times 2$ ), 126.0, 128.4, 133.0, 133.4, 151.5, 157.6, 159.4, 163.9, 167.1, 173.1; MS  $m/z$  (%) 451 ( $M + 1$ , 100); HRMS  $m/z$  calcd for  $C_{24}H_{27}N_4O_5$  [ $M + H$ ] $^+$  451.1981, found 451.2083; HPLC purity 95.7%.

**4-(2-((4-(Hydroxyamino)-4-oxobutoxy)methyl)quinazolin-4-yl)-7-methoxy-3,4-dihydroquinoxalin-2(1H)-one (6t).** A solution of **6s** (225 mg, 0.5 mmol) in  $CH_2Cl_2$  and MeOH (9 mL, 1:2, v/v) was cooled to 0 °C. Aqueous hydroxylamine, prepared from hydroxylamine hydrochloride (1.04 g, 1.5 mmol) and NaOH (0.8 g, 2.0 mmol) in 2 mL of water, was added dropwise into the solution of **6s** with stirring at 0 °C, and the mixture was stirred for an additional 1 h. After organic solvent was removed under reduced pressure, the water phase was adjusted to pH 7–8 with HOAc and extracted with EtOAc three times. The combined organic phase was washed with water and brine and dried over anhydrous  $Na_2SO_4$ . After solvent was removed in vacuo, the residue was recrystallized from EtOAc to obtain pure **6t** as a yellow solid (50 mg), mp 196–198 °C;  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.84 (2H, m,  $CH_2$ ), 2.04 (2H, t,  $J = 7.2$  Hz,  $C(O)CH_2$ ), 3.77 (3H, s,  $OCH_3$ ), 3.95 (2H, t,  $J = 7.2$  Hz,  $OCH_2$ ), 4.60 (2H, s,  $CH_2$ ), 4.61 (2H, s,  $ArCH_2O$ ), 5.20 (1H, s, OH), 6.48 (1H, dd,  $J = 8.8$  and 2.4 Hz, ArH-6'), 6.67 (1H, d,  $J = 8.8$  Hz, ArH-5'), 7.02 (1H, d,  $J = 2.4$  Hz, ArH-8'), 7.31 (1H, t,  $J = 8.0$  Hz, ArH-6), 7.36 (1H, d,  $J = 8.0$  Hz, ArH-5), 7.76 (1H, t,  $J = 8.0$  Hz, ArH-7), 7.83 (1H, d,  $J = 8.0$  Hz, ArH-8), 8.71 (1H, s, NH), 10.38 (1H, s, NH);  $^{13}C$  NMR (400 Hz,  $DMSO-d_6$ )  $\delta$  22.9, 29.6, 40.2, 51.7, 56.1, 65.4, 103.4, 108.4, 115.7, 121.2, 125.9, 126.1, 126.2, 128.6, 133.0, 133.8, 152.0, 157.4, 159.2, 165.3, 167.0, 169.1; MS  $m/z$  (%) 438 ( $M + 1$ , 100); HRMS  $m/z$  calcd for  $C_{22}H_{24}N_5O_5$  [ $M + H$ ] $^+$  438.1777, found 438.1866; HPLC purity 95.1%.

**2-Chloromethyl-3H-quinazolin-4-one (8).** Compound **8** was prepared from commercially available methyl 2-aminobenzoate (**7**) and 2-chloroacetonitrile according to the literature method.<sup>26</sup> Mp 240 °C (dec.);  $^1H$  NMR ( $CDCl_3$ )  $\delta$  4.51 (2H, s,  $CH_2$ ), 7.51 (1H, t,  $J = 8.0$  Hz, ArH-6), 7.64 (1H, d,  $J = 8.0$  Hz, ArH-8), 7.80 (1H, t,  $J = 8.0$  Hz, ArH-7), 8.08 (1H, d,  $J = 8.0$  Hz, ArH-5), 12.57 (1H, s, NH); MS  $m/z$  (%) 195 ( $M + 1$ , 100), 197 ( $M + 3$ , 33).

**2-(Chloromethyl)quinazolin-4-yl 4-methylbenzenesulfonate (9).** A mixture of **8** (500 mg 2.58 mmol), *p*-toluenesulfonyl chloride (880 mg, 4.6 mmol),  $Et_3N$  (521 mg 5.16 mmol), and DMAP (3 mg, 0.026 mmol) in  $CH_2Cl_2$  (20 mL) was stirred at rt for 3 h until the reaction was complete as judged by TLC monitoring. Solvent was removed under reduced pressure, and the residue was purified by flash column chromatography (gradient elution, EtOAc/petroleum ether, 0–15%) to obtain **9** as a pale yellow solid (790 mg, 88% yield). Mp 149–151 °C;  $^1H$  NMR ( $CDCl_3$ )  $\delta$  2.46 (3H, s,  $CH_3$ ), 4.73 (2H, s,  $CH_2$ ), 7.38 (2H, d,  $J = 8.4$  Hz, PhH), 7.68 (1H, t,  $J = 8.4$  Hz, ArH-6), 7.93 (1H, t,  $J = 8.4$  Hz, ArH-7), 7.97 (1H, d,  $J = 8.4$  Hz, ArH-5), 8.18 (2H, d,  $J = 8.4$  Hz, PhH), 8.20 (1H, d,  $J = 8.4$  Hz, ArH-8); MS  $m/z$  (%) 349 ( $M + 1$ , 100), 351 ( $M + 3$ , 33).

**2-Chloromethyl-4-(4-methoxy-2-nitrophenyl)aminoquinazoline (10).** A mixture of **9** (500 mg 1.43 mmol) and **3** (240 mg 1.43 mmol) in *i*-PrOH and  $CH_2Cl_2$  (15 mL, v/v 2:1) was stirred at 45 °C for 6 h until the reaction was complete as judged by TLC monitoring. After solvent was removed, the residue was dissolved in 50 mL of  $CH_2Cl_2$ , washed with saturated  $NaHCO_3$ , dried over anhydrous  $Na_2SO_4$ , and concentrated to yield crude product **10**, which was used in the next step without further purification. Orange solid; Mp 180–181 °C;  $^1H$  NMR ( $CDCl_3$ )  $\delta$  3.90 (3H, s,  $OCH_3$ ), 4.59 (2H, s,  $CH_2$ ), 7.38 (1H, dd,  $J = 9.2$  and 3.2 Hz, ArH-5'), 7.67 (1H, t,  $J = 8.0$  Hz, ArH-6), 7.77 (1H, d,  $J = 3.2$  Hz, ArH-3'), 7.87 (1H, t,  $J = 8.0$  Hz, ArH-7), 7.95 (1H, d,  $J = 8.0$  Hz, ArH-5), 8.64 (1H, d,  $J = 8.0$  Hz, ArH-8), 9.39 (1H, d,  $J = 9.2$  Hz, ArH-6'), 11.31 (1H, s, NH); MS  $m/z$  (%) 345 ( $M + 1$ , 100), 347 ( $M + 3$ , 33).

**2-Acetoxymethyl-4-(4-methoxy-2-nitro-phenyl)-aminoquinazoline (11).** A mixture of **10** (500 mg, 1.45 mmol) and KOAc (427 mg, 4.36 mmol) in DMF (2.0 mL) was stirred at 55 °C for 6 h. The mixture was poured into ice–water and extracted with EtOAc. The organic layer was washed with water and brine successively and dried over  $Na_2SO_4$ . After removal of solvent in vacuo, the residue was purified by flash column chromatography (gradient elution,  $CH_2Cl_2$ /EtOAc, 0–15%) to give **11** as an orange

solid (300 mg, 65% yield). Mp 159–160 °C;  $^1H$  NMR ( $CDCl_3$ )  $\delta$  2.24 (3H, s,  $CH_3$ ), 3.90 (3H, s,  $OCH_3$ ), 5.35 (2H, s,  $OCH_2$ ), 7.33 (1H, dd,  $J = 9.6$  and 3.2 Hz, ArH-5'), 7.64 (1H, t,  $J = 8.4$  Hz, ArH-6), 7.76 (1H, d,  $J = 3.2$  Hz, ArH-3'), 7.84 (1H, t,  $J = 8.4$  Hz, ArH-7), 7.93 (1H, d,  $J = 8.4$  Hz, ArH-5), 8.04 (1H, d,  $J = 8.4$  Hz, ArH-8), 9.24 (1H, d,  $J = 9.6$  Hz, ArH-6'), 11.24 (1H, s, NH); MS  $m/z$  (%) 369 ( $M + 1$ , 100).

**2-Acetoxymethyl-4-(4-methoxy-2-aminophenyl)-aminoquinazoline (12).** To a solution of **11** (154 mg, 0.54 mmol) in a mixed solvent of EtOAc and EtOH (25 mL, v/v 3:2) was added Pd/C (15 mg, 10% w/w) and hydrogen gas. The mixture was shaken under 40 psi at rt for 2 h. The Pd/C was removed by filtration and washed with EtOAc several times. After removal of the solvent in vacuo, 185 mg of light yellow solid **12** was obtained and used directly in the next step without further purification.

**2-Acetoxymethyl-4-(2-chloroacetyl-amino-4-methoxyphenyl)-aminoquinazoline (13).** Crude **12** (ca. 545 mg, 1.61 mmol) was immediately dissolved in acetone (8 mL), and  $K_2CO_3$  (667 mg, 4.83 mmol) was added. The mixture was cooled to 0 °C and chloroacetyl chloride (361 mg, 3.22 mmol) was added dropwise. Stirring was continued for an additional 1 h at 0 °C. When the reaction was complete, the mixture was poured into ice water, the pH was adjusted to neutral, and the product was extracted with EtOAc three times. The combined organic phase was washed successively with water and brine, and dried over anhydrous  $Na_2SO_4$  overnight. After removal of solvent in vacuo, **13** was obtained as a gray solid and used in the synthesis of **6q** (400 mg, 60% yield over two steps). Mp 121–122 °C; MS  $m/z$  (%) 415 ( $M + 1$ , 100), 417 ( $M + 3$ , 33).

**Aqueous Solubility Studies.** Solubility was measured at pH 7.4 by using an HPLC–UV method. Test compounds were initially dissolved in DMSO at a concentration of 1.0 mg/mL. Ten microliters of this stock solution was added to phosphate buffer (1.0 mL, pH 7.4). The mixture was stirred for 4 h at rt and then centrifuged at 3000 rpm for 10 min. The saturated supernatants were transferred to other vials for analysis by HPLC–UV. Each sample was performed in triplicate. For quantification, a model 1200 HPLC–UV (Agilent) system was used with an Agilent Eclipse XDB-C18 column (150 mm  $\times$  4.6 mm, 5  $\mu m$ ), and elution was with 50%–80% ACN in water. The flow rate was 0.8 mL/min, and injection volume was 20  $\mu L$ . Aqueous concentration was determined by comparison of the peak area of the saturated solution with a standard curve plotted peak area versus known concentrations, which were prepared by solutions of test compound in ACN at 50, 12.5, 3.13, 0.78, and 0.20  $\mu g/mL$ .

**Log P Measurements.** One to two milligrams of test compound was dissolved in 1.0–2.0 mL of *n*-octanol to obtain a 1.0 mg/mL solution. Next, the same volume of water as *n*-octanol was added to each vial. The mixture was stirred at rt for 24 h and left without stirring overnight. The aqueous and organic phases of each mixture were transferred to separate vials for HPLC analysis. The instrument and conditions were the same as those for water solubility determinations. The log *P* was calculated by the peak area ratio in *n*-octanol and in water.

**Cell-Based Assays.** Antiproliferative activities of series-6 compounds were assayed by the SRB method according to procedures described previously.<sup>27–29</sup> The panel of cell lines included A-549, MDA-MB-231, KB, KB-VIN (P-gp-overexpressing KB subline), and MCF-7. These cell lines were obtained from the Lineberger Comprehensive Cancer Center (UNC–CH) or ATCC. Cells were propagated in RPMI-1640 (Gibco) supplemented with 10% FBS (Corning), penicillin (100 IU/mL), streptomycin (1  $\mu g/mL$ ), and amphotericin B (0.25  $\mu g/mL$ ) (Corning) and were cultured at 37 °C in a humidified atmosphere of 95% air and 5%  $CO_2$ . The antiproliferative effects against tumor cell lines were expressed as  $GI_{50}$  values, which represent the molar drug concentrations required to cause 50% tumor cell growth inhibition. Flow cytometry was performed as described previously.<sup>30</sup> Briefly, A549 cells were seeded 24 h prior to treatment in a 12-well culture plate at density of 60 000 cells/well. After 24 h treatment with 15 nM **6d**, 15 nM **6e**, or 20 nM CA-4, cells were fixed and stained with PI containing RNase (BD Biosciences). DMSO (0.001%) was used as a control. Stained cells were analyzed by flow cytometry operated by FACSDiva software (BD



LSRFortessa, BD Biosciences). DMSO was used for preparation of compound, and the highest concentration of DMSO for the cell-based assay was less than 0.001% (v/v).

**Tubulin Assays.** Tubulin assembly was measured by turbidimetry at 350 nm as described previously.<sup>23</sup> Assay mixtures containing 1.0 mg/mL (10  $\mu$ M) tubulin and varying compound concentrations were preincubated for 15 min at 30 °C without guanosine 5'-triphosphate (GTP). The samples were placed on ice, and GTP to a final concentration of 0.4 mM was added. Reaction mixtures were transferred to 0 °C cuvettes, and turbidity development was followed for 20 min at 30 °C following a rapid temperature jump. Compound concentrations that inhibited increase in turbidity by 50% relative to a control sample were determined. Inhibition of the binding of [<sup>3</sup>H]colchicine to tubulin was measured as described previously.<sup>31</sup> Incubation of 1.0  $\mu$ M tubulin with 5.0  $\mu$ M [<sup>3</sup>H]colchicine and 5.0 or 1.0  $\mu$ M inhibitor was for 10 min at 37 °C, when about 40–60% of maximum colchicine binding occurred in control samples.

**Xenograft Tumor Growth Inhibition in Vivo.** Six-week-old female Balb/c-nu mice were obtained from Vital River (Beijing, China) and housed in specific-pathogen-free conditions in conformity with the Guide for the Care and Use of Laboratory Animals, as adopted and promulgated by Beijing Institute of Radiation Medicine. Human lung cancer NCI-H460 cells ( $2 \times 10^6$ ) were injected subcutaneously into the right abdominal flanks of the Balb/c-nu mice. Tumor growth was measured with a slide caliper, and volumes were estimated according to the following formula: tumor volume ( $\text{mm}^3$ ) =  $L \times W^2 \times 0.5$ , where  $L$  is length and  $W$  is width. When tumor volume reached about 100  $\text{mm}^3$ , the mice were randomly divided into five groups ( $n = 8$ ) and compound administration was started on the second day. The control group was dosed intraperitoneally daily with 7  $\mu$ L/g of vehicle (0.9% NaCl containing 5% polyethylene glycol 400 and 0.5% Tween 80). Compound 2 groups were treated at the doses of 0.25, 0.5, or 1.0 mg/kg body weight, respectively, every 5 days by iv injection for 3 weeks. The reference group received paclitaxel at a dose of 15 mg/kg body weight. Both compound 2 and paclitaxel were dissolved in vehicle solution (5% PEG400/PBS). The body weight and tumor volume were measured at each time point. At the end of the treatment, the mice were sacrificed for autopsy, and the tumors were recovered and weighed. The tumor growth inhibitory rate was calculated as follow: inhibitory rate (%) =  $[1 - (\text{mean tumor weight of treated group})/(\text{mean tumor weight of control group})] \times 100$ . Data were presented as the mean  $\pm$  SD. Statistical significance of differences between groups was compared by one-way analysis of variance test followed by Tukey's post hoc test (GraphPad Prism 5.0; GraphPad Software Inc., La Jolla, CA, USA).  $P < 0.05$  was considered to indicate a statistically significant difference. This study was approved by the Institutional Animal Care and Use Committee (IACUC) of the Beijing Institute of Radiation Medicine (Beijing, China) and performed according to the institutional guidelines.

**Immunohistochemical Staining.** The tumor tissues were formalin fixed, paraffin embedded, and sectioned into 4  $\mu$ m slices. The tissue sections were deparaffinized, rehydrated, and treated with 3%  $\text{H}_2\text{O}_2$  to quench the endogenous peroxidase activity. Antigen retrieval was performed by boiling slides in sodium citrate buffer (10 mM, pH 6.0). CD31 (Thermo Fisher Scientific; cat. no. PA5-32321), cleaved caspase-3 (Cell Signaling Technology; cat. no. 9661), or Ki67 antibody (Thermo Fisher Scientific; cat. no. MA5-14520) was used as primary antibody, and horseradish peroxidase (HRP)-conjugated antibody was used as secondary antibody. HRP was detected by diaminobenzidine (DAB) (Wuhan Boster Biological Technology, Ltd.). The immunohistochemistry reaction was developed with a DAB substrate kit (Wuhan Boster Biological Technology, Ltd.) prior to counterstaining the slides with hematoxylin. The immunostained sections were examined using an EVOS X1 microscope (Advanced Microscopy Group; Thermo Fisher Scientific, Inc.).

**Molecular Modeling.** The molecular modeling study was performed with Discovery Studio 3.0 (Accelrys, San Diego, USA), also used in our prior modeling method.<sup>13</sup> The crystal structure of tubulin in complex with CA-4 (PDB code 5LYJ) with 2.4 Å resolution

was downloaded from the RCSB Protein Data Bank (<http://www.rcsb.org/pdb>) for use in the modeling study. LIBDOCK was used to evaluate and predict in silico binding free energy of the inhibitors and for automated docking. The protein model was typed with the CHARMM force field. A binding sphere with radius of 8.5 Å was defined through the original ligand (CA-4) as the binding site for the study. The docked ligand 6e and 2 were refined using in situ ligand minimization with the Smart Minimizer algorithm by standard parameters.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.jmedchem.7b00273.

HPLC purity conditions and data of 6a–6t, HRMS data of 6a–6t, data of 2 in NIH-NCI 60 cell line panel, and representative <sup>13</sup>C spectra of target compounds (6a, 6b, 6h, 6q, 6r, 6s) (PDF)

SMILES molecular formula strings (CSV)

## ■ AUTHOR INFORMATION

### Corresponding Authors

\*For L. Xie. Phone/fax: +86-10-66931690. E-mail: [lanxieshi@yahoo.com](mailto:lanxieshi@yahoo.com); [lxie@email.unc.edu](mailto:lxie@email.unc.edu).

\*For K. H. Lee. Phone: 919-962-0066. Fax: 919-966-3893. E-mail: [khlee@unc.edu](mailto:khlee@unc.edu).

### ORCID

Kuo-Hsiung Lee: 0000-0002-6562-0070

### Notes

The content of this paper is solely the responsibility of the authors and does not necessarily reflect the official views of the National Institutes of Health.

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We thank the Molecular Pharmacology Branch, DTP, DCTD, NCI, for performing the NCI 60-cell cytotoxicity assay. This investigation was supported by Grants 81120108022 and 30930106 from the Natural Science Foundation of China (NSFC) awarded to L.X. and NIH Grant CA177584 from the National Cancer Institute awarded to K.H.L. This study was also supported in part by the Taiwan Department of Health, China Medical University Hospital Cancer Research Center of Excellence (DOH100-TD-C-111-005), and Grant 81472687 from the Natural Science Foundation of China awarded to L.L. This study was also supported in part by the Eshelman Institute for Innovation, Chapel Hill, North Carolina, awarded to M.G.

## ■ ABBREVIATIONS USED

CA-4, combretastatin A-4; DAMA-colchicine, *N*-deacetyl-*N*-(2-mercaptoacetyl)-colchicine; DMA, dimethylacetamide; DMAP, dimethylaminopyridine; DMF, dimethylformamide; DMSO, dimethyl sulfoxide; GI<sub>50</sub>, concentration that inhibits 50% human tumor cell growth; HRMS, high resolution mass spectra; HTCL, human tumor cell line; IHC, immunohistochemistry; MW, microwave; PDB, Protein Data Bank; SAR, structure–activity relationship; tumor-VDAs, tumor vascular disrupting agents

## REFERENCES

- (1) Perez-Perez, M. J.; Priego, E. M.; Bueno, O.; Martins, M. S.; Canela, M. D.; Liekens, S. Blocking Blood Flow to Solid Tumors by Destabilizing Tubulin: an Approach to Targeting Tumor Growth. *J. Med. Chem.* **2016**, *59*, 8685–8711.
- (2) Kaur, R.; Kaur, G.; Gill, R. K.; Soni, R.; Bariwal, J. Recent Developments in Tubulin Polymerization Inhibitors: an Overview. *Eur. J. Med. Chem.* **2014**, *87*, 89–124.
- (3) Tozer, G. M.; Kanthou, C.; Baguley, B. C. Disrupting Tumor Blood Vessels. *Nat. Rev. Cancer* **2005**, *5*, 423–435.
- (4) Siemann, D. W. The Unique Characteristics of Tumor Vasculature and Preclinical Evidence for Its Selective Disruption by Tumor-Vascular Disrupting Agents. *Cancer Treat. Rev.* **2011**, *37*, 63–74.
- (5) Mason, R. P.; Zhao, D.; Liu, L.; Trawick, M. L.; Pinney, K. G. A Perspective on Vascular Disrupting Agents That Interact with Tubulin: Preclinical Tumor Imaging and Biological Assessment. *Integr. Biol.* **2011**, *3*, 375–387.
- (6) Wang, X. F.; Xie, L. Vascular Disrupting Agents Targeting at Tubulin: a Novel Class of Antitumor Drug. *J. Int. Pharm. Res.* **2012**, *39*, 445–454.
- (7) Siemann, D. W.; Chaplin, D. J.; Walicke, P. A. A Review and Update of the Current Status of the Vasculature-Disabling Agent Combretastatin-A4 Phosphate (CA4P). *Expert Opin. Invest. Drugs* **2009**, *18*, 189–197.
- (8) Galbraith, S. M.; Chaplin, D. J.; Lee, F.; Stratford, M. R.; Locke, R. J.; Vojnovic, B.; Tozer, G. M. Effects of Combretastatin A4 Phosphate on Endothelial Cell Morphology In Vitro and Relationship to Tumour Vascular Targeting Activity In Vivo. *Anticancer Res.* **2001**, *21*, 93–102.
- (9) Bates, D.; Feris, E. J.; Danilov, A. V.; Eastman, A. Rapid Induction of Apoptosis in Chronic Lymphocytic Leukemia Cells by the Microtubule Disrupting Agent BNC105. *Cancer Biol. Ther.* **2016**, *17*, 291–299.
- (10) Wang, X. F.; Ohkoshi, E.; Wang, S. B.; Hamel, E.; Bastow, K. F.; Morris-Natschke, S. L.; Lee, K. H.; Xie, L. Synthesis and Biological Evaluation of *N*-Alkyl-*N*-(4-methoxyphenyl)pyridin-2-amines as a New Class of Tubulin Polymerization Inhibitors. *Bioorg. Med. Chem.* **2013**, *21*, 632–642.
- (11) Wang, X. F.; Wang, S. B.; Ohkoshi, E.; Wang, L. T.; Hamel, E.; Qian, K.; Morris-Natschke, S. L.; Lee, K. H.; Xie, L. Discovery of *N*-Aryl-6-methoxy-1,2,3,4-tetrahydro-quinolines as a Novel Class of Antitumor Agents Targeting the Colchicine Site of Tubulin. *Eur. J. Med. Chem.* **2013**, *67*, 196–207.
- (12) Wang, S. B.; Wang, X. F.; Qin, B.; Ohkoshi, E.; Hsieh, K. Y.; Hamel, E.; Cui, M. T.; Zhu, D. Q.; Goto, M.; Morris-Natschke, S. L.; Lee, K. H.; Xie, L. Optimization of *N*-Aryl-6-methoxy-1,2,3,4-tetrahydroquinolines as Tubulin Polymerization Inhibitors. *Bioorg. Med. Chem.* **2015**, *23*, 5740–5747.
- (13) Wang, X. F.; Guan, F.; Ohkoshi, E.; Guo, W.; Wang, L.; Zhu, D. Q.; Wang, S. B.; Wang, L. T.; Hamel, E.; Yang, D.; Li, L.; Qian, K.; Morris-Natschke, S. L.; Yuan, S.; Lee, K. H.; Xie, L. Optimization of 4-(*N*-Cycloamino)phenylquinazolines as a Novel Class of Tubulin Polymerization Inhibitors Targeting the Colchicine Site. *J. Med. Chem.* **2014**, *57*, 1390–1402.
- (14) Lockman, J. W.; Klimova, Y.; Anderson, M. B.; Willardsen, J. A. Synthesis of Substituted Quinazolines: Application to the Synthesis of Verubulin. *Synth. Commun.* **2012**, *42*, 1715–1723.
- (15) Yang, Z.; Wang, T.; Wang, F.; Niu, T.; Liu, Z.; Chen, X.; Long, C.; Tang, M.; Cao, D.; Wang, X.; Xiang, W.; Yi, Y.; Ma, L.; You, J.; Chen, L. Discovery of Selective Histone Deacetylase 6 Inhibitors Using the Quinazoline as the Cap for the Treatment of Cancer. *J. Med. Chem.* **2016**, *59*, 1455–1470.
- (16) Shoemaker, R. H. The NCI-60 Human Tumour Cell Line Anticancer Drug Screen. *Nat. Rev. Cancer* **2006**, *6*, 813–823.
- (17) Devambatla, R. K.; Namjoshi, O. A.; Choudhary, S.; Hamel, E.; Shaffer, C. V.; Rohena, C. C.; Mooberry, S. L.; Gangjee, A. Design, Synthesis, and Preclinical Evaluation of 4-Substituted-5-methyl-furo-[2,3-*d*]pyrimidines as Microtubule Targeting Agents that are Effective Against Multidrug Resistant Cancer Cells. *J. Med. Chem.* **2016**, *59*, 5752–5765.
- (18) The data were provided by Concord PharmaTech Co., Ltd. Nanjing, China.
- (19) Pérez-Sayáns, M.; Somoza-Martín, J. M.; Barros-Angueira, F.; Diz, P. G.; Rey, J. M. G.; García-García, A. Multidrug Resistance in Oral Squamous Cell Carcinoma: the Role of Vacuolar ATPases. *Cancer Lett.* **2010**, *295*, 135–143.
- (20) Hung, H. Y.; Ohkoshi, E.; Goto, M.; Bastow, K. F.; Nakagawa-Goto, K.; Lee, K. H. Antitumor Agents. 293. Nontoxic Dimethyl-4,4'-dimethoxy-5,6,5',6'-dimethylenedioxybiphenyl-2,2'-dicarboxylate (DDB) Analogues Chemosensitize Multidrug-resistant Cancer Cells to Clinical Anticancer Drugs. *J. Med. Chem.* **2012**, *55*, 5413–5424.
- (21) Rubinstein, L. V.; Shoemaker, R. H.; Paull, K. D.; Simon, R. M.; Tosini, S.; Skehan, P.; Scudiero, D. A.; Monks, A.; Boyd, M. R. Comparison of In Vitro Anticancer-Drug-Screening Data Generated with a Tetrazolium Assay Versus a Protein Assay Against a Diverse Panel of Human Tumor Cell Lines. *J. Natl. Cancer Inst.* **1990**, *82*, 1113–1117.
- (22) Sun, L. Q.; Zhu, L.; Qian, K.; Qin, B.; Huang, L.; Chen, C. H.; Lee, K. H.; Xie, L. Design, Synthesis, and Preclinical Evaluations of Novel 4-Substituted 1,5-Diarylanilines as Potent HIV-1 Non-Nucleoside Reverse Transcriptase Inhibitor (NNRTI) Drug Candidates. *J. Med. Chem.* **2012**, *55*, 7219–7229.
- (23) Kerns, E. H.; Di, L. *Drug-like Properties: Concepts, Structure Design and Methods From ADME to Toxicity Optimization*; Academic Press: Boston, 2008; pp 43–47.
- (24) Hamel, E. Evaluation of Antimitotic Agents by Quantitative Comparisons of Their Effects on the Polymerization of Purified Tubulin. *Cell Biochem. Biophys.* **2003**, *38*, 1–21.
- (25) Lin, C. M.; Ho, H. H.; Pettit, G. R.; Hamel, E. Antimitotic Natural Products Combretastatin A-4 and Combretastatin A-2: Studies on the Mechanism of Their Inhibition of the Binding of Colchicine to Tubulin. *Biochemistry* **1989**, *28*, 6984–6991.
- (26) Anderson, M. B. *Preparation of 4-Arylamino-quinazolines as Activators of Caspases and Inducers of Apoptosis*. U.S. Pat. Appl. Publ., 20100069383, March 18, 2010.
- (27) Boyd, M. R. Status of the NCI Preclinical Antitumor Drug Discovery Screen. In *Cancer: Principles and Practice of Oncology Updates*; Devita, V. T., Hellman, S., Rosenberg, S. A., Eds.; Lippincott Williams & Wilkins: Philadelphia, 1989; pp 1–12.
- (28) Monks, A.; Scudiero, D.; Skehan, P.; Shoemaker, R.; Paull, K.; Vistica, D.; Hose, C.; Langley, J.; Cronise, P.; Vaigro-Wolff, A.; Gray-Goodrich, M.; Campbell, H.; Mayo, J.; Boyd, M. Feasibility of a High-Flux Anticancer Drug Screen Using a Diverse Panel of Cultured Human Tumor Cell Lines. *J. Natl. Cancer Inst.* **1991**, *83*, 757–766.
- (29) Houghton, P.; Fang, R.; Techatanawat, I.; Steventon, G.; Hylands, P. J.; Lee, C. C. The Sulphorhodamine (SRB) Assay and Other Approaches to Testing Plant Extracts and Derived Compounds for Activities Related to Reputed Anticancer Activity. *Methods* **2007**, *42*, 377–387.
- (30) Nakagawa-Goto, K.; Oda, A.; Hamel, E.; Ohkoshi, E.; Lee, K. H.; Goto, M. Development of a Novel Class of Tubulin Inhibitor from Desmosmumotin B with a Hydroxylated Bicyclic B-ring. *J. Med. Chem.* **2015**, *58*, 2378–2389.
- (31) Verdier-Pinard, P.; Lai, J.-Y.; Yoo, H.-D.; Yu, J.; Márquez, B.; Nagle, D. G.; Nambu, M.; White, J. D.; Falck, J. R.; Gerwick, W. H.; Day, B. W.; Hamel, E. Structure-Activity Analysis of the Interaction of Curacin A, the Potent Colchicine Site Antimitotic Agent, with Tubulin and Effects of Analogs on the Growth of MCF-7 Breast Cancer Cells. *Mol. Pharmacol.* **1998**, *53*, 62–76.