### Accepted Manuscript

Synthesis and evaluation of the diarylthiourea analogs as novel anti-cancer agents

Shengquan Liu, Maggie C. Louie, Vanishree Rajagopalan, Guangyan Zhou, Esmeralda Ponce, Tran Nguyen, Linda Green

PII:	S0960-894X(15)00054-2				
DOI:	http://dx.doi.org/10.1016/j.bmcl.2015.01.042				
Reference:	BMCL 22382				
To appear in:	Bioorganic & Medicinal Chemistry Letters				
Received Date:	14 November 2014				
Revised Date:	17 January 2015				
Accepted Date:	20 January 2015				



Please cite this article as: Liu, S., Louie, M.C., Rajagopalan, V., Zhou, G., Ponce, E., Nguyen, T., Green, L., Synthesis and evaluation of the diarylthiourea analogs as novel anti-cancer agents, *Bioorganic & Medicinal Chemistry Letters* (2015), doi: http://dx.doi.org/10.1016/j.bmcl.2015.01.042

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**FULL TITLE:** Synthesis and evaluation of the diarylthiourea analogs as novel anti-cancer agents

### **AUTHORS:**

Shengquan Liu<sup>a,\*</sup>, Maggie C. Louie<sup>a,c</sup>, Vanishree, Rajagopalan<sup>a</sup>, Guangyan Zhou<sup>b</sup>, Esmeralda Ponce<sup>c</sup>, Tran Nguyen<sup>a</sup>, and Linda Green<sup>c</sup>

<sup>a</sup>College of Pharmacy, Touro University-California, Vallejo, CA 94592, USA <sup>b</sup>Department of Basic Science, Touro University-California, Vallejo, CA 94592, USA <sup>c</sup>Department of Natural Science and Mathematics, Dominican University of California, San Rafael, CA 94901, USA 

\*Corresponding author Phone: (707) 638-5983 Fax: (707) 638-5959 Email: shengquan.liu@tu.edu

\*Reprint requests should be addressed to the corresponding author.

**KEYWORDS:** SHetA2; diarylthiourea analogs; anticancer agents; breast cancer; prostate cancer

### **ABSTRACT:**

Ten *p*-nitrodiarylthiourea analogs were designed, synthesized and evaluated in breast (MCF-7, T-47D, MDA-MB-453) and prostate (DU-145, PC-3, LNCaP) cancer cell lines for their anticancer activities. The majority of the compounds were able to inhibit the growth of these six cancer cell lines at low micromolar concentrations. Compound **7** was found to be the most potent anticancer agent in this series with  $GI_{50}$  values of 3.16  $\mu$ M for MCF-7, 2.53  $\mu$ M for T-47D, 4.77  $\mu$ M for MDA-MB-453 breast cancer lines and 3.54  $\mu$ M for LNCaP prostate cancer cell line. These  $GI_{50}$  values were comparable to the parent compound, SHetA2.

### **GRAPHICAL ABSTRACT:**



### **ABBREVIATIONS:**

SHetA2: *N*-(2,3-dihydro-2,2,4,4-tetramethyl-6-benzothiopyranyl), *N*'-(4-nitrophenyl)thiourea; ATRA: all *trans* retinoic acid; NOAEL: no observed adverse effect level; MTT: 3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide; MTS: MTT coupled with phenazine methosulfate (soluble form of MTT); ER: estrogen receptor; AR: androgen receptor.

Previously, we synthesized *N*-(2,3-dihydro-2,2,4,4-tetramethyl-6-benzothiopyranyl), *N*'-(4nitrophenyl)thiourea (SHetA2, NSC 726189)<sup>1</sup> which has been shown to be a novel potential cancer prevention agent and is now in preclinical development for cancer prevention through the National Cancer Institute (NCI) Rapid Access to Intervention Development (RAID) program. SHetA2 evolved from a lead optimization process from all *trans*-retinoid acid (ATRA, Tretinoin<sup>®</sup>, mean

GI<sub>50</sub> is 51 µM, Figure 1),<sup>24</sup> a drug used for the treatment of acute promyelocytic leukemia. However, the exploitation of retinoic acid's full potential as a chemopreventive and/or a therapeutic drug, particularly against solid tumors, has been hampered mainly by its local and systemic toxicity and side effects often associated with its ability to activate nuclear retinoid receptors.<sup>2</sup> Thus, efforts have been made to develop novel compounds that retain retinoid anticancer activity with reduced retinoid toxicity and side effects. One such compound is SHetA2, a more effective anticancer agent than ATRA and appears to function without activating retinoid receptors.<sup>1</sup> As a result, SHetA2 does not exhibit retinoid toxicities when tested in animal models.<sup>3</sup> SHetA2 exerts its selective anticancer activity through regulating apoptosis, cell growth, differentiation and angiogenesis.<sup>5,6</sup> SHetA2 has also been shown to induce proteasomal degradation of cyclin D1,<sup>7</sup> generate mitochondrial swelling and endoplasmic reticulum stress, promote the formation of reactive oxygen species, and induce apoptosis in cancer cells while sparing normal cells.<sup>6</sup> More importantly, it was reported recently in a dog study, that no toxicity of SHetA2 was observed in any tested dose groups. The lowest observed adverse effect level (NOAEL) for ShetA2 was not established and was considered to be above 1,500 mg/kg/day.<sup>8</sup>

However, SHetA2 may not be an ideal drug candidate due to the following two limitations: its high lipophilicity and cumbersome six step synthesis. SHetA2's high lipophilicity (LogP =7.09) is higher than the upper limit of "Lipinski rule of 5" and the Log P of most marketed drugs,<sup>9,10</sup> and this might contribute to its extremely low (<1%) systemic bioavailability for all doses tested in rat and its high plasma protein binding (99.3-99.5% at low micromolar concentrations).<sup>6,11</sup> This high lipophilicity may also cause nonselectivity, liver toxicity and drug-drug interactions.<sup>12</sup> Furthermore, the synthesis of SHetA2 involves 6 steps, with low overall yield of 3%, which hinders its large scale supply and the synthesis of new analogs.<sup>13</sup> To address these issues, we have modified the SHetA2 structure based on the previous structure-activity relationship.<sup>1.2</sup> We hypothesized that the nitrophenyl group and the thiourea linker in SHetA2 structure are important for its anticancer activity. Thus, our modification strategy was to keep these two moieties intact, while replacing the thiochromane ring with another ring structure. The ring systems were selected so that designed compounds follow the Lipinski rule, are drug-like, but are not reactive and do not produce reactive metabolites. In this study, we selected aromatic ring systems so that the designed compounds resemble SHetA2 but differ in their sizes and shapes. In this way, the structure modification will

reduce the lipophilicity of the designed compounds and simplify the six-step synthesis to one step using readily available and inexpensive starting materials.

Here, we report the design, synthesis and biological activity of ten diarylthiourea analogs. These compounds were synthesized according to our previously published procedure.<sup>1</sup> The newly synthesized compounds were evaluated for their ability to inhibit the growth of breast cancer cell lines-MCF-7, T-47D, and MDA-MB-453 and prostate cancer cell lines-DU-145, LNCaP and PC-3. The potency of compounds **5** and **7** in this series were shown to be comparable to the parent compound, SHetA2.

General procedure for the preparation of compounds **1-10** with the overall yield of 19%-87% from amines and 4-nitrophenyl isothiocyanate is shown in Scheme 1. To a solution of amines in dry THF at 0°C under nitrogen, 4-nitrophenyl isothiocyanate in dry THF was added dropwise. After the reaction was completed, the solvent was evaporated; the residue was recrystallized or purified by flash column chromatography to give the corresponding thiourea product. The chemical structures of compounds **1-10** were confirmed by <sup>1</sup>H, C<sup>13</sup> NMR and LC-MS. The Log P, calculated Log P and the overall yield is summarized in Table 1. Further details of reactions, characterizations, and bioassays are provided in the Supplementary Data.

The preclinical studies of SHetA2 on cell culture and animal models showed that it is a promising agent for cancer prevention and treatment without significant toxicity and side effects.<sup>6-8</sup> However, SHetA2 is too lipophilic, with a Log P value of 7.09, which can hinder its oral bioavailability as an anticancer drug, since optimal Log P for most oral drugs is between 1-3. We have designed the SHetA2 analogs with Log P value between 2.99 and 6.36 (Table 1) so that all our compounds are less lipophilic than the lead compound SHetA2 with the premise that these compounds would have better pharmacokinetic profiles. In addition, as previously reported, the six step synthesis of SHetA2 suffers from a low overall yield of 3% and this procedure involves use of reagents that are toxic and have unpleasant odor.<sup>8</sup> Tallent's group reported a modified procedure for the synthesis of key intermediates to the SHetA2 with improved yield, but started with much more expensive chemicals and used column chromatography after every synthetic step.<sup>13</sup> We have developed a simple, one step synthetic procedure of the target compounds with cheap and commercially available chemicals with overall yield of 19%-87% (Table 1). Please note that compounds 1-10 were prepared in one step whereas SHetA2 required 6 steps. By far in the

published literature, only certain diarylurea multi-target kinase inhibitors have chemical structures related to our compounds.<sup>14,15</sup> Sorafenib is among one of these compounds that have been approved for the treatment of advanced renal cell cancer by the FDA. In addition, the wide diversity of cellular and molecular targets that can be regulated by judiciously modifying *N*, *N*'-diarylureas and ureas, suggests this part of the molecule may serve as privileged scaffolds for anticancer agents.<sup>14,15</sup>

The synthesized diarylthiourea analogs **1-10** were tested on three breast cancer cell lines — MCF7, T-47D and MDA-MB-453 cells (Table 2). While the majority of the compounds showed some effect on inhibiting growth of all three breast cancer cell lines, diarylthiourea analogs **5** and **7** showed the most significant effects. The GI<sub>50</sub> values of compounds **5** and **7** were comparable to the parent SHetA2 (2.94-6.27  $\mu$ M and 2.53-4.77  $\mu$ M vs. 3.27-4.13  $\mu$ M, respectively), with the T-47D cells displaying the greatest sensitivity to these compounds when compared to MCF7 and MDA-MB-453. Since compound **7** appeared to be the most effective in inhibiting the growth of all three breast cancer cell lines, its ability to inhibit cell growth over a longer period of time was evaluated (Figure 2 A-C). Results in figure 2 indicate that compound **7** continued to inhibit cell growth 4 days after treatment and appeared to effectively inhibit the growth of all three breast cancer cell lines.

The diarylthiourea analogs of SHetA2 were also tested on three prostate cancer cell lines— DU-145, PC-3 and LNCaP. Overall, LNCaP cells were more susceptible to the growth inhibitory effects of these agents (Table 2). Compound 7 showed the most potent growth inhibitory effect in LNCaP cells as evidenced by a low GI<sub>50</sub> value of  $3.54 \mu$ M. This value was comparable to the GI<sub>50</sub> value (2.25  $\mu$ M) of SHetA2 in the same cell-line (Table 2). Being the most potent of the diarylthiourea analogs, compound 7 was tested for its ability to inhibit cell growth over a longer period of time (Figure 2D). This experiment was carried out in LNCaP cells that were most vulnerable to the growth inhibitory effects of compound 7. Cell growth was significantly inhibited with 5  $\mu$ M compound 7 staring at day 2 and remained so until day 4. A similar trend in growth inhibition was observed at higher concentrations (10 and 20  $\mu$ M) as well (Figure 2D). Moreover, while SHetA2 inhibited the growth of all three prostate cancer cell-lines uniformly, compound **5** and **7** selectively inhibited the growth of just one of the prostate cancer cell line, the LNCaPs.

Here we report a simple, one step synthesis procedure to obtain a variety of diarylthiourea derivatives of SHetA2 and their biological evaluation in human breast and prostate cancer cells. The simple syntheses involved a coupling reaction of 4-nitrophenyl isothiocyanate in dry THF with commercially available amines. Most of these diarylthiourea derivatives of SHetA2 caused growth

inhibition of both human breast and prostate cancer cells. Among them, compounds 5 and 7 showed potency comparable to the lead compound, SHetA2. The study also validated our previous hypothesis that the 4-nitrophenyl group and the thiourea linker are important for the anticancer activity of SHetA2 because all the compounds tested have these moieties. Modification of other functional groups in SHetA2 resulted in active anticancer agents. Therefore, this study has provided a new approach to the design and synthesis of the next generation of novel anticancer agents, perhaps with novel targeted mechanisms of action.

We tested the new diarylthiourea analogs along with SHetA2 on three breast cancer and three prostate cancer cell lines and our results (Table 2) show that diarylthiourea analog 5 and 7 are effective in inhibiting cancer cell growth of four cancer cell lines-MCF7, T-47D, MDA-MB-453 (breast cancer cells) and LNCaP (prostate cancer cell). SHetA2 was previously evaluated in the panel of National Cancer Institute (NCI) human tumor cell lines by the Developmental Therapeutics Program (DTP) and results showed that SHetA2 inhibited growth of most cancer cells in the micromolar range. Specifically, reported GI<sub>50</sub> values of 4.5 µM for MCF-7, 4.8 µM for T-47D, 5.0 µM for MDA-MB-453, 4.9 µM for DU145 and 5.0 µM for PC-3,<sup>6,16</sup> are consistent with our results shown in Table 2. Our newly synthesized compounds (5 and 7) showed equivalent potency in comparison to the lead compound SHetA2 (Table 2). Noteworthy are the effects of compounds 5 and 7 on the three prostate cancer cell lines. Both these agents have shown to be more selective in inhibiting the growth of LNCaP cells in comparison to PC-3 and DU-145 cell-lines. As seen in Table 2, compound 7 is about six times more potent in inhibiting growth of LNCaP cells than PC-3 and DU145 cells and compound 5 is at least twice as potent in LNCaP cells than in PC-3 and DU-145 cells. This observation is significant since SHetA2 does not show any such selectivity (Table 2). One explanation for this selectivity could be due to differential androgen receptor signaling patterns in the three prostate cancer cell lines. Both PC-3 and DU-145 cell-lines do not express androgen receptors (AR) and are not androgen dependent for their growth, while LNCaP cells express AR (with a point mutation in the ligand binding domain) and are considered androgendependent cells<sup>17</sup>. Therefore, we hypothesize that these new analogs, compounds **5** and **7**, may induce growth inhibition in prostate cancer cell lines by acting on specific targets in these cells. Perhaps the molecular target is androgen receptor, since in addition to the LNCaP cells, all three breast cancer cell lines that responded to the growth inhibitory effects of compounds 5 and 7

express AR. Whether cell growth inhibition by compounds 5 and 7 involves the regulation of androgen receptors (AR) requires further investigation.

However, this differential response was not as significant in the breast cancer models. MCF7 and T-47D cells are both estrogen receptor (ER) positive (presence of ER $\alpha$ ) and are considered hormone-dependent, whereas MDA-MB-453 is ER negative (absence of ER $\alpha$ ). While the GI<sub>50</sub> for the ER positive cells were lower in comparison to the ER negative breast cancer cells, the difference is not statistically significant (Table 2). The differences observed in the breast and prostate cancer cells might be attributed to different mechanisms of action in the two cancer types. Although MDA-MD-453 is considered ER negative breast cancer cells, similar to MCF7 and T-47D, these cells do express ER $\beta$ .

In conclusion, this is the first report that demonstrates the growth inhibitory properties of SHetA2 and a new generation of diarylthiourea analogs on both breast and prostate cancer cells. While compounds **1-10** are not more potent than the SHetA2, we report for the first time the effects of *p*-nitrodiarylthiourea compounds on the inhibition of both breast and prostate cancer cells, along with a more simplified synthetic scheme. Data from this study indicate that these novel potential anticancer agents are promising lead compounds for further evaluation. While the mechanism of action, molecular targets and pharmacokinetic profile of these compounds remain to be elucidated in future studies, results presented in this study provide a strong foundation for further preclinical studies of these compounds as potential therapeutic agents for both breast and prostate cancers.

### Acknowledgments

This work was support by grant funding from Touro University-CA Intramural Research Award Program (IRAP) to S.L. and V.R. The authors would like to thank Dr. Michael Ellerby (Touro University-CA) for his guidance and generous gift of LNCaP, PC-3 and DU-145 prostate cancer cell lines and Dr. Miriam Gochin Touro University-CA for providing the NMR spectra.

#### **References and notes**

- Liu, S.; Brown, C. W.; Berlin, K. D.; Dhar, A.; Guruswamy, S.; Brown, D.; Gardner, G. J.; Birrer, M. J.; Benbrook, D. M. J. Med. Chem. 2004, 47, 999.
- 2. Dawson, M. I.; Eontana, J. A. Mini-Rev. Med. Chem. 2010, 10, 455.

- Abraham, D. J. (ed): Vitamins. In Burgers' Medicinal Chemistry and drug discovery; 7<sup>th</sup> ed.. Hoboken, NH, John Wiley and Sons, 2010, Vol 6, pp 500 563.
- 4. Benbrook, D. M. Mini-Rev. Med. Chem. 2002, 2, 277.
- Myers, T.; Chengedza, S.; Lightfoot, S.; Pan, Y.; Dedmond, D.; Cole, L.; Tang, Y.; Benbrook, D. M. Invest. New Drugs. 2009, 27, 304.
- Benbrook, D. M.; Kamelle, S. A.; Guruswamy, S. B.; Lightfoot, A. A.; Rutledge, T. L.; Could, N. S.; Hannafon, B. N.; Dunn, S. T.; Berlin, D. D. Invest. New Drugs. 2005, 23, 417.
- 7. Masamha, C. P.; Benbrook, D. M. Cancer Res. 2009, 69(16), 6565.
- Kabirov, K. K.; Kapetanovic, I. M.; Benbrook, D, M.; Dinger, N.; Mankovskaya, I.; Zakharov, A.; Detrisac, C.; Pereira, M.; Martin-Jimenez, T.; Onua, E.; Banerjee, A.; van Breemen, R. B.; Nikolic, D.; Chen, L.; Lyubimov, A, V. Drug Chem. Toxicol. 2013, 36(3), 284.
- 9. Lipinski, C. A.; Lombardo, F.; Dominy, B. W.; Feeney, P. J. Adv. Drug Delivery. Rev. 1997, 23, 3.
- 10. Lipinski, C. A. Drug Discov. Today Technol. 2004, 1, 337.
- 11. Leeson, P. D.; Springthorpe, B. Nat. Rev. Drug Disc. 2007, 6, 881.
- 12. Zhang, Y.; Hua, Y.; Benbrook, D. M.; Covey, J. M.; Dai, G.; Liu, Z.; Chan, K. K. Cancer Chemother. Pharmacol. 2006, 58, 561.
- Tallent, C. R.; Twine, C. E.; Risbood, P.; Seltzman, H. H. Org. Prep. Proced. Int. 2007, 39, 96.
- Denoyelle, D.; Chen, T.; Chen, L.; Wang, Y.; Klosi, E.; Halperin, J.; Aktas, B. H.; Chorev, M. Bioorg. Med. Chem. Lett. 2012, 22(1), 402.
- 15. Li, H.; Lv, P.; Yan, T.; Zhu, H. Anticancer Agents in Med. Chem. 2009, 9, 471.
- 16. Website of Developmental Therapeutics Program of NCI: http://dtp.nci.nih.gov/dtpstandard/servlet/MeanGraph?searchtype=NSC&searchlist=726189 &outputformat=HTML&outputmedium=page&chemnameboolean=AND&debugswitch=fal se&assaytype=&testshortname=NCI+Cancer+Screen+Current+Data&dataarraylength=59& endpt=GI50&button=Mean+Graph&highconc=-4.0. Accessed April 18, 2014.
- 17. Webber, M. M.; Bello, D.; Quader, S. Prostate. 1996, 29(6), 386.

**Supplementary Material:** The synthesis procedures, spectrum data of each compound and all experimental information about bioassay are provided as a separate electronic file. It can be transformed into PDF format.

### FIGURE LEGENDS

Scheme 1: General procedure for the synthesis of the compounds 1-10.

Figure 1: Evolution of SHetA2.

Figure 2: Breast cancer (A) MDA-MB-453, (B) MCF7, (C) T-47D, and prostate cancer (D) LNCaP cells were plated in 96-plates and treated with varying concentrations of SL-01-18 for 2, 3 and 4 days. Cell growth was analyzed with MTT (A-C) or MTS (D). Means and standard deviations represent at least two independent experiments done in six replicates.

MAS

Compounds	Name	Log P <sup>a</sup>	CLog P <sup>a</sup>	Yield (%)	
1	SL-01-03	3.83	3.51	83	
2	SL-01-05	ND	5.01	19	
3	SL-01-06	3.84	3.05	23	
4	SL-01-08	ND	2.83	24	
5	SL-01-09	4.11	4.52	87	
6	SL-01-15	5.34	5.39	35	6
7	SL-01-18	6.36	6.08	42	6
8	SL-01-19	5.24	4.91	70	
9	SL-01-20	3.59	3.54	60	
10	SL-01-21	2.88	2.66	73	
Parent	SHetA2	7.09	5.74	3	

### Table 1. SHetA2 analogs (1-10) produced according to Scheme 1

\_\_5.74 .iDraw.

Name and Chemical Structure	Breast cancer cell lines (μM) Mean±SD			Prostate cancer cell lines (μM) Mean±SD		
	MCF-7	T-47D	MDA-MB-453	DU-145	PC-3	LNCaP
$1 \\ \underset{H_3C}{\overset{H_3C}{\longrightarrow}} \overset{H}{\underset{S}{\longrightarrow}} \overset{H}{\underset{S}{\longrightarrow}} \overset{H}{\underset{NO_2}{\longrightarrow}} $	16.96 ±3.35 <sup>1</sup>	8.14±1.79 <sup>2</sup>	13.48±1.96 <sup>1</sup>	35.15±15.42	26.77±13.25 <sup>1</sup>	10.58±1.95 <sup>1</sup>
$2 \\ ( \mathbf{A} \mathbf{A} \mathbf{A} \mathbf{A} \mathbf{A} \mathbf{A} \mathbf{A} \mathbf{A}$	12.81±2.46 <sup>1</sup>	9.44±0.12 <sup>1</sup>	6.88±0.83 <sup>1</sup>	5.87±0.92 <sup>2</sup>	19.59±0.68 <sup>1</sup>	5.86±0.41 <sup>2</sup>
3 H <sub>9</sub> C (1) (1) (1) (1) (1) (1) (1) (1) (1) (1)	ND*	21.62±0.85 <sup>1</sup>	ND*	ND*	ND*	40.47±28.82 <sup>1</sup>
$\frac{4}{\sqrt[N_{s}]{1}} + \sqrt[H_{s}]{1} + \sqrt[H_{s}]{1} + \sqrt[N_{s}]{1} + \sqrt[N_{s}]$	ND*	15.28±3.27 <sup>1</sup>	ND*	ND*	ND*	23.53±7.11 <sup>1</sup>
5 H <sub>3</sub> C H K NO <sub>2</sub>	6.21±1.95 <sup>1</sup>	2.94 ±0.91 <sup>2</sup>	6.27±0.39 <sup>1</sup>	13.69±4.15 <sup>1</sup>	13.90±1.09 <sup>2</sup>	6.05±0.18 <sup>2</sup>
	7.81±1.19 <sup>2</sup>	3.96 ±0.58 <sup>2</sup>	6.20±2.42 <sup>2</sup>	17.48±2.20 <sup>1</sup>	16.50±1.17 <sup>1</sup>	11.32±3.90 <sup>1</sup>
	3.16±1.79 <sup>2</sup>	2.53±0.39 <sup>2</sup>	4.77±2.21 <sup>2</sup>	21.78±3.98 <sup>1</sup>	19.96±3.32 <sup>1</sup>	3.54±0.17 <sup>2</sup>
	13.30±2.73 <sup>1</sup>	11.55±3.87 <sup>1</sup>	14.49±4.58 <sup>1</sup>	6.11±1.30 <sup>2</sup>	19.27±0.41 <sup>1</sup>	7.29±2.35 <sup>2</sup>
	13.08±1.74 <sup>1</sup>	8.68±2.34 <sup>1</sup>	27.59±7.22 <sup>1</sup>	24.10±5.40 <sup>1</sup>	22.56±3.98 <sup>1</sup>	12.75±0.60 <sup>1</sup>
	9.89±1.62 <sup>2</sup>	ND*	ND*	ND*	ND*	ND*
	4.13±0.57 <sup>2</sup>	3.99±0.06 <sup>2</sup>	3.27±0.33 <sup>2</sup>	3.47±0.42 <sup>2</sup>	4.80±1.53 <sup>2</sup>	2.25±1.04 <sup>2</sup>
All Trans-RA	12.39±0.26 <sup>1</sup>	0.82±0.03 <sup>2</sup>	9.51±0.61 <sup>1</sup>	11.64±0.54 <sup>1</sup>	12.64±0.38 <sup>1</sup>	10.33±0.75 <sup>1</sup>

### Table 2. Potency (GI\_{50} values) in $\mu M$

<sup>1</sup> Based on linear model <sup>2</sup> Based on exponential model \* ND=Not determined or GI50 >>50µM



#### Figure 1





Accepter

Figure 2

