



Novel steroid sulfatase inhibitors based on *N*-thiophosphorylated 3-(4-aminophenyl)-coumarin-7-*O*-sulfamates

Mateusz Daśko¹  | Sebastian Demkowicz²  | Karol Biernacki² | Amira Harrous² | Janusz Rachon² | Witold Kozak³ | Aleksandra Martyna⁴ | Maciej Maśłyk⁴ | Konrad Kubiński⁴ | Anna Boguszevska-Czubara⁵

¹Department of Inorganic Chemistry, Faculty of Chemistry, Gdansk University of Technology, Gdansk, Poland

²Department of Organic Chemistry, Faculty of Chemistry, Gdansk University of Technology, Gdansk, Poland

³Department of Physical Chemistry, Faculty of Chemistry, University of Gdansk, Gdansk, Poland

⁴Department of Molecular Biology, Faculty of Biotechnology and Environment Sciences, The John Paul II Catholic University of Lublin, Lublin, Poland

⁵Department of Medical Chemistry, Medical University of Lublin, Lublin, Poland

Correspondence

Mateusz Daśko, Department of Inorganic Chemistry, Faculty of Chemistry, Gdansk University of Technology, Narutowicza 11/12, 80-233 Gdansk, Poland.
Email: mateusz.dasko@pg.edu.pl

Sebastian Demkowicz, Department of Organic Chemistry, Faculty of Chemistry, Gdansk University of Technology, Narutowicza 11/12, 80-233 Gdansk, Poland.
Email: sebdemko@pg.gda.pl

Funding information

Narodowe Centrum Nauki, Grant/Award Numbers: 2015/19/N/NZ7/00938, 2017/24/T/NZ7/00271; Foundation for Polish Science; National Science Centre

Abstract

In the present work, we described convenient methods for the synthesis of *N*-thiophosphorylated 3-(4-aminophenyl)-coumarin-7-*O*-sulfamates as steroid sulfatase (STS) inhibitors. To design the structures of the potential STS inhibitors, molecular modeling techniques were used. A computational docking method was used to determine the binding modes of the synthesized inhibitors as well as to identify potential interactions between specified functional groups on the inhibitors and the amino acid residues present in the active site of the enzyme. The inhibitory activities of the synthesized compounds were tested in an enzymatic assay with STS isolated from a human placenta. Within the set of newly synthesized compounds, **9e** demonstrated the highest inhibitory activity in the enzymatic assay with an IC₅₀ value of 0.201 μM (the IC₅₀ value of **667-COUMATE** in the same test was 0.062 μM). Furthermore, we tried to verify if the obtained STS inhibitors are able to pass through the cellular membrane effectively in cell line experiments. In the course of our study, we determined the STS activity in the MCF-7 cell line after incubation in the presence of the inhibitors (at 100 nM concentration). For this evaluation, we included newly synthesized compounds **9a-g** and their *N*-phosphorylated analogs **6a-h**, whose synthesis has been previously described. We found that the lowest STS activities were measured in the presence of *N*-phosphorylated derivatives **6e** (0.1% of STS activity) and **6f** (0.2% of STS activity). The measured STS activity in the presence of **667-COUMATE** (used as a reference) was 0.1%. Moreover, at concentrations up to 1 μM, the most active compounds (**6e**, **6f**, **9b**, and **9e**) did not exert any toxic effects on zebrafish embryos.

KEYWORDS

breast cancer, coumarin sulfamates, steroid sulfatase, STS inhibitors

1 | INTRODUCTION

The STS enzyme plays a crucial role in the hydrolysis of biologically inactive steroid sulfates of into their unsulfated derivatives with

biological activity. The activity of STS is responsible for the production of steroidal hormones, which induce the proliferation of cancer cells in a number of hormone-dependent cancers (Shah, Singh, Singh, Singh Jaggi, & Singh, 2016). Inhibition of the STS enzyme may be important

in the development of a new treatment strategy for hormone-dependent cancers, for example, breast cancer, which is the most common cancer and the main cause of cancer mortality in women worldwide (Ferlay et al., 2010). One of the strategies for developing new STS inhibitors is the synthesis of sulfamated analogs based on steroidal cores. For example, the steroidal derivative EMATE **1** (Figure 1) demonstrated strong inhibitory activity with an IC_{50} value of 65 μ M in the MCF-7 cell line (Purohit, Reed, Morris, Williams, & Potter, 1996). The unique activity of this compound has spawned hundreds of such sulfamate-based STS inhibitors based around the aryl sulfamate pharmacophore (Potter, 2018; Thomas & Potter, 2015). To avoid adverse side effects (estrogenic properties), several research groups have obtained new STS inhibitors based on nonsteroidal analogs, for example, coumarins. One of the first potent inhibitors based on the coumarin core was COUMATE **2**, which exhibited good activity with an IC_{50} value of 380 nM (when evaluated against placental microsomes) (Woo et al., 1998). Further modification of its structure led to a wide range of bicyclic or tricyclic coumarin derivatives that mimic the AB or ABC rings of the natural substrate (Ganeshapillai, Woo, Thomas, Purohit, & Potter, 2018). For example, 667-COUMATE **3** demonstrated potent activity toward STS with an IC_{50} value of 8 nM (Malini et al., 2000).

Recently, extensive research on the preparation of many coumarin derivatives as STS inhibitors has been carried out. Inhibitory activity against STS was determined in the case of phosphate and thiophosphate derivatives of tricyclic coumarin (Kozak et al., 2014; Kozak et al., 2015), fluorinated derivatives of 3-phenylcoumarin-7-O-sulfamate (Demkowicz et al., 2016) and fluorinated *N*-benzoyl and *N*-

phenylacetyl derivatives of 3-(4-aminophenyl)-coumarin-7-O-sulfamate (Daško et al., 2017). For example, compounds **4** and **5** demonstrated inhibitory activity with an IC_{50} value of 180 nM (for both compounds) in an assay with isolated STS.

Our previous research indicated that *N*-phosphorylated coumarin derivatives **6a-h** are also potent STS inhibitors (Daško et al., 2016). The most active derivatives, **6e** and **6f**, inhibited the STS enzyme with IC_{50} values of 190 nM and 240 nM, respectively. In the course of our research, we found a strong correlation between measured IC_{50} values and calculated $\log P$ parameters of tested STS inhibitors, which suggests that an increase in the hydrophobic properties of *N*-phosphoryl moieties causes an increase in the inhibitory activity. The higher inhibitory activities of the more hydrophobic derivatives may be explained by the establishment of stronger hydrophobic interactions with the lipophilic amino acid residues lining the cavity of STS. Encouraged by our previous research, we decided to design and synthesize a series of *N*-thiophosphorylated 3-(4-aminophenyl)-coumarin-7-O-sulfamates as compounds with potential STS inhibitory activity. We expect that the enhancement in the hydrophobic properties through replacement of the *N*-phosphoryl group with an *N*-thiophosphoryl moiety could lead to stronger STS inhibitors.

2 | MATERIALS AND METHODS

Thionyl chloride, 4-nitrophenylacetic acid, potassium carbonate, 2,4-dihydroxybenzaldehyde, sodium hydrosulfite, chlorosulfonyl isocyanate,




FIGURE 1 Chemical structures of STS inhibitors

N,N-dimethylacetamide, formic acid, dimethylchlorothiophosphate and diethylchlorothiophosphate were commercially available from Sigma-Aldrich. Diphenylchlorothiophosphate, di-*i*-propylchlorothiophosphate, di-*n*-butylchlorothiophosphate, dibenzylchlorothiophosphate, and 2-chloro-[1,3,2]dioxaphosphinane 2-sulfide were obtained according to the literature procedures (Jacobson & Nguyen, 1999). Pyridine, dichloromethane and acetone were dried and distilled using standard procedures. Melting points (uncorrected) were determined with a Stuart Scientific SMP30 apparatus. NMR spectra were recorded on a Bruker Avance III HD 400 MHz spectrometer. Chemical shifts are reported in ppm relative to the residual solvent peak (DMSO-*d*₆ 2.49 ppm for ¹H, 39.5 ppm for ¹³C) or to an external standard (85% H₃PO₄ = 0 for ³¹P). Coupling constants are given in Hertz. IR spectra were measured on a Nicolet 8700. Elemental analysis was performed using a CHNS-Carlo Erba EA-1108. Mass spectra were recorded on a TripleTOF 5600+ SCIEX mass spectrometry system. Column chromatography was performed using silica gel 60 (230–400 mesh, Merck). Preparative thin-layer chromatography was performed with Polygram SIL G/UV254 silica gel (Macherey-Nagel GmbH & Co. KG, Düren, Germany).

2.1 | General method for the synthesis of the *N*-thiophosphorylated derivatives of 3-(4-aminophenyl)-7-hydroxy-coumarin (8a-g)

The corresponding chlorothiophosphate (3.0 mmol) was added to a solution of 3-(4-aminophenyl)-7-hydroxy-coumarin **7** (380 mg, 1.5 mmol) in dry pyridine (4.5 mL). The reaction mixture was stirred under a nitrogen atmosphere for 24 hr. Next, the solvent was evaporated, and the resulting residue was purified by column chromatography using CH₂Cl₂:MeOH 100:1 as an eluent to give the desired products **8a-g**.

2.1.1 | 3-[4-(dimethoxy-thiophosphorylamino)-phenyl]-7-hydroxy-coumarin **8a**

Yield 53%, mp 165–169 °C; ν_{\max} (KBr)/cm^{−1} 3,290, 1,675, 1,609, 1,517, 1,453, 1,282, 1,226, 1,132, 1,014, 938, 634; ¹H NMR δ_{H} (400 MHz, DMSO) 10.56 (1H, s, OH), 8.59 (1H, d, *J* 15.2 Hz, NH), 8.07 (1H, s, CH), 7.62–7.54 (3H, m, Ar–H), 7.13 (2H, d, *J* 8.7 Hz, Ar–H), 6.82 (1H, dd, *J* 8.5, 2.3 Hz, Ar–H), 6.75 (1H, d, *J* 2.2 Hz, Ar–H), 3.67 (6H, d, *J* 14.0 Hz, CH₃); ¹³C NMR δ_{C} (101 MHz, DMSO) 161.4, 160.6, 155.1, 140.7, 140.3, 130.2, 129.5, 128.5, 122.4, 117.9 (d, *J*_{P-C} 7.5 Hz), 113.8, 112.6, 102.2, 53.5 (d, *J*_{P-C} 4.7 Hz); ³¹P NMR δ_{C} (162 MHz, DMSO) 69.04. Anal. calcd for: C₁₇H₁₆NO₅PS: C, 54.11; H 4.27; N, 3.71; S, 8.50. Found: C, 54.02; H, 4.33; N, 3.80; S, 8.39. HRMS (*m/z*) [M–H][−] calcd 376.0409, found 376.0616.

2.1.2 | 3-[4-(diethoxy-thiophosphorylamino)-phenyl]-7-hydroxy-coumarin **8b**

Yield 54%, mp 156–159 °C; ν_{\max} (KBr)/cm^{−1} 3,268, 1,676, 1,595, 1,514, 1,454, 1,274, 1,230, 1,118, 1,021, 927, 634; ¹H NMR δ_{H} (400 MHz, DMSO) 10.56 (1H, s, OH), 8.51 (1H, d, *J* 15.3 Hz, NH),

8.08 (1H, s, CH), 7.61–7.54 (3H, m, Ar–H), 7.14 (2H, d, *J* 8.8 Hz, Ar–H), 6.82 (1H, dd, *J* 8.5, 2.3 Hz, Ar–H), 6.75 (1H, d, *J* 2.2 Hz, Ar–H), 4.15–3.95 (4H, m, CH₂), 1.25 (6H, t, *J* 7.1 Hz, CH₃); ¹³C NMR δ_{C} (101 MHz, DMSO) 161.4, 160.6, 155.1, 141.0, 140.2, 130.2, 129.4, 128.3, 122.4, 117.8 (d, *J*_{P-C} 7.7 Hz), 113.8, 112.6, 102.1, 63.0 (d, *J*_{P-C} 4.6 Hz), 16.1 (d, *J*_{P-C} 5.0 Hz); ³¹P NMR δ_{C} (162 MHz, DMSO) 64.50. Anal. calcd for: C₁₉H₂₀NO₅PS: C, 56.29; H 4.97; N, 3.45; S, 7.91. Found: C, 56.21; H, 4.92; N, 3.55; S, 7.82. HRMS (*m/z*) [M–H][−] calcd 404.0722, found 404.1158.

2.1.3 | 3-[4-(diisopropoxy-thiophosphorylamino)-phenyl]-7-hydroxy-coumarin **8c**

Yield 74%, mp 145–148 °C; ν_{\max} (KBr)/cm^{−1} 3,269, 1,680, 1,606, 1,516, 1,449, 1,281, 1,213, 1,124, 970, 933, 633; ¹H NMR δ_{H} (400 MHz, DMSO) 10.56 (1H, s, OH), 8.43 (1H, d, *J* 16.1 Hz, NH), 8.08 (1H, s, CH), 7.62–7.55 (3H, m, Ar–H), 7.14 (2H, d, *J* 8.8 Hz, Ar–H), 6.81 (1H, dd, *J* 8.5, 2.2 Hz, Ar–H), 6.75 (1H, d, *J* 2.1 Hz, Ar–H), 4.73–4.60 (2H, m, CH), 1.32 (6H, d, *J* 6.2 Hz, CH₃), 1.18 (6H, d, *J* 6.2 Hz, CH₃); ¹³C NMR δ_{C} (101 MHz, DMSO) 161.4, 160.6, 155.1, 141.3, 140.1, 130.2, 129.1, 128.0, 122.3, 117.8 (d, *J*_{P-C} 7.7 Hz), 113.8, 112.6, 102.1, 72.1 (d, *J*_{P-C} 4.7 Hz), 23.8 (d, *J*_{P-C} 5.0 Hz), 23.5 (d, *J*_{P-C} 4.6 Hz); ³¹P NMR δ_{C} (162 MHz, DMSO) 61.31. Anal. calcd for: C₂₁H₂₄NO₅PS: C, 58.19; H 5.58; N, 3.23; S, 7.40. Found: C, 58.11; H, 5.51; N, 3.32; S, 7.48. HRMS (*m/z*) [M–H][−] calcd 432.1035, found 432.1207.

2.1.4 | 3-[4-(di-*n*-butoxy-thiophosphorylamino)-phenyl]-7-hydroxy-coumarin **8d**

Yield 46%, mp 135–136 °C; ν_{\max} (KBr)/cm^{−1} 3,277, 1,680, 1,595, 1,513, 1,452, 1,270, 1,230, 1,123, 1,019, 935, 631; ¹H NMR δ_{H} (400 MHz, DMSO) 10.56 (1H, s, OH), 8.51 (1H, d, *J* 15.5 Hz, NH), 8.08 (1H, s, CH), 7.61–7.54 (3H, m, Ar–H), 7.13 (2H, d, *J* 8.7 Hz, Ar–H), 6.82 (1H, dd, *J* 8.5, 2.2 Hz, Ar–H), 6.75 (1H, d, *J* 2.1 Hz, Ar–H), 4.11–3.85 (4H, m, CH₂), 1.59 (4H, quint, *J* 6.5 Hz, CH₂), 1.34 (4H, sext, *J* 6.7 Hz, CH₂), 0.86 (6H, t, *J* 7.4 Hz, CH₃); ¹³C NMR δ_{C} (101 MHz, DMSO) 161.4, 160.6, 155.1, 141.0, 140.1, 130.2, 129.3, 128.3, 122.3, 117.7 (d, *J*_{P-C} 7.6 Hz), 113.8, 112.6, 102.1, 66.5 (d, *J*_{P-C} 4.9 Hz), 31.9 (d, *J*_{P-C} 8.0 Hz), 18.8, 13.9; ³¹P NMR δ_{C} (162 MHz, DMSO) 65.09. Anal. calcd for: C₂₃H₂₈NO₅PS: C, 59.86; H 6.12; N, 3.03; S, 6.95. Found: C, 59.94; H, 6.03; N, 2.98; S, 7.09. HRMS (*m/z*) [M–H][−] calcd 460.1348, found 460.1775.

2.1.5 | 3-[4-(diphenoxy-thiophosphorylamino)-phenyl]-7-hydroxy-coumarin **8e**

Yield 70%, mp 215–216 °C; ν_{\max} (KBr)/cm^{−1} 3,219, 1,685, 1,592, 1,513, 1,452, 1,277, 1,209, 1,122, 1,023, 934, 627; ¹H NMR δ_{H} (400 MHz, DMSO) 10.59 (1H, s, OH), 9.33 (1H, d, *J* 17.5 Hz, NH), 8.13 (1H, s, CH), 7.70 (2H, d, *J* 8.7 Hz, Ar–H), 7.58 (1H, d, *J* 8.6 Hz, Ar–H), 7.47–7.35 (6H, m, Ar–H), 7.29–7.20 (6H, m, Ar–H), 6.82 (1H, dd, *J* 8.5, 2.3 Hz, Ar–H), 6.76 (1H, d, *J* 2.2 Hz, Ar–H); ¹³C NMR δ_{C}

(101 MHz, DMSO) 161.5, 160.6, 155.2, 150.5 (d, J_{P-C} 6.4 Hz), 140.6, 140.3 (d, J_{P-C} 3.9 Hz), 130.4, 130.3, 129.8, 129.3, 126.0, 122.2, 121.4 (d, J_{P-C} 4.8 Hz), 118.5 (d, J_{P-C} 7.8 Hz), 113.8, 112.6, 102.2; ^{31}P NMR δ_C (162 MHz, DMSO) 56.23. Anal. calcd for: $\text{C}_{27}\text{H}_{20}\text{NO}_5\text{PS}$: C, 64.67; H 4.02; N, 2.79; S, 6.39. Found: C, 64.55; H, 3.96; N, 2.70; S, 6.52. HRMS (m/z) $[\text{M}-\text{H}]^-$ calcd 500.0722, found 500.1005.

2.1.6 | 3-[4-(dibenzyloxy-thiophosphorylamino)-phenyl]-7-hydroxy-coumarin 8f

Yield 10%, mp oil; ν_{\max} (KBr)/ cm^{-1} 3,282, 1,676, 1,609, 1,516, 1,453, 1,291, 1,211, 1,130, 992, 951, 629; ^1H NMR δ_H (400 MHz, DMSO) 10.58 (1H, s, OH), 8.78 (1H, d, J 15.1 Hz, NH), 8.07 (1H, s, CH), 7.60–7.54 (3H, m, Ar–H), 7.42–7.28 (10H, m, Ar–H), 7.17 (2H, d, J 8.7 Hz, Ar–H), 6.82 (1H, dd, J 8.5, 2.2 Hz, Ar–H), 6.75 (1H, d, J 2.2 Hz, Ar–H), 5.19–4.99 (4H, m, CH_2); ^{13}C NMR δ_C (101 MHz, DMSO) 161.4, 160.6, 155.1, 140.2, 136.4 (d, J_{P-C} 9.0 Hz), 130.2, 129.4, 129.0, 128.9, 128.7, 128.3, 126.9, 122.3, 118.1 (d, J_{P-C} 7.6 Hz), 113.8, 112.6, 102.1, 68.4 (d, J_{P-C} 4.2 Hz); ^{31}P NMR δ_C (162 MHz, DMSO) 65.65. Anal. calcd for: $\text{C}_{29}\text{H}_{24}\text{NO}_5\text{PS}$: C, 65.78; H 4.57; N, 2.65; S, 6.06. Found: C, 65.91; H, 4.49; N, 2.72; S, 6.00. HRMS (m/z) $[\text{M}-\text{H}]^-$ calcd 528.1035, found 528.1502.

2.1.7 | 3-[4-(2-thioxo-2 λ^5 -[1,3,2]dioxaphosphinan-2-ylamino)-phenyl]-7-hydroxy-coumarin 8g

Yield 35%, mp >300 °C (with decomposition); ν_{\max} (KBr)/ cm^{-1} 3,255, 1,705, 1,606, 1,515, 1,451, 1,284, 1,234, 1,133, 1,020, 921, 636; ^1H NMR δ_H (400 MHz, DMSO) 10.57 (1H, s, OH), 8.70 (1H, d, J 13.4 Hz, NH), 8.08 (1H, s, CH), 7.64–7.54 (3H, m, Ar–H), 7.18 (2H, d, J 8.7 Hz, Ar–H), 6.82 (1H, dd, J 8.5, 2.3 Hz, Ar–H), 6.75 (1H, d, J 2.2 Hz, Ar–H), 4.55–4.33 (4H, m, CH_2), 2.18–1.88 (2H, m, CH_2); ^{13}C NMR δ_C (101 MHz, DMSO) 161.4, 160.6, 155.1, 140.6, 140.3, 130.2, 129.4, 128.7, 122.4, 118.4 (d, J_{P-C} 7.5 Hz), 113.8, 112.6, 102.2, 68.0 (d, J_{P-C} 6.6 Hz), 26.3 (d, J_{P-C} 6.6 Hz); ^{31}P NMR δ_C (162 MHz, DMSO) 61.26. Anal. calcd for: $\text{C}_{18}\text{H}_{16}\text{NO}_5\text{PS}$: C, 55.52; H 4.14; N, 3.60; S, 8.24. Found: C, 55.39; H, 4.21; N, 3.71; S, 8.32. HRMS (m/z) $[\text{M}-\text{H}]^-$ calcd 388.0409, found 388.0772.

2.2 | General method for the synthesis of the *N*-thiophosphorylated derivatives of 3-(4-aminophenyl)-coumarin-7-O-sulfamate (9a-g)

A mixture of formic acid (96 mg, 1.54 mmol) and *N,N*-dimethylacetamide (1.4 mg, 0.016 mmol) was added to a stirred solution of chlorosulfonyl isocyanate (212 mg, 1.50 mmol) in dry dichloromethane (0.5 mL) at 40 °C over a period of 3.5 hr. Then, a stirred solution of the appropriate *N*-thiophosphorylated derivative of 7-hydroxy-3-phenylcoumarin **8a-g** (253 mg, 1.00 mmol) in *N,N*-dimethylacetamide (3.4 mL) was added to the mixture. The resulting mixture was stirred at ambient temperature overnight and then poured into water (10 mL). Eventually, a white precipitate formed. The suspension stirred at ambient temperature for another 2 hr. The resulting precipitate was filtered, washed with water and recrystallized from acetonitrile or CH_2Cl_2 to give the desired product **9a-g**.

2.2.1 | 3-[4-(dimethoxy-thiophosphorylamino)-phenyl]-coumarin-7-O-sulfamate 9a

Yield 32%, mp 184–186 °C; ν_{\max} (KBr)/ cm^{-1} 3,284, 1,687, 1,614, 1,516, 1,456, 1,392, 1,194, 938, 774, 655; ^1H NMR δ_H (400 MHz, DMSO) 8.65 (1H, d, J 15.1 Hz, NH), 8.25 (2H, s, NH_2), 8.22 (1H, s, CH), 7.84 (1H, d, J 8.6 Hz, Ar–H), 7.64 (2H, d, J 8.6 Hz, Ar–H), 7.36 (1H, d, J 2.2 Hz, Ar–H), 7.30 (1H, dd, J 8.5, 2.3 Hz, Ar–H), 7.16 (2H, d, J 8.7 Hz, Ar–H), 3.68 (6H, d, J 14.0 Hz, CH_3); ^{13}C NMR δ_C (101 MHz, DMSO) 160.0, 153.6, 152.3, 141.4, 139.0, 130.0, 129.8, 127.8, 126.5, 119.1, 118.5, 117.9 (d, J_{P-C} 7.5 Hz), 53.6 (d, J_{P-C} 4.7 Hz); ^{31}P NMR δ_C (162 MHz, DMSO) 68.95. Anal. calcd for: $\text{C}_{17}\text{H}_{17}\text{N}_2\text{O}_7\text{PS}_2$: C, 44.73; H 3.75; N, 6.14; S, 14.05. Found: C, 44.81; H, 3.79; N, 6.21; S, 13.96. HRMS (m/z) $[\text{M}-\text{H}]^-$ calcd 455.0137, found 455.0344.

2.2.2 | 3-[4-(diethoxy-thiophosphorylamino)-phenyl]-coumarin-7-O-sulfamate 9b

Yield 58%, mp 196–198 °C (with decomposition); ν_{\max} (KBr)/ cm^{-1} 3,280, 1,685, 1,613, 1,516, 1,461, 1,389, 1,195, 929, 772, 649; ^1H NMR δ_H (400 MHz, DMSO) 8.58 (1H, d, J 15.2 Hz, NH), 8.24 (2H, s, NH_2), 8.23 (1H, s, CH), 7.84 (1H, d, J 8.6 Hz, Ar–H), 7.64 (2H, d, J 8.6 Hz, Ar–H), 7.36 (1H, d, J 2.0 Hz, Ar–H), 7.30 (1H, dd, J 8.5, 2.2 Hz, Ar–H), 7.17 (2H, d, J 8.7 Hz, Ar–H), 4.16–3.96 (4H, m, CH_2), 1.25 (6H, t, J 7.0 Hz, CH_3); ^{13}C NMR δ_C (101 MHz, DMSO) 160.0, 153.6, 152.2, 141.6, 138.9, 130.0, 129.7, 127.5, 126.5, 119.1, 118.5, 117.8, (d, J_{P-C} 7.6 Hz), 109.9, 63.1 (d, J_{P-C} 4.7 Hz), 16.1 (d, J_{P-C} 7.7 Hz); ^{31}P NMR δ_C (162 MHz, DMSO) 64.40. Anal. calcd for: $\text{C}_{19}\text{H}_{21}\text{N}_2\text{O}_7\text{PS}_2$: C, 47.10; H 4.37; N, 5.78; S, 13.24. Found: C, 47.15; H, 4.31; N, 5.90; S, 13.17. HRMS (m/z) $[\text{M}-\text{H}]^-$ calcd 483.0450, found 483.0821.

2.2.3 | 3-[4-(diisopropoxy-thiophosphorylamino)-phenyl]-coumarin-7-O-sulfamate 9c

Yield 32%, mp 190–191 °C; ν_{\max} (KBr)/ cm^{-1} 3,285, 1,685, 1,614, 1,516, 1,459, 1,386, 1,195, 933, 768, 643; ^1H NMR δ_H (400 MHz, DMSO) 8.50 (1H, d, J 16.1 Hz, NH), 8.26–8.22 (3H, m, CH, NH_2), 7.84 (1H, d, J 8.6 Hz, Ar–H), 7.64 (2H, d, J 8.7 Hz, Ar–H), 7.35 (1H, d, J 2.2 Hz, Ar–H), 7.29 (1H, dd, J 8.5, 2.3 Hz, Ar–H), 7.17 (2H, d, J 8.8 Hz, Ar–H), 4.74–4.61 (2H, m, CH), 1.32 (6H, d, J 6.2 Hz, CH_3), 1.18 (6H, d, J 6.2 Hz, CH_3); ^{13}C NMR δ_C (101 MHz, DMSO) 160.0, 153.6, 152.2, 142.0, 138.8, 130.0, 129.4, 127.2, 126.4, 119.1, 118.5, 117.9 (d, J_{P-C} 7.6 Hz), 109.9, 72.1 (d, J_{P-C} 4.8 Hz), 23.8 (d, J_{P-C} 5.0 Hz), 23.5 (d, J_{P-C} 4.7 Hz); ^{31}P NMR δ_C (162 MHz, DMSO) 61.20. Anal. calcd for: $\text{C}_{21}\text{H}_{25}\text{N}_2\text{O}_7\text{PS}_2$: C, 49.21; H 4.92; N, 5.47; S, 12.51. Found: C, 49.33; H, 4.87; N, 5.41; S, 12.44. HRMS (m/z) $[\text{M}+\text{H}]^+$ calcd 513.0919, found 513.3477.

2.2.4 | 3-[4-(di-*n*-butoxy-thiophosphorylamino)-phenyl]-coumarin-7-O-sulfamate 9d

Yield 48%, mp 175–176 °C; ν_{\max} (KBr)/ cm^{-1} 3,285, 1,693, 1,614, 1,515, 1,454, 1,375, 1,193, 936, 769, 653; ^1H NMR δ_H (400 MHz, DMSO) 8.58 (1H, d, J 15.5 Hz, NH), 8.24 (2H, s, NH_2), 8.23 (1H, s, CH),

7.85 (1H, d, J 8.6 Hz, Ar—H), 7.64 (2H, d, J 8.6 Hz, Ar—H), 7.36 (1H, d, J 2.1 Hz, Ar—H), 7.29 (1H, dd, J 8.5, 2.2 Hz, Ar—H), 7.16 (2H, d, J 8.7 Hz, Ar—H), 4.12–3.89 (4H, m, CH₂), 1.60 (4H, quint, J 7.8 Hz, CH₂), 1.34 (4H, sext, J 7.4 Hz, CH₂), 0.86 (6H, t, J 7.4 Hz, CH₃); ¹³C NMR δ_C (101 MHz, DMSO) 160.0, 153.6, 152.2, 141.6, 138.9, 130.0, 129.6, 127.5, 126.4, 119.1, 118.5, 117.7 (d, J_{P-C} 7.5 Hz), 109.9, 66.6 (d, J_{P-C} 5.0 Hz), 31.9 (d, J_{P-C} 7.9 Hz), 18.8, 13.9; ³¹P NMR δ_C (162 MHz, DMSO) 65.00. Anal. calcd for: C₂₃H₂₉N₂O₇PS₂: C, 51.10; H 5.41; N, 5.18; S, 11.86. Found: C, 51.02; H, 5.49; N, 5.29; S, 11.73. HRMS (m/z) [M+H]⁺ calcd 541.1232, found 541.3912.

2.2.5 | 3-[4-(diphenoxy-thiophosphorylamino)-phenyl]-coumarin-7-O-sulfamate 9e

Yield 43%, mp 208–210 °C (with decomposition); ν_{\max} (KBr)/cm⁻¹ 3,236, 1,691, 1,616, 1,517, 1,487, 1,392, 1,181, 916, 771, 672; ¹H NMR δ_H (400 MHz, DMSO) 9.40 (1H, d, J 17.4 Hz, NH), 8.28 (1H, s, CH), 8.25 (2H, s, NH₂), 7.85 (1H, d, J 8.6 Hz, Ar—H), 7.75 (2H, d, J 8.7 Hz, Ar—H), 7.47–7.38 (6H, m, Ar—H), 7.37 (1H, d, J 2.2 Hz, Ar—H), 7.32–7.20 (7H, m, Ar—H); ¹³C NMR δ_C (101 MHz, DMSO) 160.0, 153.7, 152.3, 150.5 (d, J_{P-C} 6.5 Hz), 140.9 (d, J_{P-C} 3.6 Hz), 139.3, 130.4, 130.1, 130.0, 128.5, 126.3, 126.0, 121.4 (d, J_{P-C} 4.8 Hz), 119.2, 118.5 (d, J_{P-C} 7.8 Hz), 118.4, 109.9; ³¹P NMR δ_C (162 MHz, DMSO) 56.15. Anal. calcd for: C₂₇H₂₁N₂O₇PS₂: C, 55.86; H 3.65; N, 4.83; S, 11.05. Found: C, 55.71; H, 3.71; N, 4.91; S, 10.94. HRMS (m/z) [M+H]⁺ calcd 581.0606, found 581.4347.

2.2.6 | 3-[4-(dibenzoyloxy-thiophosphorylamino)-phenyl]-coumarin-7-O-sulfamate 9f

Yield 38%, mp 148–149 °C; ν_{\max} (KBr)/cm⁻¹ 3,246, 1,693, 1,612, 1,513, 1,456, 1,372, 1,191, 933, 772, 656; ¹H NMR δ_H (400 MHz, DMSO) 8.85 (1H, d, J 15.0 Hz, NH), 8.24 (2H, s, NH₂), 8.22 (1H, s, CH), 7.84 (1H, d, J 8.6 Hz, Ar—H), 7.63 (2H, d, J 8.7 Hz, Ar—H), 7.39–7.32 (11H, m, Ar—H), 7.29 (1H, dd, J 8.5, 2.3 Hz, Ar—H), 7.20 (2H, d, J 8.8 Hz, Ar—H), 5.17–4.99 (4H, m, CH₂); ¹³C NMR δ_C (101 MHz, DMSO) 159.9, 153.6, 152.2, 141.4, 139.0, 136.4 (d, J_{P-C} 9.0 Hz), 130.1, 129.7, 128.9, 128.7, 128.3, 127.8, 126.4, 119.1, 118.5, 118.1 (d, J_{P-C} 7.7 Hz), 109.9, 68.42 (d, J_{P-C} 4.3 Hz); ³¹P NMR δ_C (162 MHz, DMSO) 65.55. Anal. calcd for: C₂₉H₂₅N₂O₇PS₂: C, 57.23; H 4.14; N, 4.60; S, 10.54. Found: C, 57.13; H, 4.21; N, 4.49; S, 10.61. HRMS (m/z) [M-H]⁻ calcd 607.0763, found 607.1429.

2.2.7 | 3-[4-(2-thioxo-2λ⁵-[1,3,2]dioxaphosphinan-2-ylamino)-phenyl]-coumarin-7-O-sulfamate 9g

Yield 40%, mp 267–270 °C (with decomposition); ν_{\max} (KBr)/cm⁻¹ 3,239, 1,685, 1,616, 1,517, 1,456, 1,391, 1,189, 927, 763, 651; ¹H NMR δ_H (400 MHz, DMSO) 8.76 (1H, d, J 13.3 Hz, NH), 8.25 (2H, s, NH₂), 8.23 (1H, s, CH), 7.85 (1H, d, J 8.6 Hz, Ar—H), 7.66 (2H, d, J 8.6 Hz, Ar—H), 7.36 (1H, d, J 2.1 Hz, Ar—H), 7.30 (1H, dd, J 8.5, 2.2 Hz, Ar—H), 7.22 (2H, d, J 8.7 Hz, Ar—H), 4.56–4.34 (4H, m, CH₂), 2.20–1.91 (2H, m, CH₂); ¹³C NMR δ_C (101 MHz, DMSO) 160.0, 153.6,

152.3, 141.3, 139.0, 130.1, 129.7, 128.0, 126.5, 119.1, 118.5, 118.4 (d, J_{P-C} 7.6 Hz), 109.9, 68.0 (d, J_{P-C} 6.7 Hz), 26.3 (d, J_{P-C} 6.9 Hz); ³¹P NMR δ_C (162 MHz, DMSO) 61.14. Anal. calcd for: C₁₈H₁₇N₂O₇PS₂: C, 46.15; H 3.66; N, 5.98; S, 13.69. Found: C, 46.24; H, 3.73; N, 6.05; S, 13.55. HRMS (m/z) [M+H]⁺ calcd 469.0293, found 469.2661.

2.3 | Computational studies

2.3.1 | Ligand preparation

Prior to the docking procedures, the potential inhibitors were prepared using Portable HyperChem Release 8.0.7 (Hypercube, Inc., Gainesville, FL). Each ligand was optimized using an MM+ force field and the Polak–Ribière conjugate gradient algorithm (terminating at a gradient of 0.05 kcal mol⁻¹ Å⁻¹).

2.3.2 | Protein preparation

For docking, the X-ray structure of human STS (obtained from Protein Data Bank—accession code 1P49) was prepared using the standard procedure. First, the water molecules from crystallization were removed from the structure, and the catalytic amino acid fGly75 was converted to the gem-diol form using the Protein Preparation Wizard module, delivered by Maestro (Schrödinger, LLC, New York, NY). Next, the hydrogen atoms were added onto the structure, and the prepared model of the enzyme was optimized using the OPLS-AA force field.

2.3.3 | Molecular docking

The docking of the optimized ligands to the prepared structure of human STS was carried out with AutoDock Vina 1.1.2 software (The Molecular Graphic Laboratory, The Scripps Research Institute, La Jolla, CA) (Trott & Olson, 2010). For all docking studies, a grid box of size 30 Å × 30 Å × 30 Å centered on the C β atom of amino acid 75 was used. The best poses for a particular ligand were inspected visually. Illustrations of the 3D models were generated using VMD 1.9 (University of Illinois at Urbana–Champaign, Urbana, IL).

2.3.4 | LogP calculations

LogP parameters were calculated using ChemDraw Ultra 7.0 (CambridgeSoft Corporation, Cambridge, MA).

2.4 | Biological assays

2.4.1 | Enzyme purification

STS was extracted from a human placenta and purified to homogeneity following a multistep chromatography protocol as previously described (Hernandez-Guzman, Higashiyama, Osawa, & Ghosh, 2001).

2.4.2 | In vitro activity in an enzymatic assay

The reaction mixture, at a final volume of 100 μ L, contained 20 mM Tris-HCl pH 7.4, 3 mM *p*-nitrophenyl sulfate (NPS), varied concentrations of an inhibitor (0.01–100 μ M) and 5 U of a purified enzyme (1 U is the amount of an enzyme that hydrolyzes 100 μ M NPS in 1 hr at 37 °C). The reaction was performed at 37 °C for 15 min and halted by the addition of 100 mL of 1 M NaOH. The absorbance of the released *p*-nitrophenol was measured at 405 nm using a Biotek ELx800 Microplate Reader (BioTek Instruments, Inc. Winooski, VT). The IC_{50} values were calculated using GraphPad Prism software (GraphPad Software, Inc., La Jolla, CA). All measurements were performed in triplicate.

2.4.3 | In vitro activity in a cell line experiment

The inhibition of sulfatase activity in MCF-7 breast cancer cells was measured using a method described earlier with some modifications. MCF-7 cells were maintained in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum and cultured in the above medium until 80% confluence. For experiments, cells were seeded in 24-well microplates (Nest Biotechnology) at a density of 1×10^5 cells/well. Cell number was determined using a *Bürker Counting Chamber*. Incubation of cells was conducted for 20 hr at 37 °C in a 5% CO₂ humidified incubator in serum-free medium (0.5 mL) with the addition of [³H]E1S (4×10^5 cpm, 3 nM) with or without inhibitor (100 nM). After incubation, medium (0.45 mL) was collected from each well, and the product formed by STS was isolated from the mixture by extraction with toluene (4 mL). Sulfatase activity was measured using the Radioluminometer MicroBeta (Perkin Elmer). Assays involving MCF-7 cells were carried out in triplicate.

2.4.4 | Influence of the STS inhibitors on zebrafish embryo development

The collected zebrafish embryos were transferred to a Petri dish with E3 medium (5 mM NaCl, 0.33 mM MgCl₂, 0.33 mM CaCl₂, 0.17 mM KCl; pH 7.2) and then placed in 6-well plates with 10 embryos in each well. Stock solutions of **6e**, **6f**, **9b**, and **9e** were prepared in DMSO. In the present experiments, 0.1, 0.5, and 1 μ M solutions of the compounds were employed, which were freshly prepared each time by dissolving the stock solutions in E3 medium directly before their addition to the wells. The solutions were changed once daily, and the embryos were maintained in an incubator at 28.5 °C. Zebrafish embryo development and viability were evaluated for up to 5 days using bright-field microscopy (Zeiss Axio Vert, ZEISS, Germany). Image analysis to determine the percent of dead and malformed embryos over time was performed using Zeiss software. The same embryos ($n = 10$) were followed throughout the whole study.

3 | RESULTS AND DISCUSSION

3.1 | Chemistry

In the course of our research, we synthesized all of the newly designed STS inhibitors, which are *N*-thiophosphorylated derivatives of


3-(4-aminophenyl)-coumarin-7-*O*-sulfamate (**9a-g**), using the pathway shown in Scheme 1. In the first step, we synthesized 7-hydroxy-3-(4-aminophenyl)-coumarin **7** according to the procedure that we previously described (Daško et al., 2017). Next, coumarin derivative **7** was *N*-thiophosphorylated with the corresponding chlorothiophosphate in anhydrous pyridine. The isolation of the crude products of the thiophosphorylation reaction was performed by flash column chromatography to give coumarin derivatives **8a-g**. Finally, the OH groups of compounds **8a-g** were sulfamoylated. In this cases, the solutions of stable *N*-thiophosphorylated derivatives **8a-g** in *N,N*-dimethylacetamide (DMA) were treated with H₂NSO₂Cl that was previously generated in the reaction of chlorosulfonyl isocyanate and formic acid in the presence of a catalytic amount of DMA. After the standard isolation procedure, we obtained the desired compounds **9a-g**. Furthermore, to perform the additional biological evaluation, we decided to resynthesize *N*-phosphorylated 3-(4-aminophenyl)-coumarin-7-*O*-sulfamates **6a-h** according to the procedure that we previously described (Daško et al., 2016).

3.2 | Molecular modeling

To determine the binding modes of the designed *N*-thiophosphorylated 3-(4-aminophenyl)-coumarin-7-*O*-sulfamates **9a-g**, a computational docking method was used. The docking studies also involved in identifying potential interactions between specified functional groups on the inhibitors and amino acid residues present in the active site of STS. The X-ray structure of STS was retrieved from the Protein Data Bank (Protein Data Bank accession code 1P49) and properly prepared for docking calculations. The docking procedure of the optimized ligands was performed using AutoDock Vina 1.1.2 software (Molecular Graphics Laboratory, The Scripps Research Institute, La Jolla, CA).


The calculated free binding energies of the newly designed candidates **9a-g** were in the range of -6.5 to -9.0 kcal mol⁻¹ and were significantly lower than the free binding energy value of the reference inhibitor **667-COUMATE** (-5.4 kcal mol⁻¹). The best docking result was determined for coumarin derivative **9e** (with a calculated free binding energy value of -9.0 kcal mol⁻¹). The free binding energies for resynthesized compounds **6e** and **6f** were -6.7 and -8.3 kcal mol⁻¹, respectively.

Our docking analysis for the newly designed STS inhibitor candidates **9a-g** showed that these compounds could associate well with the active site of STS in a similar manner to the mode of the reported STS inhibitor **667-COUMATE**. Figure 2 shows the structures of **9e** and **6e** and the reference inhibitor **667-COUMATE** docked into the active site of the STS enzyme. We found that the sulfamate functional groups are located very close to the catalytic amino acid residue fGly75 (~ 3.0 Å), coordinated to the Ca²⁺ ion, and surrounded by the catalytically essential residues (Asp35, Asp36, Arg79, Lys134, His136, and Lys368). The coumarin cores, with *N*-thiophosphoryl moieties in the case of the designed candidates **9a-g**, are well accommodated in the lipophilic cavity delimited by the lipophilic amino acids in the



SCHEME 1 Synthesis of *N*-thiophosphorylated derivatives of 3-(4-aminophenyl)-coumarin-7-*O*-sulfamate **9a-g**

FIGURE 2 Docked binding modes and distances to fGly75 and Arg98 for compounds **6e** (blue), **9e** (red), and the reference inhibitor **667-COUMATE** (green)



enzyme-binding region (Arg98, Leu103, Phe104, Leu167, Val 177, Phe178, Phe182, Leu185, Phe230, Phe233, Phe237, Thr484, Val486, Phe488, Trp550, and Phe553). Furthermore, in the course of our previous research with *N*-phosphorylated compounds **6a-h**, we noticed that the *N*-phosphoryl groups of compounds **6a-h** are within hydrogen bonding distance from the backbone NH group of Arg98 (Daško et al., 2016). We suggested that this additional interaction may

improve the stability of the enzyme-inhibitor complex and may influence the inhibitory activities of compounds **6a-h**. As shown in Figure 2, we detected that the *N*-thiophosphoryl groups of the *N*-thiophosphorylated 3-(4-aminophenyl)-coumarin-7-*O*-sulfamates occupied the same region of the STS active site as the *N*-phosphoryl groups of *N*-phosphorylated 3-(4-aminophenyl)-coumarin-7-*O*-sulfamates. However, it should be noted that the sulfur atom in an *N*-

thiophosphoryl group is a much weaker H-bond acceptor than the oxygen atom in an *N*-phosphoryl group.

3.3 | Biological assays

Initially, the STS inhibitory activities of synthesized compounds **9a-g** were evaluated in an enzymatic assay according to previously reported methods (Vaccaro, Salvioli, Muscillo, & Renola, 1987; Woo et al., 2010). Screening tests were performed using the STS enzyme isolated from human placenta and purified by a 3-step chromatography procedure. After purification, the obtained fraction was used directly as the enzyme source.

The summarized results of the enzymatic assay used for newly synthesized *N*-thiophosphorylated coumarin derivatives **9a-g** and the reference compounds **6e**, **6f** and **667-COUMATE** are presented in Table 1. The received data showed that all of the newly synthesized inhibitors **9a-g** are able to effectively inhibit the actions of STS. In the course of our investigations, the highest inhibitory activity among the newly synthesized inhibitors **9a-g** was exhibited by compound **9e** with an IC_{50} value of 0.201 μ M. This result is in agreement with the data from the molecular modeling studies. However, the *N*-phosphorylated derivative **6f** demonstrated a higher STS inhibitory effect (IC_{50} value of 0.083 μ M) than all of the newly designed compounds **9a-g**. This observation indicated that the possibility of the creation of stronger hydrogen bonds between the *N*-phosphoryl group of compound **6f** and the Arg98 residue compared to the *N*-thiophosphoryl group may be significant for inhibitory properties. Furthermore, the inhibitory activity of compound **6f** was similar to that of **667-COUMATE** (IC_{50} value of 0.062 μ M).

The next step of the biological evaluation was to verify whether the obtained STS inhibitors are able to pass the cellular membrane effectively and inhibit STS inside the cell. In the course of our study, we determined the STS activity in the MCF-7 cell line after incubation in the presence of inhibitor at 100 nM concentration. For this evaluation, we included the newly synthesized compounds **9a-g** and their *N*-

phosphorylated analogs **6a-h**, for which the synthesis had been previously described (Daško et al., 2016). The summarized results are presented in Table 2.

In the course of our research, we found that the lowest STS activities were measured in the presence of *N*-phosphorylated derivatives **6e** (0.1% of STS activity) and **6f** (0.2% of STS activity). The measured STS activity in the presence of **667-COUMATE** (used as a reference) was 0.1%. The inhibitory activities of *N*-phosphorylated compounds **6e** and **6f** were higher than their *N*-thiophosphorylated analogs **9e** and **9f** (despite the initial hypothesis that more lipophilic derivatives should more effectively inhibit STS activity as confirmed by molecular modeling studies). It suggests that the possibility of creating stronger hydrogen bonds between the *N*-phosphoryl groups of compounds **6e** and **6f** ($\log P$ values of 5.37 and 5.56, respectively) and the Arg98 residue in the active site of STS is more critical than increasing the hydrophobicity, as with analogs **9e** and **9f** ($\log P$ values of 6.11 and 6.30, respectively). The above observations confirm that the theoretical results of molecular modeling studies as well as predicted $\log P$ values, cannot be interpreted uncritically.

Finally, an in vivo toxicity assay of four of the most potent STS inhibitors was performed. Because zebrafish develop rapidly, this a promising model of whole-organism toxicology screening, and therefore, the morphological changes of zebrafish embryos induced by the most active STS inhibitors (**6e**, **6f**, **9b** and **9e**) were evaluated. The experiment revealed no morphological changes in the zebrafish embryos or toxic effects of the compounds used at doses of 0.1–1 μ M during 5 days of observation (Figure 3).


TABLE 1 Inhibitory activities and free binding energies calculated for the synthesized compounds **9a-g** and reference inhibitors (**6e**, **6f**, and **667-COUMATE**) in the STS enzyme assays

No.	Free binding energy (kcal Mol ⁻¹)	IC_{50} (μ M)
9a	−6.6	0.768 ± 0.04
9b	−6.5	0.399 ± 0.05
9c	−6.7	0.783 ± 0.03
9d	−6.7	0.665 ± 0.02
9e	−9.0	0.201 ± 0.02
9f	−7.9	0.802 ± 0.06
9g	−7.6	0.572 ± 0.04
6e	−6.7	0.273 ± 0.02
6f	−8.3	0.083 ± 0.007
667-COUMATE	−5.4	0.062 ± 0.006

TABLE 2 Remained STS activities in MCF-7 cell line at 100 nM concentration and $\log P$ parameters of the synthesized compounds **9a-g**, resynthesized compounds **6a-h** and reference inhibitor **667-COUMATE**

No.	STS activity (%)	$\log P$ (−)
9a	64.8	2.75
9b	1.0	3.43
9c	11.1	4.26
9d	1.0	5.16
9e	2.9	6.11
9f	2.3	6.30
9g	82.5	2.62
6a	85.9	2.01
6b	79.8	2.70
6c	56.1	3.52
6d	1.2	4.43
6e	0.1	5.37
6f	0.2	5.56
6g	55.1	1.89
6h	21.6	2.99
667-COUMATE	0.1	2.04

FIGURE 3 The influence of **6e**, **6f**, **9b**, and **9e** on zebrafish embryos on the fifth day versus the control group



4 | CONCLUSIONS

In the present work, we described our research on molecular modeling, synthesis and biological evaluation of *N*-thiophosphorylated derivatives of 3-(4-aminophenyl)-coumarin-7-*O*-sulfamate (**9a-g**) as new STS inhibitors. The data of the enzymatic assay indicated that all of the newly synthesized inhibitors **9a-g** are able to effectively inhibit the actions of STS. Among them, the highest inhibitory activity was exhibited by compound **9e** with an IC_{50} value of 0.201 μ M. However, resynthesized phosphate derivative **6f** demonstrated a higher STS inhibitory effect (an IC_{50} value of 0.083 μ M) than all of the newly designed compounds **9a-g**. The STS inhibitory properties of compound **6f** in the enzymatic assay were similar to **667-COUMATE** (an IC_{50} value of 0.062 μ M) used as a reference. In the course of the cell line experiment, we observed the highest inhibition of STS in the presence of *N*-phosphorylated derivatives **6e** (0.1% of STS activity) and **6f** (0.2% of STS activity). The measured STS activity in the presence of **667-COUMATE** was 0.1%. The results of enzymatic and cell line experiments suggest that the possibility of creating stronger hydrogen bonds between the phosphoryl groups of compounds **6e** and **6f** ($\log P$ values of 5.37 and 5.56, respectively) and the Arg98 residue in the active site of STS is more critical for inhibitory effects than increasing the hydrophobicity, as with analogs **9e** and **9f** ($\log P$ values of 6.11 and 6.30, respectively). Toxicology analysis revealed that the tested compounds are safe for zebrafish embryos at concentrations 12 times greater than the IC_{50} values.

ACKNOWLEDGMENTS

We gratefully acknowledge the National Science Centre (Poland) for financial support (PRELUDIUM grant no. 2015/19/N/NZ7/00938 and ETIUDA fellowship no. 2017/24/T/NZ7/00271). Furthermore, we gratefully acknowledge the Foundation for Polish Science (FNP) for financial support in the form of the fellowship START for one of the coauthors.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ORCID

Mateusz Daško  <https://orcid.org/0000-0002-3367-6491>

Sebastian Demkowicz  <https://orcid.org/0000-0002-4252-4297>

REFERENCES

- Daško, M., Mastyk, M., Kubiński, K., Aszyk, J., Rachon, J., & Demkowicz, S. (2016). Synthesis and steroid sulfatase inhibitory activities of *N*-phosphorylated 3-(4-aminophenyl)-coumarin-7-*O*-sulfamates. *Medicinal Chemistry Communications*, 7, 1146–1150.
- Daško, M., Przybyłowska, M., Rachon, J., Mastyk, M., Kubiński, K., Misiak, M., ... Demkowicz, S. (2017). Synthesis and biological evaluation of fluorinated *N*-benzoyl and *N*-phenylacetyl derivatives of 3-(4-aminophenyl)-coumarin-7-*O*-sulfamate as steroid sulfatase inhibitors. *European Journal of Medicinal Chemistry*, 128, 79–87.
- Demkowicz, S., Daško, M., Kozak, W., Krawczyk, K., Witt, D., Mastyk, M., ... Rachon, J. (2016). Synthesis and biological evaluation of fluorinated 3-phenylcoumarin-7-*O*-sulfamate derivatives as steroid sulfatase inhibitors. *Chemical Biology & Drug Design*, 87, 233–238.
- Ferlay, J., Shin, H. R., Bray, F., Forman, D., Mathers, C., & Parkin, D. M. (2010). Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *International Journal of Cancer*, 127, 2893–2917.
- Ganeshapillai, D., Woo, L. W. L., Thomas, M. P., Purohit, A., & Potter, B. V. L. (2018). C-3- and C-4-substituted bicyclic coumarin sulfamates as potent steroid sulfatase inhibitors. *ACS Omega*, 3, 10748–10772.
- Hernandez-Guzman, F. G., Higashiyama, T., Osawa, Y., & Ghosh, D. (2001). Purification, characterization and crystallization of human placental estrone/dehydroepiandrosterone sulfatase, a membrane-bound enzyme of the endoplasmic reticulum. *The Journal of Steroid Biochemistry and Molecular Biology*, 78, 441–450.
- Jacobson, R. M., & Nguyen, L. T. (1999). Phosphoryl hydrazine insecticides, US Pat., 6147062A.
- Kozak, W., Daško, M., Mastyk, M., Gielniewski, B., Rachon, J., & Demkowicz, S. (2015). Synthesis and biological evaluation of thiophosphate tricyclic coumarin derivatives as steroid sulfatase inhibitors. *Journal of Asian Natural Products Research*, 17, 1091–1096.

- Kozak, W., Daško, M., Maslyk, M., Pieczykolan, J. S., Gielniewski, B., Rachon, J., & Demkowicz, S. (2014). Phosphate tricyclic coumarin analogs as steroid sulfatase inhibitors: Synthesis and biological activity. *RSC Advances*, 4, 44350–44358.
- Malini, B., Purohit, A., Ganeshapillai, D., Woo, L. W. L., Potter, B. V. L., & Reed, M. J. (2000). Inhibition of steroid sulphatase activity by tricyclic coumarin sulphamates. *The Journal of Steroid Biochemistry and Molecular Biology*, 75, 253–258.
- Potter, B. V. L. (2018). Steroid sulphatase inhibition via aryl sulphamates: Clinical progress, mechanism and future prospects. *Journal of Molecular Endocrinology*, 61, 233–252.
- Purohit, A., Reed, M. J., Morris, N. C., Williams, G. J., & Potter, B. V. L. (1996). Regulation and inhibition of steroid sulfatase activity in breast cancer. *Annals of the New York Academy of Sciences*, 784, 40–49.
- Shah, R., Singh, J., Singh, D., Singh Jaggi, A., & Singh, N. (2016). Sulfatase inhibitors for recidivist breast cancers treatment: A chemical review. *European Journal of Medicinal Chemistry*, 114, 170–190.
- Thomas, M. P., & Potter, B. V. L. (2015). Discovery and development of the aryl O-sulfamate pharmacophore for oncology and Women's health. *Journal of Medicinal Chemistry*, 58, 7634–7658.
- Trott, O., & Olson, A. J. (2010). AutoDock Vina: Improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *Journal of Computational Chemistry*, 31, 455–461.
- Vaccaro, A. M., Salvioli, R., Muscillo, M., & Renola, L. (1987). Purification and properties of arylsulfatase C from human placenta. *Enzyme*, 37, 115–126.
- Woo, L. W. L., Howarth, N. M., Purohit, A., Hatem, A. M., Reed, M. J., & Potter, B. V. L. (1998). Steroidal and nonsteroidal sulfamates as potent inhibitors of steroid sulfatase, 41, 1068–1083.
- Woo, L. W. L., Jackson, T., Putey, A., Cozier, G., Leonard, P., Acharya, K. R., ... Potter, B. V. L. (2010). Highly potent first examples of dual aromatase-steroid sulfatase inhibitors based on a biphenyl template. *Journal of Medicinal Chemistry*, 53, 2155–2170.

How to cite this article: Daško M, Demkowicz S, Biernacki K, et al. Novel steroid sulfatase inhibitors based on N-thiophosphorylated 3-(4-aminophenyl)-coumarin-7-O-sulfamates. *Drug Dev Res.* 2019;1–10. <https://doi.org/10.1002/ddr.21569>