

# ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: P. G. Cozzi, A. Gualandi, G. Rodeghiero, E. Della Rocca, F. Bertoni, M. Marchini, R. Perciaccante, P. Jansen and P. Ceroni, *Chem. Commun.*, 2018, DOI: 10.1039/C8CC04048F.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



Chem. Commun.

## COMMUNICATION

## Application of Coumarin Dyes for Organic Photoredox Catalysis.

Andrea Gualandi,<sup>a\*</sup> Giacomo Rodeghiero,<sup>a,b</sup> Emanuele Della Rocca,<sup>a</sup> Francesco Bertoni,<sup>a</sup> Marianna Marchini,<sup>a</sup> Rossana Perciaccante,<sup>b</sup> Thomas Paul Jansen,<sup>b</sup> Paola Ceroni,<sup>a</sup> and Pier Giorgio Cozzi<sup>\*a</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Here we report the application of readily prepared and available coumarin dyes for photoredox catalysis able to mimic powerful reductant [Ir(III)] complexes. The coumarin derivatives **9** and **10** were employed as photoreductants in the pinacol coupling and in other reactions, in the presence of Et<sub>3</sub>N as sacrificial reducing agent. As electronic, photophysical, and sterical properties of coumarins could be varied, a wide applicability to several classes of photoredox reactions is predicted.

Photoredox catalysis has become a hot topic in catalysis, due to the facility to generate active radical species in controlled and mild conditions, via electron or energy transfer from photocatalysts in their excited states to organic molecules.<sup>1</sup> In this context, organic dyes are attracting a great interest as photoredox catalysts (PC).<sup>2,3,4</sup> Although complexes of abundant and inexpensive metals have recently found some applications in photoredox catalysis,<sup>5</sup> organic dyes have been used as valid alternatives to the expensive, but widely used Ru(II)- and Ir(III)-complexes.<sup>6</sup> Several classes of suitable dyes have been explored in photocatalytic transformations.<sup>2</sup> As a prototypical example, the Fukuzumi catalyst (Figure 1, **1**) has been exploited in many interesting photoredox reactions, thanks to its strong oxidant ability in its lowest singlet excited state ( $E_{1/2} = +2.06$  V vs SCE, MeCN).<sup>7</sup> However, organic molecules behaving as strong reductants are much less common. Murphy described powerful organic reductants able to promote radical coupling and other reactions of substrates with reduction potentials < -1.8 V (vs SCE).<sup>8</sup> Unfortunately, these molecules are rather reactive and difficult to generate. Other organic photocatalysts used as reductants of substrates with reduction potentials in the range -1.5 – -2.1 V (vs SCE) are: 10-phenylphenothiazine **2**,<sup>9a</sup> perylene **3**,<sup>9b</sup> N-aryl phenoxazines **4**<sup>9c</sup> and N,N-diaryl dihydrophenazines **5**.<sup>9d</sup> These molecules contain electron-rich motifs which stabilize the radical

cation formed in the photocatalytic process.<sup>10</sup>




Figure 1. Dyes employed in photocatalytic reactions.

Remarkably, although coumarins have been largely employed as fluorescent bio labels,<sup>11a</sup> laser dyes,<sup>11b</sup> emitting materials in organic light-emitting diodes (OLED)<sup>11c</sup> and dyes in solar cells,<sup>11d</sup> to the best of our knowledge, their systematic employment in photoredox reactions has not been yet explored. In addition, the low molecular weight and their straightforward synthesis<sup>12</sup> give the possibility to vary their photophysical and redox properties,<sup>13</sup> allowing to cover a wide range of photoredox potentials. With all these potentialities, the coumarin class attracted our interest for photoredox catalytic applications. Herein, we report the use of coumarins **9**<sup>14</sup> and **10** (see ESI for synthesis, details, and for evaluation of other coumarins in the model reaction) as powerful reductants in the photo-promoted radical coupling of carbonyls and imines,<sup>15,16</sup> and we illustrate the further possibility to apply coumarin dyes to many other photoredox reactions.

Quite recently, Rueping reported a photoredox mediated coupling of aldehydes, ketones, and imines mediated by Ir(III) photocatalyst, in the presence of Et<sub>3</sub>N as sacrificial reductant.<sup>16a,b</sup> Photoexcitation of the Ir(III) complex in the presence of Et<sub>3</sub>N yielded the reduced Ir complex ( $E_{1/2} = -1.69$  V vs Fc), which is responsible for the reduction of carbonyl and imine groups. The so-formed radical cation Et<sub>3</sub>N<sup>•+</sup>, a Lewis acid, is supposed to coordinate to the carbonyl oxygen and to activate its reduction. This transformation is

<sup>a</sup> Dipartimento di Chimica "G. Ciamician" ALMA MATER STUDIORUM, Università di Bologna, Via Selmi 2, 40126 Bologna, Italy

<sup>b</sup> Cyanagen Srl, Via Stradelli Guelfi 40/C, 40138, Bologna, Italy


† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: Screening tests. Photophysical measurements and quenching studies. Detailed procedures, description and copies of NMR spectra for all compounds. See DOI: 10.1039/x0xx00000x

## COMMUNICATION

## Journal Name

quite challenging for organic dye photocatalysts, due to the high negative potential necessary for the aldehyde reduction (e.g., benzaldehyde  $E_{1/2} = -2.11$  V vs Fc).<sup>17</sup> We commenced our investigation from the pinacol coupling of 4-chlorobenzaldehyde (**6a**, Scheme 1), as a model reaction, and investigating a series of coumarins readily prepared or commercially available.




**Scheme 1.** Reaction model for pinacol coupling (yields after chromatographic purification).

**Table 1.** Photophysical and electrochemical properties ( $E_{1/2}$  in V vs SCE) of coumarins **8**, **9** and **10** in DMF solution at 298 K.


	Absorption		Emission			Electrochemistry		
	$\lambda$ (nm)	$\epsilon$ (M <sup>-1</sup> cm <sup>-1</sup> )	$\lambda$ (nm)	$\Phi_{em}$	$\tau$ (ns)	$E_{00}$ (eV)	$E(A^+/A)$ (V)	$E(A^+/A^*)$ (V)
<b>8</b>	400	2.97	476	0.82	3.0	2.79	+0.92	-1.87
<b>9</b>	427	3.30	497	0.50	3.3	2.66	+0.79	-1.87
<b>10</b>	413	3.03	482	0.57	2.9	2.72	+0.83	-1.89

Among all the coumarins investigated (see ESI for more details) in the model reaction, only few coumarins were active, with coumarins **8**,<sup>18</sup> **9**, and **10** being the best performing photocatalysts. Coumarin **9** was readily obtained in 46% isolated yield from the reaction of diethylamino-salicylaldehyde and thiophene acetic acid, performing the reaction with acetic anhydride and triethyl amine. The preparation of **10** is a straightforward sulfonylation of **9** with sulfur trioxide N,N-dimethylformamide complex in DMF, that gave the desired product **10** in 33% yields after purification by reverse phase chromatography. The reaction was further optimized in the presence of coumarin **10** by varying solvents, reductive reagents, and conditions (see SI for more details), as **10** was proved to be the most efficient catalyst.<sup>19</sup> The optimal conditions for the reaction were reached by using DMF as reaction solvent ([substrate] = 0.2 M) in the presence of 4 equiv. of Et<sub>3</sub>N as sacrificial reductant. The generality of these conditions was investigated with different aldehydes: Scheme 2 reports the salient results. Other aldehydes were tested as well, but only low conversions or no reaction were observed (see ESI for further details). Only in the case of naphthyl aldehyde we have obtained excellent yields, and this could be due to the efficient formation of the ketyl radical coupled with an effective reaction. Our conditions could also be applied to ketones without further optimization (Scheme 3 A). We were pleased to find that benzophenone **11a** ( $E_{1/2} = -1.87$  V vs. SCE)<sup>16</sup> and its derivatives were suitable substrates for the reaction. Imines were also proper substrates and under the standard reaction conditions, different benzyl and aryl imines react in satisfactory yields (Scheme 3B). Not

only it is possible to use a cleavable benzyl as protecting group, but chiral benzylimines are suitable substrates and give access to chiral protected diamines.<sup>20</sup>




**Scheme 2.** Pinacol coupling of selected aldehydes (yields after chromatographic purification).



**Scheme 3.** Pinacol coupling of selected ketones and imines (yields after chromatographic purification).


To get insights into the mechanism of the pinacol reaction, a photochemical study was performed. The absorption spectrum of the photocatalyst is not significantly changed under the reaction conditions reported in Scheme 2 (ca. 1% degradation of coumarin **10** at the end of irradiation), demonstrating its significant photostability. Fluorescence of coumarin **10** is not quenched by Et<sub>3</sub>N (Figure S6, and comments), while it is quenched by aldehydes (Figure S7, and comments).<sup>21</sup> This behavior suggests a photoinduced oxidative quenching of the coumarin with direct formation of ketyl radicals. Indeed, energy transfer from the lowest excited state of coumarin to populate the lowest triplet excited state of aldehydes is ruled out, being endoergic.<sup>22</sup> Based on the reduction potentials of

aldehydes (Table S13) and of the  $S_1$  excited state of coumarins (Table 1), oxidative quenching is exoergonic, apart from 4-chlorobenzaldehyde **6a**. However, as evident from data reported in Table S13, the presence of Lewis acids can substantially decrease the reduction potentials of aldehydes.<sup>23</sup> Under the conditions reported in Scheme 2, the radical cation  $\text{Et}_3\text{N}^{+\bullet}$  can act as Lewis acid or Brønsted acid, as previously discussed by Rueping, making the investigated photoinduced electron transfer process exoergonic (see SI for more details). In the conditions reported by Rueping, the presence of 20 mol%  $\text{K}_2\text{CO}_3$  or  $\text{K}_3\text{PO}_4$  completely suppressed the pinacol coupling, due to the essential role of the Brønsted acidic  $\alpha$ -ammonium radical in the C=O activation event. In our case, by performing the model reaction in the presence of 20 mol%  $\text{K}_2\text{CO}_3$  and  $\text{K}_3\text{PO}_4$ , we observed a lower conversion, 16% and 12% respectively, compare to the model reaction performed under standard conditions. In our case, the improved reduction potential of coumarin dyes respect to Ir complex (-1.89 V vs -1.69 V) is probably still favoring the reaction that was, however, less efficient, confirming the importance of the Brønsted acidic  $\alpha$ -ammonium radical. We have also investigated cross-pinacol couplings, carrying out the reaction of **6a** in the presence of different aldehydes (**6b**, **6c**, **6e**, and *p*-MeOC<sub>6</sub>H<sub>4</sub>CHO, **6l**; see SI for full details). The cross products, obtained as inseparable mixtures with the homo pinacol products **7**, were observed for the reactions with **6b** and **6c** but not with **6e** and **6l**, probably due to the major stabilization of their ketyl radicals. Based on the photochemical investigation we suggest the mechanism depicted in Scheme 4: aldehydes are directly reduced by the photocatalyst in the excited state (I) to form the ketyl radical (III), at variance with the reductive quenching of the Ir(III) photocatalyst reported by Rueping.<sup>16</sup> The ketyl radical is then coupled to another ketyl radical to give the pinacol product. In the case described by Rueping, and in other examples reported in literature, the photocatalyst in its photoexcited state is quenched by the sacrificial reductant (normally a tertiary amine) and then is able to perform the electron transfer to the organic substrate in this reduced state. Remarkably, coumarins are able to directly transfer electrons to carbonyls, forming ketyl radicals.




Scheme 4. Proposed reaction mechanism for the pinacol coupling reaction.

Recently, we have introduced BODIPY dyes for the atom transfer radical addition (ATRA) reaction.<sup>24a</sup> Investigating the coumarins **9** and **10** as possible catalyst for the ATRA reaction,<sup>24b</sup> we found quite promising results (Scheme 5), without the need of sacrificial  $\text{Et}_3\text{N}$ , or other reducing agent. It is noteworthy the possibility to use ethylbromoacetate (**16d**) for the ATRA reaction promoted by coumarin, a substrate quite challenging for other photocatalysts.<sup>25</sup>



Scheme 5. ATRA reaction promoted by coumarin **9** (yields after chromatographic purification).

To further highlight the capability of coumarins **9** and **10** to promote a variety of different photoredox reactions, we have briefly investigated their use in: (i) the trifluoromethylation of alkenes by Umemoto reagent (Scheme 6, A),<sup>26</sup> (ii) the MacMillan stereoselective  $\alpha$ -alkylation of aldehydes (Scheme 6, B),<sup>27</sup> promoted by the synergistic cooperation of photo- and enamine catalysis.<sup>28</sup> The enantiomeric excesses obtained for the reactions were in line with the reported values and the organic photocatalyst does not influence or reduce the enantiomeric excess of the reactions. It was also possible to use the coumarin catalyst in the reductive protonation of bromoketones, in the presence of the Hantzsch ester (Scheme 6, c).<sup>29</sup> These selected examples demonstrate that the potentiality of coumarins in photoredox catalysis could be explored in many different reactions, allowing the replacement of ruthenium(II) or iridium(III) complexes.



Scheme 6. Application of coumarins dyes **8** and **9** in different chemical reactions.

In summary, we have introduced coumarins dyes as powerful photoreductants in the photoredox catalysis arena. These new photo-reductive catalysts shown broad applicability and further studies about their application in other photoredox reactions are in progress. The possibility of tailoring redox and photophysical properties of coumarin dyes by introducing different functional groups, their simple synthesis and their affordable cost, can be

## COMMUNICATION

## Journal Name

useful for extending the application of these dyes to new photocatalytic transformations.

## Acknowledgements

P. C. and M. M. acknowledge the European Commission ERC Starting Grant (PhotoSi, 278912). A. G. L. M. and P. G. C. are grateful to Fondazione Del Monte, Farb funds University of Bologna (project SLAMM to A.G.) and EU-Foundation through the TEC FP7 ICT-Molarnet project (318516) for the partial financial support of this research.

## Notes and references

- For selected recent reviews on photoredox catalysis, see: a) K. L. Skubi, T. R. Blum, and T. P. Yoon, *Chem. Rev.* 2016, **116**, 10035–10074; b) X. Lang, J. Zhao and X.-d. Chen, *Chem. Soc. Rev.* 2016, **45**, 3026–3038; c) M. Parasram and V. Gevorgyan, *Chem. Soc. Rev.* 2017, **46**, 6227–6240; d) K. N. Lee and M.-Y. Ngai, *Chem. Commun.* 2017, **53**, 13093–13112; e) Y. – Q. Zou, F. M. Hoermann and T. Bach, *Chem. Soc. Rev.* 2018, **47**, 278–290; f) J.-R. Chen, X.-Q. Hu, L.-Q. Lu and W.-J. Xiao, *Acc. Chem. Res.* 2016, **49**, 1911–1923; g) D. Ravelli, M. Fagnoni and A. Albini, *Chem. Soc. Rev.* 2013, **42**, 97–113.
- a) S. P. Pitre, C. D. McTiernan and J. C. Scaiano, *ACS Omega* 2016, **1**, 66–76; b) S. P. Pitre, C. D. McTiernan and J. C. Scaiano, *Acc. Chem. Res.* 2016, **49**, 1320–1330.
- a) D. A. Nicewicz and T. M. Nguyen, *ACS Catal.* 2014, **4**, 355–360; b) N. A. Romero and D. A. Nicewicz, *Chem. Rev.* 2016, **116**, 10075–10166.
- M. H. Shaw, J. Twilton and D. W. C. MacMillan, *J. Org. Chem.* 2016, **81**, 6898–6926.
- a) A. Gualandi, M. Marchini, L. Mengozzi, M. Natali, M. Lucarini, P. Ceroni and P. G. Cozzi, *ACS Catal.* 2015, **5**, 5927–5931; b) R. F. Higgins, S. M. Fatur, S. G. Shepard, S. M. Stevenson, D. J. Boston, E. M. Ferreira, N. H. Damrauer, A. K. Rappé and M. P. Shores, *J. Am. Chem. Soc.* 2016, **138**, 5451–5464; For a review: c) C. B. Larsen and O. S. Wenger, *Chem. Eur. J.* 2018, **24**, 2039–2058.
- C. K. Prier, D. A. Rankic and D. W. C. MacMillan, *Chem. Rev.* 2013, **113**, 5322–5363.
- a) K. A. Margrey and D. A. Nicewicz, *Acc. Chem. Res.* 2016, **49**, 1997–2006; For the properties of Fukuzumi's catalysts, see: S. Fukuzumi, K. Ohkubo and T. Suenobu, *Acc. Chem. Res.* 2014, **47**, 1455–1464.
- J. A. Murphy, *J. Org. Chem.* 2014, **79**, 3731–3746.
- a) E. H. Discekici, N. J. Treat, S. O. Poelma, K. M. Mattson, Z. M. Hudson, Y. Luo, C. J. Hawker and J. R. de Alaniz, *Chem. Comm.* 2015, **51**, 11705–11708; b) G. M. Miyake and J. C. Theriot, *Macromolecules* 2014, **47**, 8255–8261; c) Y. Du, R. M. Pearson, C.-H. Lim, S. M. Sartor, M. D. Ryan, H. Yang, N. H. Damrauer and G. M. Miyake, *Chem. Eur. J.* 2017, **23**, 10962–10968; d) J. C. Theriot, C.-H. Lim, H. Yang, M. D. Ryan, C. B. Musgrave and G. M. Miyake, *Science* 2016, **352**, 1082–1086. For single electron donors produced by different mechanistic pathways, see the work of König: J. I. Bardagi, I. Ghosh, M. Schmalzbauer, T. Ghosh and B. König, *Eur. J. Org. Chem.* 2018, 34–40, and ref. therein.
- Quite recently, a number of phenoxazine derivatives have been developed as visible light-absorbing, organic photoredox catalysts (PCs) with high reduction potentials (–1.42 V up to –2.01 vs SCE); see: B. McCarthy, R. Pearson, C.-H. Lim, S. M. Sartor, N. H. Damrauer and G. M. Miyake, *J. Am. Chem. Soc.* 2018, **140**, 5088–5101.
- a) H. Li, L. Cai, J. Li, Y. Hu, P. Zhou and J. Zhang, *Dyes and Pigments*, 2011, **91**, 309–316; b) B. B. Raju and T. S. Varadarajan, *Laser Chem.* 1995, **16**, 109–120; c) M. Fujiwara, N. Ishida, M. Satsuki and S. Suga, *J. Photopolym. Sci. Technol.* 2002, **15**, 237–238; d) K. Hara, Z.-S. Wang, T. Sato, A. Furube, R. Katoh, H. Sugihara, Y. Dan-oh, C. Kasada, A. Shinpo and S. Suga, *J. Phys. Chem. B* 2005, **109**, 15476–15482.
- S.-M. Yang, C.-Y. Wang, C.-K. Lin, P. Karanam and G. M. Reddy, *Angew. Chem. Int. Ed.* 2018, **57**, 1668–1672 and ref. cited therein.
- M. Ozkütük, E. Ipek, B. Aydin, S. Mamas and Z. Seferoglu, *J. Mol. Struct.* 2016, **108**, 521–532.
- Coumarin **9** is available from TCI, D5007. Price: 1g 78.00 EUR.
- For a reductive coupling by the use of perylene as photoredox catalyst, see: S. Okamoto, K. Kojima, H. Tsujioka and A. Sudo, *Chem. Comm.* 2016, **52**, 11339–11342.
- a) M. Nakajima, E. Fava, S. Loeschner, Z. Jiang and M. Rueping, *Angew. Chem. Int. Ed.* 2015, **54**, 8828–8832; b) E. Fava, A. Millet, M. Nakajima, S. Loeschner and M. Rueping, *Angew. Chem. Int. Ed.* 2016, **55**, 6776–6779; c) L. Lin, X. Bai, X. Ye, X. Zhao, C.-H. Tan, Z. Jiang, *Angew. Chem. Int. Ed.* 2017, **56**, 13842–13846; d) C.-Ming Wang, P.-J. Xia, J.-A. Xiao, J. Li, H.-Y. Xiang, X.-Q. Chen, *J. Org. Chem.* 2017, **82**, 3895–3900; e) L. J. Rono, H. G. Yayla, Y. Wang, M. F. Armstrong, R. R. Knowles, *J. Am. Chem. Soc.* 2013, **135**, 17735–17738; f) M. Zhang, W. D. Rouch, R. D. McCulla, *Eur. J. Org. Chem.* 2012, 6187–6196; g) J. Cao, G. Wang, L. Gao, X. Cheng, S. Li, *Chem. Sci.* 2018, **9**, 3664–3671.
- For electrochemical potential of organic molecule, see: H. G. Roth, N. A. Romero and D. A. Nicewicz, *Synlett* 2016, **27**, 714–723.
- Coumarin **8** was prepared in 82% yield following the literature; see: F. Jafarpour, S. Zarei, M. Barzegar Amiri Olia, N. Jalalimanesh and S. Rahiminejadan, *J. Org. Chem.* 2013, **78**, 2957–2964.
- From a photophysical point of view, coumarins **8**, **9** and **10** are quite similar: a small decrease in energy of the S<sub>0</sub>–S<sub>1</sub> electronic transition and of the redox potential corresponding to the first oxidation is observed for coumarins **9** and **10** compared to **8**. Moreover, the cyclic voltammetry analysis (SI) shows a greater stability of the oxidized species of **10** compared to that of **9**, a property that could be responsible for the better efficiency of the catalyst.
- J. Chin, F. Mancin, N. Thavarajah, D. Lee, A. Lough, D. S. Chung, *J. Am. Chem. Soc.* 2003, **125**, 15276–15277.
- The Stern-Volmer analysis was conducted varying the concentration of selected aldehyde (see SI), in the presence of tertiary amine, similarly to the reaction conditions, as the amine is not able to quench the excited state of the coumarin catalyst.
- The S<sub>1</sub> fluorescent excited states of coumarins **9** and **10** (Table 1) lie at lower energy than the T<sub>1</sub> excited states of the investigated aldehydes, see: a) H. Gerner and H. J. Kuhn, *J. Phys. Chem.* 1986, **90**, 5946–5955; b) M. J. van der Burgt, J. Jansen, A. H. Huizer and C. A.G.O. Varma, *J. Mol. Struct.* 1996, **385**, 175–183.
- E. M. Arnett and C. A. Palmer, *J. Am. Chem. Soc.* 1990, **112**, 1354–1360.
- a) G. Magagnano, A. Gualandi, M. Marchini, L. Mengozzi, P. Ceroni and P. G. Cozzi, *Chem. Commun.* 2017, **53**, 1591–1594; b) E. Arceo, E. Montroni and P. Melchiorre, *Angew. Chem. Int. Ed.* 2015, **52**, 12064–12068.
- a) J. D. Nguyen, J. W. Tucker, M. D. Konieczynska and C. R. J. Stephenson, *J. Am. Chem. Soc.* 2011, **133**, 4160–4163; b) C.-J. Wallentin, J. D. Nguyen, P. Finkbeiner, and C. R. J. Stephenson, *J. Am. Chem. Soc.* 2012, **134**, 8875–8884.
- Y. Yasu, T. Koike and M. Akita, *Angew. Chem. Int. Ed.* 2012, **51**, 9567.
- D. Nicewicz and D. W. C. MacMillan, *Science* 2008, **322**, 77–80.
- A. E. Allen and D. W. C. MacMillan, *Chem. Sci.* 2012, **3**, 633–658.
- H. S. Li and Y. Q. Li, *Chin. Chem. Lett.* 2010, **21**, 931–934.