

Communication

**Aziridinyl fluorophores demonstrate bright fluorescence  
and superior photostability through effectively  
inhibiting twisted intramolecular charge transfer**

Xiaogang Liu, Qinglong Qiao, Wenming Tian, Wenjuan Liu, Jie Chen, Matthew J Lang, and Zhaochao Xu

*J. Am. Chem. Soc.*, **Just Accepted Manuscript** • DOI: 10.1021/jacs.6b03924 • Publication Date (Web): 20 May 2016

Downloaded from <http://pubs.acs.org> on May 23, 2016

**Just Accepted**

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



**ACS Publications**

# Aziridinyl fluorophores demonstrate bright fluorescence and superior photostability through effectively inhibiting twisted intramolecular charge transfer

Xiaogang Liu,<sup>†,‡,¶,\*</sup> Qinglong Qiao,<sup>†,§,¶</sup> Wenming Tian,<sup>†</sup> Wenjuan Liu,<sup>†</sup> Jie Chen,<sup>†</sup> Matthew J. Lang,<sup>‡,⊥</sup> Zhaochao Xu<sup>†,\*</sup>

<sup>†</sup>Key Laboratory of Separation Science for Analytical Chemistry, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, 457 Zhongshan Road, Dalian 116023, China.

<sup>‡</sup>Singapore-MIT Alliance for Research and Technology (SMART), 1 CREATE Way, Singapore 138602, Singapore.

<sup>§</sup>State Key Laboratory of Fine Chemicals, Dalian University of Technology, 2 Linggong Road, Dalian 116012, China.

<sup>⊥</sup>Department of Chemical and Biomolecular Engineering and Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN 37235, United States.

## Supporting Information Placeholder

**ABSTRACT:** Replacing conventional dialkylamino substituents with a three-membered aziridine ring in naphthalimide leads to significantly enhanced brightness and photostability, via effectively suppressing twisted intramolecular charge transfer (TICT) formation. This replacement is generalizable in other chemical families of fluorophores, such as coumarin, phthalimide and nitrobenzoxadiazole dyes. In highly polar fluorophores, we show that aziridinyl dyes even outperform their azetidinyl analogues in aqueous solution. We also proposed one simple mechanism that can explain the vulnerability of quantum yield to hydrogen bond interactions in protonic solvents in various fluorophore families. Such knowledge is a critical step towards developing high-performance fluorophores for advanced fluorescence imaging.

Rapid evolution of fluorescence imaging techniques in recent years demands fluorophores with enhanced brightness and photostability.<sup>1-4</sup> This evolution, particularly in biomolecular labeling<sup>5,6</sup> and super-resolution imaging techniques,<sup>7-10</sup> has facilitated fluorescence imaging with single-molecule precision in numerous biological and biomedical studies.<sup>11-13</sup> However, many existing fluorophores lack sufficient brightness and photostability for single-molecule and live-cell imaging. Rational molecular engineering of fluorophores, based on a deep understanding of their working mechanism, is thus crucial and imperative to yield novel fluorophores with superior brightness and photostability.

On the theoretical side, it is proposed that twisted intramolecular charge transfer (TICT) acts as one of the major non-radiative de-excitation pathways in fluorophores.<sup>14</sup> In the TICT state, a dialkylamino donor twists out the fluorophore scaffold by approximately 90° upon photoexcitation, forming a non-emissive and highly reactive chemical species. Accordingly, various

experimental strategies have been applied to avoid TICT formation. Commercially available fluorophores, such as Dylight, CF and Alexa dyes, involve complex structural modifications via rigidifying amino substituents. Unfortunately, most molecular structures of these dyes are still not disclosed, constraining their further derivatization and functionalization. Song *et al.* employed a bulky 7-azabicyclo[2.2.1]heptane to replace commonly used dimethylamino substituents in rhodamine dyes.<sup>15</sup> This structural modification greatly improves the quantum yields and photostability of rhodamine dyes, but at the cost of deteriorated water solubility. In a landmark paper, Lavis *et al.* changed *N,N*-dialkylamino substituents to four-membered azetidide rings in several chemical families of fluorophores, such as rhodamines, rhodols and coumarins.<sup>2</sup> They demonstrated that the azetidide ring simultaneously improves dye brightness and photostability. Moreover, this simple structural modification retains biochemical properties of the parent compounds (such as cell-permeability and intracellular labeling), and becomes accessible even in basic chemical synthesis laboratories. Nevertheless, the degree of improvement with the azetidide ring based approach varies across different fluorophore families. For instance, the quantum yield of azetidinyl naphthalimide is markedly lower than those of coumarins and rhodamines, demanding further improvement.<sup>2</sup> In this communication, we show that a three-membered aziridine ring suppresses TICT formation more effectively over the four-membered azetidide ring and other dialkylamino substituents; consequently, the aziridine ring leads to improved brightness and superior photostability in a range of fluorophores, in conjunction with enlarged Stokes shifts.




Figure 1. The three-membered aziridine ring enhances the brightness and photostability of naphthalimide fluorophores. (a) Illustration of steric hindrance between donor and acceptor moieties of a fluorophore. (b) Three types of alignment conformations between the two amino “arms” (highlighted in yellow) and the main fluorophore scaffold (green) in the ground state ( $S_0$ ). Arrows indicate potential rotations to enter the twisted intramolecular charge transfer (TICT) state in the first excited state ( $S_1$ ). (c) The top and side views of the theoretically optimized molecular structures of 1–7 in ethanol in the ground state. (d) Calculated potential energy surfaces (PES) of the ground ( $S_0$ , in blue) and the first excited singlet ( $S_1$ , in red) states of 1–3 in ethanol, as a function of the rotational angle ( $\theta$ ) between the amino substituent and the main fluorophore scaffold.  $\theta$  is the average dihedral angle of two amino “arms” with respect to the naphthalimide scaffold. The LE state has a more planar conformation and less charge transfer compared to the TICT state. (e) Experimental spectroscopic data for 1–7 in ethanol.  $\lambda_{\text{abs}}$ , peak UV–vis absorption wavelength;  $\epsilon$ , molar extinction coefficient;  $\lambda_{\text{em}}$ , peak emission wavelength;  $\Delta\lambda$  and  $\Delta\nu$ , the Stokes shift;  $\phi$ , quantum yield. (f) Experimental quantum yields of 1–3 in various solvents. (g) Relative intensity changes of 1, 2, and Rhodamine B in DMSO/water mixture (volume ratio, 30:70) during 12 hours of strong white light exposure. (h) Relative intensity changes of 8 and 9 in water during 500 seconds of strong laser radiation (416 nm) in microfluidic channels. The insets show the molecular structures of 8 and 9.

We initially choose naphthalimide-derived 1–7 as model compounds to investigate the effect of azacyclic substituents on minimizing TICT (Figure 1). Our computational studies show that the amino “arms” and the main fluorophore scaffold of 1–7 can align in three different conformations, due to competing contributions from azacyclic strain energy, steric hindrance, and resonance effect (Figure 1a, b). The resonance effect depends on the electron-donating/withdrawing strength of both the azacyclic rings and the fluorophore scaffolds. We find that compounds 1, 2 and 3–7 possess the *up-up*, *flat* and *up-down* conformations, respectively (Figure 1c). Intuitively, the *up-up* conformation should possess the highest TICT resistance, because

one of its amino arms faces strong steric repulsion to enter the TICT state. Similarly, the *up-down* conformation should exhibit the lowest TICT resistance, because steric hindrance facilitates its TICT rotation after photoexcitation.<sup>16</sup> Indeed, our theoretical calculations show that in a polar environment (such as ethanol), the TICT state of 1 (the *up-up* conformation) is energetically unfavorable, and a significant energy barrier (0.393 eV) exists between the local excited (LE) and TICT states (Figure 1d). The TICT state of 2 (the *flat* conformation), however, becomes slightly more stable than the LE state, in conjunction with a reduced energy barrier (0.246 eV). In contrast, a strong tendency to the TICT state is most evident in 3 (the *up-down*




Figure 2. (a) Molecular structures of 10—17. (b) Experimental spectroscopic data for 10–17 in aqueous solution.  $\lambda_{\text{abs}}$ , peak UV—vis absorption wavelength;  $\epsilon$ , molar extinction coefficient;  $\lambda_{\text{em}}$ , peak emission wavelength;  $\Delta\lambda$  and  $\Delta\nu$ , the Stokes shift;  $\phi$ , quantum yield. (c). The atomic contributions to the HOMO (highest occupied molecular orbital) and LUMO (lowest unoccupied molecular orbital) electron densities of 1, 10, 14 and 16, calculated based on the optimized molecular structures in the  $S_0$  state. The blue/pink circle size represents the atomic contribution; only contributions greater than 0.02 are shown. Hydrogen bond formation sites in naphthalimide, coumarin, phthalimide, and NBD fluorophore scaffolds are indicated by arrows. Molecular sites that experience significant partial charge increase upon photoexcitation are highlighted by red arrows in 14 and 16.

conformation; Figure 1d). Nevertheless, in a non-polar environment, our calculations show that 1—3 all favor LE emission, but not TICT rotation (Supporting Information Section 1).

Experimental results are in excellent agreement with our theoretical calculations. In non-polar solvents, 1—7 all emit bright fluorescence (quantum yield,  $\phi \approx 0.750$ ). In polar solvents, only 1 and 2 demonstrate high quantum yields. We notice that 1 exhibits higher quantum yields than 2 in all tested solvents. This difference is most apparent in water ( $\phi = 0.432$  for 1 and 0.199 for 2; Figure 1e, f; Supporting Information Section 2). It is thus clear that the aziridine ring possesses higher TICT resistance than the azetidine ring and other dialkylamino substituents.

We then tested the photostability of 1 and 2 (under white light radiation; Figure 1g), and their analogues with enhanced water solubility, 8 and 9 (under 416 nm laser radiation; Figure 1h). Our tests show that aziridinyl 1 and 8 exhibit considerably higher photostability than azetidiny 2 and 9. The chemical stability of 1 was also evaluated in solutions of different pH. We found that the fluorescence intensities of 1 remained stable from pH 4.2 to 10.8. We also treated the solution of 1 with various metal ions, oxidants and nucleophiles including  $\text{H}_2\text{O}_2$ , NaHS, cysteine and glutathione. The unperturbed emission indicates that 1 has good chemical stability (Supporting Information Section 2).

Several other differences between 1 and 2 are of note. The *up-up* conformation of 1 yields a lower molar extinction coefficient and a blue shift in its UV—vis absorption peak ( $\epsilon = 9,200 \text{ M}^{-1} \text{cm}^{-1}$  and  $\lambda_{\text{abs}} = 382 \text{ nm}$ ) with respect to the planar 2 ( $\epsilon = 15,900 \text{ M}^{-1} \text{cm}^{-1}$  and  $\lambda_{\text{abs}} = 445 \text{ nm}$ ; Figure 1e). This is due to a poorer resonance effect in 1. However, a slight flattening of the aziri-

dine ring in 1 upon photoexcitation results in a large geometry relaxation in the excited state, thus leading to a favorable large Stokes shift ( $\Delta\lambda = 127 \text{ nm}$  for 1 versus 88 nm for 2).<sup>17</sup> The large Stokes shift of 1 is “indispensable in multi-colour super-resolution techniques”.<sup>18</sup> Moreover, owing to the larger quantum yield of 1, its brightness is ~27% higher than that of 2 in aqueous solution. The same patterns were observed between 8 and 9 (Supporting Information Section 2).

Inspired by the improved brightness, enhanced stability and enlarged Stokes shifts in naphthalimide dyes, we then employed the aziridine ring in coumarin, rhodamine, phthalimide and nitrobenzoxadiazole (NBD) dyes, and compared the fluorophore performance against their azetidiny and dimethylamino analogues (Figure 2a, b). Our quantum chemical calculations show that aziridinyl fluorophores demonstrate the highest TICT resistance, followed by azetidiny dyes. Experimental data also show that both aziridinyl and azetidiny fluorophores exhibit higher brightness than conventional dialkylamino fluorophores ( $\phi = 0.058$  for Coumarin 1;  $< 0.01$  for dimethylamino NBD and phthalimide in water). Between aziridinyl and azetidiny fluorophores, coumarins 10 and 11 display comparably high quantum yields ( $\phi \approx 0.900$ ) in water. Since these 4-methylcoumarin dyes have a low polarity and weak TICT tendency, the azetidine ring seems sufficient to suppress TICT formation and afford high quantum yields. The aziridine substituent in 12, however, leads to the colorless and non-fluorescent lactone form of rhodamine dyes, while azetidiny 13 are highly emissive ( $\phi = 0.459$ ). These results are consistent with Lavis and coworkers’ observations on similar rhodamine dyes, where both dialkylamino groups are replaced by aziridine or azetidine rings.<sup>2</sup> In contrast, for phthalimide and NBD dyes 14—17, the

quantum yields of aziridinyl fluorophores are approximately twice as large as those of azetidiny analogues in aqueous solution, in conjunction with enlarged Stokes shifts (Figure 2b). These results are remarkable, considering that dimethylamino phthalimide and NBD dyes are non-emissive in water.<sup>19,20</sup> Unfortunately, the absolute quantum yields of 14–17 ( $\phi = 0.011$  to 0.084) are low, largely due to intensive hydrogen bond interactions around the fluorophore scaffolds.<sup>19,21</sup>

We have found one plausible mechanism to explain the vulnerabilities of quantum yield to hydrogen bond interactions among different fluorophores. This vulnerability is closely related to the partial charge increase upon photoexcitation, at hydrogen bond formation sites in the fluorophore scaffolds (Figure 2c, highlighted by arrows). We denote the total charge change at these sites during HOMO→LUMO transition as  $\Delta$ . In naphthalimide and coumarin dyes, no significant  $\Delta$  changes are observed. Coincidentally, the quantum yields of 1 and 10 are high ( $\phi = 0.432$  for 1 and 0.899 for 10). In contrast, we notice substantial  $\Delta$  increases in phthalimide and NBD dyes 14 and 16 (Figure 2c, highlight by red arrows), indicating intensive hydrogen bond interactions after photoexcitation. Accordingly, their quantum yields are low ( $\phi = 0.020$  for 14 and 0.084 for 16). Interestingly, this mechanism also works perfectly in explaining the quantum yield differences of several other fluorophore families in protonic solvents, such as rhodol and oxazine dyes (Supporting Information Section 1). Additional quantum chemical calculations including explicit solvent molecules are required to further understand such hydrogen bond interactions, and are the subject of future work.

In conclusion, we have demonstrated that the *up-up* conformed aziridine ring is highly effective in suppressing TICT rotation. Aziridinyl fluorophores thus demonstrate considerably enhanced brightness and superior photostability, in comparison to conventional dialkylamino substituted dyes. In highly polar fluorophores with a strong TICT tendency (such as naphthalimide dyes), aziridinyl fluorophores even outperform their azetidiny analogues in aqueous solution. As another favorable feature, a slight flattening of the aziridine ring upon photoexcitation affords large Stokes shifts in aziridinyl fluorophores. Moreover, similar to that of the azetidine ring, employing the aziridine ring requires minimal structural modifications, retains advantages of the parent fluorophores, permits facile synthesis and further derivatization, and can be applied to a wide range of fluorophores to enhance dye brightness and photostability. Lastly, we show that substantial charge increase at hydrogen bond formation sites in a fluorophore is detrimental to its quantum yield in protonic solvents. Such knowledge on controlling TICT and managing hydrogen bond interactions will inspire the rational development of abundant high-performance dyes across different fluorophore families, thus enabling unprecedented fluorescence imaging applications.

## ASSOCIATED CONTENT

Supporting Information

Synthesis, characterization, and computational details. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## AUTHOR INFORMATION

Corresponding Author

\*Email: [xiaogang@smart.mit.edu](mailto:xiaogang@smart.mit.edu); Tel: +65-6516-1462.

\*Email: [zcxu@dicp.ac.cn](mailto:zcxu@dicp.ac.cn); Tel: +86-411-84379648.

Author Contributions

†X.L. and Q.Q. contributed equally to this work.

Notes

The authors declare no competing financial interests.

## ACKNOWLEDGMENT


X.L. and M.J.L. thank National Research Foundation of Singapore (through BioSyM Interdisciplinary Research Group at Singapore-MIT Alliance for Research and Technology) for supporting this project. Z.X. acknowledges the National Natural Science Foundation of China (21276251), the 100 Talents Program funded by Chinese Academy of Sciences, Dalian Cultivation Fund for Distinguished Young Scholars (2014J11JH130 and 2015J12JH205) and the National Science Fund for Excellent Young Scholars (21422606). The authors thank Dr. Jie Pan for her assistance on photostability tests.

## REFERENCES

- (1) van der Velde, J. H.; Oelerich, J.; Huang, J.; Smit, J. H.; Jazi, A. A.; Galiani, S.; Kolmakov, K.; Guoridis, G.; Eggeling, C.; Herrmann, A. *Nat. Commun.* 2016, 7, 10144.
- (2) Grimm, J. B.; English, B. P.; Chen, J.; Slaughter, J. P.; Zhang, Z.; Revyakin, A.; Patel, R.; Macklin, J. J.; Normanno, D.; Singer, R. H.; Lionnet, T.; Lavis, L. D. *Nat. Methods* 2015, 12, 244.
- (3) Dempsey, G. T.; Vaughan, J. C.; Chen, K. H.; Bates, M.; Zhuang, X. *Nat. Methods* 2011, 8, 1027.
- (4) Altman, R. B.; Terry, D. S.; Zhou, Z.; Zheng, Q.; Geggier, P.; Kolster, R. A.; Zhao, Y.; Javitch, J. A.; Warren, J. D.; Blanchard, S. C. *Nat. Methods* 2012, 9, 68.
- (5) Gautier, A.; Juillerat, A.; Heinis, C.; Corr  a, I. R.; Kindermann, M.; Beaufils, F.; Johansson, K. *Chem. Biol.* 2008, 15, 128.
- (6) Los, G. V.; Encell, L. P.; McDougall, M. G.; Hartzell, D. D.; Karassina, N.; Zimprich, C.; Wood, M. G.; Learish, R.; Ohana, R. F.; Urh, M. *ACS Chem. Biol.* 2008, 3, 373.
- (7) Planchon, T. A.; Gao, L.; Milkie, D. E.; Davidson, M. W.; Galbraith, J. A.; Galbraith, C. G.; Betzig, E. *Nat. Methods* 2011, 8, 417.
- (8) Jones, S. A.; Shim, S.-H.; He, J.; Zhuang, X. *Nat. Methods* 2011, 8, 499.
- (9) Huang, B.; Babcock, H.; Zhuang, X. *Cell* 2010, 143, 1047.
- (10) Huang, B.; Bates, M.; Zhuang, X. *Annu. Rev. Biochem.* 2009, 78, 993.
- (11) Lukinavi  ius, G.; Umezawa, K.; Olivier, N.; Honigsmann, A.; Yang, G.; Plass, T.; Mueller, V.; Reymond, L.; Corr  a Jr, I. R.; Luo, Z.-G. *Nat. Chem.* 2013, 5, 132.
- (12) Jungmann, R.; Avenda  o, M. S.; Dai, M.; Woehrstein, J. B.; Agasti, S. S.; Feiger, Z.; Rodal, A.; Yin, P. *Nat. Methods* 2016, 13, 439.
- (13) Legant, W. R.; Shao, L.; Grimm, J. B.; Brown, T. A.; Milkie, D. E.; Avants, B. B.; Lavis, L. D.; Betzig, E. *Nat. Methods* 2016, 13, 359.
- (14) Grabowski, Z. R.; Rotkiewicz, K.; Rettig, W. *Chem. Rev.* 2003, 103, 3899.
- (15) Song, X.; Johnson, A.; Foley, J. J. *Am. Chem. Soc.* 2008, 130, 17652.
- (16) Vogel, M.; Rettig, W.; Sens, R.; Drexhage, K. H. *Chem. Phys. Lett.* 1988, 147, 452.
- (17) Liu, X.; Xu, Z.; Cole, J. M. *J. Phys. Chem. C* 2013, 117, 16584.
- (18) Sednev, M. V.; Belov, V. N.; Hell, S. W. *Methods Appl. Fluoresc.* 2015, 3, 042004.
- (19) Loving, G.; Imperiali, B. *J. Am. Chem. Soc.* 2008, 130, 13630.

(20) Saha, S.; Samanta, A. *J. Phys. Chem. A* 1998, 102, 7903.  
(21) Vázquez, M. E.; Blanco, J. B.; Imperiali, B. *J. Am. Chem. Soc.* 2005, 127, 1300.

TOC Figure








Figure 1  
170x158mm (300 x 300 DPI)





Figure 2  
162x88mm (300 x 300 DPI)





TOC Figure  
85x45mm (300 x 300 DPI)