A facile microwave-assisted Diels-Alder reaction of vinylboronates†

Ariel M. Sarotti, Pablo L. Pisano and Silvina C. Pellegrinet*

Received 19th April 2010, Accepted 4th August 2010 DOI: 10.1039/c0ob00020e

The Diels–Alder reaction of vinylboronates can be easily performed using microwave irradiation giving excellent yields of the cycloadducts. Pinacol vinylboronate was the reagent of choice due to its stability towards hydrolysis, operational simplicity and yields of Diels–Alder products. To the best of our knowledge, this is the first example of microwave-assisted Diels–Alder reaction of boron-substituted dienophiles. Subsequent *in situ* oxidation of the cycloadducts with alkaline hydrogen peroxide afforded the alcohols efficiently.

Introduction

Due to their great versatility, boron-substituted dienophiles have emerged as attractive building blocks in organic synthesis.^{1,2} Unsaturated boronates exhibit poor reactivity in Diels-Alder (DA) reactions, which require temperatures as high as 200 °C and prolonged reaction times. However, unlike dialkylboranes, boronates are stable to air and moisture and can be purified by chromatography. As a result, the Diels-Alder products can be isolated, offering an interesting alternative for further elaboration of C-C, C-O and C-N bonds. Although alkynylboronates have been recently used in a number of [4+2] cycloadditions,³ examples on the use of alkenylboronates as dienophiles are scarce.⁴ In the past years, many papers on microwave-assisted reactions of boronic esters appeared in the literature.^{1,5} However, this seems to be limited to cross-coupling reactions mostly. To the best of our knowledge, the possibility of using microwave irradiation to facilitate cycloadditions of boron-functionalized compounds remained unexplored. Herein we report that the Diels-Alder reaction of vinylboronates can be easily performed using microwave irradiation.

Results and discussion

We first studied the microwave-assisted reactions of commercial vinylboronates derived from n-butanol (1a), pinacol (1b) and 2-methyl-2,4-pentanediol (1c)^{4c,6} with cyclopentadiene (2a) and 1,3-cyclohexadiene (2b), some of which have been previously carried out with conventional heating (Scheme 1).^{4a-d} We were pleased to note that the reactions could be completed in short reaction times with excellent yields (Table 1, entries 1–7). Best product yields were obtained using the pinacol derivative 1b. Also, vinylboronate 1b and its derived products offered greater stability towards hydrolysis than 1a and operational simplicity than 1c, which affords complex diastereomeric mixtures due the presence of a stereogenic center. We also observed that *endo/exo* stereoselectivities for all dienophiles were comparable



Scheme 1 Microwave-assisted Diels-Alder reaction of vinylboronates.

and similar to those reported for the thermal reactions. For pinacol vinylboronate (1b), we were able to reduce the number of equivalents of diene to three (entries 3 and 7).

In light of these encouraging results, we next performed the microwave-assisted Diels-Alder reactions with a series of structurally diverse dienes (2c-2i) (entries 8-14). All reactions were complete within a few hours and proceeded in good to excellent yields.⁷ The tolerance of the boronic ester functionality in 1b to the reaction temperatures is remarkable. The observed reactivity trend for the dienes was consistent with literature reports of Diels-Alder reactions with other electron-poor dienophiles such as tetracyanoethylene or maleic anhydride.8 Inspection of the results in Table 1 also reveals that endo/exo stereoselectivities and regioselectivities varied depending on the nature of the diene.9 Most reactions showed moderate to good exo selectivities (up to endo/exo 12:88 for 2i, entry 14). Only the reactions with 1,3cyclohexadiene (2b) showed a significant preference for the formation of the endo adduct (endo/exo 75:25, entries 6-7). In addition, for 2-substituted dienes such as isoprene (2c) and myrcene (2d) the para products were slightly favored (entries 8 and 9). 1-Substituted dienes showed different levels of regioselectivity: while 1-methoxy-1,3-cyclohexadiene (2h) showed high ortho selectivity (ortho/meta >80:20, entry 13), trans-pyperylene (2e) gave similar quantities of the ortho and meta products (entry 10).

Preliminary theoretical calculations suggest that the reactions under study are normal electron-demand Diels–Alder reactions. The low reactivity of vinylboronates appears to be related to the donating effect of the oxygens on the boron atom, which increases the energy of the LUMO. To rationalize the experimental reactivities and selectivities more calculations are currently underway,

Instituto de Química Rosario (CONICET), Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Suipacha 531, Rosario (2000), Argentina. E-mail: pellegrinet@iquir-conicet.gov.ar; Fax: +54 341 4370477; Tel: +54 341 4370477

[†] Electronic supplementary information (ESI) available: Experimental procedures, spectroscopic and analytic data and NMR spectra for all compounds. See DOI: 10.1039/c0ob00020e

Entry	Vinylboronate	Diene	Conditions ^{<i>a</i>,<i>b</i>}	Yield ^e ratios ^d	Products
1	1a	2a	10 equiv. diene dichloromethane 60 min, 150 °C	Not determined ^e	B-OBU B-OBU BuO Ba-N BuO BuO Ba-X
2	1b		10 equiv. diene 60 min, 150 °C (3 × 20 min)	100% endo/exo 37:63	B-0 3b-N 3b-X
3	1b		3 equiv. diene 60 min, 150 °C (3 × 20 min)	100% endo/exo 38:62	
4	1c		10 equiv. diene 80 min, 150 °C (4 × 20 min)	95% endo/exo 49:51	3c-N et 3c-X
5	1a	2b	15 equiv. diene 1 h, 150 °C 3 h, 220 °C	Not determined ^e	Buo 3d-N 3d-X
6	1b		10 equiv. diene 4 h, 220 °C (4 × 1 h)	91% endo/exo 75:25	B-O 3e-N 3e-X
7	1b		3 equiv. diene 4 h, 220 °C (4 × 1 h)	93% endo/exo 75:25	
8	1b	20	3 equiv. diene 3 h, 220 °C	92% meta/para 33:67	JF-M JF-P
9	1b	Zd Zd	3.4 equiv. diene 2 h, 220 °C (2 × 1 h)	94% meta/para 35:65	$\begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \\ \\ \end{array} \end{array} \\ \end{array} \\ \begin{array}{c} \\ \end{array} \\ \begin{array}{c} \\ \\ \end{array} \end{array} \\ \begin{array}{c} \\ \\ \end{array} \\ \begin{array}{c} \\ \\ \end{array} \end{array} \\ \begin{array}{c} \\ \\ \end{array} \\ \begin{array}{c} \\ \\ \end{array} \\ \begin{array}{c} \\ \\ \\ \end{array} \\ \end{array} \\ \begin{array}{c} \\ \\ \end{array} \\ \begin{array}{c} \\ \\ \end{array} \\ \end{array} \\ \end{array} \\ \begin{array}{c} \\ \\ \end{array} \\ \end{array} \\ \end{array} \\ \begin{array}{c} \\ \\ \end{array} \\ \end{array} \\ \end{array} \\ \end{array} \\ \end{array} \\ \begin{array}{c} \\ \\ \end{array} $
10	1b	2e	4 equiv. diene 4 h, 220 °C	83% endo/exo 39:61 ortho/meta 54:46	$ \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \end{array}\\ \end{array}\\ \end{array} \\ \begin{array}{c} \end{array}\\ \end{array} \\ \begin{array}{c} \end{array}\\ \end{array} \\ \begin{array}{c} \end{array} \\ \begin{array}{c} \end{array}\\ \end{array} \\ \begin{array}{c} \end{array} \\ \end{array} \\ \begin{array}{c} \end{array} \\ \begin{array}{c} \end{array} \\ \end{array} \\ \begin{array}{c} \end{array} \\ \end{array} \\ \begin{array}{c} \begin{array}{c} \end{array} \\ \end{array} \\ \begin{array}{c} \end{array} \\ \begin{array}{c} \end{array} \\ \end{array} \\ \end{array} \\ \end{array} \\ \end{array} \\ \begin{array}{c} \end{array} \\ \end{array} \\ \end{array} \\ \end{array} \\ \end{array} \\ \end{array} $ } \\ \end{array} \\ \end{array} \\ \end{array} \\ \end{array} } \\ \end{array} \\ \end{array} \\ \end{array} } \\ \end{array} \\ \end{array} } \\ \end{array} \\ \end{array} } \\ } } \\ \end{array} } } \\ \end{array} } } } \\ } } } } } } } } } }
11	1b	21	3 equiv. diene 3 h, 200 °C (3 × 1 h)	98%	J Box 3i

Table 1 Microwave-assisted Diels-Alder reaction of vinylboronates

Table 1 (Contd.)



^{*a*} All reactions were run in toluene, unless otherwise noted. ^{*b*} The reactions were monitored using ¹¹B NMR analysis in CDCl₃. Typically, the boron signal shifted 3.5 ppm downfield. As an example, the signal at δ 29.3 ppm for vinylboronate **1b** shifted to δ 33.8 ppm for the products of reaction with cyclopentadiene (**3b**). ^{*c*} Yields of isolated products after purification by chromatography on silica gel. ^{*d*} Determined by NMR integration. ^{*e*} The boronate products did not resist silica gel chromatography and hydrolysis to the boronic acids proved impractical.

though we expect that both steric and electronic effects play a significant role.

As a means to demonstrate the synthetic versatility of boronates 3, we next investigated the functionalization of the C-B bond. We studied the sequence consisting of tandem microwave-assisted Diels-Alder reaction followed by in situ oxidation with alkaline hydrogen peroxide (Scheme 2, Table 2).¹⁰ After the initial step was complete, the reaction mixtures were transferred and subjected to oxidation without isolation of the cycloadducts. It is important to remark that no special care was taken to avoid moisture and air. All reactions afforded the alcohols (4a-h) in good to excellent yields. Again, optimal yields were obtained with pinacol vinylboronate (1b). Furthermore, endo/exo stereoselectivities and regioselectivities did not change after oxidation, which confirmed that oxidation proceeded with retention of stereochemistry. These experiments demonstrate that vinylboronates can be efficiently employed as synthetic equivalents of vinyl alcohol in Diels-Alder reactions. The cyclohexenols thus obtained could be further transformed into structurally diverse compounds.



Scheme 2 Tandem microwave-assisted Diels-Alder reaction of vinylboronates-oxidation.

Conclusions

We have shown that the Diels–Alder reactions of vinylboronates can be easily performed using microwave irradiation giving excellent yields of the cycloadducts. Pinacol vinylboronate was the reagent of choice due to its stability towards hydrolysis, operational simplicity and yields of Diels–Alder products. To the best of our knowledge, this is the first example of microwave-assisted Diels– Alder reaction of boron-substituted dienophiles. Subsequent *in situ* oxidation of the cycloadducts with alkaline hydrogen peroxide afforded the alcohols efficiently, demonstrating the utility of these intermediates for C–O bond-forming reactions. We hope this investigation promotes further research on microwave-assisted boronate cycloadditions.

Experimental section

Typical procedure for the microwave-assisted Diels–Alder reaction of vinylboronates (Table 1). Synthesis of 2-bicyclo[2.2.1]hept-5-en-2-yl-4,4,5,5-tetramethyl-[1,3,2]di-oxaborolane (3b)¹¹

To an oven-dried 10 mL pressure-rated reaction vial equipped with a stirring bar were added dry toluene (0.5 mL), vinylboronate **1b** (85 μ L, 0.5 mmol) and diene **2a** (42 μ L, 0.5 mmol) under argon atmosphere. The resulting reaction mixture was stirred at 150 °C in the microwave for 20 min. Two more runs were conducted after addition of extra portions of diene (2 × 0.5 mmol of **2a** followed by 20 min at 150 °C). The solvent was removed under reduced pressure, and the crude was purified by column chromatography

Entry	Vinylboronate	Diene	Yield DA + [O] ^{<i>a</i>} ratios ^{<i>b</i>}	Products DA + [O]
1	1a	2a	91% endo/exo: 41:59	H H Ha-X
2	1b		92% endo/exo: 37 : 63¢	
3	1c		86% endo/exo: 44:56	
4	1a	D 2b	77% endo/exo: 73 : 27	4b-N 4b-X
5	1b		96% endo/exo: 73 : 27 ^d	
6	1b	20	99% meta/para 34:66	4c-M 4c-P
7	1b	Zd Zd	90% meta/para 33 : 67	Hd-M Hd-P OH
8	1b	20	68% endo/exo 39:61 ortho/meta 54:46	4e-ON 4e-OX 4e-MN 4e-MX
9	1b	21	92%	OH 4f
10	1b	Ph Ph 2g	92% endo/exo 47:53	Ph OH Ph 4g-N 4g-X
11	1b	CMe CMe	51% ortho/meta >95:5 ortho endo/ortho exo 49:51	4h-ON 4h-OX 4h-MN 4h-MX

 Table 2
 Tandem microwave-assisted Diels–Alder reaction of vinylboronates-oxidation

^{*a*} Yields of isolated products after purification by chromatography on silica gel. ^{*b*} Determined by NMR integration. ^{*c*} Conditions for the DA reaction are the same as reported in Table 1, entry 3. ^{*d*} Conditions for the DA reaction are the same as reported in Table 1, entry 7.

(hexane–AcOEt) to afford boronate **3b** (100%, 106 mg, *endo/exo* 38:62). HRMS calcd for C₁₃H₂₅BNO₂ (M+NH₄)⁺: 238.1973, found: 238.1980. **3b-N**: IR (film) v_{max} : 3058, 2976, 2934, 2874, 1412, 1371, 1310, 1273, 1146, 979, 856, 719, 690, 502. ¹H NMR (300 MHz; CDCl₃) δ : 6.00 (br s, 2H, H-5 and H-6), 3.02 (br s, 1H, H-1), 2.86 (br s, 1H, H-4), 1.83 (ddd, $J_{3n,3x} = 10.8$, $J_{2,3x} = 10.0$, $J_{3x,4} = 3.6$ Hz, 1H, H-3x), 1.40-1.32 (m, 2H, H-2 and H-7), 1.19 (s,

12H, H-9), 1.15–1.08 (m, 2H, H-3n and H-7). ¹³C NMR (75 MHz; CDCl₃) δ : 135.8 (CH, C-5), 134.4 (CH, C-6), 82.7 (2C, C-8), 50.3 (CH₂, C-7), 44.2 (CH, C-1), 42.3 (CH, C-4), 27.7 (CH₂, C-3), 24.6 (4CH₃, C-9), C-2 signal missing. ¹¹B NMR (96 MHz; CDCl₃) δ : 33.4. **3b-X**: IR (film) v_{max} : 3056, 2975, 2931, 2871, 1370, 1313, 1232, 1146, 976, 858, 719. ¹H NMR (300 MHz; CDCl₃) δ : 6.09 (dd, $J_{5,6}$ = 5.4, $J_{1,6}$ = 2.9 Hz, 1H, H-6), 5.95 (dd, $J_{5,6}$ = 5.4, $J_{4,5}$ = 3.0 Hz, 1H,

H-5), 2.88 (br s, 2H, H-1 and H-4), 1.71 (ddd, $J_{3n,3x} = 11.1$, $J_{2,3x} = 4.2$, $J_{3x,4} = 4.2$ Hz, 1H, H-3x), 1.25–1.06 (m, 15H, H-3n, H-7 and H-9), 0.68 (ddd, $J_{2,3n} = 9.7$, $J_{2,3x} = 4.9$, $J_{1,2} = 1.3$ Hz, 1H, H-2). ¹³C NMR (75 MHz; CDCl₃) δ : 137.0 (CH, C-6), 134.2 (CH, C-5), 82.8 (2C, C-8), 47.4 (CH₂, C-7), 44.0 (CH, C-1), 42.2 (CH, C-4), 27.6 (CH₂, C-3), 24.6 (4CH₃, C-9), C-2 signal missing. ¹¹B NMR (96 MHz; CDCl₃) δ : 34.1.

Typical procedure for the tandem microwave-assisted Diels–Alder reaction of vinylboronates - oxidation (Table 2). Synthesis of Bicyclo[2.2.1]hept-5-en-2-ol (4a)¹²

The microwave-assisted Diels–Alder reaction between **1b** and **2a** was performed as described above. Upon completion, the reaction mixture was diluted with THF (3 mL), and transferred to a 25 mL round-bottom flask. After the addition of Et_3N (1 mL) the solution was cooled to 0 °C and treated alternately with 3 N NaOH (3 mL) and 30% H₂O₂ (3 mL) under argon atmosphere, and then allowed to warm to room temperature and stirred overnight. The reaction mixture was diluted with water (10 mL) and extracted with Et_2O (3 × 15 mL). The combined organic layers were washed with saturated NH₄Cl (15 mL) and brine (15 mL) and dried over anhydrous Na₂SO₄. The solvent was removed under reduced pressure at 0 °C, and the crude was purified by column chromatography (pentane– Et_2O) to afford alcohol **4a** (92%, 48.3 mg, *endo/exo* 37:63) with NMR spectra identical to the commercial material.

Acknowledgements

We thank CONICET, U.N.R. and ANPCyT for financial support. A.M.S. thanks CONICET for the award of a fellowship. P.L.P. thanks ANPCyT for the award of a fellowship.

Notes and references

- 1 Boronic Acids, ed. D. G. Hall, Wiley-VCH: Weinheim, 2005.
- 2 G. Hilt and P. Bolze, Synthesis, 2005, 2091–2115.
- 3 (a) J. E. Moore, M. York and J. P. A. Harrity, *Synlett*, 2005, 860–862;
 (b) M. D. Helm, J. E. Moore, A. Plant and J. P. A. Harrity, *Angew. Chem., Int. Ed.*, 2005, 44, 3889–3892; (c) P. M. Delaney, J. E. Moore and J. P. A. Harrity, *Chem. Commun.*, 2006, 3323–3325; (d) M. D. Helm, A. Plant and J. P. A. Harrity, *Org. Biomol. Chem.*, 2006, 4, 4278–4280; (e) E. Gomez-Bengoa, M. D. Helm, A. Plant and J. P. A.

Harrity, J. Am. Chem. Soc., 2007, **129**, 2691–2699; (f) M. D. Helm, A. Plant and J. P. A. Harrity, Synlett, 2007, 2885–2887; (g) D. L. Browne, M. D. Helm, A. Plant and J. P. A. Harrity, Angew. Chem., Int. Ed., 2007, **46**, 8656–8658; (h) P. M. Delaney, D. L. Browne, H. Adams, A. Plant and J. P. A. Harrity, Tetrahedron, 2008, **64**, 866–873; (i) P. M. Delaney, J. Huang, S. J. F. Macdonald and J. P. A. Harrity, Org. Lett., 2008, **10**, 781–783; (j) D. L. Browne, J. F. Vivat, A. Plant, E. Gomez-Bengoa and J. P. A. Harrity, J. Am. Chem. Soc., 2009, **131**, 7762–7769; (k) R. S. Foster, J. Huang, J. F. Vivat, D. L. Browne and J. P. A. Harrity, Org. Lett., 2008, Chem., 2009, **7**, 4052–4056; (l) D. L. Browne and J. P. A. Harrity, Tetrahedron, 2019, **66**, 553–568.

- 4 (a) D. S. Matteson and J. Wasserbillig, J. Org. Chem., 1963, 28, 366–369; (b) K. Kiyoshi, G. W. Willcockson, I. S. Bengelsdorf, and W. G. Woods, US Pat. 3135781, 1964; (c) W. G. Woods and I. S. Bengelsdorf, J. Org. Chem., 1966, 31, 2769–2772; (d) D. S. Matteson and M. L. Talbot, J. Am. Chem. Soc., 1967, 89, 1123–1126; (e) D. A. Evans, W. L. Scott and L. K. Truesdale, Tetrahedron Lett., 1972, 13, 121–124; (f) P. Martinez-Fresneda and M. Vaultier, Tetrahedron Lett., 1989, 30, 2929–2932; (g) K. Narasaka and I. Yamamoto, Tetrahedron, 1992, 48, 5743–5754; (h) C. Rasset and M. Vaultier, Tetrahedron, 1994, 50, 3397–3406; (i) J. D. Bonk and M. A. Avery, Tetrahedron: Asymmetry, 1997, 8, 1149–1152; (j) G. Lorvelec and M. Vaultier, Tetrahedron Lett., 1998, 39, 5185–5188; (k) R. A. Batey, A. N. Thadani and A. J. Lough, J. Am. Chem. Soc., 1999, 121, 450–451.
- P. Lidström, J. Tierney, B. Wathey and J. Westman, *Tetrahedron*, 2001, **57**, 9225–9283; (b) C. O. Kappe, *Angew. Chem., Int. Ed.*, 2004, **43**, 6250–6284; (c) C. O. Kappe and D. Dallinger, *Mol. Diversity*, 2009, **13**, 71–193; (d) S. Caddick and R. Fitzmaurice, *Tetrahedron*, 2009, **65**, 3325–3355; (e) *Microwaves in Organic Synthesis*, ed. A. Loupy, Wiley-VCH: Weinheim, 2002.
- 6 (a) A. P. Lightfoot, G. Maw, C. Thirsk, S. J. R. Twiddle and A. Whiting, *Tetrahedron Lett.*, 2003, 44, 7645–7648; (b) A. P. Lightfoot, S. J. R. Twiddle and A. Whiting, *Org. Biomol. Chem.*, 2005, 3, 3167–3172.
- 7 Several reactions were run to determine the optimal reaction time and temperature for each diene.
- 8 J. Sauer and R. Sustmann, Angew. Chem., Int. Ed. Engl., 1980, 19, 779–807.
- 9 The structure of the products were determined by a combination of spectroscopic methods. See the Supporting Information.
- 10 Conditions for DA reactions are the same as reported in Table 1.
- 11 Boronates 3b, 3f and 3i have been previously synthezised by Diels-Alder reactions of dichlorovinylborane and the corresponding dienes followed by hydrolysis and esterification with pinacol: N. Noiret, A. Youssofi, B. Carboni and M. Vaultier, J. Chem. Soc., Chem. Commun., 1992, 1105–1107.
- 12 Alcohols 4a–4f have been previously synthezised by other methods, including the Diels–Alder reactions of vinylboranes and the corresponding dienes followed by oxidation: (a) D. A. Singleton, J. P. Martinez, J. V. Watson and G. M. Ndip, *Tetrahedron*, 1992, 48, 5831–5838; (b) D. A. Singleton, J. P. Martinez and G. M. Ndip, J. Org. Chem., 1992, 57, 5768–5771; (c) M. Zaidlewicz, J. R. Binkul and W. Sokól, J. Organomet. Chem., 1999, 580, 354–362.