Accepted Manuscript

Stereoselective synthesis of dehydroamino acids using malonic acid half oxyester and aromatic aldehydes

Yuttapong Singjunla, Silvia Colombano, Jérôme Baudoux, Jacques Rouden

PII: S0040-4020(16)30174-0

DOI: 10.1016/j.tet.2016.03.043

Reference: TET 27583

To appear in: *Tetrahedron*

Received Date: 8 February 2016

Revised Date: 2 March 2016

Accepted Date: 13 March 2016

Please cite this article as: Singjunla Y, Colombano S, Baudoux J, Rouden J, Stereoselective synthesis of dehydroamino acids using malonic acid half oxyester and aromatic aldehydes, *Tetrahedron* (2016), doi: 10.1016/j.tet.2016.03.043.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Graphical Abstract

To create your abstract, type over the instructions in the template box below. Fonts or abstract dimensions should not be changed or altered.



Chilling and a second s



Tetrahedron journal homepage: www.elsevier.com

Stereoselective synthesis of dehydroamino acids using malonic acid half oxyester and aromatic aldehydes

Yuttapong Singjunla, Silvia Colombano, Jérôme Baudoux,* Jacques Rouden*

Laboratoire de Chimie Moléculaire et Thio-organique, ENSICAEN, Université de Normandie, CNRS, 6 boulevard du Maréchal Juin, 14050 Caen, France.

ARTICLE INFO

ABSTRACT

Article history: Received Received in revised form Accepted Available online

Keywords: Amino acid Doebner condensation Malonic half oxyester Organocatalysis Decarboxylation

1. Introduction

 α,β -Dehydroamino acids (Δ AAs) are a new class of nonproteinogenic amino acids occurring in recently characterized natural peptides.¹ Several bioactive products are composed of trisubstituted dehydroamino acids derived from valine (Δ Val), isoleucine (Δ Ile), leucine (Δ Leu) and phenylalanine (Δ Phe) such as Tentoxin.^{1,2} The incorporation of Δ AAs into a peptide could modify the backbone geometry of the chain by increasing its rigidity. Moreover, α,β dehydroamino acids may be involved in biological processes as biochemical intermediates to bring better stability in biological media.³ As a result, the development of new approaches for the synthesis of α,β -dehydroamino acid derivatives is essential to evaluate their biological reactivity and to facilitate the preparation of new synthetic peptides. In synthetic chemistry, dehydroamino acids have been used as starting materials for several reactions, mainly in asymmetric hydrogenations⁴ but also epoxidations,⁵ conjugated additions,⁶ cyclopropanations⁷ and Diels-Alder reactions.⁸

The most commonly used strategy for the one-pot synthesis of α,β -dehydroamino acids involves Wittig reaction⁹ and alternative methods.¹⁰ Ito et al. used the Schöllkopf method, isolating an oxazoline intermediate which was opened in the presence of Pd(0).¹¹ Alvarez-Ibarra et al. used a Schiff base in the presence of ethyl propynoate and potassium tert-butoxide to induce a [1.3]-hydride shift followed by a double bond migration to the α,β -position of the amino acid moiety.¹² Alternatively, Henry's reaction followed by dehydration was used to access α -nitroesters- α,β -unsaturated esters from nitroacetate.¹³ Other transformations reported for the synthesis of α,β -dehydroamino acids involve the elimination of β -hydroxy- α -amino esters is well described, easy access to these compounds is still challenging. Furthermore, the dehydration step sometimes

Corresponding authors. E-mail address: jerome.baudoux@ensicaen.fr (J. Baudoux), Jacques.rouden@ensicaen.fr (J. Rouden).

An efficient and direct approach was developed for the synthesis of α , β -dehydroamino acid derivatives with a broad range of substrates. Amido-substituted Malonic Acid Half Oxyesters (MAHOs) have proven to be excellent partners of various aromatic aldehydes in the presence of secondary amine, trifluoromethanesulfonic acid to perform a Knoevenagel-Doebner condensation under mild conditions with good to excellent yields. A mechanistic study revealed that the sequence involved the formation of an iminium intermediate to provide stereoselectively Z- α , β -unsaturated amino acids.

2016 Elsevier Ltd. All rights reserved.

requires restrictive conditions with the use of EDC/Cu(OTf)₂,¹⁵ TsCl/DCC/CuCl₂, EDC/CuCl₂,¹⁶ Ts₂O/DABCO¹⁷ or PPh₃/DEAD¹⁸ and very recently an interesting study has been realized by Ess and Castle to selectively generate Z or *E*- Δ Ile by using Martin sulfurane.¹⁹ In recent years, malonic acid half oxyesters (MAHOs) have been utilized with aliphatic aldehyde to perform decarboxylative aldol reactions followed by a subsequent dehydration step but very low overall yields of α, β -dehydroamino esters were observed in all cases.²⁰

In this context, the Knoevenagel-Doebner condensation²¹ represents a convenient and mild method to create a double bond from unsubstituted malonic acid or malonic acid half oxyesters (MAHOs) and aldehydes *via* a one-pot domino decarboxylative aldol-elimination process. The recent report by List and co-workers who developed a set of mild conditions for this transformation is particularly attractive, but only unsubstituted malonates were used (scheme 1a).^{21d} To the best of our knowledge, only one example reported by Xu and co-workers involves this methodology starting from a substituted 2-amido MAHO with an alkyl aldehyde to synthesize a ΔAA derivative in moderate yield and as a mixture of stereoisomers (Z-dehydroamino acid as the major).²²

With the aim of developing a short and flexible methodology to synthesize α,β -dehydroamino esters of various structural and electronic properties and with our experience in the use of 2-amido MAHO,²³ we report hereafter a straightforward synthesis of Δ AAs under mild conditions using the Doebner reaction (scheme 1b).

Tetrahedron



Scheme 1: Previous work and our strategy using MAOHs in the Doebner reaction.

2. Results and discussion

2

Our first attempt to transform 2-benzamido MAHO **1a** into a Δ AA derivative was carried out by using the efficient conditions developed by List and co-worker with unsubstituted malonates.^{21d} The reaction led quantitatively to decarboxylative aldol product **4** and no traces of Doebner compound **3a** was detected even after a prolonged reaction time (entry 1, table 1). Compared to unsubstituted malonates, amido-substituted MAHO appeared to behave quite differently and the Doebner reaction with these starting materials therefore represent an interesting synthetic challenge.

Reassessment of these conditions started using a secondary amine such as pyrrolidine to transform benzaldehyde 2 into a more reactive iminium intermediate as Xu recently developed with an alkyl aldehyde.²² Again, the reaction led to a mixture of three different compounds, the expected Z-dehydroamino ester 3a,²⁴ the decarboxylative-aldol ester 4 and the decarboxylative-protonated ester 5, all in very poor yields (entry 2, table 1). Based on previous works including ours, we envisaged a double activation of the substrates, basic for MAHO and nucleophilic for the aldehyde. We attempted the reaction by adding triethylamine to deprotonate MAHO but also a strong acid as co-catalyst with pyrrolidine to facilitate the iminium formation from benzaldehyde 2. Compared to alkyl aldehyde, a careful optimization had to be performed in order to determine the best catalyst combination working for all types of aromatic aldehydes.

Table 1

Optimization of the Knoevenagel-Doebner condensation



Tetrahedron 19% 53% MA8 D trace E 34% 19% 35% 10 А TfOH 4 Å MS 70% <2% 11% 11^{j} TfOH 4 Å MS^{e} 74% <2% <2% А $12^{d,g}$ A TfOH 4 Å MS ' 59% trace trace 13^{d} TfOH 4 Å MS e F 49% trace trace 14^h TfOH 4 Å MS ^e А 82% 8% 8%

^{*a*} Reactions performed on 0.2 mmol scale at 0.33 M **1a**. ^{*b*} Isolated yields. ^{*c*¹}H NMR conversion. ^{*d*} Without Et₃N. ^{*e*} Molecular sieves. ^{*f*} Performed at 10 °C. ^{*g*} Performed at 10 °C with 1 eq. of benzaldehyde, 1 eq. of TfOH and 2 eq. of pyrrolidine. ^{*h*} Performed at 10 °C with 2 eq. of benzaldehyde, pyrrolidine and TfOH.

Three strong acids, trifluoroacetic acid (TFA), trifluoromethane sulfonic acid (TfOH) and 4-nitrobenzoic acid, were tested in stoichiometric amounts (entries 3-5). TfOH initially provided the best improvement allowing the isolation of Doebner product **3a** in 61% yield along with **4** and **5**. With TfOH, the use of piperidine or morpholine, two amines with similar nucleophilicity to pyrrolidine,²⁵ afforded product **3a** in low yield (entries 6-7).

To enhance cooperation between the acid and the pyrrolidine in order to increase the concentration of iminium in the reaction mixture, we selected proline **D** and pyrrolidine-NHTf **E** as bifunctionnal catalysts. In the case of proline, only trace of **3a** was observed, probably due to the poor solubility of this catalyst in THF (entry 8). It is noteworthy that proline in ethanol as solvent afforded 8% yield of **3a** while the use of proline-Boc in THF with *in-situ* deprotection by TfOH gave 18% yield of **3a** (data not shown). Pyrrolidine *N*-triflimide **E** gave only a slight improvement (entry 9) compared to proline with 19% yield. In all cases when changing the nucleophilic catalyst, glycine was observed as a major product (product of protodecarboxylation).

To shift the equilibrium toward the formation of iminium, we added molecular sieves and a significant enhancement was observed to afford **3a** in 70% yield (entry 10). Performing the reaction at 10 °C, a 74% yield of Doebner product was achieved with very low amounts of side products **4** and **5** (entry 11). The selected combination of acid, tertiary and secondary amines to simultaneously activate MAHO **1a** and aldehyde **2** was confirmed during additional experiments. Without deprotonation of MAHO (no Et₃N) but with optimal iminium formation, the reaction lead to **3a** in only 59% yield (entry 12). Diamine **F** containing the base and the nucleophile on the same molecule afforded 49% yield (entry 13). Finally, the yield of **3a** was improved to 82% with a stoichiometric amount of Et₃N, two equivalents of pyrrolidine, TfOH and aldehyde (entry 14). All these experiments afforded the same stereoisomer and exclusively the *Z*-dehydroamino ester **3a**.



Scheme 2: Assessment of various protecting groups.

With these optimized conditions in hand, the effect on the reaction of various protecting groups at the nitrogen of MAHO was investigated (Scheme 2). *N*-acetyl **1b** afforded a similar yield of 85%. The reaction with *N*-(4-methylbenzoyl) **1c** and *N*-(4-methoxybenzoyl) **1d** gave slightly lower yields, **3c** in 74% and **3d** in 70% yield respectively. Benzyl carbamate *N*-Cbz **1e** generated Doebner product **3e** in 65% yield.

The scope of the reaction with various aromatic aldehydes was then evaluated (table 2). We started with 2-naphthyl-carboxaldehyde and the hindered 2-phenylbenzaldehyde. The corresponding dehydroamino esters **6** and **7** were obtained in 71% and 77% yields, respectively. With benzaldehydes bearing an electron donating group (4-MeO, 3,4-diMeO and 3-OMe), the reaction afforded products **8**, **9** and 10 in very good yields (up to 89%). In the presence of a hydroxyl group, 4-hydroxybenzaldehyde reacted slowly to provide the compound 11 in 42% yield. Even with 4-nitrobenzaldehyde (one equivalent) which is difficult to transform into an iminium intermediate, the desired compound 12 was isolated, albeit in moderate yield (38%). This low yield was due to a competitive decarboxylative aldol reaction.²⁶ Halogen-substituted benzaldehydes were successfully tested with MAHO 1a at 0 °C. At this temperature, Doebner condensation prevails to provide the compounds 13 (58%), 14 (50%) and 15 (72%). Furthermore, we extended our study to heterocyclic such furfural, aldehydes as 2and 3thiophenecarboxaldehyde, 2and 4-pyridinecarboxaldehyde. Electron-rich or -poor aromatic aldehydes reacted similarly in our conditions and the π -conjugated heterocyclic products 16-20 were obtained in very good yields between 75% and 85%.

Table 2

Synthesis of various substituted Z-dehydroamino esters



 a performed with 1 eq. of aldehyde, TfOH, and pyrrolidine. 26b Performed at 0 $^\circ C.$

Indole carboxaldehyde proposed a significant challenge. In our first attempts, indole-3-carboxaldehyde or 1-methylindole-3-carboxaldehyde were unreactive substrates (data not shown). However, with a Cbz group the reaction led to the product **21** in 59% NMR yield.²⁷ Finally the Doebner condensation with cinnamaldehyde afforded diene-amino ester **22** in 33% yield. Enolizable aldehydes such as crotonaldehyde or butyraldehyde did

not lead to the desired products, only glycine **5** was observed after 72 h of reaction (data not shown). However, the conditions developed by Xu and co-workers should provide the Doebner products with these aliphatic aldehydes.²²

Next, we investigated the mechanism operating under our conditions by performing a few additional experiments. Our initial strategy was based on a nucleophilic addition of an amine to aldehyde forming an iminium species. After the addition of the α -amido MAHO 1a, a spontaneous decarboxylative β -elimination reaction would occur from the intermediate β -ammonium carboxylate **B** to generate the double bond under mild conditions. Thus, the reaction begins simultaneously by deprotonation of MAHO with triethylamine (activated MAHO A) and by the iminium formation in presence of TfOH (scheme 3). The key step could be the addition of the secondary amine to the aldehyde in order to generate the highly electrophilic iminium species. To confirm this hypothesis, we tested Eschenmoser's salt with MAHO 1a and Et₃N, the reaction was complete in less than 20 minutes to give product 23 in a quantitative yield (scheme 4a). Similarly, the isolated iminium triflate 24^{23} reacted completely with 1a after only 2 h and compound 3a was obtained in an excellent yield without side products (Scheme 4b). These experiments prove a rapid addition of the activated MAHO 1a to the electrophilic iminium followed by the spontaneous decarboxylative β -elimination to afford the Doebner product. Therefore, the limiting step of this overall process is the slow condensation of the aldehyde with pyrrolidine to form the highly electrophilic iminium which was confirmed by the reaction of 1a and 4-nitrobenzaldehyde affording the expected dehydroamino esters 12 in a low yield (table 2).



Scheme 3: Plausible mechanism for the Doebner condensation.

Moreover, adding pyrrolidine and trifluoromethanesulfonic acid to β -hydroxy- α -amino ester 25 did not generate the dehydrated product 12 even after 72 h (scheme 4c). This result completes the previous experiments insofar as, during the Doebner condensation, dehydroamino esters are not formed by dehydration of putative hydroxyl amino ester intermediates such as 25 or 4.



Scheme 4: Additional experiments for mechanistic studies.

According to the literature, the reaction between unsubstituted hemimalonates and aldehydes in presence of pyridine (or DMAP) and Ac₂O gives a mixture of *E*- and *Z*-alkenes. With α -amido MAHOs, Xu et al. have observed a significant evolution of the ratio in favor of the *Z*-alkene (a ratio similar to Horner-Wadsworth-Emmons condensation) but the *E*-isomer is always present in small amounts and contaminates the crude reaction.²² Hengartner et al. were able to convert *E*-alkene into the more stable *Z*-isomer by treatment with NaOEt.²⁸ Under our conditions, the Doebner condensation of α -amido MAHOs gives exclusively the thermodynamically more stable *Z*-dehydroamino esters and the minor diastereomers were not observed during the process (¹H NMR monitoring).

3. Conclusion

In summary, we have developed an efficient and direct approach to prepare Z-dehydroamino acid derivatives according to the Knoevenagel-Doebner reaction. A variety of α -amido MAHOs and aromatic aldehydes in the presence of pyrrolidine, triethylamine and trifluoromethanesulfonic acid afforded various dehydroamino esters in moderate to excellent yields. The developed Doebner reaction using 2-amido MAHOs with aromatic aldehydes complements well the already known conditions. The methodology displays interesting features such as readily available starting materials and a simple experimental procedure which could be useful for large-scale applications. It is therefore a practical tool to access new scaffolds for making synthetic peptides.

4. Experimental section

4.1. General

All reagents were purchased from Acros Organics, Sigma Aldrich, Alfa Aesar, TCI or Fluka and were used without further purification and used as received. Solvents were used in RPE grade without further purification. Anhydrous solvents were obtained from a PURESOLV SPS400 apparatus developed by Innovative Technology Inc. ¹H and ¹³C NMR spectra were recorded on a Bruker DRX 400 MHz or a Bruker DRX 500 MHz spectrometer. Samples were dissolved in an appropriate deuterated solvent (CDCl₃, acetone-d6, D₂O). The chemical shifts (δ) are expressed in ppm relative to internal tetramethylsilane for ¹H and ¹³C nuclei, and coupling constants are indicated in Hz. Abbreviations for signal coupling are as follows: s = singlet; d = doublet; dd = doublet of doublets; t = triplet; q = quartet; quin = quintet; m = multiplet; br = broad signal. To assign the signals to the different proton and carbon atoms, as well as the relative stereochemistry of the cycloadducts, additional 2D NMR experiments (COSY, HSQC, HMBC) and NOESY

experiments were performed. Mass spectra were obtained on a GC/MS Saturn 2000 spectrometer. High-resolution mass spectra (HRMS) were performed on Q-TOF Micro WATERS by electrospray ionisation (ESI). Infrared (IR) spectra were recorded with a Perkin Elmer 16 PC FT-IR ATR spectrometer, using the pure product (oil or solid). Thin Layer Chromatography (TLC) was run on pre-coated aluminum plates of silica gel 60 F-254 (Merck). Flash chromatography was performed on silica gel column (Merck silica gel, 40-63 µm) using air pressure. Compounds $1b^{29}$, $1e^{30}$ were known in literature and synthesized as described below.

4.2. General procedure for the preparation of malonic half oxyesters (MAOHs)

Diethylmalonate (1 eq.) was diluted in EtOH. A KOH solution (1 eq.) in 10:1 EtOH/water was added dropwise at 0 °C. The mixture was stirred 15 h. A 1M HCl solution was added dropwise until pH = 1, then the mixture was saturated with NaCl and extracted twice with ethyl acetate. The organic layer was dried over MgSO₄ and solvents removed under reduced pressure. All malonic half oxyesters were easily detected by TLC (Rf = 0 in 60/40 cyclohexane/EtOAc)

4.3. General procedure for Doebner reaction

Procedure I: With aldehyde

Under inert atmosphere, triethylamine (0.2 mmol) was added to a solution of malonic acid half ester (0.2 mmol) in dry THF (0.6 ml) in the presence of molecular sieve (50 mg). At 0 $^{\circ}$ C, pyrrolidine (0.4 mmol) was added dropwise followed by trifluoromethanesulfonic acid (0.4 mmol) and aldehyde (0.4 mmol). The mixture was stirred at 10 $^{\circ}$ C for 24-120 h. The solution was concentrated under reduced pressure. The residue was directly purified by flash chromatography.

Procedure II: With iminium

Triethylamine (0.2 mmol) was added to a solution of malonic acid half ester 2 (0.2 mmol) in dry THF (0.6 ml). Iminium (0.4 mmol) was added to the solution and the reaction mixture was stirred at room temperature for 0.3-2 h. The solution was concentrated under reduced pressure. The residue was directly purified by flash chromatography.

4.3.1. 3-Ethoxy-2-benzamido-3-oxopropanoic acid (1a). White solid, mp 105-106 °C, 95% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.83 (2H, t, *J* = 7.2 Hz), 7.51-7.55 (1H, m), 7.39-7.45 (3H, m), 5.37 (d, *J* = 6.8 Hz, 1H), 4.30 (q, *J* = 7.2 Hz, 2H), 1.30 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 168.8, 168.3, 166.6, 132.8, 132.6, 129.0, 127.7, 63.5, 57.2, 14.2. IR (neat) 3299, 2926, 1748, 1709, 1639, 1520, 1248, 1197, 1172, 711, 691 cm⁻¹. HRMS *m*/*z* (ESI): calcd. for C₁₂H₁₃NO₅Na [M+Na]⁺: 274.0691, found: 274.0703.

4.3.2. 3-*Ethoxy*-2-(4-*methylbenzamido*)-3-oxopropanoic acid (**1c**). White solid, mp 114-115 °C, 61% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.81 (d, *J* = 8.8 Hz, 2H), 7.26 (b, 1 H), 6.93 (d, *J* = 8.8 Hz, 2 H), 5.81 (b, 1H), 5.33 (d, *J* = 6.4 Hz, 1H), 3.04 (s, 3H), 4.32 (q, *J* = 7.2 Hz, 2H), 1.31 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 168.4, 167.5, 166.6, 162.9, 129.4, 124.5, 113.9, 63.1, 57.0, 55.5, 13.9. IR (neat) 3397, 2982, 1738, 1605, 1497, 1257, 1183, 1024, 767 cm⁻¹. HRMS *m*/z (ESI) calcd for C₁₃H₁₄NO₆ [M-H]: 280.0821, found: 280.0814.

4.3.3. 3-Ethoxy-2-(4-methoxybenzamido)-3-oxopropanoic acid (**1d**). White solid, mp 109-112 °C. 52% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.74 (d, *J* = 8.2 Hz, 2H), 7.26 (d, *J* = 8.2 Hz, 2 H), 6.71 (b, 1H), 5.35 (d, *J* = 6.4 Hz, 1H), 4.33 (q, *J* = 7.2 Hz, 2H), 1.33 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 168.4, 168.0, 166.3, 143.2, 129.5, 129.4, 127.4, 63.1, 56.9, 21.5, 13.9. IR (neat) 3305, 3131, 1749, 1726, 1634, 1523, 1499, 1238, 1185, 1164, 759, 669 cm⁻¹. HRMS *m*/*z* (ESI) calcd for C₁₃H₁₆NO₅ [M+H]⁺: 266.1028, found: 266.1035.

4.3.4. (*Z*)-*Ethyl* 2-*benzamido-3-phenylacrylate* (**3***a*). White solid, mp 128-130 °C, 48.6 mg, 82% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.87 (d, *J* = 8.2 Hz, 2H), 7.77 (b, 1H), 7.56 (t, *J* = 8.4 Hz, 1H), 7.47-7.51 (m, 5H), 7.30-7.36 (m, 3H), 4.33 (q, *J* = 7.2 Hz, 2H), 1.36 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.9, 165.7, 134.2, 133.9, 132.4, 131.6, 129.9, 129.6, 129.0, 128.8, 127.7, 124.7, 62.2, 14.5. IR (neat) 3262, 2924, 1709, 1644, 1635, 1518, 1486, 1239, 1141, 1090, 712 cm⁻¹. HRMS *m*/z (ESI) calcd. for C₁₈H₁₈NO₃ [M+H]⁺: 296.1287; found 296.1274.

4.3.5. (*Z*)-*Ethyl* 2-acetamido-3-phenylacrylate (**3b**). Solid, mp 92-93 °C, 39.5 mg, 85% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.46 (d, *J* = 6.8 Hz, 2H), 7.35-7.38 (m, 4H), 6.97 (b, 1H), 4.31 (q, *J* = 7.2 Hz, 2H), 2.14 (s, 3H), 1.35 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 169.1, 165.6, 134.1, 132.1, 129.9, 129.6, 128.8, 124.8, 62.1, 23.7, 14.5. IR (neat) 3242, 2983, 1716, 1665, 1642, 1513, 1490, 1371, 1251, 1203, 691 cm⁻¹. HRMS *m*/z (ESI) calcd. for C₁₃H₁₅NO₃Na [M+Na]⁺: 256.0950; found 256.0943.

4.3.6. (*Z*)-*Ethyl* 2-(4-*methylbenzamido*)-3-*phenylacrylate* (3*c*). Solid, mp 120-123 °C, 45.8 mg, 74% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.92 (s, 1H), 7.72 (d, *J* = 8.0 Hz, 2H), 7.46 (d, *J* = 7.8 Hz, 2H), 7.39 (s, 1H), 7.23-7.30 (m, 3H), 7.19 (d, *J* = 8.0 Hz, 2H), 4.24 (q, *J* = 7.2 Hz, 2H), 2.36 (s, 3H), 1.29 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.5 (2C), 142.8, 134.1, 130.9, 130.8, 129.6, 129.4, 129.3, 128.6, 127.5, 124.5, 61.9, 21.5, 14.3. IR (neat) 3304, 2981,1724, 1636, 1525, 1497, 1490, 1250, 1189, 752, 667 cm⁻¹. HRMS *m/z* (ESI) calcd. for C₁₉H₂₀NO₃ [M+H]⁺: 310.1443; found 310.1451.

4.3.7. (*Z*)-*Ethyl* 2-(4-methoxybenzamido)-3-phenylacrylate (**3d**). Solid, mp 117-119 °C, 45.5 mg, 70% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.83 (d, *J* = 8.8 Hz, 2H), 7.69 (s, 1H), 7.49 (d, *J* = 8.0 Hz, 2H), 7.43 (s, 1H), 7.29-7.34 (m, 3H), 6.95 (d, *J* = 8.8 Hz, 2H), 4.32 (q, *J* = 7.2 Hz, 2H), 3.86 (s, 3H), 1.35 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.5, 165.1, 162.8, 134.1, 130.6, 129.6, 129.4, 129.3, 128.6, 126.0, 124.6, 113.9, 61.9, 55.5, 14.3. IR (neat) 3286, 2972, 1709, 1643, 1603, 1479, 1246, 1175, 1026, 761, 690 cm⁻¹. HRMS *m*/z (ESI) calcd. for C₁₉H₁₉NO₄Na [M+Na]⁺: 348.1212; found 348.1214.

4.3.8. (*Z*)-*Ethyl* 2-(((*benzyloxy*)*carbonyl*)*amino*)-3-*phenylacrylate* (*3e*). Colorless oil, 42.3 mg, 65% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.43-7.53 (m, 2H), 7.25-7.27 (m, 8H), 6.29 (b, 1H), 5.04 (s, 2H), 4.21 (q, *J* = 7.2 Hz, 2H), 3.86 (s, 3H), 1.25 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.3, 153.9, 136.0, 133.8, 136.0, 133.8, 131.4, 129.8, 129.4, 128.6, 128.5, 128.3, 128.2, 124.5, 67.5, 61.8, 14.2. IR (neat) 3296, 2977, 1699, 1498, 1312, 1254, 1221, 1201, 1039, 1024, 734, 690 cm⁻¹. HRMS *m/z* (ESI) calcd. for C₁₉H₁₉NO₄Na [M+Na]⁺: 348.1212; found 348.120.

4.3.9. (Z)-Ethyl 2-benzamido-3-(naphthalen-2-yl)acrylate (6). White solid, mp 99-102 °C, 49.1 mg, 71% yield. ¹H NMR (400 MHz, CDCl₃) δ 8.03 (d, J = 8.2 Hz, 1H), 7.78-7.89 (m, 3H), 7.68-7.71 (m, 3H), 7.46-7.61 (m, 4H), 7.36-7.43 (m, 3H), 4.39 (q, J = 7.2 Hz, 2H), 1.39 (t, J = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.6, 165.0, 133.0, 133.5, 132.1, 131.2, 131.1, 129.5, 128.8, 128.7, 127.5, 127.4, 127.3, 126.7, 126.3, 126.2, 125.3, 124.3, 62.0, 14.3. IR (neat) 3229, 2977, 1636, 1509, 1481, 1292, 1241, 768, 689 cm⁻¹. HRMS *m*/z (ESI) calcd. for C₂₂H₂₀NO₃ [M+H]⁺: 346.1443; found: 346.1437.

4.3.10. (*Z*)-*Ethyl* 3-([1,1'-*biphenyl*]-2-*yl*)-2-*benzamidoacrylate* (7). White solid, mp 138-140 °C, 57.2 mg, 77% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.58-7.63 (m, 3H), 7.44 (d, *J* = 7.7 Hz, 1H), 7.33-7.37 (m, 3H), 7.06-7.28 (m, 9H), 4.07 (q, *J* = 7.2 Hz, 2H), 1.07 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.3, 165.2, 141.9, 140.4, 133.9, 132.7, 132.0, 130.6, 130.2, 129.9, 129.0, 128.7, 128.3, 128.1, 128.0, 127.5, 127.4, 127.1, 124.8, 61.7, 14.2. IR (neat) 3281, 2967, 1709, 1651, 1509, 1476, 1266, 1244, 730, 705 cm⁻¹. HRMS *m*/z (ESI) calcd. for C₂₄H₂₂NO₃ [M+H]⁺: 372.1600; found: 372.1594.

4.3.11 S(Z)-Ethyl 2-benzamido-3-(4-methoxyphenyl)acrylate (8). Solid, mp 136-137 °C, 58.0 mg, 89% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.89 (d, J = 7.4 Hz, 2H), 7.77 (s, 1H), 7.54-7.58 (m, 1H), 7.47-7.50 (m, 5H), 6.85 (d, J = 8.8 Hz, 2H), 4.30 (q, J = 7.2 Hz, 2H), 3.79 (s, 3H), 1.34 (t, J = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.7, 165.6, 160.5, 133.9, 132.2, 131.8, 131.7, 128.8, 127.5, 126.5, 121.9, 114.1, 61.7, 55.3, 14.3. IR (neat) 3289, 2927, 1713, 1649, 1603, 1509, 1477, 1244, 1173, 728, 706 cm⁻¹. HRMS m/z (ESI) calcd. for C₁₉H₂₀NO₄ [M+H]⁺: 326.1392; found 326.1386.

4.3.12. (Z)-Ethyl 2-benzamido-3-(3,4-dimethoxyphenyl)acrylate (9). White solid, mp 112-114 °C, 50.5 mg, 71% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.90 (d, J = 7.4 Hz, 2H), 7.74 (t, J = 7.2 Hz, 1H), 7.46-7.50 (m, 3H), 7.10-7.14 (m, 2H), 6.83 (d, J = 8.2 Hz, 2H), 4.32 (q, J = 7.2 Hz, 2H), 3.87 (s, 1H), 3.68 (s, 1H), 1.36 (t, J = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.6 (2C), 150.2, 148.6, 133.8, 132.2, 132.1, 128.8, 127.4, 127.0, 126.7, 124.1, 112.2, 110.8, 61.8, 55.8, 55.6, 14.3. IR (neat) 3309, 2980, 1713, 1649, 1513, 1480, 1246, 1140, 1023, 721 cm⁻¹. HRMS m/z (ESI) calcd. for C₂₀H₂₁NO₅Na [M+Na]⁺: 378.1317; found: 378.1318.

4.3.13. (*Z*)-*Ethyl* 2-*benzamido*-3-(3-*methoxyphenyl*)*acrylate* (10). White solid, mp 82-84 °C, 47.5 mg, 73% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.69 (d, *J* = 7.4 Hz, 2H), 7.59 (s, 1H), 7.37-7.40 (m, 1H), 7.28-7.32 (m, 2H), 7.05-7.09 (m, 2H), 6.92 (d, *J* = 7.7 Hz, 1H), 6.87 (s, 1H), 6.69 (dd, *J* = 2.2, 8.2 Hz, 1H), 4.16 (q, *J* = 7.2 Hz, 2H), 3.53 (s, 3H), 1.19 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.6, 165.3, 135.2, 133.6, 132.2, 131.1, 129.6, 128.8, 127.4, 124.7, 122.2, 115.5, 114.4, 61.9, 55.1, 14.3. IR (neat) 3279, 2982, 1716, 1651, 1511, 1478, 1295, 1271, 1236, 709, 689 cm⁻¹. HRMS *m*/z (ESI) calcd, for C₁₉H₁₉NO₄Na [M+Na]⁺: 348.1212; found 348.1219.

4.3.14. (Z)-Ethyl 2-benzamido-3-(4-hydroxyphenyl)acrylate (11). White solid, mp 100-103 °C, 26.2 mg, 42% yield. ¹H NMR (400 MHz, CD₃OD) δ 7.97 (d, J = 7.4 Hz, 2H), 7.53-7.56 (m, 6H), 7.79 (d, J = 8.6 Hz, 2H), 4.28 (q, J = 7.2 Hz, 2H), 1.32 (t, J = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.5, 165.7, 159.2, 135.7, 133.6, 131.8, 128.3, 127.3, 124.7, 122.4, 115.2, 61.8, 13.2. IR (neat) 3264, 2928, 1695, 1644, 1602, 1508, 1478, 1202, 1251, 1165, 708 cm⁻¹. HRMS m/z (ESI) calcd. for C₁₈H₁₈NO₄ [M+H]⁺: 312.1236; found: 312.1225.

4.3.15. (*Z*)-*Ethyl* 2-benzamido-3-(4-nitrophenyl)acrylate (**12**). White solid, mp 165-166 °C, 25.9 mg, 38% yield. ¹H NMR (400 MHz, CDCl₃) δ 8.21 (br, 1H), 8.16 (d, *J* = 8.0 Hz, 2H), 7.84 (d, *J* = 8.0 Hz, 2H), 7.56-7.59 (m, 3H), 7.47-7.51 (m, 3H), 4.38 (q, *J* = 8.0 Hz, 2H), 1.40 (t, *J* = 8.0 Hz, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 165.1, 164.8, 147.3, 141.6, 133.4, 132.8, 130.0, 129.1, 127.6, 126.4, 126.1, 123.7, 62.8, 14.4. IR (neat) 3273, 2984, 1719, 1660, 1517, 1478, 1343, 1305, 1253 cm⁻¹. HRMS *m*/*z* (ESI) calcd. for C₁₈H₁₆N₂O₅Na [M+Na]⁺: 363.0957; found: 363.0947.

4.3.16. (*Z*)-*Ethyl* 2-benzamido-3-(3-fluorophenyl)acrylate (**13**). The experiment was performed at 0 °C.White solid, mp 135-136 °C, 36.4 mg, 58% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.86 (d, *J* = 7.6 Hz, 3H), 7.57 (t, *J* = 7.6 Hz, 1H), 7.48 (t, *J* = 7.6 Hz, 2H), 7.43 (s, 1H), 7.27-7.32 (m, 2H), 7.19 (d, *J* = 9.9 Hz, 1H), 6.98 (t, *J* = 7.6 Hz, 1H), 4.34 (q, *J* = 7.2 Hz, 2H), 1.36 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (125 MHz, CDCl₃) δ 165.4, 165.2, 162.2 (d, *J* = 246.4 Hz), 136.3 (d, *J* = 8.2 Hz), 133.5, 132.3, 130.0 (d, *J* = 8.3 Hz), 129.4, 128.8, 127.4, 125.4 (d, *J* = 2.8 Hz), 124.9, 116.2 (d, *J* = 21.3 Hz), 116.1 (d, *J* = 22.4 Hz), 62.2, 14.2. ¹⁹F NMR (376 MHz, CDCl₃) δ -112.7. IR (neat) 3272, 2985, 1719, 1652, 1509, 1479, 1291, 1230, 1153, 709, 686 cm⁻¹. HRMS *m*/z (ESI) calcd. for C₁₈H₁₆NO₃FNa [M+Na]⁺: 336.1012; found: 336.1004.

4.3.17. (Z)-Ethyl 2-benzamido-3-(4-fluorophenyl)acrylate (14). The experiment was performed at 0 °C. White solid, mp 120-121 °C, 31.4 mg, 50% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.62-6.66 (m, 3H), 7.36 (t, *J* = 7.4 Hz, 1H), 7.27-7.29 (m, 4H), 7.04 (s, 1H), 6.80 (t,

J = 8.6 Hz, 2H), 4.12 (q, *J* = 7.2 Hz, 2H), 1.15 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (125 MHz, CDCl₃) δ 165.4 (2C), 162.6 (d, *J* = 250.0 Hz), 133.6, 132.3, 131.7 (d, *J* = 8.4 Hz), 130.3 (d, *J* = 3.3 Hz), 130.2, 128.8, 127.5, 123.6, 115.6 (d, *J* = 21.8 Hz), 62.0, 14.3. ¹⁹F NMR (376 MHz, CDCl₃) δ -110.2. IR (neat) 3273, 2986, 1717, 1652, 1601, 1507, 1479, 1251, 1230, 708 cm⁻¹. HRMS *m/z* (ESI) calcd. for C₁₈H₁₆NO₃FNa [M+Na]⁺: 336.1012; found: 336.1013.

4.3.18. (*Z*)-*Ethyl* 2-benzamido-3-(2-bromophenyl)acrylate (**15**). The experiment was performed at 0 °C. White solid, mp 106-107 °C, 53.9 mg, 72% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.78 (d, *J* = 7.1 Hz, 3H), 7.62 (dd, *J* = 1.1, 8.0 Hz, 1H), 7.51-7.55 (m, 2H), 7.42-7.47 (m, 3H), 7.21 (td, *J* = 0.8, 7.6 Hz, 1H), 7.14 (td, *J* = 1.6, 7.6 Hz, 1H), 4.36 (q, *J* = 7.2 Hz, 2H), 1.38 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.1, 165.0, 135.1, 133.6, 132.9, 132.2, 130.1, 129.3, 128.8, 128.5, 127.4, 127.2, 126.0, 124.4, 62.2, 14.2. IR (neat) 3286, 2977, 1714, 1648, 1509, 1479, 1287, 1249, 1024, 705 cm⁻¹. HRMS *m*/*z* (ESI) calcd. for C₁₈H₁₆NO₃BrNa [M+Na]⁺: 396.0211; found: 396.0208.

4.3.19. (Z)-Ethyl 2-benzamido-3-(furan-2-yl)acrylate (16). White solid, mp 124-125 °C, 45.1 mg, 79% yield. ¹H NMR (400 MHz, CDCl₃) δ 8.23 (s, 1H), 7.93 (d, J = 6.8 Hz, 2H), 7.56-7.59 (m, 1H), 7.48-7.52 (m, 3H), 7.08 (s, 1H), 6.57 (d, J = 3.4 Hz, 1H), 6.47 (dd, J = 1.8, 3.4 Hz, 1H), 4.31 (q, J = 7.2 Hz, 2H), 1.33 (t, J = 7.2 Hz, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 165.6, 164.9, 150.1, 144.4, 133.7, 132.2, 128.8, 127.6, 123.2, 115.9, 115.3, 112.3, 61.7, 14.2. IR (neat) 3292, 2982, 1715, 1650, 1509, 1480, 1286, 1259, 1210, 1182, 1022, 716 cm⁻¹. HRMS *m*/z (ESI) calcd. for C₁₆H₁₅NO₄Na [M+Na]⁺: 308.0899; found: 308.0907.

4.3.20. (*Z*)-*Ethyl* 2-*benzamido-3*-(*thiophen-2-yl*)*acrylate* (**17**). White solid, mp 170-173 °C, 45.2 mg, 75% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.96 (d, *J* = 7.2 Hz, 2H), 7.87 (s, 1H), 7.42-7.60 (m, 5H), 7.35 (d, *J* = 3.4 Hz, 1H), 7.07 (dd, *J* = 3.4, 4.9 Hz, 1H), 4.30 (q, *J* = 7.2 Hz, 2H), 1.33 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 166.7, 164.9, 136.5, 133.9, 133.2, 132.2, 130.7, 128.8, 128.7, 127.6, 127.2, 121.6, 61.7, 14.3. IR (neat) 3281, 2972, 1714, 1648, 1628, 1512, 1471, 1261, 1198, 705 cm⁻¹. HRMS *m*/z (ESI) calcd. for C₁₆H₁₅NO₃SNa [M+Na]⁺: 324.0670; found: 324.0679.

4.3.21. (Z)-Ethyl 2-benzamido-3-(thiophen-3-yl)acrylate (18). White solid, mp 177-179 °C, 51.3 mg, 85% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.90 (d, J = 7.4 Hz, 2H), 7.68 (s, 1H), 7.46-7.57 (m, 5H), 7.26-7.28 (m, 2H), 4.29 (q, J = 7.2 Hz, 2H), 1.33 (t, J = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.9, 165.5, 135.1, 133.7, 132.4, 129.3, 128.8, 127.9, 127.5, 126.9, 126.1, 122.6, 61.8, 14.3. IR (neat) 3260, 2987, 1699, 1630, 1514, 1476, 1256, 1203, 1140, 1087, 705, 690 cm⁻¹. HRMS m/z (ESI) calcd. for C₁₆H₁₅NO₃SNa [M+Na]⁺: 324.0670; found: 324.0685.

4.3.22. (*Z*)-*Ethyl* 2-*benzamido*-3-(*pyridin*-4-*yl*)*acrylate* (**19**). White solid, mp 117-120 °C, 45.0 mg, 76% yield. ¹H NMR (400 MHz, CDCl₃) δ 8.57 (d, J = 3.7 Hz, 2H), 8.17 (s, 1H), 7.83-7.85 (m, 2H), 6.57 (d, J = 3.4 Hz, 1H), 7.59 (t, J = 7.4 Hz, 1H), 7.49 (t, J = 7.4 Hz, 2H), 7.32-7.34 (m, 3H), 4.37 (q, J = 7.2 Hz, 2H), 1.39 (t, J = 7.2 Hz, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 164.7, 164.6, 148.8, 143.6, 133.2, 132.7, 128.9, 127.5, 127.2, 125.5, 123.4, 62.7, 14.2. IR (neat) 3265, 2967, 1721, 1633, 1514, 1476, 1287, 1249, 1016, 710, 685 cm⁻¹. HRMS *m*/*z* (ESI) calcd. for C₁₇H₁₇N₂O₃ [M+H]⁺: 297.1239; found: 297.1245.

4.3.23. (Z)-Ethyl 2-benzamido-3-(pyridin-2-yl)acrylate (20). Colorless oil, 44.5 mg, 75% yield. ¹H NMR (400 MHz, CDCl₃) δ 12.9 (s, 1H), 8.05 (d, J = 4.6 Hz, 1H), 7.71 (td, J = 1.8, 7.8 Hz, 1H), 7.50-7.60 (m, 3H), 7.26 (d, J = 7.8 Hz, 1H), 7.18 (dd, J = 4.6, 7.8 Hz, 1H), 6.34 (s, 1H) 4.40 (q, J = 7.2 Hz, 2H), 1.38 (t, J = 7.2 Hz, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 165.4, 165.1, 155.3, 148.2, 137.2, 135.0, 133.4, 132.3, 128.7, 127.9, 125.4, 121.9, 113.3, 61.8, 14.1. IR (neat) 3063, 1724, 1673, 1635, 1469, 1421, 1309, 1279,

 $J = 8.6 \text{ Hz}, 2\text{H}, 4.12 \text{ (q}, J = 7.2 \text{ Hz}, 2\text{H}, 1.15 \text{ (t}, J = 7.2 \text{ Hz}, 3\text{H}). [A254, J 1208, [1145, 698 \text{ cm}^{-1}]. \text{ HRMS } m/z \text{ (ESI) calcd. for C NMR (125 MHz, CDCl_3) } \delta 165.4 \text{ (2C)}, 162.6 \text{ (d}, J = 250.0 \text{ Hz}), C_{17}H_{16}N_2O_3Na [M+Na]^+: 319.1059; \text{ found: } 319.1073.$

4.3.24. (Z)-Benzyl 3-(2-benzamido-3-ethoxy-3-oxoprop-1-en-1-yl)-1H-indole-1-carboxylate (21). Yellow solid, mp 122-124 °C, 54.2 mg, 59% NMR yield. ¹H NMR (400 MHz, CDCl₃) δ 8.15 (d, J = 7.5 Hz, 1H), 7.68-7.83 (m, 6H), 7.53 (t, J = 7.3 Hz, 1H), 7.41 (t, J = 7.5 Hz, 2H), 7.22-7.34 (m, 7H), 5.31 (s, 2H), 4.31 (q, J = 7.2 Hz, 2H), 1.34 (t, J = 7.2 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.2 (2C), 150.3, 134.9, 134.5, 133.8, 132.1, 129.5, 128.8, 128.74, 128.72, 128.3, 127.4, 127.3, 125.4, 123.6, 122.3, 119.0, 115.3, 115.1, 69.0, 61.9, 14.3. IR (neat) 3255, 2977, 1742, 1711, 1666, 1481, 1453, 1339, 1233, 1216, 1082, 753, 698 cm⁻¹. HRMS *m*/z (ESI) calcd. for C₂₈H₂₄N₂O₅Na [M+Na]⁺: 491.1583; found; 491.1591.

4.3.25. (2Z,4*E*)-*Ethyl* 2-*benzamido*-5-*phenylpenta*-2,4-*dienoate* (22). Yellow solid, mp 95-96 °C, 21.2 mg, 33% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.90 (d, *J* = 7.2 Hz, 2H), 7.82 (br s, 1H), 7.54 (t, *J* = 7.2 Hz, 1H), 7.42-7.53 (m, 4H), 7.23-7.29 (m, 4H), 6.87-7.01 (m, 2H), 4.27 (q, *J* = 7.2 Hz, 2H), 1.32 (t, *J* = 7.2 Hz, 3H).¹³C NMR (101 MHz, CDCl₃) δ 165.9, 165.4, 140.1, 136.7, 134.3, 132.3, 131.7, 129.1, 129.0, 128.9, 127.7, 127.6, 124.5, 123.3, 62.0, 14.5 IR (neat) 3288, 2924, 1713, 1647, 1508, 1478, 1276, 1230, 967, 705 cm⁻¹. HRMS *m*/*z* (ESI) calcd. for C₂₀H₂₀NO₃ [M+H]⁺: 322.1443; found: 322.1440.

4.3.26. *Ethyl 2-benzamidoacrylate* (**23**). Colorless oil, 43.0 mg, 98% yield. ¹H NMR (400 MHz, CDCl₃) δ 8.55 (br s, 1H), 7.83 (d, *J* = 8.0 Hz 2H), 7.45-7.56 (m, 3H), 6.78 (s, 1H), 5.99 (s, 1H), 4.33 (q, *J* = 8.0 Hz, 2H), 1.37 (t, *J* = 8.0 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 166.1, 164.6, 134.6, 132.3, 131.5, 129.1, 127.2, 108.9, 62.7, 14.4. IR (neat) 3355, 2983, 1730, 1651, 1604, 1244, 1095, 1024, 710, 691 cm⁻¹. HRMS *m*/*z* (ESI) calcd. for C₁₂H₁₄NO₃ [M+H]⁺: 220.0974; found: 220.0974.

Acknowledgements

The Ministery of Higher Education and Research, the "Region Basse Normandie" (fellowships to Y.S.), CNRS, and the European Union (FEDER) are gratefully acknowledged for funding this work. We thank the LABEX SynOrg (ANR-11-LABX-0029) for financial support.

Supplementary data

Supplementary data associated with this article can be found in the online version, at

References and notes

- (a) Jiang, J.; Ma, Z.; Castle, S. L. *Tetrahedron.* 2015, 71, 5431-5451; (b) Bonauer, C.; Walenzyk, T.; König, B. *Synthesis* 2006, 1-20
- Jiménez, J. C.; Chavarria, B.; Lopez-Macia, A.; Royo, M.; Giralt, E.; Albericio, F. Org. Lett. 2003, 5, 2115-2118.
- 3. Thibodeaux, G. N.; van der Donk, W. A. Chem. Commun. 2012, 48, 10615-10617.
- (a) Noyori, R. Asymmetric Catalysis in Organic Synthesis; Wiley Interscience: New York, **1994**; 16-28; (b) MacNeil, P. A.; Roberts, N. K.; Bosnich, B. J. Am. Chem. Soc. **1981**, 103, 2273-2280.
- (a) Bartels, A.; Jones, P. G.; Liebscher, J. *Tetrahedron Lett.* 1995, *36*, 3673-3674; (b) Bartels, A.; Jones, P. G.; Liebscher, J. *Synthesis* 2003, 67-72.
- (a) Ferreira, P. M. T.; Maia, H. L. S.; Monteiro, L. S.; Sacramento, J.; Sebastiao, J. J. Chem. Soc., Perkin Trans. 1 2000, 3317-3324; (b) Ferreira, P. M. T.; Maia, H. L. S.; Monteiro, L. S.; Sacramento, J. Tetrahedron Lett. 2000, 41, 7437-7441; (c) Javidan, A.; Schafer, K.; Pyne, S. G. Synlett. 1997, 100-102; (d) Pyne, S. G.; Javidan, A.; Skeleton, B. W.;

White, A. H. *Tetrahedron* **1995**, *51*, 5157-5168; (e) Bull, S. D.; M.A. Davies, S. G.; O'Shea, M. D. J. Chem. Soc., Perkin Trans. 1 **1998**, 3657-3658; (f) Jin, S. D.; Wessig, P.; Liebscher, J. *Eur. J. 2 Org. Chem.* **2000**, *10*, 1993-1999.

- (a) Adams, L. A.; Aggarwal, V. K.; Bonnert, R. V.; Bressel, B.; Cox, R. J.; Shepherd, J.; de Vicente, J.; Walter, M.; Whittingham, W. G.; Winn, C. L. J. Org. Chem. 2003, 68, 9433-9440; (b) Adams, L. A.; Charmant, J. P. H.; Cox, R. J.; Walter, M.; Whittingham, W. G. Org. Biomol. Chem. 2004, 2, 542-553; (c) Aggarwal, V. K.; Alonso, E.; Fang, G. Y.; Ferrar, M.; Hynd,G.; Porcelloni, M. Angew. Chem. Int. Ed. 2001, 40, 1433-1436.
- (a) Horikawa, H.; Nishitani, T.; Iwasaki, T.; Mushika, Y.; Inoue, I.; Miyoshi, M. *Tetrahedron Lett.* **1980**, *21*, 4101-4104; (b) Avenoza, A.; Cativiela, C.; Fernandez-Rico, M. A.; Peregrina, J. M. *J. Chem. Soc., Perkin Trans. 1* **1999**, 3375-3379; (c) Crossley, M. J.; Stamford, A. W. *Aust. J. Chem.* **1993**, *46*, 1443-1446.
- Mazurkiewicz, R.; Kuznik, A.; Grymel, M.; Kuznik, N. Monatsh. Chem. 2004, 135, 807-815.
- (a) Schmidt, U.; Lieberknecht, A.W.J. Synthesis 1988, 159-172;
 (b) Schmidt, U.; Griesser, H.; Leitenberger, V.; Lieberknecht, A.; Mangold, Meyer, R.; Riedl, B. Synthesis 1992, 487-490; (c) Schmidt U.; Riedl, B. Chem. Commun. 1992, 1186-1187; (d). Schmidt, U. Riedl, B. Synthesis 1993, 809-814; (e) Schmidt, U.; Riedl, B. Synthesis 1993, 815-818; (f) Yasuno, Y.; Hamada, M.; Yamada, T.; Shinada T.; Ohfune, Y. Eur. J. Org. Chem. 2013, 1884-1888; (g) Hamada, M.; Yamada, T.; Shinada T.; Ohfune, Y. Org. Lett. 2009, 11, 4664-4667.
- Ito, Y.; Matsuura, T.; Saegusa, T. Tetrahedron Lett. 1985, 26, 5781-5784.
- Alvarez-Ibarra, C.; Csaky, A. G.; Ortega, E. M.; De la Morena, M. J.; Quiroga, M. L. *Tetrahedron Lett.* **1997**, *38*, 4501-4502.
- (a) Lehnert, W. *Tetrahedron* **1972**, 28, 663-666; (b) Fornicola, R.
 S.; Oblinger, E.; Montgomery, J. J. Org. Chem. **1998**, 63, 3528-3529.
- 14. (a) Ross, A. C.; Liu, H.; Pattabiraman, V. R.; Vederas, J. C. J. Am. Chem. Soc. 2010, 132, 462-463; (b) Ferreira, P. M. T.; Maia, H. L. S.; Monteiro, L. S. Tetrahedron Lett. 1998, 39, 9575-9578; (c) Goodall, K.; Parsons, A. F. Tetrahedron Lett. 1995, 36, 3259-3260; (d) Li, K. W.; Wu, J.; Xing, W.; Simon, J. A. J. Am. Chem. Soc. 1996, 118, 7237-7238; (e) Cherney, R. J.; Wang, L. J. Org. Chem. 1996, 61, 2544-2546; (f) Somekh, L.; Shanzer, A. J. Org. Chem. 1983, 48, 907-908; (g) Sai, H.; Ogiku, T.; Ohmizu, H. Synthesis 2003, 201-204; (h) Sai, H.; Ohmizu, H. Tetrahedron Lett. 1999, 40, 5019-5022; (i) Bonauer, C.; Walenzyk, T.; König, B. Synthesis 2006, 1-20.
- Shangguan, N.; Joullié, M. Tetrahedron Lett. 2009, 50, 6748-6750.
- (a) Sai, H.; Ogiku, T.; Ohmizu, H. Synthesis 2003, 201-204. (b) Sai, H.; Ohmizu, H. Tetrahedron Lett. 1999, 40, 5019-5022.
- 17. Li, K. W.; Wu, J.; Xing, W.; Simon, J. A. J. Am. Chem. Soc. 1996, 118, 7237-7238.
- 18. Cherney, R. J.; Wang, L. J. Org. Chem. 1996, 61, 2544-2546.
- (a) Ma, Z.; Naylor, B. C.; Loertscher, B. M.; Hafen, D. D.; Li, J. M.; Castle, S. L. J. Org. Chem. 2012, 77, 1208-1214. (b) Ma, Z.; Jiang, J.; Luo, S.; Cai, Y.; Cardon, J. M.; Kay, B. M.; Ess, D. H.; Castle, S. L. Org. Lett. 2014, 16, 4044-4047.
- (a) Faucher, A- M.; Bailey, M. D.; Beaulieu, P. L.; Brochu, C.; Duceppe, J- S.; Ferland, J- M.; Ghiro, E.; Gorys, V.; Halmos, T.; Kawai, S. H.; Poirier, M.; Simoneau, B.; Tsantrizos, Y. S.; Llinas-Brunet, M. Org. Lett. 2004, 6, 2901-2904; (b) Velazquez, F.; Venkatraman, S.; Wu, W.; Blackman, M.; Prongay, A.; Girijavallabhan, V.; Shih, N.-Y.; Njoroge, F.G. Org. Lett. 2007, 9, 3061-3064.
- (a) Doebner, O. Ber. Dtsch. Chem. Ges. 1900, 33, 2141-2142; (b) Doebner, O. Ber. Dtsch. Chem. Ges. 1902, 35, 3311 -3347; (c) Rodriguez, J.; Waegell, B. Synthesis 1988, 534–535; (d) List, B.; Doehring, A.; Hechaverria Fonseca, M. T.; Job, A.; Rios Torres,

- A R. Tetrahedron 2006, 62, 476–482; (e) Heravi M. M.; Asadi S.;
- Azarakhshi F. Curr. Org. Chem. 2014, 11, 701-731.
- Xu, F.; Zacuto, M.; Yoshikawa, N.; Desmond, R.; Hoerrner, S.; Itoh, T.; Journet, M.; Humphrey, G. R.; Cowden, C.; Strotman, N.; Devine, P. J. Org. Chem. 2010, 75, 7829-7841.
- For enantioselective decarboxylative protonations of MAHOs see: (a) Amere, M.; Lasne, M.-C.; Rouden, J. Org. Lett. 2007, 9, 2621–2624; (b) Seitz, J.; Baudoux, J.; Bekolo, H.; Cahard, D.; Plaquevent, J.-C.; Lasne, M.-C.; Rouden, J. Tetrahedron 2006, 62, 6155–6165; (c) Blanchet, J.; Baudoux, J.; Amere, M.; Lasne, M.-C.; Rouden, J. Eur. J. Org. Chem. 2008, 5493–5506. For decarboxylative aldol reactions of MAHOs see: (d) Baudoux, J.; Lefevre, P.; Lasne, M.-C.; Rouden, J. Green Chem. 2010, 12, 252–259; (e) Blaquiere, N.; Shore, D. G.; Rousseaux, S.; Fagnou, K. J. Org. Chem. 2009, 74, 6190-6198; (f) Li, X.; Xiong, H.; Hua, M.; Nie, J.; Ma, J.-A. Tetrahedron Lett. 2012, 53, 2117-2120. For diastereoselective decarboxylative aldol reaction of MAHOs see: (g) Singjunla, Y.; Baudoux, J.; Rouden, J. Org. Lett. 2013, 15, 5770-5773.
- 24. The correct stereochemistry of the dehydroamino ester was determined by a NOE experiment (see Supporting Information for details).
- Appel, R.; Chelli, S.; Tokuyasu, T.; Troshin, K.; Mayr, H. J.Am.Chem.Soc. 2013, 135, 6579-6587.
- 26. The following hydroxyamino ester **25** was isolated in 50% yield as the major product.



- 27. N-CBz pyrrolidine was found as side product.
- Hengartner, U.; Valentine Jr. D.; Johnson, K. K.; Larscheid, M. M. Pigott, F.; Scheidl, F.; Scott, J. W.; Sun, R. C.; Townsend J. M.; Williams, T. H. J. Org. Chem. 1979, 44, 3741-3747.
- 29. Faucher, A-M.; Bailey, M.D.; Beaulieu, P.L.; Brochu, C.; Duceppe, J-S.; Ferland, J-M.; Ghiro, E.; Gorys, V.; Halmos, T.; Kawai, S.H.; Poirier, M.; Simoneau, B.; Tsantrizos, Y.S.; Llina`s-Brunet, M. Org. Lett. **2004**, *6*, 2901-2904.
- 30. Mai, A.H.; Pawar, S.; De Borggraeve, W.M. *Tetrahedron Lett.* **2014**, *55*, 4664-4666.