

Articles

Electrophilic Methylplatinum Complexes: First Structure of a Hydroxytris(pentafluorophenyl)borate Complex

Geoffrey S. Hill,[†] Ljubica Manojlovic-Muir,[‡] Kenneth W. Muir,^{*,‡} and Richard J. Puddephatt^{*,†}

Departments of Chemistry, The University of Western Ontario, London, Ontario, Canada N6A 5B7, and The University of Glasgow, Glasgow G12 8QQ, Scotland


Received September 30, 1996[⊗]

Treatment of $[\text{PtMe}_2(\text{bu}_2\text{bpy})]$ (bu_2bpy = 4,4'-di-*tert*-butyl-2,2'-bipyridine) with $\text{B}(\text{C}_6\text{F}_5)_3$ in the presence of the ligand L gives, under anhydrous conditions, $[\text{PtMeL}(\text{bu}_2\text{bpy})][\text{MeB}(\text{C}_6\text{F}_5)_3]$ {L = CO (**1a**), C_2H_4 (**2a**), PPh_3 (**3a**)}. Similar reactions performed in the presence of H_2O afford $[\text{PtMeL}(\text{bu}_2\text{bpy})][\text{HOB}(\text{C}_6\text{F}_5)_3]$ {L = CO (**1b**), C_2H_4 (**2b**), PPh_3 (**3b**)}. In the absence of L, the treatment of $[\text{PtMe}_2(\text{bu}_2\text{bpy})]$ with $\text{B}(\text{C}_6\text{F}_5)_3$ and H_2O gives $[\text{Pt}\{\text{HOB}(\text{C}_6\text{F}_5)_3\}\text{Me}(\text{bu}_2\text{bpy})]$ (**4**), the first published example of a hydroxytris(pentafluorophenyl)borate complex. All of the complexes are fully characterized by NMR and IR spectroscopy and, in the case of complex **4**, by an X-ray analysis which confirms the attachment of the $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ ligand to the square-planar platinum atom *via* a Pt–O bond of 2.062(2) Å. We propose that, in the presence of water, $\text{B}(\text{C}_6\text{F}_5)_3$ forms an adduct $[\text{H}_2\text{OB}(\text{C}_6\text{F}_5)_3]$ (which acts as a strong acid $\text{H}[\text{HOB}(\text{C}_6\text{F}_5)_3]$) and this then protonates a Pt–Me bond of $[\text{PtMe}_2(\text{bu}_2\text{bpy})]$ forming CH_4 and $[\text{PtMe}(\text{bu}_2\text{bpy})]^+[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$. This protonolysis methodology provides an alternative route to the well-established electrophilic methyl-ligand abstraction by $\text{B}(\text{C}_6\text{F}_5)_3$ for the production of late transition-metal cations.

Introduction

There has been increasing interest in electrophilic cationic transition-metal complexes as they often possess unique and highly desirable catalytic properties. For example, early transition-metal metallocenium and *ansa*-metallocenium cations¹ as well as late transition-metal cations of the type $[\text{MMe}(\text{N}=\text{N})]^+$ (M = Ni, Pd; N=N = diimine)² are highly active catalysts for the polymerization of ethylene and α -olefins.

One of the most common methods used to produce transition-metal cations is by direct electrophilic attack on a M–Me bond by the powerful Lewis acid $\text{B}(\text{C}_6\text{F}_5)_3$, to produce $[\text{M}]^+[\text{MeB}(\text{C}_6\text{F}_5)_3]^-$. This methodology was introduced several years ago for creating vacant sites in early transition-metal complexes³ but has only been recently applied to late transition-metal systems.⁴ For example, we have shown that under anhydrous condi-



tions, methyl-ligand abstraction from $[\text{PtMe}_2(\text{bu}_2\text{bpy})]$ by $\text{B}(\text{C}_6\text{F}_5)_3$ in the presence of the ligand L affords the complexes $[\text{PtMeL}(\text{bu}_2\text{bpy})][\text{MeB}(\text{C}_6\text{F}_5)_3]$ {L = CO (**1a**), C_2H_4 (**2a**); bu_2bpy = 4,4'-di-*tert*-butyl-2,2'-bipyridine}⁴ and we now show that in the presence of PPh_3 it leads to the formation of $[\text{PtMe}(\text{PPh}_3)(\text{bu}_2\text{bpy})][\text{MeB}(\text{C}_6\text{F}_5)_3]$ (**3a**) (Scheme 1).

In this paper we also report that in the presence of water $\text{B}(\text{C}_6\text{F}_5)_3$ does not directly attack a Pt–Me bond of $[\text{PtMe}_2(\text{bu}_2\text{bpy})]$ but forms an adduct $[\text{H}_2\text{OB}(\text{C}_6\text{F}_5)_3]$ (which acts as a strong acid $\text{H}[\text{HOB}(\text{C}_6\text{F}_5)_3]$) which rapidly protonates a Pt–Me bond to form CH_4 and the proposed cationic intermediate $[\text{PtMe}(\text{bu}_2\text{bpy})]^+[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$. This protonation methodology provides an alternative synthetic route to transition-metal cations which still uses $\text{B}(\text{C}_6\text{F}_5)_3$ as the key reagent. The

[†] The University of Western Ontario.

[‡] The University of Glasgow.

[⊗] Abstract published in *Advance ACS Abstracts*, January 1, 1997.


(1) Brintzinger, H. H.; Fischer, D.; Mulhaupt, R.; Rieger, B.; Waymouth, R. M. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 1143.

(2) (a) Rix, F. C.; Brookhart, M.; White, P. S. *J. Am. Chem. Soc.* **1996**, *118*, 4746. (b) Johnson, L. K.; Mecking, S.; Brookhart, M. *J. Am. Chem. Soc.* **1996**, *118*, 267. (c) Johnson, L. K.; Killian, C. M.; Brookhart, M. *J. Am. Chem. Soc.* **1995**, *117*, 6414. (d) Rix, F. C.; Brookhart, M. *J. Am. Chem. Soc.* **1995**, *117*, 1137. (e) Brookhart, M.; Rix, F. C.; DeSimone, J. C.; Barborak, J. C. *J. Am. Chem. Soc.* **1992**, *114*, 5895.

(3) Yang, X.; Stern, C. L.; Marks, T. J. *J. Am. Chem. Soc.* **1991**, *113*, 3623.

(4) Hill, G. S.; Rendina, L. M.; Puddephatt, R. J. *J. Chem. Soc., Dalton Trans.* **1996**, 1809.

Scheme 2



closest analogies to this system of which we are aware employ either the trialkylammonium acid⁵ $[\text{HNBu}_3]^+[\text{B}(\text{C}_6\text{F}_5)_4]^-$ or the oxonium acid⁶ $[\text{H}(\text{OEt}_2)_2]^+[\{3,5-(\text{CF}_3)_2\text{C}_6\text{H}_3\}_4\text{B}]^-$. The protonation of a zirconium–methyl bond in $[(\text{Cp}^*)_2\text{ZrMe}_2]$ by $[\text{HNEt}_3][\text{HOB}(\text{C}_6\text{F}_5)_3]$ gives *not* a cationic species but the zirconoxyborane $[(\text{Cp}^*)_2\text{ZrOB}(\text{C}_6\text{F}_5)_3]$.⁷

Results and Discussion

Under anhydrous conditions, electrophilic methyl-ligand abstraction from $[\text{PtMe}_2(\text{bu}_2\text{bpy})]$ by the powerful Lewis acid $\text{B}(\text{C}_6\text{F}_5)_3$ in the presence of the ligand L gives the complexes $[\text{PtMeL}(\text{bu}_2\text{bpy})][\text{MeB}(\text{C}_6\text{F}_5)_3]$ $\{\text{L} = \text{CO} \text{ (1a)}, \text{C}_2\text{H}_4 \text{ (2a)}, \text{PPh}_3 \text{ (3a)}\}$; $\text{bu}_2\text{bpy} = 4,4'$ -di-*tert*-butyl-2,2'-bipyridine. The synthesis and characterization of complexes **1a** and **2a** have been reported previously.⁴ Complex **3a** is an orange hygroscopic solid that can be isolated in good yield and has been characterized on the basis of NMR spectroscopic (¹H, ¹¹B, and ³¹P) and microanalytical data. The ¹H NMR spectrum shows the expected two *tert*-butyl resonances and six sets of aromatic resonances due to the inequivalent nature of the two pyridyl moieties of the bu_2bpy ligand. The Pt–Me ligand resonates at $\delta = 0.71$ with $^2J(\text{PtH}) = 72.0$ Hz, typical of other cationic platinum(II) complexes.⁴ The ³¹P NMR spectrum shows the expected singlet at $\delta = 20.0$. This resonance is flanked by ¹⁹⁵Pt satellite signals with $^1J(\text{PtP}) = 4343$ Hz, typical of other cationic platinum(II)–phosphine complexes.⁹ The ¹¹B NMR spectra of complexes **1a**, **2a**, and **3a** show a singlet at $\delta = -15.1$, indicative of the $[\text{MeB}(\text{C}_6\text{F}_5)_3]^-$ anion.^{3,7,10}

During the initial preparations of complexes **1a**, **2a**, and **3a**, it was noted that some of the corresponding complex, $[\text{PtMeL}(\text{bu}_2\text{bpy})][\text{HOB}(\text{C}_6\text{F}_5)_3]$ $\{\text{L} = \text{CO} \text{ (1b)}, \text{C}_2\text{H}_4 \text{ (2b)}, \text{PPh}_3 \text{ (3b)}\}$, was often produced (Scheme 2). The cause of this is most certainly the introduction of adventitious H₂O into the reaction vessel since these $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ complexes are formed exclusively if 1 equiv of H₂O is added during the initial stages of the reaction. Complexes **1b**, **2b**, and **3b** give very similar NMR spectra to complexes **1a**, **2a**, and **3a**, respectively,

with the exception of the ¹¹B NMR which shows a singlet at $\delta = -4.4$, indicative of the $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ anion.⁷ The infrared spectra of complexes **1b**, **2b**, and **3b** show a strong O–H stretching band at 3684 cm^{-1} that is absent in the otherwise very similar infrared spectra of **1a**, **2a**, and **3a**, respectively.

In the absence of the ligand L, the low-temperature reaction of $[\text{PtMe}_2(\text{bu}_2\text{bpy})]$ with $\text{B}(\text{C}_6\text{F}_5)_3$ and 1 equiv of water produces CH₄ and the {hydroxytris(pentafluorophenyl)borato}platinum(II) complex **4** in high yield (Scheme 2).¹¹ To our knowledge, the only other example of a transition-metal complex incorporating the $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ group is $[(\text{Cp}^*)_2\text{Ta}(\text{OH})\text{Me}][\text{HOB}(\text{C}_6\text{F}_5)_3]$, which is formed upon treatment of $[(\text{Cp}^*)_2\text{TaMe}_2]\text{SO}_3\text{CF}_3$ with $[\text{NaB}(\text{C}_6\text{F}_5)_4]$ and in which the $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ group is not coordinated.⁸ Thus, complex **4** provides the first published example of a transition-metal complex containing a coordinated $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ ligand, though such species have been mentioned in the patent literature.⁷ The ¹H NMR spectrum of complex **4** shows the expected two *tert*-butyl resonances and six sets of aromatic resonances due to the inequivalent nature of the two pyridyl moieties of the bu_2bpy ligand. The Pt–Me ligand resonates at $\delta = 0.81$ with $^2J(\text{PtH}) = 78.9$ Hz. The PtO(*H*)B group gives a broad resonance at $\delta = 3.21$ which does not show any resolvable splitting even at -90°C . No coupling of the PtO(*H*)B group to ¹⁹⁵Pt is observed, and we assume that this is due to a rapid exchange process. This PtO(*H*)B resonance disappears upon treatment of an Et₂O solution of complex **4** with D₂O and is absent in the otherwise identical ¹H NMR spectrum of $[\text{Pt}\{\text{DOB}(\text{C}_6\text{F}_5)_3\}\text{Me}(\text{bu}_2\text{bpy})]$ prepared from $[\text{PtMe}_2(\text{bu}_2\text{bpy})]$, $\text{B}(\text{C}_6\text{F}_5)_3$, and D₂O. The infrared spectrum of complex **4** shows a sharp O–H stretching band at 3640 cm^{-1} . This band is replaced by a weak absorption at 2673 cm^{-1} in the infrared spectrum of $[\text{Pt}\{\text{DOB}(\text{C}_6\text{F}_5)_3\}\text{Me}(\text{bu}_2\text{bpy})]$. At room temperature, rapid rotation about the O–B and B–C bonds renders the three pentafluorophenyl moieties of **4** equivalent, and the ¹⁹F NMR spectrum shows only three sets of resonances, one set for each of the *ortho*-, *meta*-, and *para*-fluorine atoms. However, at -90°C these three sets resolve into eight separate resonances, indicating that the three pentafluorophenyl moieties are no longer equivalent. This low-temperature phenomenon is certainly a result of the steric interference introduced upon coordination of the $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ ion to platinum, since none of the compounds which contain either the free $[\text{MeB}(\text{C}_6\text{F}_5)_3]^-$ ion (in complexes **1a**, **2a**, **3a**) or $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ ion (in complexes **1b**, **2b**, **3b**) shows any further resolution in the ¹⁹F NMR at -90°C due to inequivalent pentafluorophenyl groups.

The structure of **4** has been confirmed by X-ray analysis (Figure 1). The *cis*-N₂CO donor set which surrounds the Pt atom departs significantly from ideal square-planar coordination: as expected, the N–Pt–N angle narrows to accommodate the restricted bite of the bu_2bpy ligand and there is also a slight tetrahedral distortion, with donor atoms lying alternately 0.04 Å above and below the coordination plane. The Pt–N bond lengths (Table 1) differ by 0.117(3) Å, reflecting the high *trans*-influence of $\sigma\text{-CH}_3$ and the low *trans*-

(5) Yang, X.; Stern, C. L.; Marks, T. L. *Organometallics* **1991**, *10*, 840.

(6) (a) Brookhart, M.; Grant, B.; Volpe, Jr., A. F. *Organometallics* **1992**, *11*, 3920. (b) Holtcamp, M. W.; Labinger, J. A.; Bercaw, J. E. Personal communication.

(7) (a) Siedle, A. R.; Newmark, R. A.; Lamanna, W. M.; Huffman, J. C. *Organometallics* **1993**, *12*, 1491. (b) Siedle, A. R.; Lamanna, W. M. *PCT Int. Appl. WO* **1993**, 21,238 (*Chem. Abs.* **1994**, *121*, 36424x).

(8) Schaefer, W. P.; Quan, R. W.; Bercaw, J. E. *Acta Crystallogr.* **1993**, *C49*, 878.

(9) Appleton, T. G.; Clark, H. C.; Manzer, L. E. *Coord. Chem. Rev.* **1973**, *10*, 335.

(10) Siedle, A. R.; Newmark, R. A. *J. Organomet. Chem.* **1995**, *497*, 119.

(11) CH₃D was detected by GC-MS when the reaction was performed in the presence of D₂O.




Figure 1. Molecular structure of $[\text{Pt}\{\text{HOB}(\text{C}_6\text{F}_5)_3\}\text{Me}(\text{bu}_2\text{bpy})]$, **4**. Displacement ellipsoids of 30% are shown except for hydrogen atoms which are displayed as spheres of arbitrary size. The disorder of the methyl carbon atoms attached to C11 is not illustrated. Phenyl carbon atoms are numbered cyclically C21–C26, C31–C36, and C41–C46, in each case starting from the atom bonded to B1. The atoms F1–F5 are attached in sequence to C22–C26, F6–F10 to C32–C36, and F11–F15 to C42–C46.

influence of oxygen donor ligands,^{9,12} thereby confirming less precise results of the only previous structural study of a platinum(II) compound with a *cis*-($\text{sp}^2\text{-N}$)₂CO donor set.¹³ The Pt–O bond length in **4** [2.062(2) Å] is a little longer than the mean Pt(II)–OH distance obtained from a database survey [2.006 Å],¹⁴ while the reverse is true of the Pt–CH₃ distance [2.037(3) Å in **4** compared with a database value of 2.107 Å].

The bu₂bpy ligand contains two rigorously planar pyridyl rings (rms, $\Delta < 0.004$ Å) twisted about the central single bond [C5–C6 = 1.476(3) Å] so that the C4–C5–C6–C7 torsion angle is 6.9(4)°. This twist still leaves a short H4⋯H7 contact of 2.16 Å; it may explain the tetrahedral distortion of the Pt coordination. The acuteness of the endocyclic C–C–C angles at C3 and C8 [115.7(3) and 115.8(2)°] can be attributed to electron release by the *tert*-butyl substituents.¹⁵ Each of the three methyl groups attached to C11 is disordered over two sites with occupancy ratio *ca.* 4:1. The methyl groups attached to C15 show physically unreasonable displacement parameters [e.g. $U_{\text{max}} = 0.27$ Å² for C17] suggestive of disorder which is not explicitly accounted for in the final structural model.

The $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ ligand in **4** is much the most novel feature of the structure. The analysis allowed the hydrogen atom attached to O1 to be successfully, though not very accurately, located. The central boron atom has a somewhat distorted tetrahedral coordination and forms B–C bonds whose lengths [1.636(4)–1.659(4) Å] cluster about the database mean¹⁴ for the corresponding bonds in BPh_4^- (1.643 Å). The B–O bond length [1.526(3) Å] may be slightly lengthened by coordination

Table 1. Selected Distances (Å) and Angles (deg) in $[\text{Pt}\{\text{HOB}(\text{C}_6\text{F}_5)_3\}\text{Me}(\text{bu}_2\text{bpy})]$, **4**

(a) Bond Lengths			
Pt1–O1	2.062(2)	Pt1–N1	1.980(2)
Pt1–N2	2.097(2)	Pt1–C19	2.037(3)
C21–B1	1.659(4)	O1–B1	1.526(3)
C41–B1	1.636(4)	C31–B1	1.654(3)
(b) Bond Angles			
O1–Pt1–N1	172.6(1)	O1–Pt1–N2	93.3(1)
O1–Pt1–C19	90.6(1)	N1–Pt1–N2	80.1(1)
N1–Pt1–C19	96.1(1)	N2–Pt1–C19	175.7(1)
Pt1–O1–N10	115(3)	Pt1–O1–B1	128.2(2)
H10–O1–B1	111(3)	O1–B1–C21	108.3(2)
O1–B1–C31	111.7(2)	O1–B1–C41	103.2(2)
C21–B1–C31	104.3(2)	C21–B1–C41	115.2(2)
C31–B1–C41	114.3(2)		
(c) Torsion Angles			
C19–Pt1–O1–B1	73.8(2)	Pt1–O1–B1–C21	132.5(2)
Pt1–O1–B1–C31	–113.2(2)	Pt1–O1–B1–C41	10.0(3)
C22–C21–B1–O1	–140.5(2)	C26–C21–B1–O1	46.2(3)
C32–C31–B1–O1	–14.0(4)	C36–C31–B1–O1	170.5(2)
C42–C41–B1–O1	70.2(3)	C46–C41–B1–O1	–100.6(3)
(d) Hydrogen Bonds			
bond	O⋯F	O–H	H⋯F
O1–H10⋯F5	2.722(3)	0.67(4)	2.32(4)
O1–H10⋯F6	2.692(3)	0.67(4)	2.18(4)
			O–H⋯F
			120(4)
			135(5)

to Pt: corresponding values in structures containing the uncomplexed $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ anion are 1.487(3) Å in $[\text{Et}_3\text{NH}][\text{HOB}(\text{C}_6\text{F}_5)_3]$ and 1.490(10) Å in $[(\text{Cp}^*)_2\text{TaMe}(\text{OH})][\text{HOB}(\text{C}_6\text{F}_5)_3]$.^{7,8} It has been shown by a crystal structure analysis of $[\text{H}_2\text{OB}(\text{C}_6\text{F}_5)_3] \cdot 2\text{H}_2\text{O}$ that only one water molecule is directly attached to boron, forming the conjugate acid of $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$.¹⁶ However in $[\text{Re}^{\text{VO}_2}(\text{C}_{10}\text{H}_{24}\text{N}_4)]\text{Cl} \cdot 2[\text{H}_2\text{OBPh}_3]$ the B–O distance [1.602(6) Å] is much longer than in **4**.¹⁷ The only other structure related to **4** of which we are aware is $[(\text{Cp}^*)_2\text{Zr}^{\text{IV}}(\text{OB}(\text{C}_6\text{F}_5)_3)]$ in which the conjugate base of $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ is bonded to zirconium through an *ortho* fluorine atom as well as through oxygen [B–O 1.460(6) Å]; the B–O–Zr angle of 151.2(3)° in this structure is substantially more obtuse than the B–O–Pt angle of 128.2(2)° in **4**.⁷ However, it may also be worth noting here that there is a single structural report of the attachment of OBPh_3^{2-} to a tungsten(VI) cation.¹⁸

The conformation of the $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ ligand in **4** is such that B1 lies far from the coordination plane of the metal atom (C19–Pt1–O1–B1 = 74°). The Pt1–O1–B1–C41 chain is nearly planar (torsion angle 10°) so that the C41 ring lies above the coordination plane of Pt1; however, the shortest contact between the C41 ring and the metal [Pt1⋯C41 = 3.185(2) Å] is clearly nonbonding. This arrangement leaves the face of the Pt1 coordination plane opposite to the C41 ring relatively exposed, and in consequence there is a short intermolecular Pt⋯Pt contact of 4.1201(4) Å with a centrosymmetrically related molecule.

One further feature of the conformation adopted by the $[\text{HOB}(\text{C}_6\text{F}_5)_3]^-$ ligand in **4** requires comment. The OH group is contained in a cleft between the C21 and C31 C_6F_5 rings so that a weak bifurcated intramolecular

(12) Manojlovic-Muir, Lj.; Muir, K. W. *Inorg. Chim. Acta* **1974**, *10*, 47.

(13) Dieck, H. T.; Fendesak, G.; Munz, C. *Polyhedron* **1991**, *10*, 255.

(14) *International Tables for Crystallography*, Vol. C, International Union of Crystallography; Kluwer Academic Publishers: Dordrecht, The Netherlands, 1992; Vol. C, Tables 9.5.1.1 and 9.6.3.3.

(15) *Accurate Molecular Structures*; Domenicano, A., Hargittai, I., Eds.; International Union of Crystallography; OUP: Oxford, U.K., 1992; Chapter 18 and references therein. In particular: Domenicano, A.; Vaciago, A.; Coulson, C. A. *Acta Crystallogr.* **1975**, *B31*, 221.

(16) Siedle, A. R.; Lamanna, W. M.; Newmark, R. A.; Stevens, J.; Richardson, D. E.; Ryan, M. *Makromol. Chem. Macromol. Symp.* **1993**, *66*, 215.

(17) Blake, A. J.; Greig, J. A.; Schroder, M. *J. Chem. Soc., Dalton Trans.* **1988**, 2645.

(18) Schreiber, P.; Wieghardt, K.; Nuber, B.; Weiss, J. *Z. Naturforsch. B* **1990**, *45*, 619.


hydrogen bond is formed involving F5 and F6 as acceptors (Table 1d). Interestingly, the C–F5 and C–F6 bond lengths [1.366(4) and 1.362(3) Å] are very slightly longer than the other 13 C–F bond lengths [1.343(4)–1.355(3) Å]. Although the 11 atoms of each C₆F₅ ring barely deviate from planarity (rms $\Delta < 0.02$ Å) the displacements of the boron atom from the ring planes [respectively 0.113(3), 0.026(3), and 0.186(3) Å for the C21, C31, and C41 rings] are irregular and in two cases highly significant. The acuteness of the endocyclic C–C–C angles subtended by the atoms C21, C31, and C41 [113.0(3)–113.5(3)°] is compensated by an increase to 123.9(3)–124.5(3)° of the adjacent angles at carbon atoms *ortho* to boron. These features have been noted previously in the free [HOB(C₆F₅)₃][–] anion⁸ and may be ascribed to the difference in the electro-negativities of B and F.¹⁵

The [HOB(C₆F₅)₃][–] ligand in complex **4** is only weakly coordinated to platinum as treatment of **4** with CO, C₂H₄, and PPh₃ gives the complexes **1b**, **2b**, and **3b**, respectively (Scheme 2). Similar complexes of the type [PtMeX(bu₂bpy)] (X = O₃SCF₃, O₂CCF₃) do not react with C₂H₄, illustrating that the [HOB(C₆F₅)₃][–] ligand is an even poorer ligand than triflate or trifluoroacetate for platinum.⁴

In the synthesis of complexes **1a**, **2a**, and **3a**, B(C₆F₅)₃ acts as a powerful Lewis acid and directly attacks a Pt–Me bond in [PtMe₂(bu₂bpy)] to form [PtMe(Bu₂bpy)]⁺[MeB(C₆F₅)₃][–]. However, in the presence of water a different mechanism must be followed because the corresponding [PtMe(bu₂bpy)]⁺[HOB(C₆F₅)₃][–] complex is preferentially formed. Although B(C₆F₅)₃ has often been described as a “water-tolerant Lewis acid”,¹⁹ it has been shown that, in the presence of H₂O, B(C₆F₅)₃ forms the hydrate [H₂OB(C₆F₅)₃]·2H₂O with one molecule of H₂O directly coordinated to boron.¹⁶ In the presence of a suitable base, Nu, this hydrate can be deprotonated to form the acid [HNu]⁺[HOB(C₆F₅)₃][–].^{7,16} It has been shown that the highly nucleophilic complex [PtMe₂(bu₂bpy)] reacts rapidly with a variety of protic acids, resulting in protonation of the Pt–Me bond to initially form CH₄ and [PtMe(bu₂bpy)]⁺.²⁰ Thus, in the synthesis of complexes **1b**, **2b**, and **3b**, [PtMe₂(bu₂bpy)] most probably acts as Nu and deprotonates H₂OB(C₆F₅)₃ to form CH₄ and [PtMe(bu₂bpy)]⁺[HOB(C₆F₅)₃][–]. This complex can either react with L to form [PtMeL(bu₂bpy)][HOB(C₆F₅)₃][–] {L = CO (**1b**), C₂H₄ (**2b**), PPh₃ (**3b**)} or, in the absence of L, form the hydroxytris(pentafluorophenyl)borate complex (**4**) (Scheme 2). In the reaction with D₂O in place of H₂O, the organic products contained CH₃D but no CH₂D₂, indicating that exchange between PtH and PtCH₃ protons does not occur to a significant extent in this reaction.²¹

It has been well established that the protonation of the Pt–C bond follows an oxidative addition/reductive elimination mechanism proceeding through an alkyl-(hydrido)platinum(IV) intermediate.^{20,21} The most definitive evidence for the proposed protonation mechanism for the production of complexes **1b**, **2b**, **3b**, and **4**

Scheme 3



would be the detection of the methyl(hydrido)platinum(IV) intermediate **5** (Scheme 3). In an attempt to detect this intermediate, the reaction of B(C₆F₅)₃ and water with [PtMe₂(bu₂bpy)] in CD₂Cl₂ was monitored by low-temperature ¹H NMR spectroscopy (–90 °C). A resonance was detected at $\delta = -18.5$, suggestive of a (hydrido)platinum species.^{20,21} Unfortunately, the stability of this hydrido species was poor and at any given time the concentration was not high enough to discern the ¹⁹⁵Pt satellite signals.²² Since the order of stability toward CH₄ reductive elimination from the methyl-(hydrido)platinum(IV) complexes [PtX(H)Me₂(bu₂bpy)] (X = Cl, Br, I, O₂CCF₃, SO₃CF₃) follows the order of the Pt–X bond strength, i.e. Cl > Br > I > O₂CCF₃, SO₃CF₃, it is not unexpected that complex **5**, containing the weakly bound [HOB(C₆F₅)₃][–] ion, would be highly unstable.²⁰ Nevertheless, this low-frequency resonance in the ¹H NMR spectrum is highly suggestive of a transient (hydrido)platinum intermediate and, thus, provides further evidence for the protonation mechanism proposed in Scheme 3.

It has been suggested that the reaction of H₂O with [PtMe₂(N–N)] (N–N = 2,2′-bipyridine, 1,10-phenanthroline) proceeds through the hydroxy(hydrido)platinum(IV) intermediate [PtH(OH)Me₂(N–N)]²³ (Scheme 4). It is thus conceivable that complex **5** is formed by an initial reaction of H₂O with [PtMe₂(bu₂bpy)], forming [PtH(OH)Me₂(bu₂bpy)], which then reacts with B(C₆F₅)₃ to give complex **5**. However, the reaction of H₂O with [PtMe₂(N–N)] is slow, taking several hours to reach completion,²³ whereas the reaction of Scheme 2 is very rapid, so this mechanism can immediately be eliminated. Since H[HOB(C₆F₅)₃] is a strong acid, it is expected to react with [PtMe₂(bu₂bpy)] much more rapidly than water itself would.

Conclusions

In addition to the well-established methyl-ligand abstraction by B(C₆F₅)₃ to form [M]⁺[MeB(C₆F₅)₃][–], we have shown that, in the presence of water, B(C₆F₅)₃ protonates a Pt–Me bond of [PtMe₂(bu₂bpy)] to form [PtMe(bu₂bpy)]⁺[HOB(C₆F₅)₃][–]. In the presence of the

(19) (a) Ishihara, K.; Hanaki, N.; Funahashi, M.; Miyata, M.; Yamamoto, H. *Bull. Chem. Soc. Jpn.* **1995**, *68*, 1721. (a) Yang, X.; Stern, C. L.; Marks, T. J. *J. Am. Chem. Soc.* **1994**, *116*, 10015.


(20) Hill, G. S.; Rendina, L. M.; Puddephatt, R. J. *Organometallics* **1995**, *14*, 4966.

(21) (a) Stahl, S. S.; Labinger, J. A.; Bercaw, J. E. *J. Am. Chem. Soc.* **1995**, *117*, 9371; **1996**, *118*, 5961. (b) De Felice, V.; De Renzi, A.; Panunzi, A.; Tesaro, D. *J. Organomet. Chem.* **1995**, *488*, C13.

(22) Monitoring similar reactions by low-temperature ¹H NMR (–90 °C) with either CH₃OH or C₆H₅OH in place of H₂O resulted in no observable hydride resonances.

(23) (a) Canty, A. J.; Fritsche, S. D.; Jin, H.; Honeyman, R. T.; Skelton, B. W.; White, A. H. *J. Organomet. Chem.* **1996**, *510*, 281. (b) Canty, A. J.; Fritsche, S. D.; Jin, H.; Skelton, B. W.; White, A. H. *J. Organomet. Chem.* **1995**, *490*, C18. (c) Canty, A. J.; Honeyman, R. T.; Roberts, A. S.; Traill, P. R.; Colton, R.; Skelton, B. W.; White, A. H. *J. Organomet. Chem.* **1994**, *471*, C8. (d) Appleton, T. G.; Hall, J. R.; Neale, D. W.; Williams, M. A. *J. Organomet. Chem.* **1984**, *276*, C73. (e) Monaghan, P. K.; Puddephatt, R. J. *Organometallics* **1984**, *3*, 444.

Scheme 4



ligand L (L = CO, C₂H₄, PPh₃) the corresponding cationic complex, [PtMeL(bu₂bpy)][HOB(C₆F₅)₃], is formed. However, in the absence of L, the [HOB(C₆F₅)₃][−] anion coordinates to platinum to form the first published example of a hydroxytris(pentafluorophenyl)borate complex (**4**), which has been fully characterized by NMR spectroscopy and X-ray crystallography. This procedure provides an additional route to the production of late transition-metal cations by the powerful Lewis acid B(C₆F₅)₃.

Experimental Section

General Procedures. All reactions were performed under a N₂ atmosphere using standard Schlenk techniques, unless otherwise stated. All solvents were freshly distilled, dried, and degassed prior to use. NMR spectra were recorded using a Varian Gemini spectrometer (¹H at 300.10 MHz, ¹¹B at 64.17 MHz, ¹⁹F at 282.32 MHz, ³¹P at 121.44 MHz, and ¹⁹⁵Pt at 42.99 MHz). Chemical shifts are reported in ppm with respect to TMS (¹H), BF₃·Et₂O (¹¹B), CFCl₃ (¹⁹F), H₃PO₄/D₂O (³¹P), or K₂[PtCl₄]/D₂O (¹⁹⁵Pt). The ¹H, ¹¹B, ¹⁹F, ³¹P, and ¹⁹⁵Pt NMR spectra are referenced to the residual protons of the deuterated solvents or to BF₃·Et₂O, CFCl₃, H₃PO₄/D₂O, or K₂[PtCl₄]/D₂O contained in a coaxial insert, respectively. IR spectra (Nujol mull) were recorded in the range 4000–400 cm^{−1} using a Perkin-Elmer 2000 FT-IR instrument. Elemental analyses were determined by Guelph Chemical Laboratories, Guelph, Canada.

[PtMe₂(bu₂bpy)],²⁴ [PtMe(CO)(bu₂bpy)][MeB(C₆F₅)₃],⁴ and [Pt(Me)(C₂H₄)(bu₂bpy)][MeB(C₆F₅)₃]⁴ were prepared by the literature methods. B(C₆F₅)₃ was obtained commercially (Strem).

Preparation of Complexes. **4,4′-Di-*tert*-butyl-2,2′-bipyridine{hydroxytris(pentafluorophenyl)borato}-methylplatinum(II), [Pt{HOB(C₆F₅)₃}Me(bu₂bpy)] (**4**).** All glassware must be flame-dried under vacuum prior to use. To a 100 mL Schlenk tube charged with [PtMe₂(bu₂bpy)] (10.0 mg, 0.020 mmol) and B(C₆F₅)₃ (10.4 mg, 0.020 mmol) was added Et₂O (5.0 mL) and distilled H₂O (0.4 μL, 0.02 mmol) immediately producing an orange to yellow color change. Removal of the solvent *in vacuo* and recrystallization from CH₂Cl₂/*n*-pentane gave a bright yellow microcrystalline product. Yield: 18.4 mg (92%). ¹H NMR (CD₂Cl₂): δ = 8.64 [d, 1H, ³J(H⁶H⁵) = 6.3 Hz, ³J(PtH) = *ca.* 60 Hz, H⁶], 8.50 [d, 1H, ³J(H⁶H⁵) = 5.6 Hz, H⁶], 7.91 [d, 1H, ⁴J(H³H⁵) = 1.9 Hz, H³], 7.84 [d, 1H, ⁴J(H³H⁵) = 2.2 Hz, H³], 7.58 [dd, 1H, ⁴J(H⁵H³) = 1.9 Hz, ³J(H⁵H⁶) = 5.7 Hz, H⁵], 7.35 [dd, 1H, ⁴J(H⁵H³) = 2.2 Hz, ³J(H⁵H⁶) = 6.3, H⁵], 3.21 [br s, 1H, Pt–O(H)B], 1.41 (s, 9H, *tert*-bu), 1.39 (s, 9H, *tert*-bu), 0.81 [s, 3H, ²J(Pt–H) = 78.9 Hz, Pt–Me]. ¹¹B NMR (CD₂Cl₂): δ = 0.1 (br s). ¹⁹F NMR (CD₂-

Cl₂, 25 °C): δ = −124.5 [br s, 6F, *o*-F], −160.8 [m, 3F, *p*-F], −165.8 [m, 6F, *m*-F]. ¹⁹F NMR (CD₂Cl₂, −90 °C): δ = −132 (br m, 3F, *o*-F), −133 (s, 1F, *o*-F), −139 (s, 1F, *o*-F), −143 (s, 1F, *o*-F), −159 (s, 1F, *p*-F), −160 (br m, 2F, *p*-F), −164 (br m, 3F, *m*-F), −165 (br m, 3F, *m*-F). ¹⁹⁵Pt NMR (CD₂Cl₂): δ = −1330 [br s (Δν^{1/2} = *ca.* 400 Hz)]. IR (Nujol, cm^{−1}): 3640 [s, ν(OH)]. Anal. Calcd for C₃₇H₂₈BF₁₅N₂O₂Pt: C, 44.0; H, 3.1; N, 2.7. Found: C, 44.1; H, 2.8; N, 2.8.

Alternate Method for the Preparation of **4.** To a suspension of [PtMe₂(bu₂bpy)] (0.100 g, 0.203 mmol) in Et₂O (5.0 mL) at −78 °C was added a solution of B(C₆F₅)₃ (0.105 g, 0.203 mmol) in Et₂O (2.0 mL). The solution was stirred at −78 °C until the color had completely changed from orange to bright-yellow (*ca.* 2.0 h). The mixture was then allowed to warm to room temperature, and the solvent was removed *in vacuo* to give a yellow product. The product could be recrystallized from CH₂Cl₂/*n*-pentane to afford a yellow microcrystalline solid. Yield: 0.189 g (93%).

(4,4′-Di-*tert*-butyl-2,2′-bipyridine)methyl(triphenylphosphine)platinum(II) Methyltris(pentafluorophenyl)borate, [PtMePPh₃(bu₂bpy)][MeB(C₆F₅)₃] (3a**).** All glassware must be flame-dried under vacuum prior to use. To a 50 mL Schlenk tube charged with [PtMe₂(bu₂bpy)] (0.20 g, 0.405 mmol), B(C₆F₅)₃ (0.208 g, 0.405 mmol), and PPh₃ (0.106 g, 0.405 mmol) was added Et₂O (10.0 mL). This bright orange mixture was stirred at room temperature for *ca.* 24 h giving a pale orange solution. Removal of the solvent *in vacuo* gave a pale orange powder. Yield: 0.510 g (99%). ¹H NMR (CD₂Cl₂): δ = 8.89 [d, 1H, ³J(H⁶H⁵) = 6.4 Hz, ³J(PtH) = *ca.* 35 Hz, H⁶], 8.88 [d, 1H, ³J(H⁶H⁵) = 6.2 Hz, ³J(PtH) = *ca.* 35 Hz, H⁶], 8.20 (br s, 1H, H³), 8.11 [d, 1H, ⁴J(H³H⁵) = 2.2 Hz, H³], 7.78 [m, 6H, PPh₃], 7.55 [dd, 1H, ⁴J(H⁵H³) = 1.9 Hz, ³J(H⁵H⁶) = 6.4 Hz, H⁵], 7.50 (m, 9H, PPh₃), 7.00 [dd, 1H, ⁴J(H⁵H³) = 2.2 Hz, ³J(H⁵H⁶) = 6.1, H⁵], 1.49 [s, 9H, *tert*-bu], 1.32 (s, 9H, *tert*-bu), 0.71 [s, 3H, ²J(PtH) = 72.0 Hz, ³J(PtP) = 3.7 Hz, Pt–Me]. ¹¹B NMR (CD₂Cl₂): δ = −15.1 (br s). ³¹P NMR (CD₂Cl₂): δ = 20.0 [s, ¹J(PtP) = 4343 Hz]. Anal. Calcd for C₅₆H₄₅BF₁₅N₂-PPh₃: C, 53.0; H, 3.6; N, 2.2. Found: C, 52.7; H, 3.5; N, 2.8.

Synthesis of Complexes **1b, **2b**, and **3b**.** The addition of 1 equiv of H₂O during the initial stages of the preparation of complexes **1a**, **2a**, and **3a** afforded exclusively complexes **1b**, **2b**, and **3b**, respectively. The ¹H NMR spectra of the cationic platinum complexes are identical to those of the analogous [MeB(C₆F₅)₃][−] complexes (**1a**, **2a**, and **3a**, respectively). ¹¹B NMR (CD₂Cl₂): δ = −15.1. IR (Nujol, cm^{−1}): 3684 [s, ν(OH)].

Crystal structure analysis of **4:** C₃₇H₂₈BF₁₅N₂O₂Pt, fw = 1007.51, triclinic, space group *P* $\bar{1}$, *a* = 10.7613(7) Å, *b* = 12.2174(11) Å, *c* = 15.9389(7) Å, α = 110.855(5)°, β = 96.347(4)°, γ = 98.985(6)°, *V* = 1902.7(2) Å³ [from setting angles of 25 reflections with 20.7 ≤ θ(Mo Kα) ≤ 22.3°], *Z* = 2, *D*_{calc} = 1.759 Mg/m³, μ = 3.794 mm^{−1}, *F*(000) = 980.

The intensities of 12 786 reflections with 2.3 ≤ θ(Mo Kα) ≤ 30° and −15 ≤ *h* ≤ 15, −17 ≤ *k* ≤ 2, and −21 ≤ *l* ≤ 22 were measured at 20 °C from ω/2θ scans on an Enraf-Nonius CAD4 diffractometer with Mo Kα X-rays, λ = 0.710 73 Å, using a yellow, needle-shaped crystal of dimensions 0.66 × 0.36 × 0.21 mm. The mean intensity of the three standard reflections decreased by 9.4% during the experiment. Correction for this decomposition and for absorption²⁵ (analytical method,²⁶ transmission factors 0.32–0.53) and subsequent merging gave 11 037 independent reflections (*R*_{int} = 0.015). Refinement of *F*² on 536 parameters using 11 036 observations converged (Δ/*σ* < 0.001) at *R*₁ = 0.025, *wR*₂ = 0.059 for 9203 reflections with *I* > 2σ(*I*) and *R*₁ = 0.040, *wR*₂ = 0.065 for all 11 037 data with

(25) Computer programs used for data processing were as follows.

(a) GX: Mallinson, P. R.; Muir, K. W. *J. Appl. Crystallogr.* **1985**, *18*, 53. (b) PLATON: Spek, A. L. *Acta Crystallogr.* **1990**, *A46*, 31.

(26) de Meulenaer, J.; Tompa, H. *Acta Crystallogr.* **1965**, *19*, 1014.

(24) Achar, S.; Scott, J. D.; Vittal, J. J.; Puddephatt, R. J. *Organometallics* **1993**, *12*, 4592.

$|\Delta\rho| < 0.96 \text{ e}\cdot\text{\AA}^{-3}$ and $w = 1/[\sigma^2(F_o^2) + (0.0342P)^2 + 0.72P]$, where $P = (F_o^2 + 2F_c^2)/3$.²⁷

In the final calculations the positional and anisotropic displacement parameters of 57 non-hydrogen atoms (C₃₇-BF₁₅N₂OPt), a scale factor, an extinction parameter, and orientation parameters for seven methyl groups were refined (Supporting Information). The C12–C14 methyl carbon atoms were found to be disordered with site occupancy, α , of 0.794–(5). The alternative sites C12*, C13*, and C14* were assigned occupancies $1 - \alpha$ and U^{ij} parameters equal respectively to those of C13, C12, and C14; only their positional parameters were refined. Hydrogen atoms attached to C12*–C14* were not included in the calculations. Despite the disorder of the methyl groups, the hydrogen atom bonded to O1 was located

in a difference map. Its positional and isotropic thermal parameters were refined freely. Other hydrogen atoms rode on their parent carbon atoms with C–H and $U(\text{H})_{\text{iso}}$ set at SHELXL default values.²⁷ Atomic scattering factors and dispersion corrections were taken from ref 28.

Acknowledgment. We thank the NSERC (Canada) for financial support to R.J.P. and for a Post-graduate Scholarship to G.S.H., EPSRC (U.K.) and the University of Glasgow for support to K.W.M., and Dr. Louis Farrugia for supplying PC versions of GX and PLATON.

Supporting Information Available: For **4**, tables of atomic parameters and a complete geometry listing (9 pages). Ordering information is given on any current masthead page. X-ray data have also been deposited with the Cambridge Crystallographic Data Centre.

OM960834J

(27) Sheldrick, G. M. SHELXL-93. A Program for the Refinement of Crystal Structures, University of Göttingen, Germany, 1993.

(28) *International Tables for Crystallography*; International Union of Crystallography; Kluwer Academic Publishers: Dordrecht, The Netherlands, 1992; Vol. C, Tables 4.2.6.8. and 6.2.1.1.