REGULAR ARTICLE



Special Issue on 150 years of the Periodic Table

Solvent-assisted monomeric molecular structure of the phosphate diester and the synthesis of menthol-based phosphate diesters

DEBDEEP MANDAL^a, VIVEK GUPTA^a, BISWAJIT SANTRA^a, NICOLAS CHRYSOCHOS^b, VIVEK W BHOYARE^c, AVIJIT MAITI^a, ABHISEK KAR^a, SUMAN^a, ANKU GUHA^a, PALLAVI THAKUR^a, RAMAKIRUSHNAN SURIYA NARAYANAN^a, NITIN T PATIL^{c,*}, CAROLA SCHULZKE^{b,*}, VADAPALLI CHANDRASEKHAR^{a,d,*} and ANUKUL JANA^{a,*}, ^aTata Institute of Fundamental Research Hyderabad, Gopanpally, Hyderabad 500 107, Telangana, India ^bInstitut für Biochemie, Universität Greifswald, Felix-Hausdorff-Straße 4, 17487 Greifswald, Germany ^cDepartment of Chemistry, Indian Institute of Science Education and Research (IISER)-Bhopal, Bhopal 462 066, Madhya Pradesh, India

^dDepartment of Chemistry, Indian Institute of Technology Kanpur, Kanpur 208 016, Uttar Pradesh, India E-mail: npatil@iiserb.ac.in; carola.schulzke@uni-greifswald.de; vc@iitk.ac.in; ajana@tifrh.res.in

MS received 5 August 2019; revised 7 November 2019; accepted 7 November 2019

Abstract. Phosphate diesters are well known to form intermolecular H-bonded dimeric structures in their solid-state. Recently, we reported 2,6-(CHPh)₂-4-*i*Pr-phenyl substituted phosphate diester exists as H-bonded monomeric molecular structure along with water dimer in the solid-state. Herein we report 2,6-(CHPh)₂-4-*i*Pr-phenyl substituted phosphate diester forms a monomeric molecular structure in the solid-state upon cocrystallization with dimethylformamide, DMF (Me₂NCHO). The -CHO group of DMF simultaneously acts as an H-bond acceptor to P-OH and an H-bond donor to P=O moieties. We also used the alcohols, ROH (R = Me, Et, *i*Pr, and *t*Bu), for crystallisation of 2,6-(CHPh)₂-4-*i*Pr-phenyl substituted phosphate diester. In these instances, solvent-incorporated dimeric structures are found in the solid-state. We also report the syntheses and molecular structures of anionic phosphate diesters of 2,6-(CHPh)₂-4-*i*Pr-phenyl substituted phosphate diester to structures of phosphate diesters based on (–)-menthol, (+)-menthol and (+)/(–)-menthol. These exist as H-bonded dimers in the solid-state.

Keywords. Phosphate monoesters; phosphate diesters; phosphonic acids; phosphinic acids; X-ray diffraction.

1. Introduction

In the solid-state, the existence of an entirely free terminal P(O)(OH) moiety in phosphate monoesters, phosphate diesters, phosphonic acids and phosphinic acids is *not* known.^{1–9} Such compounds are present as H-bonded structures either through intermolecular association or through interaction with solvent molecules through the donor (P–OH) and acceptor (P=O) moieties.^{10–25} These hydrogen bonding interactions in phosphates lead to the formation of various types of aggregates in the solid-state.^{26–28} Structures of these

aggregates depend on the solvent of crystallization and the steric bulk of the substituents.⁶ Aggregation behaviour of *para*-substituted monoarylphosphates have been very well studied and the formation of H-bonded non-porous and porous 3-D framework structures was reported.^{29–32} Recently Murugavel *et al.*,³³ and we³⁴ have shown that phosphate monoesters, 2,6-(CHPh)₂-4-R-C₆H₂O-P(O)(OH)₂ form solvent assisted dimeric molecular structures in the solidstate. The corresponding anionic phosphate monoesters derived from these bulky phosphate monoesters exhibit structural diversity in supramolecular

^{*}For correspondence

Electronic supplementary material: The online version of this article (https://doi.org/10.1007/s12039-019-1712-8) contains supplementary material, which is available to authorized users.

organization based on counter cations³⁵ as well as modulation of the nuclearity of molecular Mg(II)phosphates.³⁶ We further disclosed that the phosphate diester, (2,6-(CHPh)₂-4-*i*Pr-C₆H₂O)₂P(O)OH, encapsulates a water dimer and as a result is present as a monomer in the solid-state.³⁷ In order to examine the influence of solvents on the formation of monomeric or dimeric structures of phosphate diester, (2.6- $(CHPh)_2-4-iPr-C_6H_2O)_2P(O)OH$, we have carried out further studies which are reported here. The syntheses of anionic phosphate diesters were also considered and their solid-state molecular structures examined. Moreover, we report the syntheses and molecular structures of phosphate diesters based on stereochemically distinct (-)-menthol, (+)-menthol and (+)/(-)-menthol.

2. Experimental

2.1 Materials and general procedures

Reactions involving nBuLi and POCl₃ were carried out under an argon atmosphere using standard Schlenk techniques and all remaining reactions were carried out under an open atmosphere inside a fume hood. THF was dried by Innovative Technology solvent purification system. Compounds $1 \cdot \text{Et}_2 \text{O}_{,37}^{,37} 2 \cdot \text{Et}_3 \text{N}^{37}$ and $(RO)_2 POCl^{38}$ with R = (-)-menthol and/or (+)-menthol were prepared according to literature procedures. NMR spectra were recorded on a BrukerNanoBay 300 MHz NMR spectrometer. ¹H and ¹³C{¹H} NMR spectra were referenced to the peaks of residual protons of the deuterated solvent (¹H) or the deuterated solvent itself $({}^{13}C{}^{1}H{})$. ${}^{31}P$ NMR spectra were referenced to external H₃PO₄. FT-IR spectra were recorded on a Bruker-Alpha spectrometer. Melting points were determined in Stuart melting point apparatus SMP10 and are uncorrected. Electrospray ionization mass spectrometry (ESI-MS) spectra were recorded on a Waters-Q-Tof Premier-HAB213 spectrometer.

2.2 Synthesis of 1-4

2.2a Isolation of $1 \cdot DMF$ Slow evaporation of a dimethylformamide (DMF) (0.5 mL) solution of $1 \cdot Et_2O$ (0.100 g) at room temperature leads to an almost quantitative formation of $1 \cdot DMF$.

2.2b Isolation of $1 \cdot \text{MeOH}$ Slow evaporation of a CH₃OH (5 mL) solution of $1 \cdot \text{Et}_2 O$ (0.100 g) at room temperature leads to the almost quantitative formation of $1 \cdot \text{MeOH}$.

2.2c Isolation of $1 \cdot \text{EtOH}$ Slow evaporation of an ethanol (5 mL) solution of $1 \cdot \text{Et}_2 O$ (0.100 g) at room temperature leads to the almost quantitative formation of $1 \cdot \text{EtOH}$.

2.2d Isolation of 1-iPrOH

Method I: Slow evaporation of an *iso* propanol (5 mL) solution of $1 \cdot Et_2O$ (0.100 g) at room temperature leads to the almost quantitative formation of $1 \cdot iPrOH$.

Method II: Slow evaporation of an acetone (1 mL), acetonitrile (1 mL) and *iso* propanol (1 mL) solvent mixture of $1 \cdot Et_2O$ (0.100 g) at room temperature leads to the almost quantitative formation of $1 \cdot iPrOH$.

2.2e Isolation of $1 \cdot tBuOH$ Slow evaporation of an acetone (1 mL), acetonitrile (1 mL) and *tert*butanol (0.5 mL) solvent mixture of $1 \cdot Et_2O$ (0.100 g) at room temperature leads to the almost quantitative formation of $1 \cdot tBuOH$.

2.2f Synthesis of 2·iPr₂NEt; [(iPrAr'O)₂P(O)(O)]⁻·H- $N(Et)iPr_2^+$ To a solution of $1 \cdot Et_2O$ (0.107 g, 0.1 mmol) in acetonitrile (5 mL) was added EtNiPr₂ (0.02 mL, 0.1 mmol) and stirred for 5 h. During this period a colorless residue was formed, which was dissolved by addition of DMF (2 mL) followed by warming. The reaction mixture was filtered and the filtrate kept for crystallization at room temperature affording colorless crystals. Yield: 0.072 g, 64%. M.p.: >250 °C. ¹H NMR (300 MHz, CDCl₃, 25 °C): δ = 11.25 (s, br, 1H, N–H), 6.99–6.82 (m, 40H, Ar-H), 6.65–6.60 (m, 8H, Ar-H + CHPh₂), 2.94 (br, 1H,), 2.66 (sept, 2H,CH(CH₃)₂), 2.48 (br, 1H,), 2.26 (br, 2H, N(CH(CH₃)₂)₂, 1.02–0.93 (m, 27H, CH(CH₃)₂ + $NCH_2CH_3 + N(CH(CH_3)_2)_2)$ ppm. ¹³C{¹H} NMR (75.431 MHz, CDCl₃, 25 °C,): δ = 146.92 (Ar–Cquart), 146.80 (Ar-Cquart), 144.66 (Ar-Cquart), 143.52 (Ar-Cquart), 136.73 (Ar-Cquart), 129.73 (Ar-CH), 129.41 (Ar-CH), 127.91 (Ar-CH), 127.62 (Ar-CH), 125.35 (Ar-CH), 52.46 (N(CH(CH₃)₂)₂), 49.32 (CHPh₂), 41.0 (NCH₂CH₃), 33.32 (CH(CH₃)₂), 23.98 (CH(CH₃)₂), 18.16 (N(CH(CH₃)₂)₂), 17.16 (NCH₂CH₃), 11.14 (NCH₂*C*H₃) ppm. ³¹P{¹H} NMR (121.442 MHz, CDCl₃, 25 °C): $\delta = -11.0$ ppm. FT-IR (KBr pellet, cm⁻¹): $\bar{v} = 3420$ (w, br), 3082 (w), 3057 (m), 3024 (m), 2957 (s), 2870 (w), 2387 (w), 2299 (w), 1943 (w), 1888 (w), 1801 (w), 1599 (m), 1493 (s), 1465 (s), 1445 (s), 1399 (w), 1384 (w), 1362 (w), 1318 (w), 1292 (w), 1244 (s), 1205 (m), 1177 (w), 1159 (m), 1120 (m), 1083 (s), 1031 (m), 920 (m), 896 (s), 850 (m), 829 (w), 796 (w), 763 (m), 744 (w), 721 (s), 698 (s), 666 (w), 646 (w), 628 (w), 606 (m), 580 (w), 516 (w).

2.2g Synthesis of $2 \cdot \text{NHC}^{i\text{Pr2Me2}}$; $[(i\text{PrAr'O})_{2} - P(O)(O)]^{-1} \cdot H - NHC^{i\text{Pr2Me2}+1}$ In a 100 mL Schlenk

flask, 1·Et₂O (0.072 g, 0.07 mmol) and NHC^{iPr2Me2} (0.013 g, 0.07 mmol) were mixed. Then dry THF (20 mL) was added and the resulting reaction mixture was stirred for 5 h at room temperature. On evaporation of all volatiles followed by washing with diethyl ether (2 x 5 mL), a colorless solid was obtained as desired product. Single crystals suitable for X-ray diffraction studies were grown by slow evaporation of acetonitrile solution of 2.NHC^{iPr2Me2} at room temperature. Yield: 0.048 g, 59%. M.p.: >250 °C. ¹H NMR (25 °C, 300 MHz, CDCl₃): $\delta = 10.54$ (s, 1H, N– CH-N), 7.06–6.71 (m, 43H, Ar-H + CHPh₂), 6.61 (s, 4H, Ar-H), 5.91 (s, 1H, CHPh₂), 4.30 (sept, 2H, (N-CH(CH₃)₂), 2.65 (sept, 2H, CH(CH₃)₂), 2.16 (s, 6H, $C(4,5)-CH_3$, 1.29 (d, 12H, (N-CH(CH_3)_2), 1.01 (d, 12H, CH(CH₃)₂) ppm. ¹³C{¹H} NMR (25 °C, 75.431 MHz, CDCl₃): $\delta = 144.95$ (Ar– C_{quart}), 142.44 (Ar– C_{quart}), 137.30 (Ar-C_{quart}), 137.26 (N-CH-N), 134.89 $(Ar-C_{quart}), 129.80 (Ar-C_{quart}), 127.38 (Ar-CH),$ 126.83 (Ar-CH), 125.23 [C(4,5), Im], 125.03 (Ar-CH), 51.36 (CHPh₂), 49.86 (N-CH(CH₃)₂), 33.36 (CH(CH₃)₂), 24.08 (N–CH(CH₃)₂), 22.36 (CH(CH₃)₂), 8.76 $[C(4,5)-CH_3]$ ppm. ³¹P{¹H} NMR (25 °C, 121.442 MHz, CDCl₃): $\delta = -9.2$ ppm. FT-IR (KBr pellet, cm⁻¹): $\bar{v} = 3421$ (m, br), 3117 (w), 3081 (w), 3057 (m), 3024 (m), 2958 (s), 2865 (m), 2801 (w), 1947 (w), 1884 (w), 1807 (w), 1629 (w), 1599 (m), 1581 (w), 1555 (m), 1493 (s), 1464 (s), 1445 (s), 1393 (w), 1379 (w), 1359 (w), 1319 (w), 1289 (w), 1249 (s), 1202 (m), 1156 (m), 1118 (s), 1096 (s), 1030 (m), 1002 (w), 986 (w), 967 (w), 916 (w), 890 (m), 878 (s), 857 (w), 841 (s), 794 (w), 764 (m), 719 (s), 701 (s), 659 (s), 633 (w), 623 (w), 606 (w), 598 (m), 588 (m), 574 (m), 546 (m). ESI-MS: Calcd (m/z) for $[C_{11}H_{21}N_2^+]$: 181.1699; found: 181.1613.

Synthesis of 3; $[(iPrAr'O)_2P(O)(O)]_2^{2-}$. 2.2h $Zn(MeOH)_4(H_2O)_2^{2+}$ To a solution of $2 \cdot Et_3N$ (0.031 g, 0.028 mmol) in methanol (5 mL) was added a solution of ZnCl₂ (0.004 g, 0.028 mmol) in methanol (5 mL) with stirring followed by addition of DMF (0.03 mL) at room temperature. The resulting reaction mixture was heated at 70 °C for 5 h and during this period a colorless residue was formed which was filtered-off and the filtrate kept for crystallization. Colorless crystals suitable for X-ray diffraction studies were obtained by slow evaporation at room temperature. Yield: 0.012 g, 38%. M.p.: 224–228 °C. ¹H NMR (300 MHz, CDCl₃, 25 °C): $\delta = 6.78-6.98$ (m, 40H, Ar-H), 6.62 (s, 4H, Ar-H), 6.32 (s, 4H, CHPh₂), 2.57–2.71 (m, 16H. CH(CH₃)₂+4CH₃OH), 1.01 (d, 12H, CH(CH₃)₂), 0.83 (s, 4H, H_2 O) ppm. ¹³C{¹H} NMR (75.431 MHz, CDCl₃, 25 °C): $\delta = 143.69$ (Ar–C_{quart}), 129.59

(Ar–CH), 127.85 (Ar–CH), 127.43 (Ar–CH), 125.75 (Ar–CH), 49.76 (CHPh₂), 45.54 (CH₃OH), 33.37 (CH(CH₃)₂), 23.93 (CH(CH₃)₂) ppm. ³¹P{¹H} NMR (121.442 MHz, CDCl₃, 25 °C): δ = -10.6 ppm. FT-IR (KBr pellet, cm⁻¹): \bar{v} = 3565 (s, br), 3083 (w), 3059 (m), 3026 (m), 2958 (s), 2869 (w), 1948 (w), 1885 (w), 1804 (w), 1657 (m), 1599 (m), 1493 (s), 1466 (m), 1445 (s), 1383 (w), 1363 (w), 1319 (w), 1291 (w), 1258 (m), 1200 (s), 1157 (m), 1117 (s), 1078 (s), 1031 (m), 1003 (w), 966 (w), 925 (s), 856 (m), 828 (w), 797 (w), 761 (m), 746 (w), 722 (s), 698 (s), 672 (w), 647 (w), 633 (w), 621 (w), 605 (m), 584 (w), 544 (w).

2.2i Synthesis of 4a; (-) $Mt_2P(O)OH$ (-)-Menthol (5.0 g, 32 mmol) was dissolved in 10 mL of THF; an equal volume of diethyl ether was added and the mixture was cooled to -78 °C. The dropwise addition of nBuLi (1.6 M in hexane, 20 mL, 32 mmol) resulted in the formation of a colorless suspension. After the mixture was stirred for an additional 10 min at -78 °C, POCl₃ (1.5 mL, 16 mmol) was slowly added and the solution was warmed to ambient temperature and stirred overnight. All volatiles were removed under vacuum affording a colorless oily compound with LiCl. The resulting reaction mixture was dissolved in a mixture of CH₃CN/water (40 mL/10 mL) and refluxed for 6 h. During this period a colorless clear solution was formed, which was kept for two days and led to the formation of the colorless crystalline compound. This colorless crystalline compound was filtered followed by washing with acetonitrile and dried under vacuum at 50 °C for 3 h to give 4a as colorless compound. Single crystals suitable for X-ray diffraction studies were grown by slow evaporation from a CH₃CN/water (4/1) mixture solution of 4a at room temperature. Yield: 2.85 g, 48%. M.p.: 102-104 °C. ¹H NMR (25 °C, 300 MHz, CDCl₃): $\delta = 8.78$ (s, br, 1H, P-OH), 4.10-4.06 (m, 2H, H1), 2.27-2.15 (m, 4H), 1.67–1.63 (m, 4H), 1.40–1.29 (m, 4H), 1.20– 1.12 (m, 2H), 1.0-0.97 (m, 2H), 0.92-0.88 (m, 14H), 0.81–0.79 (m, 6H) ppm. ¹³C{¹H} NMR (25 °C, 75.431 MHz, CDCl₃): δ = 78.99 (C1), 78.90 (C1), 48.56, 48.47, 42.39 (C6), 34.09 (CH₂), 31.61, 30.95, 25.42 (CH(CH₃)₂), 22.89 (CH₂), 22.01 (CH₃), 20.97 (CH₃), 15.74 (CH₃) ppm. ³¹P{¹H} NMR (25 °C, 121.442 MHz, CDCl₃): $\delta = 0.55$ ppm. FT-IR (KBr pellet, cm⁻¹): $\bar{v} = 3417$ (m, br), 2954 (s), 2927 (s), 2870 (m), 2722 (w), 2385 (w), 1639 (w), 1456 (s), 1387 (m), 1371 (m), 1349 (w), 1232 (w), 1180 (m), 1154 (m), 1061 (s), 1014 (s), 977 (s), 928 (m), 898 (m), 835 (m), 818 (w), 801 (w), 594 (w), 566 (w), 552 (m). ESI-MS: Calcd (m/z) for $[C_{20}H_{39}O_4P + Na^+]^+$: 397.2478; found: 397.2489.

2.2j Synthesis of 4b; (+)Mt₂P(O)OH Compound 4b was prepared starting from (+)-menthol as colorless crystals following the same procedure as described above for the synthesis of **4a**. Quantities: (+)-menthol (5.0 g, 32 mmol), nBuLi (1.6 M in hexane, 20 mL, 32 mmol), POCl₃ (1.5 mL, 16 mmol), THF (10 mL), diethyl ether (10 mL). Single crystals suitable for Xray diffraction studies were grown by slow evaporation from a CH₃CN/water (4/1) mixture solution of 4b at room temperature. Yield: 2.72 g, 45%. M.p.: 108-110 °C. ¹H NMR (25 °C, 300 MHz, CDCl₃): $\delta = 9.24$ (s, br, 1H, P-OH), 4.10-4.06 (m, 2H, H1), 2.27-2.15 (m, 4H), 1.67–1.63 (m, 4H), 1.39–1.29 (m, 4H), 1.20– 1.08 (m, 2H), 1.0-0.97 (m, 2H), 0.92-0.88 (m, 14H), 0.81–0.79 (m, 6H) ppm. ¹³C{¹H} NMR (25 °C, 75.431 MHz, CDCl₃): δ = 78.98 (C1), 78.90 (C1), 48.57, 48.47, 42.40 (C6), 34.10 (CH₂), 31.61, 25.43 (CH(CH₃)₂), 22.90 (CH₂), 22.0 (CH₃), 20.96 (CH₃), 15.75 (CH₃) ppm. ³¹P{¹H} NMR (25 °C, 121.442) MHz, CDCl₃): $\delta = 0.55$ ppm. FT-IR (KBr pellet, cm^{-1}): $\bar{v} = 3420$ (m, br), 2954 (s), 2927 (s), 2870 (m), 2722 (w), 2385 (w), 1654 (m), 1456 (s), 1387 (m), 1371 (m), 1349 (w), 1231 (s), 1180 (m), 1154 (m), 1061 (s), 1014 (s), 977 (s), 928 (m), 898 (m), 835 (m), 818 (w), 801 (w), 772 (w), 594 (w), 566 (s), 552 (s), 506 (w). ESI-MS: Calcd (m/z) for $[C_{20}H_{39}O_4P +$ Na⁺]⁺: 397.2478; found: 397.2489.

2.2k Synthesis of 4c; (+)-Mt(-)-MtP(O)OH nBuLi (1.6 M in hexane, 20 mL, 32 mmol) was added dropwise to a solution of (+)-menthol (5.0 g, 32 mmol, in 10 mL of dry diethyl ether) at -78 °C resulting in the formation of a colorless suspension. Afterwards it was stirred for an additional 10 min at -78 °C, the suspension was added dropwise to a solution containing POCl₃ (9 mL, 96 mmol, in 30 mL of dry diethyl ether) at -78 °C and stirring was continued for overnight. All volatiles were removed under vacuum affording a colorless oily compound of RO-POCl₂; R = (+)-Mt with LiCl. In another Schlenk flask, nBuLi (1.6 M in hexane, 20 mL, 32 mmol) was added dropwise to a solution of (-)-menthol (5.0 g, 32 mmol, in 10 mL of dry diethyl ether) at -78 °C. After completion of addition, the reaction mixture was stirred for an additional 10 min at -78 °C resulting in the formation of a colorless suspension of $R'O^-Li^+$; R = (-)-Mt. This colorless suspension of R'O⁻Li⁺ was added to a stirred solution of RO-POCl₂ in THF (20 mL) at -78 °C by cannula within 1 h. Then the reaction mixture was slowly allowed to reach room temperature and stirring was continued for overnight. During this period a colorless suspension was formed, and then the volatiles were removed under vacuum affording a colorless oily compound of (RO)(R'O)POC1 with LiCl. The resulting reaction mixture was dissolved in a mixture of CH₃CN/water (80 mL/20 mL) and refluxed for 6 h. During this period a colorless clear solution was formed, which was kept for two days and lead to the formation of a colorless crystalline compound. This colorless crystalline compound was filtered followed by washing with acetonitrile and drying under vacuum at 50 °C for 3 h to give 4c as colorless compound. Single crystals suitable for X-ray diffraction studies were grown by slow evaporation from a CH₃CN/water (4/1) mixture solution of **4c** at room temperature. Yield: 1.92 g, 16%. M.p.: 118–120 °C. ¹H NMR (25 °C, 300 MHz, CDCl3): δ = 8.75 (s, br, 1H, P– OH), 4.15–4.04 (m, 2H, H1), 2.27–2.13 (m, 4H), 1.67– 1.63 (m, 4H), 1.43-1.29 (m, 4H), 1.20-1.08 (m. 2H). 1.01-0.97 (m, 2H), 0.93-0.89 (m, 14H), 0.82-0.79 (m, 6H) ppm. ¹³C{¹H} NMR (25 °C, 75.431 MHz, CDCl₃): $\delta = 79.09$ (C1), 79.0 (C1), 48.52, 48.43, 42.40 (C6), 34.09 (CH₂), 31.6, 25.42 (CH(CH₃)₂), 22.92 (CH₂), 22.02 (CH₃), 20.96 (CH₃), 15.76 (CH₃) ppm. ${}^{31}P{}^{1}H{}$ NMR (25 °C, 121.442 MHz, CDCl₃): $\delta = 0.53$ ppm. FT-IR (KBr pellet, cm⁻¹): $\bar{v} = 3414$ (m, br), 2954 (s), 2925 (s), 2868 (s), 2656 (w), 2383 (w), 1684 (w), 1453 (s), 1385 (m), 1368 (m), 1346 (w), 1219 (s), 1180 (m), 1159 (m), 1079 (m), 1061 (s), 1020 (s), 976 (s), 927 (m), 896 (m), 835 (m), 814 (w), 799 (w), 771 (w), 593 (w), 557 (m), 541 (m), 506 (w). ESI-MS: Calcd (m/z) for $[C_{20}H_{30}O_4P + Na^+]^+$: 397.2478; found: 397.2487.

2.3 X-ray crystallography

Single-crystal X-ray diffraction data of 1.DMF, 1. MeOH, 1. iPrOH (solvent of crystallization: acetone, acetonitrile and iso-propanol), 1.tBuOH, 2.*i*Pr₂NEt, 2.NHC^{*i*Pr₂Me²}, 3, 4a, 4b and 4c were collected on a Rigaku XtaLAB X-ray Diffractometer system equipped with a CCD area detector and operated at 30 W power (50 kV, 0.6 mA) to generate Mo K\alpha radiation ($\lambda = 0.71073$ Å) at 120(2) K. Singlecrystal X-ray diffraction data of 1. EtOH and 1. iPrOH (solvent of crystallization: iso-propanol) were collected at low temperature (170 K) on a STOE-IPDS II diffractometer equipped with a normal-focus, 2.4 kW, sealed tube X-Ray source with graphite-mono-chromated Mo K α radiation ($\lambda = 0.71073$ Å). The integration of diffraction profiles was performed with the program XArea; numerical absorption correction was carried out with the programs X-shape and X-red32: all from $STOE^{\odot}$. The structures were solved with the SHELXT structure solution program using Intrinsic Phasing and refined with the SHELXL refinement package using Least Squares minimisation in the Olex-2 software.^{39–43} All non-hydrogen atoms were refined with anisotropic thermal parameters. All hydrogen atoms were placed in geometrically calculated positions or found in the Fourier difference map and included in the refinement process using a riding model. Crystal data and structure refinement of all these compounds are given in Tables S5 and S6 (Supplementary Information).

3. Results and Discussion

We have synthesized $1 \cdot Et_2O$ according to the procedure shown in Scheme $1.^{37}$ Subsequently, compound $1 \cdot Et_2O$ was crystallized using different solvents capable of hydrogen bonding such as DMF, CH₃OH, EtOH, and *i*PrOH as well as from the mixtures of *i*PrOH/CH₃COCH₃/CH₃CN and *t*BuOH/CH₃COCH₃/ CH₃CN solvents to study the H-bonding interaction. Crystallization of the phosphate diester in CH₃-COCH₃, CH₂Cl₂, CHCl₃ or DMSO did not succeed. Analysis of the single crystals of $1 \cdot DMF$ obtained from DMF indicates two types of H-bond between P– O–H…O=CH and P=O…H–C=O moieties. This results in the stabilization of the monomer in the solid-state. On the other hand, analysis of crystals obtained from CH₃OH, EtOH, *i*PrOH/CH₃COCH₃/CH₃CN indicated the formation of an alcohol-assisted dimer (Figure 1). The important bond distances and angles of these compounds are given in Tables S1 and S2 (Supplementary Information).

The molecular structures of the phosphate diesters. 1.DMF, 1.MeOH, 1.EtOH and 1.iPrOH (solvent of crystallization: acetone, acetonitrile and iso-propanol) reveal three types of P-O bond distances (Tables S1 and S2, Supplementary Information). The shortest distance is 1.469(3) Å (1.DMF), 1.471(11) Å (1·MeOH), 1.471(4) Å (1·EtOH) and 1.473(11) Å (1.*i*PrOH) which is consistent with the P=O distance of similar compounds. A total of six O-P-O angles found in each structure: 109.05 (17)°, were 107.49(16)°, 95.06(15)°, 113.30(17)°, 115.20(17)°, $114.82(17)^{\circ}$ $(1 \cdot \mathbf{DMF}).$ 103.64(6)°. 105.31(6)°. 103.0(6)°, 112.01(6)°, 114.57(6)°, 116.95(7)° (1·MeOH); 116.95(4)°, 112.01(2)°, 114.28(23)°, 105.71(24)°, 103.14(3)°, 103.40(33)° (1·EtOH) and 103.45(6)°, 102.78(6)°, 115.12(6)°, 105.67(6)°, $112.22(6)^{\circ}$, $116.21(7)^{\circ}$ (**1**·*i***PrOH**). In all these instances, the central prochiral phosphorus centre is present in a distorted tetrahedral geometry. The



Scheme 1. Synthesis of 1.DMF, 1.MeOH, 1.EtOH, 1.iPrOH and 1.tBuOH.



Figure 1. H-bonded molecular structures of (a) 1·DMF, (b) 1·MeOH, (c) 1·EtOH and (d) 1·*i*PrOH; all hydrogen atoms except O–H, C–H of DMF and one disordered solvent molecule in 1·DMF were omitted for clarity reasons and thermal ellipsoids represent the 50% probability level in all cases.

molecular structure of 1.DMF shows a non-planar seven-membered ring (Figure 1), and in the case of 1.MeOH, 1.EtOH and 1.iPrOH non-planar twelvemembered rings (Figure 1). The crystal structure of 1.DMF also shows two types of intermolecular $(P-OH\cdots O=C = 1.843(7) \text{ Å}$ hydrogen-bonds 1 $P=O\cdots H-C = 2.733 \text{ Å}$) with an $O\cdots O$ separation of only 2.418(4) Å. On the other hand, crystal structures of 1.MeOH, 1.EtOH and 1.iPrOH reveal intermolecular hydrogen-bonding between one molecule of phosphate diester and one molecule of alcohol (P-OH···OH-R/P=O···H-OR) via two short O-H···O [P-OH-OH-R = 1.584(3)]Å bonds $(1 \cdot MeOH),$ 1.532(7) Å (1·EtOH) and 1.51(3) Å (1·*i*PrOH); $P=O\cdots H-OR = 1.75(3) \text{ Å}$ (1·MeOH), 1.762(7) Å (1.EtOH) and 1.91 Å (1.iPrOH)] with O...O separations of 2.435(16) Å/2.671(16) Å (1·MeOH), 2.433(11) Å/2.652(10) Å (1·EtOH) and 2.413(18) Å/

2.703(17)Å (1·*i***PrOH**). 1·*i***PrOH** when crystallized in pure *iso*-propanol solvent results in severe disorder in the *iso*-propanol and therefore the H-bonding interactions are not discussed. The metric parameters of 1·*i***PrOH** and 1·*t***BuOH** are given in Table S2, Supplementary Information.

Previously we reported the reaction of $1 \cdot \text{Et}_2 O$ with Et₃N affording anionic phosphate diester, $2 \cdot \text{Et}_3 N$.³⁷ Subsequently, we also performed related reactions of $1 \cdot \text{Et}_2 O$ with one equivalent of $i Pr_2 NEt$ or $NHC^{iPr2Me2}$, resulting in the formation of colourless compounds $2 \cdot i Pr_2 NEt$ and $2 \cdot NHC^{iPr2Me2}$, respectively. The characterization of $2 \cdot i Pr_2 NEt$ and $2 \cdot NHC^{iPr2Me2}$ in solution was carried out by NMR and ESI-MS methods. The most prominent features were the appearance of resonances of N–H in $2 \cdot i Pr_2 NEt$ or C2–H in $2 \cdot NHC^{iPr2Me2}$ in the deshielded region (11.25 ppm in CDCl₃ for $2 \cdot i Pr_2 NEt$ and 10.54 ppm in CDCl₃ for



Scheme 2. Synthesis of $2 \cdot i Pr_2 NEt$ and $2 \cdot NHC^{iPr2Me2}$.

Table 1. ${}^{31}P{}^{1}H$ NMR and selected ${}^{1}H$ NMR data of newly synthesized compounds.

Compound	$\delta^{31} P{^1H} NMR$ (ppm)	δ ¹ HNMR of P-OH (ppm)
2· <i>i</i> Pr ₂ NEt 2·NHC ^{<i>i</i>Pr2Me2} 3	-11.0 -9.2 -10.6	
4a 4b 4c	0.55 0.55 0.53	8.78 9.24 8.75

All NMR spectra were measured using CDCl₃.

2·**NHC**^{*i***Pr2Me2**}) (Scheme 2, Figures S1 and S4, Supplementary Information). The ³¹P{¹H} solution NMR spectra of **2**·*i***Pr₂NEt** or **2**·**NHC**^{*i***Pr2Me2**} show each a singlet at $\delta = -11.0$ or -9.2ppm suggesting the

presence of only one type of phosphorus centre in the dissolved state (Table 1). The base peak in the ESI-MS spectrum for $2 \cdot \text{NHC}^{i\text{Pr2Me2}}$ recorded in positive ion mode corresponds to the imidazolium cation. Furthermore, $2 \cdot i\text{Pr}_2\text{NEt}$ and $2 \cdot \text{NHC}^{i\text{Pr2Me2}}$ were characterized in their solid-state by single-crystal X-ray diffraction analysis.

The asymmetric units of $2 \cdot i Pr_2 NEt$ and $2 \cdot NHC^{iPr2Me2}$ are shown in Figure 2. The crystallographic parameters for these compounds are given in Table S6 (Supplementary Information). The negative charge of the phosphate diester ligand in $2 \cdot i Pr_2 NEt$ and $2 \cdot NHC^{iPr2Me2}$ is delocalized as confirmed from the P–O distances [P1–O1 1.622(7), P1–O2 1.618(6), P1–O3 1.490(7), P1–O4 1.467(3) in $2 \cdot i Pr_2 NEt$ and P1–O1 1.625(1), P1–O2 1.628(6), P1–O3 1.468(5), P1–O4 1.486(3) in $2 \cdot NHC^{iPr2Me2}$]. The phosphorus atom exhibits a distorted tetrahedral geometry with the



Figure 2. Molecular structures of $2 \cdot i \Pr_2 NEt$ and $2 \cdot NHC^{i \Pr_2 Me2}$. Thermal ellipsoids are at the 50% probability level (hydrogen atoms except N–H for $2 \cdot i \Pr_2 NEt$ and C71–H71 for $2 \cdot NHC^{i \Pr_2 Me2}$ are omitted for clarity).



Scheme 3. Synthesis of 3.

angles varying from $98.09(9)^{\circ}-119.37(11)^{\circ}$ in $2 \cdot i Pr_2$. **NEt** and $93.84(8)^{\circ}-118.47(10)^{\circ}$ in $2 \cdot NHC^{iPr2Me2}$ (Table S3, Supplementary Information). Ethyl diisopropyl ammonium ($2 \cdot i Pr_2 NEt$) and imidazolium ($2 \cdot NHC^{iPr2Me2}$) cations are involved in strong hydrogen bonding interaction with the anionic phosphate diester [N1-H1N···O3, 1.61(3) Å in $2 \cdot i Pr_2 NEt$ and C71-H71·O4 2.00(3) Å in $2 \cdot NHC^{iPr2Me2}$].

Subsequently, we were interested to obtain zincphosphate diester complex using $2 \cdot \text{Et}_3 \text{N}$ and ZnCl_2 . However, the reaction of $2 \cdot \text{Et}_3 \text{N}$ with ZnCl_2 in methanol afforded the ion-pair **3** (Scheme 3). The ³¹P{¹H} NMR spectrum of **3** revealed the presence of a singlet at $\delta = -10.6$ ppm suggesting the presence of only one type of phosphorus centre in the dissolved state (Table 1). In comparison to the ion pair $2 \cdot \text{Et}_3$. N ($\delta = -9.8$), the chemical shift for **3** appears in the upfield region.

The solid-state structure of 3 is shown in Figures 3and S28, Supplementary Information. The crystallographic parameters are given in Table S6 (Supplementary Information). The asymmetric unit contains two anionic diester phosphates and the cationic hexacoordinated Zn(II) complex, $[Zn(H_2O)_2(CH_3OH)_4]^{2+}$. The negative charge of the phosphate diester in 3 is delocalized as confirmed from the P-O distances [P1-O1 1.611(3), P1-O2 1.603(2), P1-O3 1.486(3), P1-O4 1.491(3) in **3**] and the phosphorous atom is present in a distorted tetrahedral geometry with angles around phosphorus varying from 100.59(13)^o-117.23(15)° (Table S3, Supplementary Information). The coordinating methanol and methoxide ligands in the hexa-coordinated cationic moiety [Zn(H₂O)₂(- $(CH_3OH)_4]^{2+}$ and, however, disordered in two crystallographic special positions (Figure 3; only one part is shown) and therefore the metric parameters are not discussed.

Subsequently, in order to explore the assembly of chiral phosphate ligands, menthol-based compounds **4a**, **4b** and **4c** were prepared (Scheme 4). While hydrolysis of the symmetric phosphoryl dichlorides afforded **4a** and **4b**, the corresponding hydrolysis of the unsymmetrical derivative afforded **4c** (Scheme 4).



Figure 3. Molecular structure of **3** revealing the hydrogen bonding interactions. Thermal ellipsoids are at the 50% probability level (hydrogen atoms except O–H are omitted for clarity).

The ¹HNMR spectra of **4a**, **4b** and **4c** reveal the presence of a broad singlet in each corresponding to POH at $\delta = 8.78$, 9.24 and 8.75 ppm, respectively (Table 1). The ³¹P{¹H}NMR spectra of **4a**, **4b** and **4c** revealed singlets at $\delta = 0.55$, 0.55 and 0.53 ppm, respectively, which are upfield shifted compared to the resonances observed in the corresponding phosphoryldichlorides (δ : 3.3 ppm) (Table 1). Single crystals of **4a**, **4b**, and **4c** were obtained by slow evaporation from CH₃CN/water mixture solutions and were subjected to X-ray diffraction studies (Figure 4, Figures S29–S30, Supplementary Information).

The important bond distances and angles of **4a**, **4b** and **4c** are given in Table S4, Supplementary Information. The molecular structures of **4a**, **4b** and **4c**, show three types of P–O bond distances (Table S4, Supplementary Information). The shortest distance is 1.492(5) Å (**4a**), 1.489(4) Å (**4b**) and 1.495(1) Å (**4c**), which is consistent with the P=O distance of known similar compounds. A total of six O–P–O angles were found in each structure: [101.9(3), 109.3(3), 110.4(3), 111.5(3), 107.2(3), 115.6(3) and 111.3(4) (**4a**)], [102.0(2), 106.9(3), 108.9(2), 116.0(3), 111.5(3) and 110.6(3) (**4b**)] and [103.40(6), 108.70(7), 112.26(6), 115.31(7), 109.50(6) and 106.99(7) (**6c**). The molecular structures of **4a** and **4c** reveal a non-planar eight-



Scheme 4. Syntheses of 4a, 4b and 4c.



Figure 4. Molecular structure of **4a** showing the hydrogen bonding interactions in the crystal. Thermal ellipsoids are at the 50% probability level (hydrogen atoms except O– H are omitted for clarity).

membered ring (Figures 4 and S30, Supplementary Information) as a result of an intermolecular hydrogen-bonding network between the two molecules of phosphate diesters (P=O/P–OH) *via* two short O–H···O bonds [P=O···HO–P = {1.611(8) Å and 1.700(5) Å (4a)} and 1.779(3) Å (4c)] with a O···O separation of {2.472(5) Å and 2.527(3) Å (4a)}, and 2.533(1) Å (4c)]. In the case of 4b, these hydrogen-bonding interactions are missing; this can be attributed to the orientation of the hydroxide moiety of the P–OH group which leads to the monomeric structure in the solid state.

4. Conclusions

In summary, we report the monomeric solid-state molecular structure of 2,6-(CHPh)₂-4-*i*Pr-phenyl substituted phosphate diester as crystallized from DMF solvent. With alcohols as the crystallizing solvents, solvent-bridged dimeric structures are the result. The reactions of phosphate diesters with bases such as amine and NHC afforded ion-pairs which are present as monomers in the solid-state. We also obtained the anionic phosphate diester with a cationic Zn(II) complex as the counter cation. In addition, we have also synthesized phosphate diesters containing (-)-menthol, (+)-menthol and (+)/(-)-menthol substituents.

Supplementary Information (SI)

CCDC 1892137–1892148 contains the supplementary crystallographic data for 1·DMF, 1·MeOH, 1·EtOH, 1·iPrOH (Solvent of crystallization: *iso*-propanol), 1·iPrOH (Solvent of crystallization: acetone, acetonitrile and *iso*-propanol), 1·tBuOH (Solvent of crystallization: acetone, acetonitrile and *tert*-butanol), 2·iPr₂NEt and 2·NHC^{iPr2Me2}, 3, 4a, 4b, and 4c. These data can be obtained free of charge *via* http://www.ccdc.cam.ac.uk/ conts/retrieving.html, or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: (+44) 1223-336-033; or e-mail: deposit@ccdc.-cam.ac.uk. Supplementary data associated with this article is available at www.ias.ac.in/chemsci.

Acknowledgements

This work is supported by the TIFR Centre for Interdisciplinary Science Hyderabad, Hyderabad, India. V. G. acknowledges financial support from the Science and Engineering Research Board (SERB-NPDF) (Project No. PDF/2017/001902), Government of India. VC is thankful to the Department of Science and Technology, New Delhi, India, for National J. C. Bose fellowships.

References

1. Murugavel R, Choudhury A, Walawalkar M G, Pothiraja R and Rao C N R 2008 Metal Complexes of Organophosphate Esters and Open-Framework Metal Phosphates: Synthesis, Structure, Transformations, and Applications *Chem. Rev.* **108** 3549

- Roesky H W, Walawalkar M G and Murugavel R 2001 Is Water a Friend or Foe in Organometallic Chemistry? The Case of Group 13 Organometallic Compounds Acc. Chem. Res. 34 201
- 3. Murugavel R, Walawalkar M G, Dan M, Roesky H W and Rao C N R 2004 Transformations of Molecules and Secondary Building Units to Materials: A Bottom-Up Approach *Acc. Chem. Res.* **37** 763
- Guillou N, Gao Q, Forster P M, Chang J-S, Nogus M, Park S-E, Frey G and Cheetham 2001 A K, Nickel(ii) Phosphate VSB-5: A Magnetic Nanoporous Hydrogenation Catalyst with 24-Ring Tunnels Angew. Chem. 113 2913
- 5. Walawalkar M G, Roesky H W and Murugavel R 1999 Molecular Phosphonate Cages: Model Compounds and Starting Materials for Phosphate Materials *Acc. Chem. Res.* **32** 117
- 6. Goura J and Chandrasekhar V 2015 Molecular Metal Phosphonates *Chem. Rev.* **115** 6854
- Murugavel R, Kuppuswamy S, Gogoi N and Steiner A 2010 Assembling Discrete D4R Zeolite SBUs through Noncovalent Interactions. 3. Mediation by Butanols and 1,2-Bis(dimethylamino)ethane *Inorg. Chem.* 49 2153
- Murugavel R, Kuppuswamy S, Maity A N and Singh M P 2009 Di-, Tri-, Tetra-, and Hexanuclear Copper(II) Mono-organophosphates: Structure and Nuclearity Dependence on the Choice of Phosphorus Substituents and Auxiliary N-Donor Ligands *Inorg. Chem.* 48 183
- 9. Murugavel R and Shanmugan S 2008 Asymmetric Pentameric and Tetrameric Organooxotin Clusters: Insights into Their Formation through Partial Dearylation *Organometallics* **27** 2784
- 10. Cheng W, Feng Z-Q and Tang J-M 2011 Bis(3,5dimethoxyphenyl)phosphinic acid Acta Crystallogr., Sect. E 67 0896
- 11. Székely G, Farkas V, Párkányi L, Tóth T, Hollósi M. and Huszthy P 2010 Crystal structures of crown ethers containing an alkyl diarylphosphinate or a diarylphosphinic acid unit *Struct. Chem.* **21** 277
- Lyssenko K A, Grintselev-Knyazev G V and Antipin M Y 2002 Nature of the P–O bond in diphenylphosphonic acid: experimental charge density and electron localization function analysis *Mendeleev Commun.* 12 128
- Fenske D, Mattes R, Löns J and Tebbe K-F 1973 Die Kristallstruktur von Diphenylphosphinsäure Chem. Ber. 106 1139
- Xu B, Zhu S-F, Zhang Z-C, Yu Z-X, Ma Y. and Zhou Q-L 2014 Highly enantioselective S–H bond insertion cooperatively catalyzed by dirhodium complexes and chiral spiro phosphoric acids *Chem. Sci.* 5 1442
- 15. Nayak S K, Chandrasekhar S and Row T N G 2008 1,1'-Bi-naphthalene-2,2'-diyl hydrogen phosphate *Acta Crystallogr., Sect. E* **64** 0256
- 16. Swamy K C K, Kumaraswamy S and Kommana P 2001 Very Strong C-H···O, N-H···O, and O-H···O Hydrogen Bonds Involving a Cyclic Phosphate J. Am. Chem. Soc. **123** 12642
- 17. Klussmann M, Ratjen L, Hoffmann S, Wakchaure V, Goddard R and List B 2010 Synthesis of TRIP and Analysis of Phosphate Salt Impurities *Synth. Lett.* 2189

- DeFord J, Chu F and Anslin E V 1996 Dimerization constants for phosphoric acid diesters *Tetrahedron Lett.* 37 1925
- Peppard D F, Ferraro J R and Mason G W 1957 Possible hydrogen bonding in certain interactions of organic phosphorus compounds *J. Inorg. Nucl. Chem.* 4 371
- 20. Peppard D F, Ferraro J R and Mason G W 1958 Hydrogen bonding in organophosphoric acids J. Inorg. Nucl. Chem. 7 231
- Dar A A, Mallick A and Murugavel R 2015 Synthetic strategies to achieve further-functionalised monoaryl phosphate primary building units: crystal structures and solid-state aggregation behaviour *New J. Chem.* **39** 1186
- 22. Kalita A C, Sharma K and Murugavel R 2014 Pseudopolymorphism leading and two different supramolecular aggregations in a phosphate monoester: role of a rare water-dimer *CrystEngComm* **16** 51
- 23. Kuczek M, Bryndal I. and Lis T 2006 4-Nitrophenyl phosphoric acid and its four different potassium salts: a solid state structure and kinetic study *CrystEngComm* **8** 150
- 24. Onoda A, Yamada Y, Okamura T-A, Yamamoto H and Ueyama N 2002 Mononuclear Ca(II)-Bulky Aryl-Phosphate Monoanion and Dianion Complexes with Ortho-Amide Groups *Inorg. Chem.* **41** 6038
- 25. Onoda A, Okamura T, Yamamoto H and Ueyama N 2001One-dimensional P–OH…O=P hydrogen bonds restricted by the bulky molecule 2,6-diiso-propyl-phe-nyl di-hydrogen phosphate *Acta Crystallogr., Sect. E* o1022
- 26. Mehring M, Schurmann M and Ludwig R 2003 *tert*-Butylphosphonic Acid: From the Bulk to the Gas Phase *Chem. Eur. J.* **9** 837
- 27. Chandrasekhar V, Sasikumar P, Boomishankar R and Anantharaman G 2006 Assembly of Lipophilic Tetranuclear (Cu₄ and Zn₄) Molecular Metallophosphonates from 2,4,6-Triisopropylphenylphosponic Acid and Pyrazole Ligands *Inorg. Chem.* **45** 3344
- 28. Weakley T J R 1976 Benzenephosphonic acid Acta Cryst. B32 2889
- 29. Parmar D, Sugiono E, Raja S and Rueping M 2014 Complete Field Guide to Asymmetric BINOL-Phosphate Derived Brønsted Acid and Metal Catalysis: History and Classification by Mode of Activation; Brønsted Acidity, Hydrogen Bonding, Ion Pairing, and Metal Phosphates *Chem. Rev.* **114** 9047
- 30. Minyaev M E, Nifantev I E, Tavtorkin A N, Korchagina S A and Zeynalova S S 2015 Crystal structure of [bis(2,6-diisopropylphenyl) phosphato- κO]tris(methanol- κO)lithium methanol monosolvate *Acta Cryst.* **E71** 443
- 31. Murugavel R, Sathiyendiran M, Pothiraja R, Walawalkar M G, Mallah T and Rivire E 2004 Monomeric, Tetrameric, and Polymeric Copper Di-*tert*-butyl Phosphate Complexes Containing Pyridine Ancillary Ligands *Inorg. Chem.* **43** 945
- 32. Murugavel R and Shanmugan S 2007 Seeking tetrameric transition metal phosphonate with a D4R core and organising it into a 3-D supramolecular assembly *Chem. Commun.* 1257

- 33. Saxena P, Mandal S K, Sharma K and Murugavel R 2018 Delineating factors that dictate the framework of a bulky phosphate derived metal complexes: Sterics of phosphate, anion of the metal salt and auxiliary N-donor ligand *Inorg. Chim. Acta* 469 353
- 34. Mandal D, Santra B, Kalita P, Chrysochos N, Malakar A, Narayanan R S, Biswas S, Schulzke C, Chandrasekhar V and Jana A 2017 2,6-(Diphenylmethyl)-Aryl-Substituted Neutral and Anionic Phosphates: Approaches to H-Bonded Dimeric Molecular Structures *Chem. Select* 2 8898
- 35. Santra B, Mandal D, Gupta V, Kalita P, Kumar V, Narayanan R S, Dey A, Chrysochos N, Mohammad A, Singh A, Zimmer M, Dalapati R, Biswas S, Schulzke C, Chandrasekhar V, Scheschkewitz D and Jana A 2019 Structural Diversity in Supramolecular Organization of Anionic Phosphate Monoesters: Role of Cations ACS Omega 4 2118
- 36. Santra B, Narayanan R S, Kalita P, Kumar V, Mandal D, Gupta V, Zimmer M, Huch V, Chandrasekhar V, Scheschkewitz D, Schulzke C and Jana A 2019 Modulation of the Nuclearity of Molecular Mg(II)-Phosphates: Solid-State Structural Change Involving Coordinating Solvents *Dalton Trans.* 48 8853

- 37. Gupta V, Santra B, Mandal D, Das S, Narayanan R S, Kalita P, Rao D K, Schulzke C, Pati S K, Chandrasekhar V and Jana A 2018 Neutral and anionic phosphate-diesters as molecular templates for the encapsulation of a water dimer *Chem. Commun.* 54 11913
- Korb M and Lang H 2014 Planar Chirality from the Chiral Pool: Diastereoselective Anionic Phospho-Fries Rearrangements at Ferrocene Organometallics 33 6643
- SMART & SAINT Software Reference manuals, Version 6.45; Bruker Analytical X-ray Systems, Inc.: Madison, WI, 2003
- 40. Sheldrick G M 1996 SADABS, Program for Empirical Absorption Correction, University of Gottingen, Germany
- 41. Sheldrick G M 2015 *SHELXT* Integrated space-group and crystal structure determination *Acta Crystallogr*. **A71** 3
- 42. Sheldrick G M 2015 Crystal structure refinement with *SHELXL Acta Crystallogr.* C71 3
- 43. Dolomanov O V, Bourhis L J, Gildea R J, Howard J A K and Puschmann H 2009 *OLEX2*: a complete structure solution, refinement and analysis program *J. Appl. Cryst.* **42** 339