

This article was downloaded by: [Erciyes University]


On: 25 December 2014, At: 04:18

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T

3JH, UK



Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for
authors and subscription information:

<http://www.tandfonline.com/loi/lsc20>

Synthesis of 4-Arylpiperazine Derivatives of Moclobemide: Potential Antidepressants with a Dual Mode of Action

Hernán Pessoa-Mahana^a, Gonzalo Recabarren
Gajardo^a, Ramiro Araya-Maturana^a, Johann
Kosche Cárcamo^a & C. David Pessoa-Mahana^b

^a Departamento de Química Orgánica y
Fisicoquímica, Facultad de Ciencias Químicas y
Farmacéuticas, Universidad de Chile, Casilla 233,
Santiago, 1, Chile

^b Departamento de Farmacia, Pontificia
Universidad Católica de Chile, Santiago, Chile
Published online: 10 Jan 2011.

To cite this article: Hernán Pessoa-Mahana, Gonzalo Recabarren Gajardo, Ramiro Araya-Maturana, Johann Kosche Cárcamo & C. David Pessoa-Mahana (2004) Synthesis of 4-Arylpiperazine Derivatives of Moclobemide: Potential Antidepressants with a Dual Mode of Action, *Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry*, 34:14, 2513-2521, DOI: [10.1081/SCC-200025581](https://doi.org/10.1081/SCC-200025581)

To link to this article: <http://dx.doi.org/10.1081/SCC-200025581>

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Synthesis of 4-Arylpiperazine Derivatives of Moclobemide: Potential Antidepressants with a Dual Mode of Action

Hernán Pessoa-Mahana,^{1,*} Gonzalo Recabarren Gajardo,¹
Ramiro Araya-Maturana,¹ Johann Kosche Cárcamo,¹
and C. David Pessoa-Mahana²

¹Departamento de Química Orgánica y Fisicoquímica, Facultad de Ciencias Químicas y Farmacéuticas, Universidad de Chile, Santiago, Chile

²Departamento de Farmacia, Pontificia Universidad Católica de Chile, Santiago, Chile

ABSTRACT

We report here the synthesis of substituted 4-chloro-*N*-[3-*oxo*-3-(4-aryl)-1-piperaziny]-propyl] benzamides (**5–9**), as potential new antidepressants, incorporating in a single molecule structural moieties related to a dual pharmacological profile: MAO-A inhibitor and 5-HT_{1A} receptor affinity.

*Correspondence: Hernán Pessoa-Mahana, Departamento de Química Orgánica y Fisicoquímica, Facultad de Ciencias Químicas y Farmacéuticas, Universidad de Chile, Casilla 233, Santiago 1, Chile; E-mail: hpessoa@ciq.uchile.cl.

2513



DOI: 10.1080/SCC-200025581
Copyright © 2004 by Marcel Dekker, Inc.

0039-7911 (Print); 1532-2432 (Online)
www.dekker.com

Request Permissions / Order Reprints
powered by **RIGHTS LINK**
COPYRIGHT CLEARANCE CENTER, INC.

INTRODUCTION

Depression is one of the most frequent psychiatric disorders, affecting up to one third of all people at the same time. The causes of depression are multifactorial, although it is accepted that neurochemical disorders reflected by alterations in the levels of some central neurotransmitters are ultimately responsible for the appearance of the depressive symptoms.^[1–3] The neurotransmitter serotonin (5-hydroxytryptamine, 5-HT) modulates the activity of central nervous system and peripheral tissues by interaction with seven classes of receptors (5-HT_{1–7}), containing 15 distinct subpopulations of receptors^[4,5] where the 5-HT_{1A} subtype is the best studied due to its involvement in psychiatric disorders such as anxiety and depression.^[6,7] A well-known class of 5-HT_{1A} receptor ligands are the “long chain” arylpiperazine derivatives. Among these Buspirone a partial agonist at 5-HT_{1A} receptors is an effective antianxiety and was the first arylpiperazine approved for clinical use. On the other hand Moclobemide, a reversible MAO-A inhibitor exerts its antidepressant therapeutic effect by preventing the degradation of neurotransmitters and xenobiotic amines (oxidative deamination), modulating the level of these biogenic amines.



Although, a variety of method for the synthesis of Moclobemide and phenylpiperazine derivatives have been developed,^[8–11] there is no report dealing with the obtention of 4-aryl piperazine, connected with the alkyl-*p*-chlorobenzamide framework of Moclobemide in a single chemical entity.

In this article, we describe the synthesis of a series of molecular templates of 4-chloro-*N*-[3-oxo-3-(4-aryl-1-piperazinyl)-propyl] benzamides, as potential antidepressants of dual action, based on coupling structural moieties related to reversible MAO-A inhibitor Moclobemide and typical 5-HT_{1A} ligands (Fig. 1).

The synthesis of the title benzamides started by reaction of *p*-chlorobenzoyl chloride with β -alanine hydrochloride, in the presence of anhydrous pyridine under inert atmosphere to give the ester (**2**) (93.2%), which was further hydrolyzed in basic medium to provide the carboxylic acid (**3**) (91.5%) (Sch. 1). The hydrolysis of the ester function of (**2**) had to be carried out with special care,





Figure 1.



Scheme 1. Reagents: (a) $\text{NH}_2(\text{CH}_2)_2\text{-COOEt}$ hydrochloride/anhydrous THF, dry pyridine, N_2 atmosphere; (b) KOH-EtOH (30 min); (c) SOCl_2 , 40°C (3 hr); (d) substituted 4-phenylpiperazines/anhydrous THF, dry pyridine, N_2 atmosphere.

Table 1. The synthesis of 4-arylpiperazine derivatives of Moclobemide (**5–9**) in excellent yields.

Compound	R	Yield (%)
5	Hydrogen	95.0
6	4-Nitro	96.2
7	2- Fluor	94.7
8	4-Fluor	93.6
9	2-Methoxy	98.3

in order to avoid competitive hydrolysis of the amide, which was detected as a side reaction.

Reaction of the carboxylic acid (**3**) with thionyl chloride under smooth conditions of heating gave the corresponding acid chloride (**4**), in near quantitative yield, which was rapidly identified by its IR spectrum (strong carbonylic band at 1798 cm^{-1}), and reacted under inert atmosphere with different 4-arylpiperazines to afford the expected 4-arylpiperazino benzamides (**5–9**) in high yields (Table 1).

In conclusion, we have described the synthesis of new 4-arylpiperazine derivatives of Moclobemide (**5–9**) in excellent yields, through a four step sequence, starting from commercially available reagents. The potential dual pharmacological profile of the synthesized products could lead to a new class of antidepressants. The corresponding biological assays of these compounds are in progress.

EXPERIMENTAL

All reagents were obtained commercially and used without further purification. Melting points were measured on a Kofler hot-stage apparatus and are uncorrected. IR spectra were recorded with a FT Bruker spectrophotometer for KBr. $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra were obtained with a Bruker DRX-300 spectrophotometer. The chemical shifts are expressed in ppm (δ scale) downfield from TMS, J values are given in Hertz for solutions in CDCl_3 unless otherwise indicated. Microanalysis was determined on a Fisons EA 1108 analyzer. Silica gel Merck 60 (70–230 mesh) and DC-alufolien 60 F_{254} were normally used for column and TLC chromatography, respectively.

Ethyl 3-[(4-chlorobenzoyl) amino] propanoate (2). To a stirred solution of 3-amino-propanoate ethyl ester hydrochloride (β -alanine) (500 mg,

3.25 mmol), in dry THF (50 mL) containing anhydrous pyridine (381.3 mg, 4.88 mmol) at 0°C, was successively added 4-chloro benzoyl chloride (569.7 mg, 3.25 mmol) under nitrogen atmosphere, the whole was then stirred for 4 hr, upon this time the solution was poured into water (100 mL) and extracted with ethyl acetate (3 × 50 mL). The combined organic layers were dried (Na₂SO₄) and evaporated to dryness. The crude was purified by silica gel column chromatography (CH₂Cl₂/AcOEt = 1 : 1) to afford (2) (775.8, 93.2%). M.p. 84–85°C white crystals (ethanol). Anal. Calcd. for C₁₂H₁₄ClNO₃: C, 56.46; H, 5.53; N, 5.49. Found: C, 56.22; H, 5.47; N, 5.32%; IR ν_{\max} : 3338 (N-H), 1728 (COOEt), 1633 (CONH). ¹H-NMR (CDCl₃) δ : 1.28 (t, 3H, *J* = 7.1, –CH₂–CH₃), 2.64 (t, 2H, *J* = 5.9, –CH₂–COOEt), 3.72 (q, 2H, *J* = 6.0, NH–CH₂–CH₂), 4.17 (q, 2H, *J* = 7.1, COOCH₂CH₃), 6.97 (br. t, 1H, CONH–), 7.4 (d, 2H, *J* = 8.4, *p*-Cl–C₆H₄–, 3-H and 5-H), 7.71 (d, 2H, *J* = 8.4, *p*-Cl–C₆H₄–, 2-H and 6-H). ¹³C-NMR (CDCl₃) δ : 14.2, 33.8, 35.4, 60.9, 128.4 (2C), 128.8 (2C), 132.7, 137.7, 166.3, 172.9.

3-[(4-Chlorobenzoyl) amino] propionic acid (3). A solution of the ester (2) (347 mg, 1.36 mmol) in 0.5 N KOH : EtOH (1 : 1 v/v), (50 mL) was stirred at room temperature for 30 min. The mixture was concentrated in vacuo and the aqueous residue was cooled at 10°C and acidified with 0.1 N HCl to pH 1, the resulting precipitate was filtered off to provide (3) (263 mg, 91.5%). Recrystallization from ethanol/petroleum ether (1 : 2) furnished pure carboxylic acid (3) a white needles.

M.p. 168–169°C (ethanol/petroleum ether, 1 : 2). Anal. Calcd. for C₁₀H₁₀Cl NO₃: C, 52.86; H, 4.44; N, 6.17. Found: C, 52.75; H, 4.28; N, 6.11%; IR ν_{\max} : 3600–2870 (O-H), 3311 (N-H), 1702 (COOH), 1633 (NHC=O). ¹H-NMR (DMSO-*d*₆) δ : 2.31 (t, 2H, *J* = 7.0, –CH₂–COOH), 3.24 (q, 2H, *J* = 6.2, NH–CH₂–CH₂), 7.31 (d, 2H, *J* = 8.6, *p*-Cl–C₆H₄–, 3-H and 5-H), 7.65 (d, 2H, *J* = 8.6, *p*-Cl–C₆H₄–2-H and 6-H), 8.40 (br.t, 1H, ArCONH–), 12.2 (s, 1H, COOH). ¹³C-NMR (DMSO-*d*₆) δ : 34.1, 36.0, 128.8 (2C), 129.6 (2C), 133.6, 136.4, 165.63, 173.3.

3-[(4-Chlorobenzoyl) amino] propanoyl chloride (4). A solution of carboxylic acid (3) (312 mg, 1.47 mmol) in thionyl chloride (40 mL) was smoothly refluxed in an oil silicone bath, at 40°C for 3 hr. The solvent was then removed in vacuo, to afford (357.7 mg, 98.6%) of propanoyl chloride (4) as a yellow pale oil, being immediately characterized by spectroscopic analysis and reacted. IR ν_{\max} : 1797 (–COCl), 1664 (HO–C=N–). ¹H-NMR (CDCl₃) δ : 3.33 (t, 2H, *J* = 6.2, CH₂–CH₂–COCl), 3.97 (t, 2H, *J* = 6.2, C=N–CH₂–CH₂), 7.36 (d, 2H, *J* = 8.6, *p*-Cl–C₆H₄–, 3-H and 5-H), 7.90 (d, 2H, *J* = 8.6, *p*-Cl–C₆H₄–, 2-H and 6-H). ¹³C-NMR (CDCl₃) δ : 46.9, 49.2, 128.6 (2C), 130.2 (2C), 133.5, 138.2, 143.8, 172.2.

Preparation of 4-Chloro-*N*-[3-oxo-3-(4-aryl-1-piperazinyl)-propyl]-benzamide Derivatives (5–9). Typical Procedure.
(5) As an Example

Propanoyl chloride (**4**) (292 mg, 1.19 mmol) in dry THF (10 mL) was slowly added to a stirred solution at 0°C of 4-phenyl-1-piperazine (193 mg, 1.19 mmol), pyridine (94 mg, 1.19 mmol) and dry THF (50 mL) in nitrogen atmosphere. The mixture was maintained with stirring for 6 hr at room temperature and then poured into a water solution (100 mL). The mixture was extracted with ethyl acetate (3 × 50 mL) and the organic layers dried over MgSO₄. Removal of the solvent afforded 4-chloro-*N*-[3-oxo-3-(4-phenyl-1-piperazinyl)-propyl]-benzamide (**5**) (420 mg, 95%), as a white powder which was purified by silica gel column chromatography (CH₂Cl₂/AcOEt = 1:1). m.p. 172–173°C (ethanol). Anal. Calcd for C₂₀H₂₂ClN₃O₂: C, 64.60; H, 5.96; N, 11.30. Found: C, 64.52; H, 5.94; N, 11.26%; IR ν_{max} : 3295 (N-H), 1639 (C-O), 1632 (C=O), 1542 (N-H). ¹H-NMR (CDCl₃) δ : 2.67 (t, 2H, *J* = 5.5, -CH₂-CO-Pip.), 3.14–3.18 (m, 4H, (2 × -CH₂), Pip.), 3.60 (m, 2H, CONH-CH₂-), 3.75–3.80 (m, 4H, (2 × -CH₂), Pip.), 6.89–6.93 (m, 3H, C₆H₅), 7.25–7.31 (m, 3H, C₆H₅ and -CONH-), 7.37 (d, 2H, *J* = 8.6, *p*-Cl-C₆H₄-, 3-H and 5-H), 7.40 (d, 2H, *J* = 8.6, *p*-Cl-C₆H₄ -2-H and 6-H). ¹³C-NMR δ : 32.7, 35.6, 41.6, 45.3, 49.4, 49.6, 116.8 (2C), 121.0, 128.5 (2C), 128.8 (2C), 129.3 (2C), 132.8, 137.6, 150.8, 166.2, 170.3.

4-Chloro-*N*-[3-oxo-3-(4-(4-nitrophenyl)-1-piperazinyl)-propyl]-benzamide (6). Compound (**6**) (318 mg, 96.2%) yellow powder (column chromatographed, AcOEt/CH₂Cl₂ = 1:1). was prepared from (**4**) (195 mg, 0.793 mmol) and 4-nitrophenyl piperazine (164 mg, 0.793 mmol); m.p. 165–166°C (ethanol/petroleum ether = 3:1). Anal. Calcd for C₂₀H₂₁Cl N₄O₄: C, 57.63; H, 5.08; N, 13.44. Found: C, 57.36; H, 5.00; N, 13.24%; IR ν_{max} : 3300 (N-H), 1642 (CONH), 1635 (CO-Pip.). ¹H-NMR (CDCl₃) δ : 2.70 (t, 2H, *J* = 5.5, -CH₂-CO-Pip.), 3.41–3.50 (m, 4H, (2 × CH₂)-Pip.), 3.64–3.68 (m, 2H, *p*-Cl-CONH-CH₂-), 3.75–3.83 (m, 4H, (2 × CH₂)-Pip.), 6.82 (d, 2H, *J* = 9.4, *p*-NO₂-C₆H₄-2'-H and 6'-H), 7.21 (br t, 1H, ArCONH-), 7.39 (d, 2H, *J* = 8.6, *p*-Cl-C₆H₄-3-H and 5-H), 7.73 (d, 2H, *J* = 8.6, *p*-Cl-C₆H₄-, 2-H and 6-H), 8.13 (d, 2H, *J* = 9.4, *p*-NO₂-C₆H₄-, 3'-H and 5'-H). ¹³C-NMR δ : 32.8, 35.5, 40.9, 44.5, 46.7, 46.8, 113.0 (2C), 126.0 (2C), 128.5 (2C), 128.8 (2C), 132.7, 137.8, 139.1, 154.3, 166.3, 170.5.

4-Chloro-*N*-[3-oxo-3-(4-(2-fluorophenyl)-1-piperazinyl)-propyl]-benzamide (7). Compound (**7**) (795.3 mg, 94.7%) white powder (column chromatographed, AcOEt/CH₂Cl₂ = 1:1). was prepared from (**4**) (530 mg, 2.15 mmol) and 2-fluorophenyl piperazine (388 mg, 2.15 mmol); m.p. 184–185°C (ethanol–petroleum ether 1:1). Anal. Calcd for C₂₀H₂₁ClFN₃O₂: C,

61.68; H, 5.44; N, 10.80. Found: C, 61.46; H, 5.43; N, 10.82%; IR ν_{\max} : 3308 (N-H), 1639 (C=O), 1635 (C=O), 1547 (N-H). $^1\text{H-NMR}$ (DMSO- d_6) δ : 2.65 (t, 2H, $J = 7.1$, $-\text{CH}_2\text{-CO-Pip.}$), 2.91–3.02 (m, 4H, $(2 \times \text{CH}_2)\text{-Pip.}$), 3.46–3.54 (m, 2H, $\text{CONH-CH}_2\text{-}$), 3.59–3.65 (m, 4H, $(2 \times \text{CH}_2)\text{-Pip.}$), 6.93–7.15 (m, 4H, $o\text{-F-C}_6\text{H}_4\text{-}$), 7.53 (d, 2H, $J = 8.6$, $p\text{-Cl-C}_6\text{H}_4\text{-}$, 3-H and 5-H), 7.86 (d, 2H, $J = 8.6$, $p\text{-Cl-C}_6\text{H}_4\text{-}$, 2-H and 6-H), 8.61 (br t, 1H, ArCONH-). $^{13}\text{C-NMR}$ (DMSO- d_6) δ : 32.7, 36.5, 41.5, 45.5, 50.6, 50.9, 116.5 (d, $^2J_{\text{C-F}} = 21$ Hz), 120 (d, $^4J_{\text{C-F}} = 2.7$ Hz), 123.3 (d, $^3J_{\text{C-F}} = 7.9$ Hz), 125.3 (d, $^2J_{\text{C-F}} = 3.5$ Hz), 128.9 (2C), 129.5 (2C), 133.6, 136.5, 140.0 (d, $^3J_{\text{C-F}} = 8.3$ Hz), 155.5 (d, $^1J_{\text{C-F}} = 244$ Hz), 165.6, 169.6.

4-Chloro-N-[3-oxo-3-(4-(4-fluorophenyl)-1-piperazinyl)-propyl]-benzamide (8). Compound (8) (365 mg, 93.6%) white crystals was prepared from (4) (246 mg, 1.0 mmol) and 4-fluorophenyl piperazine (180 mg, 1.0 mmol); m.p. 148–149°C (ethanol/petroleum ether 2:1). Anal. Calcd for $\text{C}_{20}\text{H}_{21}\text{ClFN}_3\text{O}_2$: C, 61.68; H, 5.44; N, 10.80. Found: C, 61.67; H, 5.35; N, 10.81%; IR ν_{\max} : 3292 (N-H), 1638 (C-O), 1632 (C=O), 1538 (N-H). $^1\text{H-NMR}$ (CDCl_3) δ : 2.67 (t, 2H, $J = 5.5$, $-\text{CH}_2\text{-CO-Pip.}$), 3.04–3.10 (m, 4H, $(2 \times \text{CH}_2)\text{-Pip.}$), 3.57–3.61 (m, 2H, $\text{NH-CH}_2\text{-}$), 3.74–3.80 (m, 4H, $(2 \times \text{CH}_2)\text{-Pip.}$), 6.84–7.01 (m, 4H $o\text{-F-C}_6\text{H}_4\text{-}$), 7.24 (br t, 1H, ArCONH-), 7.38 (d, 2H, $J = 8.5$, $p\text{-Cl-C}_6\text{H}_4\text{-}$, 3-H and 5-H), 7.72 (d, 2H, $J = 8.5$, $p\text{-Cl-C}_6\text{H}_4\text{-}$, 2-H and 6-H). $^{13}\text{C-NMR}$ (CDCl_3) δ : 32.7, 35.4, 41.5, 45.3, 50.4, 50.7, 115.8 (d, 2C, $^2J_{\text{C-F}} = 22$ Hz), 118.7 (d, 2C, $^3J_{\text{C-F}} = 7.8$ Hz), 128.5 (2C), 128.8 (2C), 132.7, 137.7, 147.5 (d, $^4J_{\text{C-F}} = 2.3$ Hz), 157.7 (d, $^1J_{\text{C-F}} = 240$ Hz), 166.2, 170.3.

4-Chloro-N-[3-oxo-3-(4-(2-methoxy phenyl)-1-piperazinyl)-propyl] benzamide (9). Compound (9) (526 mg, 98.3%) was prepared from (4) (328 mg, 1.33 mmol) and 2-methoxyphenyl piperazine (256 mg, 1.33 mmol); m.p. 109–110°C white crystals (column chromatographed, $\text{AcOEt/CH}_2\text{Cl}_2 = 2:1$). Anal. Calcd for $\text{C}_{21}\text{H}_{24}\text{ClN}_3\text{O}_3$: C, 62.82; H, 6.03; N, 10.47. Found: C, 62.46; H, 6.15; N, 10.33%; IR ν_{\max} : 3308 (N-H), 1656 (C=O), 1621 (C=O), 1538 (N-H). $^1\text{H-NMR}$ (CDCl_3) δ : 2.68 (t, 2H, $J = 5.5$, $-\text{CH}_2\text{-CO-Pip.}$), 3.01–3.06 (m, 4H, $(2 \times \text{CH}_2)\text{-Pip.}$), 3.60–3.63 (m, 2H, $\text{NH-CH}_2\text{-}$), 3.72–3.81 (m, 4H, $(2 \times \text{CH}_2)\text{-Pip.}$), 3.87 (s, 3H, $o\text{-OCH}_3\text{-C}_6\text{H}_4\text{-}$), 6.83–7.06 (m, 4H $o\text{-OCH}_3\text{-C}_6\text{H}_4\text{-}$), 7.38 (d, 3H, $J = 8.5$, $p\text{-Cl-C}_6\text{H}_4$ 3-H, 5-H and $p\text{-Cl-C}_6\text{H}_4\text{CONH-}$), 7.74 (d, 2H, $J = 8.5$, $p\text{-Cl-C}_6\text{H}_4$, 2-H, 6-H). $^{13}\text{C-NMR}$ (CDCl_3) δ : 32.7, 35.7, 41.9, 45.6, 50.5, 50.8, 55.4, 111.3, 118.4, 121.0, 123.7, 128.5 (2C), 128.7 (2C), 132.8, 137.6, 140.5, 152.2, 166.2, 170.3.

ACKNOWLEDGMENTS

This work was supported by the Research Project CEPEDEQ- FACUL-TAD of the Chemical and Pharmaceutical Sciences Faculty, University of Chile, Santiago, Chile, and FONDECYT 1030916.

REFERENCES

1. Blier, P. The pharmacology of putative early-onset antidepressant strategies. *Eur. Neuropsychopharmacol.* **2003**, *13*, 57–66.
2. Pessoa-Mahana, H.; Araya-Maturana, R.; Saitz, B.C.; Pessoa-Mahana, C. David. A synthetic overview of new molecules with 5-HT_{1A} binding affinities. *Minirev. Med. Chem.* **2003**, *3*, 77–93.
3. Oficialdegui, A.M.; Martinez, J.; Perez, S.; Heras, B.; Irurzun, M.; Palop, J.A.; Tordera, R.; Lasheras, B.; del Río, J.; Monge, A. Design, synthesis and biological evaluation of new 3-[(4-aryl)piperazin-1-yl]-1-arylpropane derivatives as potential antidepressants with a dual mode of action. Serotonin reuptake inhibition and 5-HT_{1A} receptor antagonism. *Il Farmaco* **2000**, *55*, 345–353.
4. Glennon, R.A. Higher-end serotonin receptors: 5-HT₅, 5-HT₆, and 5-HT₇. *J. Med. Chem.* **2003**, *46* (14), 2795–2812.
5. Mattson, R.J.; Catt, J.D.; Sloan, C.P.; Gao, Q.; Carter, R.B.; Gentile, A.; Mahle, C.D.; Mattos, F.F.; McGvern, R.; VanderMaelen, C.P.; Yocca, F.D. Development of a presynaptic 5-HT_{1A} antagonist. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 285–288.
6. Orús, L.; Sáinz, Y.; Pérez, S.; Oficialdegui, A.M.; Martínez, J.; Lasheras, B.; del Río, J.; Monge, A. New 3-[4-(aryl) piperazin-1-yl] -1-(benzo [b] thio-phen-2-yl)propane derivatives with dual action at 5-HT_{1A} serotonin receptors and serotonin transporter as a new class of antidepressants. *Pharmazie* **2002**, *57* (6), 355–357.
7. Takeuchi, K.; Kohn, T.J.; Honigsmid, N.A.; Rocco, P.V.; Spinazze, P.G.; Koch, D.J.; Nelson, D.L.; Wainscott, D.B.; Ahmad, L.J.; Shaw, J.; Threlkeld, P.G.; Wong, D.T. Advances toward new antidepressants beyond SSRIs: 1-aryloxy-3-piperidinylpropan-2-ols with dual 5-HT_{1A} receptor antagonism/SSRI activities. Part 1. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 1903–1905.
8. Ghambarpour, A.; Hadizadeh, H.; Piri, F.; Rashidi-Ranjbar, P. Synthesis, conformational analysis and antidepressant activity of moclobemide new analogues. *Pharm. Acta Helv.* **1997**, *72*, 119–122.

9. Pessoa-Mahana, H.; Astudillo, C.I.; Araya-Maturana, R. Synthesis of *N*-(morpholinomethyl)benzamides as moclobemide analogues. *Synth. Commun.* **2002**, 32 (9), 1437–1445.
10. López-Rodríguez, M.L.; Ayala, D.; Benhamú, B.; Morcillo, M.J.; Viso, A. Arylpiperazine derivatives acting at 5-HT_{1A} receptors. *Curr. Med. Chem.* **2002**, 9 (4), 443–469.
11. López-Rodríguez, M.L.; Morcillo, M.J.; Fernández, E.; Benhamú, B.; Tejada, I.; Ayala, D.; Viso, A.; Olivella, M.; Pardo, L.; Delgado, M.; Manzanares, J.; Fuentes, J.A. Design and synthesis of *S*-(–)-2[[4-(naphth-1-yl)piperazin-1-yl]-methyl]-1,4-dioxoperhydropyrrolo[1,2-*a*]pyrazine (CSP-2503) using computational stimulation. A 5-HT_{1A} receptor agonist. *Bioorg. Med. Chem. Lett.* **2003**, 13, 1429–1432.

Received in the USA March 3, 2004