This article was downloaded by: [University of Arizona] On: 24 December 2012, At: 02:01 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lsyc20</u>

# A Green Preparation of N - Chloro- and N - Bromosaccharin

Soraia P. L. de Souza<sup>a</sup>, Joaquim F. M. da Silva<sup>a</sup> & Marcio C. S. de Mattos<sup>a</sup> <sup>a</sup> Instituto de Química, Departamento de Química Orgânica, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil Version of record first published: 17 Aug 2006.

To cite this article: Soraia P. L. de Souza , Joaquim F. M. da Silva & Marcio C. S. de Mattos (2003): A Green Preparation of N -Chloro- and N -Bromosaccharin, Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry, 33:6, 935-939

To link to this article: <u>http://dx.doi.org/10.1081/SCC-120016353</u>

### PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: http://www.tandfonline.com/page/terms-and-conditions

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

SYNTHETIC COMMUNICATIONS<sup>®</sup> Vol. 33, No. 6, pp. 935–939, 2003

## A Green Preparation of *N*-Chloro- and *N*-Bromosaccharin

Soraia P. L. de Souza, Joaquim F. M. da Silva, and Marcio C. S. de Mattos<sup>\*</sup>

Instituto de Química, Departamento de Química Orgânica, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

#### ABSTRACT

The reaction of the sodium salt of saccharin, potassium chloride or bromide and oxone<sup>®</sup> in water at room temperature for 24 h produced pure *N*-chloro- and *N*-bromosaccharin in 58% and 64% yield, respectively.

Key Words: Green chemistry; Saccharin; N-Halosaccharin.

935

DOI: 10.1081/SCC-120016353 Copyright © 2003 by Marcel Dekker, Inc. 0039-7911 (Print); 1532-2432 (Online) www.dekker.com

<sup>\*</sup>Correspondence: Marcio C. S. de Mattos, Instituto de Química, Departamento de Química Orgânica, Universidade Federal do Rio de Janeiro, Caixa Postal 68545 21945-970, Rio de Janeiro, Brazil; Fax: +55-21-25627133; E-mail: mmattos@iq.ufrj.br.

M

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

936

#### de Souza, da Silva, and de Mattos

#### INTRODUCTION

Although *N*-halosaccharins are more electrophilic than the structurally analogue *N*-haloimides,<sup>[1]</sup> they have found little attention in the literature. Whereas *N*-iodo-<sup>[1,2]</sup> and *N*-fluorosaccharin<sup>[3]</sup> are scarcely mentioned, *N*-chloro- and *N*-bromosaccharin are mostly described as analytical<sup>[4,5]</sup> and/or oxidizing reagents. These compounds are mild oxidants of aldehydes,<sup>[6,7]</sup> alcohols,<sup>[8,9]</sup> glycols,<sup>[9]</sup>  $\alpha$ -amino-<sup>[10]</sup> and  $\alpha$ -hydroxyacids,<sup>[9]</sup> etc, and their application as halogenating reagent for aromatic compounds<sup>[11,12]</sup> and allylic,<sup>[13]</sup> benzylic,<sup>[13,14]</sup> and  $\alpha$ -carbonylic <sup>[14,15]</sup> positions is very useful.

With few exceptions,<sup>[16,17]</sup> the classical <sup>[18]</sup> procedure for preparation of *N*-chloro- and *N*-bromosaccharin is reaction of sodium salt of saccharin with the halogen. However, the handling and manipulation of the toxic and corrosive halogens is problematic at multigram-scale.

Recently, Curini published the *N*-chlorination of amides and carbamates by the reaction with sodium chloride and oxone<sup>®</sup> (2 KHSO<sub>5</sub>·KHSO<sub>4</sub>·K<sub>2</sub>SO<sub>4</sub>) in a suspension of wet alumina in chloroform at  $45^{\circ}$ C.<sup>[19]</sup> The role of oxone<sup>®</sup> in these reactions is to oxidize the chloride anion and generate chlorine *in situ*.<sup>[20]</sup>

The utilization of environmentally acceptable, healthy, safe, and readily available materials in a simple process is very attractive nowadays and consistent with green chemistry challenges.<sup>[21]</sup> Based on this precedent we present here our results on an alternate and safe route to N-chloro and N-bromosaccharin.

#### RESULTS

The reaction of the sodium salt of saccharin with potassium halide (1 equiv. mol) and oxone<sup>®</sup> (1 equiv. mol) was performed in water at room temperature for 24 h. Using these conditions on a 10 mmol scale, the corresponding *N*-halosaccharin precipitates at the end of the reaction. The products were obtained in high purity (no need of recrystallization), albeit in moderate yield after washing with cold water (Sch. 1 and Table 1). The purity of the *N*-halosaccharins obtained was determined by iodometric titration<sup>[1]</sup> and they were characterized by spectroscopic methods and their melting point. The reaction was performed at 50-mmol scale and the products were obtained in similar yield and purity.

In summary we propose a new preparation of *N*-chloro- and *N*-bromosaccharin that does not use or handle toxic and corrosive halogens. The reaction conditions are mild, the yield is moderate and

937

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

N-Chloro- and N-Bromosaccharin



Table 1. Preparation of N-halosaccharin.

Х	Yield/%	Purity/% <sup>a</sup>	M.p. [lit.]/ $^{\circ}C$	$v_{\mathrm{KBr}}/\mathrm{cm}^{-1}$
Cl	58	>96	145–148	3094; 1739 (C=O); 1361 (SO <sub>2</sub> ); 1235
Br	64	>99	$[152^{-17}]$ 165–167 $[170–172^{[17]}]$	$\begin{array}{c} (N-C1); 1194 (SO_2); 955, 748; 757 \\ 3094; 1709 (C=O); 1352 (SO_2); 1231 \\ (N-Br); 1191 (SO_2); 949; 748; 574 \end{array}$

<sup>*a*</sup>Determined by iodometric titration.<sup>[1]</sup>

the isolation of the products in high purity is simple. Our methodology is easier and safer than the predecessors, the reaction can be scaled up and water is the reaction solvent, all of which is in accordance with the green chemistry philosophy.<sup>[22]</sup>

#### EXPERIMENTAL

Typical procedure for the preparation of *N*-halosaccharin: To a wellstirred solution of sodium saccharin (10.25 g, 50 mmol),  $Na_2CO_3$  (2.65 g, 25 mmol) and KCl or KBr (3.78 g or 6.00 g, respectively—50 mmol) in water (250 mL), a solution of oxone<sup>®</sup> (30.75 g, 50 mmol) in water (30 mL) was added slowly at 0°C. After stirring for 24 h at room temperature, the solid was filtered off, washed with cold water and dried to give the pure *N*-halosaccharin. The results obtained are shown in Table 1.

#### ACKNOWLEDGMENTS

SPLS thanks CNPq for a fellowship. We thank Prof. W. Bruce Kover and Joel Jones Jr. for helpful discussions. We also thank Prof. Carlos A.S. Riehl for technical assistance in the *N*-halosaccharin titration.

XX

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

938

#### de Souza, da Silva, and de Mattos

#### REFERENCES

- 1. Dolenc, D. N-Iodosaccharin—A new reagent for iodination of alkenes and activated aromatics. Synlett **2000**, (4), 544–546.
- Fairbanks, A.J.; Aloui, M. N-Iodosaccharin: A potent new activator of thiophenylglycosides. Synlett 2001, (6), 797–799.
- Gakh, A.A.; Romaniko, S.V.; Ugrak, B.I.; Fainzilberg, A.A. N-Fluorination with cesium fluoroxysulfate. Tetrahedron 1991, 47 (35), 7447–7458.
- Chauhan, R.P.S.; Ahmed, I.; Kumar, A. Determination of milligram amounts of thiourea, allylthiourea and phenylthiourea using N-chlorosaccharin in acetic-acid media. Asian J. Chem. 1995, 7 (3), 667–668. (Chem. Abstr. 1995, 123, 245771p)
- Das, C.M. Determination of some sulfur-compounds with N-bromoimides. Talanta 1991, 38 (3), 347–350. (Chem. Abstr. 1991, 114, 258834a)
- Khan, M.U.; Verma, J.K.; Nigam, S.K.; Parihar, S.S.; Dwiverdi, H.P. Kinetics and mechanistic approach on N-chlorosaccharin oxidation of some aliphatic aldehydes. Oxid. Commun. 1998, 21 (3), 362–368.
- Sharma, V.K.; Sharma, K.; Gupta, H.D.; Gupta, P.P. Kinetics of oxidation of benzaldehyde and substituted benzaldehydes by N-bromosaccharin in aqueous acetic acid-perchloric acid medium. Oxid. Commun. 1995, 18 (4), 395–406.
- Manoharan, V.; Venkatasubramanian, N. Kinetics of oxidation of aliphatic secondary alcohols by N-bromosaccharin. J. Indian Chem. Soc. 1986, 63 (6), 613–615.
- Bachhawat, J.M.; Mathur, N.K. Oxidation of secondary alcohols, glycols, α-hydroxy acids, amino acids and hydrocarbons by N-bromosaccharin. Indian J. Chem. 1971, 9 (12), 1335–1336.
- Manikyamba P.; Sundaram, E.V. Kinetics of oxidation of aminoacids by N-bromosaccharin. Indian J. Chem. 1980, 19A (11), 1122–1124.
- Zajc, B. Synthesis of (+/-)-trans-7,8-Dihydrodiol of 6-fluorobenzo[a] pyrene via hydroxyl-directed regioselective functionalization of Substituted Pyrene. J. Org. Chem. 1999, 64 (6), 1902–1907.
- Mozek, I.; Sket, B. Chlorination and bromination of molecules by an N-halosaccharin/pyridinium poly(hydrogen fluoride) system. Synth. Commun. 1992, 22 (17), 2513–2520.
- 13. Bachhawat, J.M.; Koul, A.K.; Prashad, B.; Ramegowd, N.S.; Narang, C.K.; Mathur, N.K. Allylic and Benzylic chlorination

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

#### N-Chloro- and N-Bromosaccharin

and oxidation of organic substrate using N-chlorosaccharin. Indian J. Chem. **1973**, *11* (6), 609–610.

- Sanchez, E.I.; Fumarola, M.J. N-Bromosaccharin: Benzylic and α-carbonylic bromination. J. Org. Chem. 1982, 47 (8), 1588–1590.
- Panchatsharam, V.S.; Sundaram, S.; Venkatasubramanian, N. Chlorination of ketones with N-chlorosaccharin. Indian J. Chem. 1981, 20A (5), 469–472.
- 16. Zajc, B. A convenient preparation of N-bromosaccharin, source of electrophilic bromine. Synth. Commun. **1999**, *29* (10), 1779–1784.
- Sanchez, E.I.; Fumarola, M.J. A new synthesis of N-bromosaccharin using bromine chloride. Synthesis 1976, (11), 736–737.
- Chattaway, F.D. A contribution to the chemistry of o-Benzoic Sulphinida. J. Chem. Soc. 1905, 87 (2), 1882–1887.
- Curini, M.; Epifani, F.; Marcotullio, M.C.; Rosati, O.; Tsadjout, A. N-chlorination of amides and carbamates by oxone<sup>®</sup> and sodium chloride. Synlett **2000**, (6), 813–814.
- 20. Dieter, R.K.; Nice, L.E.; Velu, S.E. Oxidation of  $\alpha$ , $\beta$ -enones and alkenes with oxone and sodium halides: A convenient laboratory preparation of chlorine and bromine. Tetrahedron Lett. **1996**, *37* (14), 2377–2380.
- 21. Clark, J.H. Green chemistry: Challenges and opportunities. Green Chem. **1999**, *1* (1), 1–8.
- 22. Anastas, P.T.; Warner, J.C. *Green Chemistry: Theory and Practice*; Oxford University Press: Oxford, 2000.

Received in the USA April 3, 2002



©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.