pubs.acs.org/joc

## Oxidation of Organotrifluoroborates via Oxone

Gary A. Molander\* and Livia N. Cavalcanti

Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6323, United States

gmolandr@sas.upenn.edu

Received November 5, 2010



A method for the oxidation of organotrifluoroborates using Oxone was developed. A variety of aryl-, heteroaryl-, alkenyl-, and alkyltrifluoroborates were converted into the corresponding oxidized products in excellent yields. This method proved to be tolerant of a broad range of functional groups, and in secondary alkyl substrates it was demonstrated to be completely stereospecific.

## Introduction

Phenols are found in numerous natural products, pharmaceuticals, and polymers and have been widely used as versatile synthetic intermediates.<sup>1</sup> Traditional methods for the synthesis of these compounds include various protocols for substitution of aryl halides. Unfortunately, the harsh reaction conditions required for direct nucleophilic substitution make this approach incompatible with a variety of common functional groups.<sup>2</sup> Copper- and palladium-catalyzed hydroxylation of aryl halides has been reported in the literature and offers an alternative to the former method.<sup>3</sup> In this case, however, the use of transition metals with attendant ligands and/or high reaction temperatures leads to processes that are less than ideal. Various boron species can be incorporated into aromatic and heteroaromatic systems in a variety of complementary ways, and recently the oxidation of boronic acids in a copper-catalyzed reaction has been reported.<sup>4</sup> Although perhaps useful on a small scale in the laboratory, the use of a copper reagent, ligand, and excess base significantly reduces its appeal on a larger scale. As an alternative to this method, several nonmetal-promoted oxidations of boronic acids and their derivatives have been described.<sup>5</sup> Among the oxidants used for this transformation, hydrogen peroxide, <sup>5a,c,e,f</sup> hydroxylamine, <sup>5b</sup> and Oxone <sup>5d,g</sup> are most often employed. The first two reagents require long reaction times, and the products are obtained in moderate yields for arylboronic acids containing electron-withdrawing groups. On the other hand, the use of Oxone provides an extremely rapid and generally efficient reaction. <sup>5d,g</sup>

Over the past years organotrifluoroborates have been used in many contexts as a valuable alternative to boronic acids and boronate esters.<sup>6</sup> Because of their tetracoordinate nature, these organoboron species do not undergo undesirable side reactions with commonly employed organic reagents such as bases, organic acids, and nucleophiles. Thus, organotrifluoroborates,

 <sup>(1) (</sup>a) Rappoport, Z. *The Chemistry of Phenols*; Wiley-VCH: Weinheim, Germany, 2003. (b) Tyman, J. H. P. *Synthetic and Natural Phenols*; Elsevier: New York, 1996.

<sup>(2) (</sup>a) Fyfe, C. A. In *The Chemistry of the Hydroxyl Group*; Patai, S., Ed.; Wiley-Interscience: New York, 1971; Vol. 1. (b) Hoarau, C.; Pettus, T. R. R. *Synlett* **2003**, 127. (c) Hanson, P.; Jones, J. R.; Taylor, A. B.; Walton, P. H.; Timms, A. W. J. Chem. Soc., Perkin Trans. 2 **2002**, 1135. (d) George, T.; Mabon, R.; Sweeney, G.; Sweeney, J. B.; Tavassoli, A. J. Chem. Soc., Perkin Trans. 1 **2000**, 2529 and references therein.

<sup>(3) (</sup>a) Schulz, T.; Torborg, C.; Schaffner, B.; Huang, J.; Zapf, A.; Kadyrov, R.; Borner, A.; Beller, M. Angew. Chem., Int. Ed. 2009, 48, 918.
(b) Tlili, A.; Xia, N.; Monnier, F.; Taillefer, M. Angew. Chem., Int. Ed. 2009, 48, 9725.
(c) Zhao, D.; Wu, N.; Zhang, S.; Xi, P.; Su, X.; Lan, J.; You, J. Angew. Chem., Int. Ed. 2009, 48, 8729.
(d) Anderson, K. W.; Ikawa, T.; Tundel, R. E.; Buchwald, S. L. J. Am. Chem. Soc. 2006, 128, 10694.

<sup>(4)</sup> Xu, J.; Wang, X.; Shao, C.; Su, D.; Cheng, G.; Hu, Y. Org. Lett. 2010, 12, 1964.

<sup>(5) (</sup>a) Prakash, G. K. S.; Chacko, S.; Panja, C.; Thomas, T. E.; Gurung, L.; Rasul, G.; Mathew, T.; Olah, G. A. Adv. Synth. Catal. **2009**, 351, 1567. (b) Kianmehr, E.; Yahyaee, M.; Tabatabai, K. Tetrahedron Lett. **2007**, 48, 2713. (c) Clay, J. M.; Vedejs, E. J. Am. Chem. Soc. **2005**, 127, 5766. (d) Maleczka, R. E.; Shi, F.; Holmes, D.; Smith, M. R., III. J. Am. Chem. Soc. **2003**, 127, 7792. (e) Benjamin, R.; Travis, B. R.; Ciaramitaro, B. P.; Borhan, B. *Eur. J. Org. Chem.* **2002**, 3429. (f) Simon, J.; Salzbrunn, S.; Prakash, G. K. S.; Petasis, N. A.; Olah, G. A. J. Org. Chem. **2001**, 66, 633. (g) Webb, K. S.; Levy, D. Tetrahedron Lett. **1995**, 36, 5117.

<sup>(6)</sup> For reviews, see: (a) Darses, S.; Genêt, J.-P. Chem. Rev. 2008, 108, 288.
(b) Doucet, H. Eur. J. Org. Chem. 2008, 2013. (c) Molander, G. A.; Ellis, N. M. Acc. Chem. Res. 2007, 40, 275. (d) Stefani, H. A.; Cella, R.; Vieira, A. S. Tetrahedron 2007, 63, 3623. (e) Molander, G. A.; Figueroa, R. Aldrichim. Acta 2005, 38, 49. (f) Darses, S.; Genêt, J.-P. Eur. J. Org. Chem. 2003, 4313.

<sup>(7)</sup> For selected examples, see: (a) Molander, G. A.; Raushel, J.; Ellis, N. M. J. Org. Chem. 2010, 75, 4304. (b) Molander, G. A.; Canturk, B.; Kennedy, L. E. J. Org. Chem. 2009, 74, 973. (c) Molander, G. A.; Febo-Ayala, W.; Ortega-Guerra, M. J. Org. Chem. 2008, 73, 6000. (d) Molander, G. A.; Cooper, D. J. J. Org. Chem. 2008, 73, 3885. (e) Molander, G. A.; Ellis, N. M. J. Org. Chem. 2006, 71, 7491. (f) Molander, G. A.; Felix, L. A. J. Org. Chem. 2005, 70, 3950.



Fg = Functional group

in contrast to most other boron species, can be functionalized to build molecular complexity, while leaving the boron-carbon bond intact (Scheme 1).<sup>7</sup> The organotrifluoroborates can then be conveniently converted in a variety of ways, depending on the conditions utilized.<sup>8</sup> Of further importance, heteroaryltrifluoroborates exhibit enhanced stability, and thus they can be prepared and stored for long periods of time, whereas many of the corresponding boronic acids rapidly decompose.<sup>8a,9</sup> Thus, many advantages to using organotrifluoroborates as surrogates for boronic acids are beginning to emerge.

To the best of our knowledge, although organotrifluoroborates present all of the benefits mentioned above, reports of the oxidation of these species into the corresponding phenol, alcohol, or related derivatives are currently limited.<sup>4,10</sup> For example, phenyltrifluoroborate has been oxidized to phenol in a copper-catalyzed procedure requiring 8 h at room temperature.<sup>4</sup> In another isolated example, an alkyltrifluoroborate has been oxidized by utilizing an excess of a hypervalent iodonium complex to afford an excellent yield of the desired alcohol in 3 h.<sup>10a</sup> Finally, during the preparation of this paper Fensterbank et al. reported the 2,2,6,6-tetramethylpiperidin-1oxyl (TEMPO)-promoted oxidation of several alkyltrifluoroborates, which required high temperatures and long reaction times (120 °C, 20 h) with excess TEMPO (1.2 equiv) and copper salts (1.2 equiv) in DMSO, or 3 equiv of TEMPO in the presence of Dess-Martin periodinane in Et<sub>2</sub>O at room temperature for 1-2 days. The products isolated were the corresponding TEMPO derivatives, which were accessed in modest to good yields depending on the substrates. Importantly, these reactions were demonstrated to proceed via radical intermediates, and thus they would not be expected to be stereospecific for enantiomerically enriched alkyltrifluoroborates, although this latter aspect was not specifically addressed in an unambiguous manner.

The lack of examples detailing a practical method for the oxidation of organotrifluoroborates and the limitations associated with other boron derivatives prompted us to investigate a general, efficient, mild, rapid, and convenient method for this transformation. Herein, we report the oxidation of a broad range of aryl-, heteroaryl-, alkenyl-, and alkyltrifluoroborates using Oxone at room temperature.

## **Results and Discussion**

On the basis of previous research in which a variety of arylboronic acids and pinacol boronates were transformed to

the corresponding phenols,<sup>5d,g</sup> investigations were initiated on the oxidation of potassium naphthalen-1-yltrifluoroborate. Very rapidly it was evident that the protocol developed by Maleczka and Smith for pinacol boronates, using 1 equiv of Oxone (as a 0.2 M solution in  $H_2O$ ) in acetone at room temperature, open to the air, was also ideal for the rapid (<5 min) oxidation of organotrifluoroborates. The desired phenol was obtained by an aqueous workup and simple filtration in excellent yield (99%). Because Oxone is an inexpensive,<sup>11</sup> widely available, and industrially acceptable reagent, all subsequent reactions were performed using this material.

With optimal conditions in hand, the investigation continued to outline the full scope of this process. The requisite potassium organotrifluoroborates needed for the study were readily prepared by previously reported procedures.<sup>6,12</sup> Initially, a broad range of electron-rich and electron-neutral aryltrifluoroborates were investigated (Table 1). In all cases, quantitative conversion to the desired products was accomplished in only 2 min. The reaction provided excellent yields for all substrates containing para, meta, or ortho substituents (entries 3-5). Sterically hindered compounds also afforded the products in high yields (entry 7). Importantly, we demonstrated that the reaction could be scaled to 55 mmol, providing product 1a in 96% yield (Table 1, entry 1). In contrast to the metal-catalyzed processes utilizing other organoboron species, the use of organotrifluoroborates and Oxone in every case afforded the desired products with little or no formation of organic side products. Therefore, a simple aqueous workup followed, at most, by filtration through a short plug of silica with charcoal afforded the desired products in virtually quantitative yield. Throughout the course of this study, no chromatography was required to obtain pure products.

To demonstrate the ability of organotrifluoroborates to serve in the role detailed in Scheme 1, in which a functionalized substrate could be elaborated and then converted into the

<sup>(8)</sup> For selected examples see: (a) Molander, G. A.; Cavalcanti, L. N.; Canturk, B.; Pan, P.-S.; Kennedy, L. E. J. Org. Chem. 2009, 74, 7364. (b) Kabalka, G. W.; Mereddy, A. R. Tetrahedron Lett. 2004, 45, 343. (c) Kabalka, G. W.; Mereddy, A. R. Organometallics **2004**, *23*, 4519.
 (9) Gillis, E. P.; Burke, M. D. J. Am. Chem. Soc. **2007**, *129*, 6716.

<sup>(10) (</sup>a) Ochiai, M.; Miyamoto, K.; Yokota, Y.; Suefuji, T.; Shiro, M. Angew. Chem., Int. Ed. 2005, 44, 75. (b) Sorin, G.; Mallorquin, R. M.; Contie, Y.; Baralle, A.; Malacria, M.; Goddard, J.-P.; Fensterbank, L. Angew. Chem., Int. Ed. 2010, 49, 8721.

<sup>(11)</sup> Catalog price: US\$ 7.00/mol.

 <sup>(12) (</sup>a) Dreher, S. D.; Lim, S.-E.; Sandrock, D. L.; Molander, G. A.
 J. Org. Chem. 2009, 74, 3626. (b) Gerbino, D. C.; Mandolesi, S. D.; Schmalz, H.-G.; Podesta, J. C. Eur. J. Org. Chem. 2009, 3964. (c) Molander, G. A.; Sandrock, D. L. J. Am. Chem. Soc. 2008, 130, 15792. (d) Dreher, S. D.; Dormer, P. G.; Sandrock, D. L.; Molander, G. A. J. Am. Chem. Soc. 2008, 130, 9257. (e) Molander, G. A.; Petrillo, D. E. J. Am. Chem. Soc. 2006, 128, 9634. (f) Molander, G. A.; Figueroa, R. J. Org. Chem. 2006, 71, 6135. (g) Smoum, R.; Rubinstein, A.; Srebnik, M. Org. Biomol. Chem. 2005, 3, 941. (h) Yuen, A. K. L.; Hutton, C. A. Tetrahedron Lett. 2005, 46, 7899. (i) Navarre, L.; Darses, S.; Genet, J.-P. Eur. J. Org. Chem. 2004, 69. (j) Molander, G. A.; Biolatto, B. J. Org. Chem. 2003, 68, 4302. (k) Molander, G. A.; Biolatto, B. Org. Lett. 2002, 4, 1867. (1) Molander, G. A.; Ito, T. Org. Lett. 2001, 3, 393. (m) Vedejs, E.; Chapman, R. W.; Fields, S. C.; Lin, S.; Schrimpf, M. R. J. Org. Chem. 1995, 60, 3020.

 TABLE 1.
 Oxidation of Electron-Rich and Electron-Neutral Potassium

 Aryltrifluoroborates<sup>a</sup>



<sup>*a*</sup>All reactions were carried out using 1 mmol of aryltrifluoroborate and Oxone (5 mL, 0.2 M in  $H_2O$ ) in 5 mL of acetone for 2 min at room temperature. <sup>*b*</sup>55 mmol scale.

phenol in a late synthetic stage, potassium (*Z*)-(4-(4-cyanobut-1-en-1-yl)phenyl)trifluoroborate was synthesized through a Wittig reaction of (4-formylphenyl)trifluoroborate and the nitrile-containing ylide<sup>12f</sup> (Scheme 2). The organotrifluoroborate was then converted into the phenol, using the oxidation





protocol with Oxone. The desired product was obtained in excellent yield in only 2 min without affecting either the nitrile or the double bond of the molecule.<sup>13</sup>

Next, attention was turned to electron-poor aryltrifluoroborates. The conditions developed worked equally well for these substrates, providing the phenols in excellent yields (Table 2). Substrates such as aldehyde-, ester-, keto-, nitrile-, and nitro-containing aryltrifluoroborates afforded the hydroxylated product without affecting the pendant functional groups (entries 7-11). Halogenated trifluoroborates also provide the corresponding phenol in excellent yields (entries 1-6).

To expand the utility of the developed conditions further, heteroaryl systems were investigated, including dibenzothiophenyl, dibenzofuranyl, pyridinyl, benzothiophenyl, benzofuranyl, thiophenyl, and furanyl derivatives (Table 3). Heteroaryls, such as dibenzo[*b*,*d*]furan-4-yl-, dibenzo[*b*,*d*]thiophen-4-yl-, 6-chloropyridin-3-yl-, and 6-fluoro-5-methylpyridin-3-yltrifluoroborate, afforded the desired phenol in excellent yields in only 5 min (entries 1–4). For substrates where the most stable tautomers are the carbonyl isomers,<sup>14</sup> only the keto tautomer was observed (entries 5–8). 2-Trifluoroboratofuran afforded the  $\beta$ , $\gamma$ -unsaturated lactone (entry 9), along with 10% of the  $\alpha$ , $\beta$ -unsaturated product. Interestingly, under the reaction conditions, no oxidation of the nitrogen or sulfur atom of these heterocycles was observed.<sup>15</sup>

The scope of the general reaction conditions was further extended to alkyl- and alkenyltrifluoroborates (Table 4). We were pleased to find that the desired alcohol was obtained in excellent yields from primary, secondary, and benzylic alkyltrifluoroborates (entries 1-8). Primary alcohols containing halogens or ester groups were generated without affecting these incorporated functional groups. Alkenyltrifluoroborates were converted into the corresponding aldehydes, also in excellent yields (entries 9-11). However, under the reaction conditions, minor amounts of the hydrated aldehyde were observed (entries 9 and 10). Oxidation of both alkyltrifluoroborates and alkenyltrifluoroborates was complete in only 2 min.

To investigate the stereochemical integrity of the process, the enantiomerically enriched  $\beta$ -trifluoroboratoamide was prepared from the corresponding  $\alpha$ , $\beta$ -unsaturated amide and bis(pinacolato)diboron in a copper-catalyzed process using (*R*)-(*S*)-Josiphos as the chiral ligand, according to the

<sup>(13)</sup> For selected examples see: (a) Zhu, W.; Ford, W. T. J. Org. Chem.
1991, 56, 7022. (b) Adam, W.; Hadjiarapoglou, L.; Smerz, A. Chem. Ber.
1991, 124, 227. (c) Corey, P. F.; Ward, F. E. J. Org. Chem.
1986, 51, 1925. (d) Bloch, R.; Abecassis, J.; Hassan, D. J. Org. Chem.
1985, 50, 1544. (e) Jeyaraman, R.; Murray, R. W. J. Am. Chem. Soc.
1984, 106, 2462. (f) Curci, R.; Fiorentino, M.; Troisi, L.; Edwards, J. O.; Pater, R. H. J. Org. Chem.
1980, 45, 4758.

<sup>(14) (</sup>a) Elguero, J.; Marzin, C.; Katritzky, A. R.; Linda, P. *The Tautomerism of Heterocyclic Compounds*; Academic Press: New York, 1976.
(b) Capon, B.; Kwok, F.-C. *Tetrahedron Lett.* **1986**, *27*, 3275.

<sup>(15)</sup> For selected examples see: (a) Greehalgh, R. P. Synlett 1992, 235.
(b) Itahara, T. Chem. Lett. 1991, 57, 1591. (c) Kettani, A. E.; Bernadou, J.; Meunier, B. J. Org. Chem. 1989, 54, 3213. (d) Murray, R. W.; Jeyaraman, R. J. Org. Chem. 1985, 50, 2847. (e) Evans, T. L.; Grade, M. M. Synth. Commun. 1986, 16, 1207. (f) Trost, B. M.; Curran, D. P. Tetrahedron Lett. 1981, 22, 1287.

TABLE 2. Oxidation of Electron-Poor Potassium Aryltrifluoroborates<sup>a</sup>



<sup>*a*</sup>All reactions were carried out using 1 mmol of aryltrifluoroborate and Oxone (5 mL, 0.2 M in  $H_2O$ ) in 5 mL of acetone for 2 min at room temperature.

procedures previously reported by Yun et al. (Scheme 3).<sup>16</sup> With the enantioeriched organotrifluoroborate in hand, this material was subjected to the oxidation conditions with Oxone. We were pleased to observe that the transformation transpired with complete retention of configuration. The desired alcohol was obtained in 97% yield in only 2 min, in a 94:6 (*R*:*S*) ratio.

TABLE 3. Hydroxylation of Potassium Heteroaryltrifluoroborates<sup>a</sup>

(HetAr)-BF <sub>3</sub> K <u>conditions</u> (HetAr)-OH						
entry	substrate	product		isolated yield (%)		
1	BF <sub>3</sub> K	ОН	3a	97		
2	BF <sub>3</sub> K	ОН	3b	95		
3	CI N BF <sub>3</sub> K	CI N OH	3c	91		
4	F N BF <sub>3</sub> K	F N OH	3d	94		
5	BF <sub>3</sub> K		3e	99		
6	BF <sub>3</sub> K		3f	97		
7	Br	Br	3g	94		
8	S BF3K	k s <sup>0</sup>	3h	89		
9	€ OBF <sub>3</sub> K		3i	97		

<sup>*a*</sup>All reactions were carried out using 1 mmol of aryltrifluoroborate and Oxone (5 mL, 0.2 M in  $H_2O$ ) in 5 mL of acetone for 2 min at room temperature.

In summary, we have successfully developed a general, rapid, and efficient method for the oxidation of aryl-, heteroaryl-, alkyl-, and alkenyltrifluoroborates. The reactions were complete in only 2–5 min at room temperature, affording the desired products in virtually quantitative yield. The use of Oxone as the oxidant makes the process both economical and environmentally sound. Numerous attendant functional groups were tolerated in this process, and no chromatography was required—a simple aqueous workup followed at most by filtration through a plug of silica gel afforded pure material. The oxidation of enantiomerically enriched secondary alkyltrifluoroborates affords the desired alcohols with complete retention of configuration.

## **Experimental Section**

General Experimental Procedure for the Preparation of Phenols (55 mmol Scale). Preparation of Phenol (1a)<sup>4</sup>. To a 50 mL round-bottom flask containing a mixture of potassium phenyltrifluoroborate (10.12 g, 55.0 mmol) and acetone (275 mL, 0.2 M) was added Oxone (275 mL of a 0.2 M solution in H<sub>2</sub>O, 1 equiv) in one portion. The reaction mixture was stirred at room temperature until <sup>11</sup>B NMR indicated completion of the reaction (~2 min). To the crude mixture was added H<sub>2</sub>O (30 mL) and

<sup>(16)</sup> Chea, H.; Sim, H.-S.; Yun, J. Adv. Synth. Catal. 2009, 351, 855.

	R BF₃K —	conditions R OH		
	ŔŔ	R R		
entry	substrate	product		isolated yield (%)
1	BzO	BzO ()2 OH	4a	99
2	BzO ()_BF <sub>3</sub> K	BzO ( )_OH	4b	99
3	Br BF <sub>3</sub> K	Br OH	4c	98
4	BF3K	ОН	4d	95
5	₩ <sup>™</sup> BF <sub>3</sub> K		4e	96
6	BF <sub>3</sub> K	-ОС-он	4f	98
7	BF <sub>3</sub> K	ОН	4g	99
8	BF <sub>3</sub> K	ОН	4h	99
9	OH H BF <sub>3</sub> K		4i	99
10	MeO BF <sub>3</sub> K	MeO H	4j	99
11			4k	99

TABLE 4. Hydroxylation of Potassium Alkyl- and Alkenyltrifluoroborates  $^a$ 

 $^{a}All$  reactions were carried out using 1 mmol of aryltrifluoroborate and Oxone (5 mL, 0.2 M in H<sub>2</sub>O) in 5 mL of acetone for 2 min at room temperature.

aqueous HCl (0.1 M, 20 mL), and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>( $3\times50$  mL). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated, and dried in vacuo. The crude extract was filtered through a small plug of silica topped with charcoal, with CH<sub>2</sub>Cl<sub>2</sub> as eluent, to afford the desired pure product in 96% yield (5.0 g, 53 mmol) as a light yellow solid, mp 41–43 °C (lit. mp 40–42 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.24 (t, *J*=7.5 Hz, 2H), 6.93 (t, *J*=7.5 Hz, 1H), 6.83 (d, *J*=8.5 Hz, 2H), 4.88 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  155.4, 129.7, 120.8, 115.3.

General Experimental Procedure for the Preparation of Phenols (1 mmol Scale). Preparation of Naphthalen-1-ol (1b)<sup>4</sup>. To a 50 mL round-bottom flask containing a mixture of potassium naphthalen-1-yltrifluoroborate (0.23 g, 1.0 mmol) and acetone (5 mL, 0.2 M) was added Oxone (5 mL of a 0.2 M solution in H<sub>2</sub>O, 1 equiv) in one portion. The reaction mixture was stirred at room temperature until <sup>11</sup>B NMR indicated completion of the reaction ( $\sim 2$  min). To the crude mixture was added H<sub>2</sub>O (5 mL) and aqueous HCl (0.1 M, 3 mL), and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>(3×15 mL). The combined organic layers SCHEME 3



were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated, and dried in vacuo. The crude extract was filtered through a small plug of silica topped with charcoal, with CH<sub>2</sub>Cl<sub>2</sub> as eluent, to afford the desired pure product in 99% yield (0.14 g, 0.99 mmol) as a white solid, mp 91–93 °C (lit.<sup>3a</sup> mp 92–94 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  8.17 (m, 1H), 7.81 (m, 1H), 7.50–7.48 (m, 2H), 7.44 (d, *J* = 8.5 Hz, 1H), 7.30 (t, *J* = 7.5 Hz, 1H), 6.81 (d, *J* = 7.5 Hz, 1H), 5.21 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  151.3, 134.8, 127.7, 126.4, 125.8, 125.2, 124.3, 121.5, 120.7, 108.6.

**Naphthalen-2-ol** (1c).<sup>4</sup> The general procedure was employed using potassium naphthalen-2-yltrifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 97% yield (0.14 g, 0.97 mmol) as a white solid, mp 121–123 °C (lit. mp 122–124 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.75 (t, J=8.9 Hz, 2H), 7.67 (d, J=8.2 Hz, 1H), 7.42 (m, 1H), 7.32 (m, 1H), 7.13 (m, 1H), 7.08 (m, 1H), 5.01 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  134.4, 134.3, 133.1, 130.9, 128.8, 127.9, 126.9, 126.6, 126.0.

**4-Methoxyphenol (1d).**<sup>4</sup> The general procedure was employed using potassium (4-methoxyphenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 98% yield (0.12 g, 0.98 mmol) as a white solid, mp 57–58 °C (lit. mp 57–58 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  6.79–6.75 (m, 4H), 5.31 (s, 1H), 3.76 (s, 3H). <sup>13</sup>C NMR (125.8 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  161.6, 137.6, 136.0, 114.0, 55.8.

**3-Methoxyphenol (1e).**<sup>4</sup> The general procedure was employed using potassium (3-methoxyphenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 93% yield (0.11 g, 0.93 mmol) as a light yellow oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.12 (m, 1H), 6.50 (m, 1H), 6.44–6.41 (m, 2H), 5.62 (br s, 1H), 3.75 (s, 3H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  160.8, 156.6, 130.2, 107.9, 106.5, 101.6, 55.3.

**2-Methoxyphenol (1f).**<sup>4</sup> The general procedure was employed using potassium (2-methoxyphenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 97% yield (0.12 g, 0.97 mmol) as a light yellow oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  6.93 (m, 1H), 6.88–6.84 (m, 3H), 5.61 (br s, 1H), 3.89 (s, 3H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  146.6, 145.7, 121.5, 120.1, 114.5, 110.7, 55.9. **2,4-Dimethoxyphenol (1g).**<sup>17</sup> The general procedure was em-

**2,4-Dimethoxyphenol (1g).**<sup>17</sup> The general procedure was employed using potassium (2,4-dimethoxyphenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 95% yield (0.15 g, 0.95 mmol) as a light yellow oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  6.75 (d, *J* = 9.0 Hz, 1H), 6.56 (d, *J* = 3.0 Hz, 1H), 6.37 (m, 1H), 5.78 (br s, 1H), 3.81 (s, 3H), 3.73 (s, 3H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  154.5, 146.4, 141.0, 111.5, 104.5, 101.8, 56.5, 55.6.

<sup>(17)</sup> Gonzalez, R. R.; Gambarotti, C.; Liguori, L.; Bjorsvik, H.-R. J. Org. Chem. 2006, 71, 1703.

<sup>(18)</sup> Ohkubo, M.; Mochizuki, S.; Sano, T.; Kawaguchi, Y.; Okamoto, S. Org. Lett. 2007, 9, 773.

**2,6-Dimethylphenol** (1h).<sup>4</sup> The general procedure was employed using potassium (2,6-dimethylphenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 94% yield (0.11 g, 0.94 mmol) as a white solid, mp 43–45 °C (lit. mp 42–43 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  6.96 (d, *J*=7.5 Hz, 2H), 6.75 (t, *J*=7.5 Hz, 1H), 4.61 (br s, 1H), 2.23 (s, 6H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  152.1, 128.6, 123.0, 120.2, 15.8.

152.1, 128.6, 123.0, 120.2, 15.8. **4-tert-Butylphenol** (**1i**).<sup>3d</sup> The general procedure was employed using potassium (4-tert-butylphenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.15 g, 0.99 mmol) as a white solid, mp 97–99 °C (lit. mp 96–99 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.25 (d, *J*=6.5 Hz, 2H), 7.17 (d, *J*=6.5 Hz, 2H), 4.66 (br s, 1H), 1.29 (s, 9H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  153.1, 143.6, 126.4, 114.7, 34.1, 31.5. **4-(Benzyloxy)phenol (1j).**<sup>18</sup> The general procedure was em-

**4-(Benzyloxy)phenol (1j).**<sup>18</sup> The general procedure was employed using potassium (4-(benzyloxy)phenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 97% yield (0.19 g, 0.97 mmol) as a light yellow solid, mp 117–121 °C (lit. mp 118–122 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.42–7.36 (m, 4H), 7.32 (m, 1H), 6.85 (d, J=9 Hz, 2H), 6.75 (d, J=9 Hz, 2H) 5.00 (s, 2H), 4.46 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  149.6, 137.2, 128.5, 127.9, 127.5, 116.1, 116.0, 70.8.

(*Z*)-5-(4-Hydroxyphenyl)pent-4-enenitrile (1k). The general procedure was employed using potassium ((*Z*)-4-(4-cyanobut-1-enyl)phenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 98% yield (0.17 g, 0.98 mmol) as a white solid, mp 67–69 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.12 (d, *J*=8.5 Hz, 2H), 6.81 (d, *J*=8.0 Hz, 2H) 6.52 (d, *J* = 11.5 Hz, 1H), 5.55 (m, 1H), 5.00 (br s, 1H), 2.69 – 2.65 (m, 2H), 2.44 (t, *J* = 7.0 Hz, 2H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  154.8, 131.6, 130.1, 129.2, 126.0, 119.2, 115.2, 24.4, 17.6. IR (neat) 3325, 2257, 1608, 1514, 1228, 836 cm<sup>-1</sup>. HRMS (ESI): *m/z* calcd for C<sub>11</sub>H<sub>11</sub>NONa (M + Na)<sup>+</sup> 196.0738, found 196.0741.

**4-Iodophenol** (2a).<sup>19</sup> The general procedure was employed using potassium (4-iodophenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 85% yield (0.19 g, 0.85 mmol) as a light yellow solid, mp 90–93 °C (lit. mp 92–95 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.50 (d, *J*=9.0 Hz, 2H), 6.61 (d, *J*=9.0 Hz, 2H), 5.00 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  155.2, 138.4, 117.8, 82.8.

**4-Bromophenol (2b).**<sup>20</sup> The general procedure was employed using potassium (4-bromophenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 98% yield (0.17 g, 0.98 mmol) as a light yellow solid, mp 55–58 °C (lit. mp 54–64 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.31 (d, J=8.5 Hz, 2H), 6.70 (d, J=9.0 Hz, 2H), 5.64 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  154.4, 132.5, 117.2, 112.9.

**4-Chlorophenol (2c).**<sup>4</sup> The general procedure was employed using potassium (4-chlorophenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.13 g, 0.99 mmol) as a white solid, mp 43–46 °C (lit. mp 44–45 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.18 (d, *J*=9.0 Hz, 2H), 6.75 (d, *J*=8.5 Hz, 2H), 5.52 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  153.9, 129.5, 125.7, 116.7.

**4-Fluorophenol (2d).**<sup>4</sup> The general procedure was employed using potassium (4-fluorophenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.11 g, 0.99 mmol) as a white solid, mp

45–47 °C (lit. mp 46–47 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 6.95–6.90 (m, 2H), 6.79–6.74 (m, 2H), 4.88 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>): δ 157.2 (d, J = 235.9 Hz), 151.4, 116.2 (d, J = 10.0 Hz, 2C), 116.0 (d, J = 22.9, 2C). <sup>19</sup>F NMR (470.8 MHz, CDCl<sub>3</sub>): δ –124.3.

**2,4-Difluorophenol (2e).**<sup>5f</sup> The general procedure was employed using potassium (2,4-difluorophenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 97% yield (0.13 g, 0.97 mmol) as a light yellow oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  6.95 (m, 1H), 6.85 (m, 1H), 6.76 (m, 1H), 5.75 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  155.9 (dd, J = 240.9, 10.3 Hz), 150.4 (dd, J = 240.1, 12.2 Hz), 139.9 (dd, J = 14.1, 3.5 Hz), 117.3 (d, J = 6.4 Hz), 111.7–110.6 (m), 104.5–103.3 (m). <sup>19</sup>F NMR (470.8 MHz, CDCl<sub>3</sub>):  $\delta$  –121.0, –135.7.

**4-(Trifluoromethyl)phenol (2f).**<sup>21</sup> The general procedure was employed using potassium (4-(trifluoromethyl)phenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 98% yield (0.16 g, 0.98 mmol) as a light yellow oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.50 (d, *J*=8.5 Hz, 2H), 6.90 (d, *J*=9 Hz, 2H) 5.45 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  158.1, 127.2 (q, *J* = 3.8 Hz), 125.4, 123.7 – 122.7 (m), 115.4. <sup>19</sup>F NMR (470.8 MHz, CDCl<sub>3</sub>):  $\delta$  -61.5.

122.7 (m), 115.4. <sup>19</sup>F NMR (470.8 MHz, CDCl<sub>3</sub>):  $\delta$  –61.5. **4-Hydroxybenzonitrile (2g).**<sup>5f</sup> The general procedure was employed using potassium (4-cyanophenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 98% yield (0.12 g, 0.98 mmol) as a light yellow solid, mp 107–109 °C (lit. mp 108–109 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.57 (d, *J*=9.0 Hz, 2H), 6.94 (d, *J*= 8.5 Hz, 2H), (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  160.0, 134.3, 119.2, 116.4, 103.3.

**4-Hydroxybenzaldehyde** (2h).<sup>4</sup> The general procedure was employed using potassium (4-formylphenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.12 g, 0.99 mmol) as a white solid, mp 114–117 °C (lit. mp 114–116 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  9.87 (s, 1H), 7.82 (d, J = 8.5 Hz, 2H), 6.97 (d, J = 9.0 Hz, 2H), 6.73 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  191.1, 161.4, 132.4, 129.9, 115.9.

Methyl 4-Hydroxybenzoate (2i).<sup>22</sup> The general procedure was employed using potassium 4-(methoxycarbonyl)phenyltrifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.15 g, 0.99 mmol) as a white solid, mp 121–123 °C (lit. mp 127–129 °C). <sup>1</sup>H NMR (500 MHz, acetone- $d_6$ ):  $\delta$  9.12 (br s, 1H), 7.89 (d, J=9.0 Hz, 2H), 6.92 (d, J=9.0 Hz, 2H), 3.82 (s, 3H). <sup>13</sup>C NMR (125.8 MHz, acetone- $d_6$ ):  $\delta$  167.8, 163.4, 133.2, 123.3, 116.8, 52.6. 1-(4-Hydroxyphenyl)ethanone (2j).<sup>20</sup> The general procedure

**1-(4-Hydroxyphenyl)ethanone (2j).**<sup>20</sup> The general procedure was employed using potassium (4-acetylphenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.13 g, 0.99 mmol) as a white solid, mp 105–107 °C (lit. mp 108–109 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  8.24 (br s, 1H), 7.92 (d, J = 9.0 Hz, 2H), 6.97 (d, J = 9.0 Hz, 2H), 2.59 (s, 3H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  199.0, 161.6, 131.3, 129.4, 115.6, 26.3.

CDCl<sub>3</sub>):  $\delta$  199.0, 161.6, 131.3, 129.4, 115.6, 26.3. **3-Nitrophenol** (2k).<sup>4</sup> The general procedure was employed using potassium (3-nitrophenyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 98% yield (0.14 g, 0.98 mmol) as a light yellow solid, mp 96–99 °C (lit. mp 98–99 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.82 (m, 1H), 7.71 (t, *J* = 2.0 Hz, 1H), 7.41 (t, *J* = 8.0 Hz, 1H), 7.19 (m, 1H), 5.61 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  156.3, 130.3, 122.0, 115.9, 110.6.

<sup>(19)</sup> Zhou, C.-Y.; Li, J.; Peddibhotla, S.; Romo, D. Org. Lett. 2010, 12, 2104.

<sup>(20)</sup> Magano, J.; Chen, M. H.; Clark, J. D.; Nussbaumer, T. J. Org. Chem. 2006, 71, 7103.

<sup>(21)</sup> Sergeev, A. G.; Schulz, T.; Torborg, C.; Spannenberg, A.; Neumann, H.; Beller, M. Angew. Chem., Int. Ed. 2009, 48, 7595.

<sup>(22)</sup> Sivakumar, S.; Reddy, M. L. P.; Cowleyb, A. H.; Vasudevanb, K. V. Dalton Trans. 2010, 39, 776.

**Dibenzo**[*b,d*]**furan-4-ol** (3a). The general procedure was employed using potassium (dibenzo[*b,d*]**furan-4-y**])trifluoroborate, and the reaction was complete in 5 min. The desired pure product was obtained in 97% yield (0.18 g, 0.97 mmol) as a white solid, mp 98–100 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.93 (d, *J* = 7.7 Hz, 1H), 7.57 (d, *J* = 8.3 Hz, 1H), 7.51 (d, *J* = 7.7 Hz, 1H), 7.46 (t, *J* = 7.8 Hz, 1H), 7.35 (t, *J* = 7.5 Hz, 1H), 7.23 (m, 1H), 7.02 (d, *J* = 7.9 Hz, 1H), 5.39 (s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  156.1, 144.0, 141.1, 127.3, 125.7, 124.6, 123.7, 1249, 744 cm<sup>-1</sup>. HRMS (ESI): *m*/*z* calcd for C<sub>12</sub>H<sub>7</sub>O<sub>2</sub> (M – H)<sup>-1</sup> 183.0446, found 183.0454.

**Dibenzo**[*b,d*]**thiophen-4-ol** (**3b**). The general procedure was employed using potassium (dibenzo[*b,d*]thiophen-4-yl)trifluoroborate, and the reaction was complete in 5 min. The desired pure product was obtained in 97% yield (0.18 g, 0.97 mmol) as a white solid, mp 158–160 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  8.13 (m, 1H), 7.89 (m, 1H), 7.78 (d, *J* = 7.9 Hz, 1H), 7.50–7.42 (m, 2H), 7.34 (t, *J* = 7.8 Hz, 1H), 6.88 (d, *J* = 7.7 Hz, 1H), 5.35 (s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  150.5, 139.6, 137.9, 135.9, 126.9, 126.5, 125.7, 124.4, 123.1, 122.0, 114.5, 111.7. IR (neat): 3230, 1569, 1444, 1253, 744 cm<sup>-1</sup>. HRMS (ESI): *m/z* calcd for C<sub>12</sub>H<sub>7</sub>OS (M – H)<sup>-</sup> 199.0218, found 199.0222.

**6-Chloropyridin-3-ol (3c).** The general procedure was employed using potassium (6-chloropyridin-3-yl)trifluoroborate, and the reaction was complete in 5 min. The desired pure product was obtained in 91% yield (0.12 g, 0.97 mmol) as a white solid, mp 155–157 °C. <sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>):  $\delta$  9.04 (br s, 1H), 7.99 (d, J=2.9 Hz, 1H), 7.40–7.14 (m, 2H). <sup>13</sup>C NMR (125.8 MHz, acetone-*d*<sub>6</sub>):  $\delta$  154.9, 142.5, 138.9, 127.6, 126.1. IR (neat): 3004, 1573, 1464, 1278, 1228, 826 cm<sup>-1</sup>. HRMS (ESI): *m/z* calcd for C<sub>5</sub>H<sub>3</sub>NOCl (M – H)<sup>-</sup> 127.9903, found 127.9902.

**6-Fluoro-5-methylpyridin-3-ol** (**3d**). The general procedure was employed using potassium (6-chloropyridin-3-yl)trifluoroborate, and the reaction was complete in 5 min. The desired pure product was obtained in 94% yield (0.12 g, 0.94 mmol) as a white solid, mp 120–122 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.69 (br s, 1H), 7.61 (s, 1H), 7.22 (m, 1H), 2.25 (s, 3H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  157.1, 155.2, 151.1, 129.8 (dd, *J* = 44.4, 9.5 Hz), 120.7 (d, *J* = 34.1 Hz), 14.4. <sup>19</sup>F NMR (470.8 MHz, CDCl<sub>3</sub>):  $\delta$  –83.9. IR (neat): 3048, 1471, 1232, 770 cm<sup>-1</sup>. HRMS (ESI): *m*/*z* calcd for C<sub>6</sub>H<sub>7</sub>NOF (M + H)<sup>+</sup> 128.0512, found 128.0510.

**Benzo**[*b*]thiophen-2(3*H*)-one (3e).<sup>23</sup> The general procedure was employed using potassium (benzothiophen-2-yl)trifluoroborate, and the reaction was complete in 5 min. The desired pure product was obtained in 99% yield (0.15 g, 0.99 mmol) as a light yellow oil. <sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>):  $\delta$  7.44 (m, 1H), 7.38 (d, *J* = 7.5 Hz, 1H), 7.33 (m, 1H), 7.25 (m, 1H), 4.09 (s, 2H). <sup>13</sup>C NMR (125.8 MHz, acetone-*d*<sub>6</sub>):  $\delta$  203.7, 138.3, 134.7, 129.8, 127.7, 126.7, 124.5, 48.4.

**Benzofuran-2**(*3H*)**-one** (**3f**).<sup>24</sup> The general procedure was employed using potassium (benzofuran-2-yl)trifluoroborate, and the reaction was complete in 5 min. The desired pure product was obtained in 97% yield (0.13 g, 0.97 mmol) as a light yellow oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.32–7.24 (m, 2H), 7.13 (m, 1H), 7.08 (d, *J* = 8.1 Hz, 1H), 3.71 (s, 2H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  174.0, 154.6, 128.8, 124.5, 123.9, 123.0, 110.6, 32.8.

**5-Bromobenzo**[*b*]**thiophen-2**(*3H*)**-one** (**3g**). The general procedure was employed using potassium (5-bromobenzo[*b*]thiophen-2-yl)trifluoroborate, and the reaction was complete in

5 min. The desired pure product was obtained in 94% yield (0.21 g, 0.94 mmol) as a light yellow solid, mp 117–120 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.49–7.37 (m, 2H), 7.21 (d, *J*=8.9 Hz, 1H), 3.97 (s, 2H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  201.5, 136.1, 133.9, 131.4, 127.9, 124.3, 119.7, 47.1. FT-IR (neat): 1711, 1017, 819 cm<sup>-1</sup>. HRMS (ESI): *m/z* calcd for C<sub>8</sub>H<sub>4</sub>OSBr (M – H)<sup>-</sup> 226.9166, found 226.9158.

**4-Methylthiophen-2**(*3H*)-one (3h). The general procedure was employed using potassium (4-methylthiophen-2-yl)trifluoroborate, and the reaction was complete in 5 min. The desired pure product was obtained in 89% yield (0.10 g, 0.89 mmol) as a light yellow oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  6.09 (m, 1H), 3.97 (s, 2H), 2.21 (s, 3H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  199.7, 167.7, 129.0, 40.8, 18.9. FT-IR (neat) 2918, 1676, 1096, 653 cm<sup>-1</sup>. HRMS (ESI): *m*/*z* calcd for C<sub>5</sub>H<sub>7</sub>OS (M + H)<sup>+</sup> 115.0218, found 115.0215.

115.0218, found 115.0215.
Furan-2(3H)-one (3i).<sup>25</sup> The general procedure was employed using potassium (thiophen-3-yl)trifluoroborate, and the reaction was complete in 5 min. The desired pure product was obtained in 98% yield (0.10 g, 0.98 mmol) as a light yellow oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 6.77 (m, 1H), 5.55 (m, 1H), 3.14 (s, 2H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>): δ 176.5, 143.8, 105.4, 32.3.

2H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  176.5, 143.8, 105.4, 32.3. **4-Hydroxybutyl Benzoate** (4a).<sup>26</sup> The general procedure was employed using potassium (4-(benzoyloxy)butyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.19 g, 0.99 mmol) as a colorless oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  8.04–8.02 (m, 2H), 7.53 (m, 1H), 7.44–7.40 (m, 2H), 4.35 (t, *J*=6.5 Hz, 2H), 3.70 (t, *J*=6.5 Hz, 2H), 2.60, (br s, 1H), 1.88–1.83 (m, 2H), 1.74–1.69 (m, 2H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  166.7, 132.8, 130.2, 129.4, 128.3, 64.8, 62.0, 29.1, 25.1.

**6-Hydroxyhexyl Benzoate (4b).**<sup>27</sup> The general procedure was employed using potassium (7-(benzoyloxy)hexyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.23 g, 0.99 mmol) as a colorless oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  8.05–8.03 (m, 2H), 7.53 (m, 1H), 7.44–7.41 (m, 2H), 4.31 (t, *J*=6.5 Hz, 2H), 3.63 (t, *J*=6.5 Hz, 2H), 2.38, (br s, 1H), 1.80–1.75 (m, 2H), 1.62–1.56 (m, 2H), 1.48–1.43 (m, 4H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  166.6, 132.7, 130.2, 129.4, 128.2, 64.8, 62.4, 32.4, 28.5, 25.7, 25.3.

**3-Bromopropan-1-ol** (4c).<sup>28</sup> The general procedure was employed using potassium (3-bromopropyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 98% yield (0.14 g, 0.98 mmol) as a colorless oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  3.78 (t, J = 6.0 Hz, 2H), 3.54 (t, J = 6.5 Hz, 2H), 2.69 (br s, 1H), 2.11–2.07 (m, 2H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  60.1, 34.9, 30.3.

**Cyclopentanol** (4d).<sup>29</sup> The general procedure was employed using potassium cyclopentyltrifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.09 g, 0.99 mmol) as a colorless oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  4.33 (m, 1H), 1.78–1.76 (m, 4H), 1.59 (br s, 1H), 1.58–1.56 (m, 4H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  74.0, 35.5, 23.2.

**2-Methylcyclohexanol** (**4e**).<sup>30</sup> The general procedure was employed using potassium (2-methylcyclohexyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 96% yield (0.11 g, 0.96 mmol) as a

<sup>(23)</sup> Chen, S.; Zhao, X.; Chen, J.; Chen, J.; Kuznetsova, L.; Wong, S. S.; Ojima, I. *Bioconjugate Chem.* 2010, *21*, 979.

<sup>(24)</sup> Kim, M.-J.; Kim, J.-J.; Won, J.-E.; Kang, S.-E; Park, S.-E.; Jung, K.-J.; Lee, S.-G.; Yoon, Y.-J. *Bull. Korean Chem. Soc.* **2008**, *29*, 2247.

<sup>(25)</sup> Yanai, H.; Takahashi, A; Taguchi, T. Tetrahedron Lett. 2007, 48, 2993.

<sup>(26)</sup> Ochiai, M.; Miyamoto, K.; Yokota, Y.; Suefuji, T.; Shiro, M. Angew. Chem., Int. Ed. 2005, 44, 75.

<sup>(27)</sup> Saburi, H.; Tanaka, S.; Kitamura, M. *Angew. Chem., Int. Ed.* **2005**, 44, 1730.

 <sup>(28)</sup> Shah, S. T. A.; Singh, S.; Guiry, P. J. J. Org. Chem. 2009, 74, 2179.
 (29) Abraham, R. J.; Koniotou, R.; Sancassan, F. J. Chem. Soc., Perkin Trans. 2 2002, 2025.

<sup>(30)</sup> Brown, H. C.; Weissman, S. A.; Perumal, P. T.; Dhokte, U. P. J. Org. Chem. **1990**, 55, 1217.

colorless oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  3.11 (m, 1H), 1.95–1.93 (m, 1H), 1.75–1.59 (m, 4H), 1.36–1.18 (m, 4H), 1.01 (d, J = 6.5 Hz, 3H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  76.5, 40.2, 35.5, 33.6, 25.7, 25.2, 18.5.

**3,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol** (**4f**).<sup>31</sup> The general procedure was employed using potassium (3,6,6-trimethylbicyclo[3.1.1]heptan-2-yl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 98% yield (0.15 g, 0.98 mmol) as a white solid, mp 49–52 °C (lit. mp 51–53 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  4.05 (m, 1H), 2.64 (br s, 1H), 2.50 (m, 1H), 2.37 (m, 1H), 2.00–1.90 (m, 2H), 1.79 (m, 1H), 1.73 (m, 1H), 1.21 (s, 3H), 1.13 (d, *J*=7.4 Hz, 3H), 1.06 (d, *J* = 9.7 Hz, 1H), 0.92 (s, 3H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  71.3, 47.7, 47.4, 41.67, 38.8, 38.1, 34.2, 27.6, 23.6, 20.6. IR (neat): 3260, 2905, 1469, 1043, 1006, 923 cm<sup>-1</sup>. HRMS (ESI): *m/z* calcd for C<sub>10</sub>H<sub>17</sub> (M – OH)<sup>+</sup> 137.1330, found 137.1326. **2-Phenylethanol (4g).**<sup>27</sup> The general procedure was employed

**2-Phenylethanol (4g).**<sup>27</sup> The general procedure was employed using potassium phenethyltrifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.12 g, 0.99 mmol) as a colorless oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.30–7.27 (m, 2H), 7.21–7.18 (m, 3H), 3.78 (t, J = 6.5 Hz, 2H), 2.81 (t, J = 6.5 Hz, 2H), 2.20 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  138.6, 129.0, 128.5, 126.4, 63.5, 39.1. **Phenylmethanol (4h).**<sup>32</sup> The general procedure was employed

**Phenylmethanol (4h).**<sup>32</sup> The general procedure was employed using potassium benzyltrifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.11 g, 0.99 mmol) as a colorless oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.32–7.25 (m, 5H), 4.56 (s, 2H), 2.90 (br s, 1H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  140.8, 128.4, 127.4, 126.9, 64.9.

**11-Hydroxydodecanal** (**4i**).<sup>33</sup> The general procedure was employed using potassium ((*E*)-11-hydroxydodec-1-en-1-yl)-trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.2 g, 0.99 mmol) as a colorless oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  9.76 (t, J=1.9 Hz, 1H), 3.79 (m, 1H), 2.44–2.40 (m, 2H), 1.69–1.50 (m, 2H), 1.41 (d, J=14.2 Hz, 2H), 1.29 (s, 12H), 1.19 (d, J=6.2 Hz, 3H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  68.2, 43.9, 39.3, 29.6, 29.3, 29.1, 25.7, 22.1.

**Methyl 6-Oxohexanoate** (4j).<sup>34</sup> The general procedure was employed using potassium ((*E*)-6-methoxy-6-oxohex-1-en-1-yl)-trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.14 g, 0.99 mmol) as a colorless oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  9.77 (t,

J = 1.2 Hz, 1H), 3.67 (s, 3H), 2.55–2.25 (m, 4H), 1.67 (m, 4H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  202.0, 174.0, 173.7, 51.5, 43.5, 33.7, 24.3, 21.5.

**2-(2,2-Dimethyl-4-oxo-4***H***-benzo[***d***][1,3]dioxin-5-yl)acetaldehyde (4k). The general procedure was employed using potassium ((***E***)-2-(2,2-dimethyl-4-oxo-4***H***-benzo[***d***][1,3]dioxin-5-yl)vinyl)trifluoroborate, and the reaction was complete in 2 min. The desired pure product was obtained in 99% yield (0.22 g, 0.99 mmol) as a light yellow solid, mp 55–57 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): \delta 9.84 (s, 1H), 7.50 (t,** *J* **= 8 Hz, 1H), 6.96 (d,** *J* **= 8.3 Hz, 1H), 6.91 (d,** *J* **= 7.5 Hz, 1H), 4.22 (s, 2H), 1.75 (s, 6H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>): \delta 198.1, 160.8, 157.1, 137.0, 135.6, 126.4, 117.0, 112.6, 105.7, 49.0, 25.6. IR (neat): 1722, 1292, 1052 cm<sup>-1</sup>. HRMS (ESI):** *m/z* **calcd for C<sub>12</sub>H<sub>12</sub>O<sub>4</sub>Na (M + Na)<sup>+</sup> 243.0633, found 243.0633.** 

(*R*)-3-Hydroxy-*N*-(4-methoxyphenyl)butanamide (5).<sup>35</sup> The general procedure was employed using 0.1 mmol of potassium ((*R*)-4-((4-methoxyphenyl)amino)-4-oxobutan-2-yl)trifluoroborate (*R*: *S* 97:3), and the reaction was complete in 2 min. The desired pure product was obtained in 97% yield (0.020 g, 0.097 mmol) as a white solid, mp 136–138 °C (lit. mp 135 °C). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.75 (br s, 1H), 7.39 (d, *J*=9.0 Hz, 2H), 6.85 (d, *J*=9.0 Hz, 2H), 4.28 (m, 1H), 3.78 (s, 3H), 3.43 (d, *J*=2.9 Hz, 1H), 2.52–2.41 (m, 2H), 1.27 (d, *J*=6.5 Hz, 3H). <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>):  $\delta$  170.3, 156.5, 130.5, 122.0, 114.1, 64.9, 55.5, 44.8, 22.9.

Acknowledgment. This work was generously supported by the NIGMS (Grant Nos. R01 GM086209, GM035249) and a Conselho Nacional de Desenvolvimento Cientítico e Tecnológico (CNPq) Graduate Research Fellowship to L.N.C. We also acknowledge Aldrich, BoroChem, and Frontier Scientific for their donation of the boronic acids and Dr. Deidre Sandrock for providing the enantiomerically enriched organotrifluoroborate sample. Dr. Rakesh Kohli (University of Pennsylvania) is acknowledged for obtaining the HRMS data.

**Supporting Information Available:** Figures giving <sup>1</sup>H, <sup>13</sup>C, and <sup>19</sup>F spectra for all compounds prepared by the method described. This material is available free of charge via the Internet at http://pubs.acs.org.

<sup>(31)</sup> Brown, H. C.; Gupta, A. K. J. Organomet. Chem. 1988, 341, 73.
(32) Jagdale, A. R.; Paraskar, A. S.; Sudalai, A. Synthesis 2009, 660.

<sup>(33)</sup> Li, X.-Q.; Zhang, C. Synthesis 2009, 1163.

<sup>(34)</sup> Hickmann, V.; Alcarazo, M.; Furstner, A. J. Am. Chem. Soc. 2010, 132, 11042.

<sup>(35)</sup> Chea, H.; Sim, H.-S.; Yun, J. Adv. Synth. Catal. 2009, 351, 855.