

# **CHEMISTRY** A European Journal



Direct oxidation c trifluoromethylate	of Csp3-H bo	ndo using in			
	a dioxirane	in flow	situ generated		
<b>)rs:</b> Mathieu Lesie Jacq, Christoj Pasau	eur, Claudio phe Genicot	Battilocchio, , Steven Victo	Ricardo Labes, or Ley, and Pat	, Jérôme rick	
manuscript has be pted Article online e final Version of g the Digital Obje shed online in Ea is Accepted Artic /oR from the jou isure accuracy of ent of this Accept	een accepter e prior to edi Record (Vo ct Identifier rly View as s le as a resu rnal website information. ed Article.	d after peer re iting, proofing oR). This wor (DOI) given oon as possii ult of editing. e shown belo . The authors	view and appe , and formal pu k is currently o below. The Vo ble and may be Readers shou w when it is p are responsib	ars as an ublication itable by R will be different Id obtain published le for the	
cited as: Chem. Eu	r. J. 10.1002/	/chem.20180	5657		
o VoR: http://dx.do	oi.org/10.100	)2/chem.2018	805657		
				S	upport
					ACE
	Pasau manuscript has be epted Article onlin e final Version of g the Digital Obje shed online in Ea is Accepted Artic VoR from the jou sure accuracy of ent of this Accepte e cited as: <i>Chem. Eu</i> to VoR: http://dx.de	Pasau manuscript has been accepter epted Article online prior to edi e final Version of Record (Vo g the Digital Object Identifier shed online in Early View as s is Accepted Article as a resu- VoR from the journal website isure accuracy of information. ent of this Accepted Article. e cited as: <i>Chem. Eur. J.</i> 10.1002, to VoR: http://dx.doi.org/10.100	manuscript has been accepted after peer re epted Article online prior to editing, proofing e final Version of Record (VoR). This worl g the Digital Object Identifier (DOI) given to shed online in Early View as soon as possit is Accepted Article as a result of editing. VoR from the journal website shown belo issure accuracy of information. The authors ent of this Accepted Article. e cited as: <i>Chem. Eur. J.</i> 10.1002/chem.201805 to VoR: http://dx.doi.org/10.1002/chem.2018	manuscript has been accepted after peer review and apperent of the Digital Object Identifier (DOI) given below. The Vo shed online in Early View as soon as possible and may be is Accepted Article as a result of editing. Readers shou VoR from the journal website shown below when it is presure accuracy of information. The authors are responsible ent of this Accepted Article.	add, Christophe Genicol, Steven Victor Ley, and Patrick Pasau manuscript has been accepted after peer review and appears as an opted Article online prior to editing, proofing, and formal publication e final Version of Record (VoR). This work is currently citable by g the Digital Object Identifier (DOI) given below. The VoR will be shed online in Early View as soon as possible and may be different is Accepted Article as a result of editing. Readers should obtain VoR from the journal website shown below when it is published usure accuracy of information. The authors are responsible for the ent of this Accepted Article. ecited as: Chem. Eur. J. 10.1002/chem.201805657 to VoR: http://dx.doi.org/10.1002/chem.201805657

COMMUNICATION

## WILEY-VCH

# Direct oxidation of C<sub>sp</sub><sup>3</sup>-H bonds using *in-situ* generated trifluoromethylated dioxirane in flow

Mathieu Lesieur,\*<sup>[a]</sup> Claudio Battilocchio,<sup>[b][c]</sup> Ricardo Labes,<sup>[b]</sup> Jérôme Jacq,<sup>[a]</sup> Christophe Genicot,<sup>[a]</sup> Steven V. Ley,\*<sup>[b]</sup> and Pasau Patrick\*<sup>[a]</sup>

**Abstract:** A fast, scalable and safer  $C_{sp}^{3}$ -H oxidation of activated and un-activated aliphatic chain can be enabled by methyl(trifluoromethyl)dioxirane (TFDO). The continuous flow platform allows the *in situ* generation of TFDO gas and its rapid reactivity toward tertiary and benzylic  $C_{sp}^{3}$ -H bonds. The process exhibits a broad scope and good functional group compatibility (28 examples, 8 - 99 %). The scalability of this methodology is demonstrated on 2.5 g scale oxidation of adamantane.

The development of new and powerful synthetic methodology for direct C<sub>sp</sub><sup>3</sup>-H oxidation of un-activated methylene and methine systems is of considerable importance.<sup>1</sup> More specifically, the site-specific C<sub>sp</sub><sup>3</sup>-H hydroxylation of valuable scaffolds bearing further functional groups is of major current interest for both academia and industry.<sup>2</sup> Over the past decade, direct Csp<sup>3</sup>-H oxidation has been investigated using transition metal complexes (Ru, Fe, V, Co, Mn or Ir)<sup>3</sup> and some organocatalysts<sup>4</sup>. Unfortunately, specific challenges including cost of the catalyst, complex synthetic design of the ligand, use of directing group, harsh reaction conditions, functional group incompatibility and site selectivity remain as major drawbacks to the field (Figure 1).<sup>5</sup> Nowadays, new chemical methods such as chemo-catalytic approaches,<sup>6</sup> electrochemical oxidation<sup>7</sup> or photochemical C<sub>sp</sub><sup>3</sup>-H hydroxylation in pure oxygen<sup>8</sup> or using hydrogen peroxide<sup>9</sup> offer alternative approaches (Figure 1). Nevertheless, regio-selective and stereo-selective oxidation of an un-activated Csp3-H bond remains a challenging task.

Among existing  $C_{sp}^{3}$ -H oxidation strategies, the use of strong oxidants such as dimethyldioxirane (DMDO, 1)<sup>10</sup> and methyl(trifluoromethyl)dioxirane (TFDO, 2)<sup>11</sup>, have a long history of success on complex systems without the need for a directing group. Regrettably, their use has been limited due to a long list of disadvantages such as instability, volatility, laborious preparation, safety requirements and scale limitations.<sup>12</sup> Moreover, the use of dioxirane for the direct  $C_{sp}^{3}$ -H oxidation requires precise control of the temperature below 10 °C, a large excess of reagent and long reaction times.<sup>13</sup> For all these reasons, dioxiranes have only been deployed on batch mode and limited to small scale application. Recently, Hilinski published a catalytic batch method for the *in situ* formation of a less volatile but expensive reactive dioxirane (3-

 [a] Dr Mathieu Lesieur, Dr Jérôme Jacq, Dr Christophe Genicot and Dr Patrick Pasau
 UCB Biopharma, Avenue de l'industrie, 1420 Braine l'Alleud, Belgium
 E-mail: patrick.pasau@ucb.com

[b] Dr Claudio Battilocchio, Dr Ricardo Labes and Pr Steven V. Ley Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, United Kingdom E-Mail: svl1000@cam.ac.uk

 [c] Dr Claudio Battilocchio
 Syngenta Crop Protection AG, Schaffhauserstrasse, CH-4332 Stein, Switzerland butyl-3-(trifluoropentyl)dioxirane, **3**). Specifically, limitations of this method include low temperature (4 °C), long reaction time (72 hours), the use of HFIP (hexafluoro-2-propanol) as co-solvent and iterative addition of reagents to achieve satisfactory yield.<sup>14</sup>



Figure 1. Methods for the direct C<sub>sp</sub><sup>3</sup>-H oxidation.

In recent years, continuous flow technologies have been recognized as a powerful tool to generate *in situ* hazardous reagents or unstable intermediates with an efficient mass and heat transfer.<sup>15</sup> Additionally, it offers other advantages in making procedures more reproducible and scalable. With an interest in further developing methods for  $C_{sp}^3$ -H oxidation, we sought to develop an alternative direct  $C_{sp}^3$ -H oxidation using TFDO in a rapid and safer continuous flow process.

A flow set up was investigated using adamantane (5a) as a model substrate for the C<sub>sp</sub><sup>3</sup>-H oxidation (Table 1). A solution of adamantane (0.05 M) in dichloromethane containing 1,1,1trifluoroacetone (4, 1.0 M) was mixed with an aqueous solution of NaHCO<sub>3</sub> (1.5 M) and Oxone (0.5 M) in a sequential manner in order to generate in situ the TFDO. The desired 1-adamantanol (6a) can be efficiently obtained with a conversion of 93 % (Table 1, Entry 1) in an optimal residence time of 80 seconds. A better mass transfer of the reaction mixture was observed with the use of a coil reactor packed with glass beads (425 - 600  $\mu m$  diameter) compared to the conventional reactor (Table 1, Entry 2 vs. Entry 1). Decreasing the amount of the 1,1,1-trifluoroacetone (4, 0.5 M) or increasing the quantity of adamantane (5a, 0.1 M) have, respectively, an negative effect on the conversion (60 %, Table 1, Entry 3 vs. Entry 1) and (62 %, Table 1, Entry 4 vs. Entry 1). An excess of NaHCO<sub>3</sub> (1.5 M) compared to Oxone (0.5 M) is required to reach higher conversion (Table 1, Entry 5 vs. Entry 1), whereas decreasing their flow rates showed inferior results (Table 1, Entry 6 vs. Entry 1). Moreover, the reactions proceeded at room temperature, and under pressure, rendering the reaction conditions simple and more reliable than in batch mode<sup>12d</sup> (Table 1, Entry 7 vs. Entry 1). Reducing the residence time to 32 seconds, led to a lower conversion (79 %) (Table 1, Entry 8 vs. Entry 1).

## COMMUNICATION

Finally, the use of the less volatile 1,1,1-trifluorobutanone (7) did not provide a substantial improvement on the reaction outcome (Table 1, Entry 9 vs. Entry 1) and no reaction was observed with acetone as a source of dioxirane due to the formation of a precipitate (Table 1, Entry 10 vs. Entry 1).

Table 1. Optimization for the  $C_{\text{sp}}{}^3\text{-}\text{H}$  oxidation of adamantane (5a).



[a] Reaction condition: 0.25 mmol scale reaction. Reaction carried out at 0.05 M concentration of **5a** and 1.0 M of **4** in DCM, 1.5 M concentration of NaHCO<sub>3</sub> in water and 0.5 M concentration of Oxone in water, 5 mL glass beads reactor, residence time: 80 seconds, 25 °C. [b] Conversion determined by <sup>1</sup>H NMR, average of two reactions.

To confirm the *in situ* formation of TFDO (**2**) in the reaction, a solution of CDCl<sub>3</sub> containing 1,1,1-trifluoroacetone (1.0 M) was mixed with an aqueous solution of NaHCO<sub>3</sub> (1.5 M) and Oxone (0.5 M) using the flow set up depicted in Scheme 1. After a continuous phase separation, using the Zaiput unit, and collection of the organic phase, <sup>19</sup>F and HMBC <sup>1</sup>H/<sup>13</sup>C NMR spectroscopy revealed the formation of TFDO (**2**) (<sup>19</sup>F NMR = -87.3 ppm).<sup>16</sup> Furthermore, titration of the organic phase, revealed a molarity of TFDO (**2**) around 0.05 M (5 % yield)<sup>17,18</sup> This results emphasise the benefit of the flow procedure to achieve a safe, faster and higher TFDO concentration than reported in batch mode (2 % yield).<sup>12d</sup>

With the optimized conditions in hand, the scope for this transformation was explored as shown in Scheme 2. As expected, investigations of the substrate scope of this new flow method revealed a strong preference for hydroxylation of tertiary centers.





Scheme 1. Production of TFDO in flow.

In the same way, trans-decaline (5g) and trans-hydrindane (5h) afforded the  $C_{sp}^{3}$ -H oxidation products 6g and 6h at the tertiary position with an isolated yield of 74 % and 98 %, respectively. Interestingly, product of tertiary hydroxylation remote position are strongly preferred over proximal oxidable position. For example, Csp3-H oxidation products of 5i can be obtained with an isolated yield of 68 % (6i) and 7 % (6i') with a ratio of (91:9) toward the remote  $C_{sp}^3$ -H bond. Lower yields were obtained for substrates (5j) and (5k) bearing only one hindered proximal tertiary carbon (6j, 21 % and 6k, 8 %). Importantly, the retention of configuration was achieved on the L-menthyl-acetate scaffold (51). As previously described, the  $C_{sp}^{3}$ -H oxidation occurred, preferentially, at the two tertiary remote positions with an isolated yield of 18 % (6I) and 21 % (6I'), without racemization, suggesting no deviation from the current mechanistic understanding of dioxirane hydroxylation.<sup>18</sup> Product **6m** can be selectively oxidized on the tertiary position with an isolated yield of 31 %. Notably, increasing the residence time and the concentration of 1,1,1-trifluoroacetone (4) by a factor of 2 showed higher isolated yield in the desired product (75 %). Activated methylenes, such as benzylic Csp3-H bonds bearing cyclohexyl (5n) and cyclobutyl (5o) ring can be oxidized, at the tertiary carbon, with an isolated yield of 42 % and 51 %, respectively. investigated with Regio-selectivity was substrate (4methylcyclohexyl)benzene (5p) bearing an activated benzylic position and an un-activated C<sub>sp</sub><sup>3</sup>-H bond.

## COMMUNICATION



Scheme 2. Substrate scope for the direct Csp<sup>3</sup>-H oxidation using TFDO.<sup>[a]</sup> [a] Reaction conditions: 0.25 mmol scale reaction. Reaction carried out at 0.05 M concentration of **5** and 2.0 M of **4** in DCM, 1.5 M concentration of NaHCO<sub>3</sub> in water and 0.5 M concentration of Oxone in water, 5 mL glass beads reactor, residence time: 80 seconds, 25 °C. Isolated yield after chromatography, average of two reactions. [b] Reaction carried out at 0.05 M concentration of **5** and 4.0 M of **4** in DCM, 1.5 M concentration of Oxone in water, 20 mL helicoidal static-mixer coil reactor, residence time: 160 seconds, 25 °C. [c] 0.125 mmol scale reaction. Reaction carried out at 0.025 M concentration of **5** and 2.0 M of **4** in DCM. [d] 1.25 mmol scale reaction. Reaction carried out at 0.025 M concentration of **5** and 2.0 M of **4** in DCM. [d] 1.25 mmol scale reaction. Reaction carried out at 0.025 M concentration of **5** and 4.0 M of **4** in DCM.

The benzylic position was preferentially oxidized with an isolated yield of 27 % toward the un-activated methine position (**6p**', 9 %) with a ratio of (**6p:6p'** = 3:1). Interestingly, indole derivative bearing a sulfonyl chloride functionality (**5q**) can be tolerated. Hydroxylation reaction occurred on the benzylic position with an isolated yield of 24 %. The reaction was also achieved on a 1.25 mmol scale reaching an isolated yield of 56 % with an increase of the 1,1,1-trifluoroacetone (**4**) by a factor of 2.

Secondary  $C_{sp}^{3}$ -H bond oxidation was evaluated using few substrates bearing un-activated and activated positions. The reaction on ibuprofen methyl ester (**5r**) led to oxidation on both benzylic and tertiary  $C_{sp}^{3}$ -H in a 3:2 ratio favouring the tertiary position. In the case of protected phthalimide-methyl ester scaffold bearing a cyclohexyl ring (**5s**), a mixture (1/1) of two regio-isomer ketone **6s** and **6s'** was obtained in low yield 8 % in both cases. The excellent potential of TFDO to oxidize secondary

# COMMUNICATION

alcohols was also demonstrated with camphor (95 %, **6t**) and on 1-adamantanone (98 %, **6u**). In general, a variety of functionalities are well tolerated under the reaction conditions. For example, ester, nitrile, carbamate, phthalimide, ketone and sulfonyl chloride remain intact during the reaction, except for non-protected primary amine (**6e**) which are oxidized to the nitro derivative, quantitatively which is an interesting reaction in its own right. Natural products such as sclareolide (**5v**) and cholestane (**5w**) bearing multiple oxidative sites were studied. In the case of the sclareolide, the C<sub>sp</sub><sup>3</sup>-H oxidation was selectively observed at the secondary carbon position C2 whereas for cholestane the transformation occurred preferentially at the terminal tertiary position.



Figure 2. Retrosynthetic step for the synthesis of Adapalene<sup>18</sup>

The scalability of this flow procedure has been demonstrated on 2.7 g synthesis of the alcohol (**6a**), a common starting material for the preparation of Adapalene, a retinoid drug (Figure 2).<sup>20</sup> The initial flow set-up has been scale out by increasing the size of the reactor from 5 mL to 40 mL to accommodate the critical mixing requirement at high flow rate with the use of two "helicoidal static-mixer coil". A 2.3 hours run at steady state in standard conditions provided the desired product **6a** with similar yield (96 %) to the smaller scale reaction with this particular reactor design (Scheme 3).



Scheme 3. The 2.5 grams scale C<sub>sp</sub><sup>3</sup>-H oxidation of adamantane (5a).

In summary, the development of a new general trifluoromethylated dioxirane-mediated  $C_{sp}^3$ -H oxidation method has been enabled in a fast, efficient and simple fashion through a continuous flow process. This practical method employs inexpensive reagents and can be applied to a wide panel of substrates bearing a large range of functionalities. The flow conditions provide a scalable and safer solution for transforming simple aliphatic compounds to complex materials.



#### Acknowledgements

Acknowledgements Text The authors gratefully acknowledge the Walloon region (Belgium) – DG06 (Convention  $n^{\circ}7240$ ) for funding.

Keywords: oxidation • Flow • TFDO • dioxirane • Oxone

- W. Lu, L. Zhou in Oxidation of C-H Bonds; John Wiley & Sons; Hoboken, New Jersey, 2017.
- [2] a) T. Newhouse, P. S. Baran, Angew. Chem. Int. Ed. 2011, 50, 3362; b)
  M. C. White, Science 2012, 335, 807; c) M. Canta, M. Rodríguez, M. Costas, Top. Curr. Chem. 2016, 372, 27; d) D. Font, M. Canta, M. Milan,
  O. Cussó, X. Ribas, R. J. M. Klein Gebbink, M. Costas, Angew. Chem. Int. Ed. 2016, 55, 5776; e) K. Kamata, K. Yonehara, Y. Nakagawa, K. Uehara, N. Mizuno, Nat. Chem. 2010, 2, 478; f) D. P. Hruszkewycz, K. C. Miles, O.R. Thiel, S. S. Stahl, Chem. Sci. 2017, 8, 1282; g) N. Sauermann, T. Meyer, C. Tian, L. Ackermann, J. Am. Chem. Soc. 2017, 139, 18452; h) G. Olivo, G. Farinelli, A. Barbieri, O. Lanzalunga, S. Di Stefano, M. Costas, Angew. Chem. Int. Ed. 2017, 56, 16347; i) E. M. Simmons, J. F. Hartwig, Nature 2012, 482, 70; j) M. Zhou, N. D. Schley, R. H. Crabtree, J. Am. Chem. Soc. 2010, 132, 12550.
- [3] a) E. McNeill, J. Du Bois, *Chem. Sci.* 2012, 3, 1810; b) E. McNeill, J. Du Bois, *J. Am. Chem. Soc.* 2010, *132*, 10202.
- [4] a) D. Wang, W. G. Shuler, C. J. Pierce, M. K. Hilinski, Org. Lett. 2016, 18, 3826; b) C. J. Pierce, M. K. Hilinski, Org. Lett. 2014, 16, 6504; c) A. M. Adams, J. Du Bois, Chem. Sci. 2014, 5, 656; d) B. H. Brodsky, J. Du Bois, J. Am. Chem. Soc. 2005, 127, 15391; e) M. Lee, M. S. Sandford, Org. Lett. 2017, 3, 572; f) X. Li, X. Che, G.-H. Chen, J. Zhang, J.-L. Yan, Y.-F. Zhang, L.-S. Zhang, C.-P. Hsu, Y. Q. Gao, Z.-J. Shi, Org. Lett. 2016, 6, 1234.
- [5] P. E. Gormisky, M. C. White, J. Am. Chem. Soc. 2013, 135, 14052.
- a) X. Ren, J. A. Yorke, E. Taylor, T. Zhang, W. Zhou, L. L. Wong, *Chem. Eur. J.* 2015, *21*, 15039; b) J. Genovino, D. Sames, L. G. Hamann, B. B. Touré, *Angew. Chem. Int. Ed.* 2016, *55*, 14218.
- [7] Y. Kawamata, M. Yan, Z. Liu, D.-H. Bao, J. Chen, J. T. Starr, P. S. Baran, J. Am. Chem. Soc. 2017, 139, 7448.
- [8] a) G. Laudadio, S. Govaerts, Y. Wang, D. Ravelli, H. F. Koolman, M. Fagnoni, S. W. Djuric, T. Noël, *Angew. Chem. Int. Ed.* 2018, *15*, 4078.
  [9] D. M. Schultz, F. Lévesque, D. A. Di Rocco, M. Reibarkh, Y. Ji, L. A. Dropinski, J. F. Joyce, H. Sheng, B. D. Sherry, I. W. Davies, *Angew. Chem. Int. Ed.* 2017, *48*, 15476.
- [10] a) J. O. Edwards, R. H. Pater, P. R. Curci, F. Di Furia, *Photochem. Photobiol.* **1979**, *30*, 63; b) J. K. Crandall, R. Curci, L. D'Accolti, C. Fusco, Dimethyldioxirane. In *Encyclopedia of reagents for Organic Synthesis*; John Wiley & Sons; New York, **2005**.
- [11] a) R. Mello, M. Fiorentino, C. Fusco, R. Curci, J. Am. Chem. Soc. 1989, 111, 6749; b) R. W. Murray, R. J. Jeyaraman, Org. Chem. 1985, 50, 2847; c) G. Asensio, M. E. Gonzalez-Nunez, C. Boix Bernardini, R. Mello, W. Adam, J. Am. Chem. Soc, 1993, 115, 7250; d) J. K. Crandall, R. Curci, L. D'Accolti, C. Fusco, Methyl(trifluoromethyl)dioxirane. In Encyclopedia of reagents for Organic Synthesis; John Wiley & Sons; New York, 2005.
- [12] a) H. Mikula, D. Svatunek, D. Lumpi, F. Glöcklhofer, C. Hametner, J. Fröhlich, Org. Process Res. Dev. 2013, 17, 313; b) M. Chabanas, Teles, G. Heydrich, R. S. Sanderson, US 2008/0177092 A1, Jul. 24, 2008; c) R. Mello, M. E. González-Núñez, G. Asensio, Synlett 2007, 1, 47; d) W. Adam, C. Van Barneveld, D. Golsh, Tetrahedron, 1996, 52, 2377.
- [13] a) J. S. Lee, P. L. Fuchs, *Org. Lett.* 2003, *5*, 2247; b) S. Kasuya, S. Kamijo, M. Inoue, *Org. Lett.* 2009, *11*, 3630; c) K. Chen, P. S. Baran, *Nature* 2009, *459*, 824; d) K. Chen, A. Eschenmoser, P. S Baran, *Angew. Chem. Ed.* 2009, *48*, 9705.
- [14] W. G. Shuler, S. L. Johnson, M. K. Hilinski, Org. Lett. 2017, 19, 4790.
- [15] a) M. Movsisyan, E. I. P. Delbeke, J. K. E. T. Berton, C. Battilocchio, S. V. Ley, C. V. Stevens, *Chem. Soc. Rev.* **2016**, *45*, 4892; b) D. Webb, T.

# COMMUNICATION

F. Jamison, *Chem. Sci.* **2010**, *1*, 675; c) B. Gutmann, D. Cantillo, C. O. Kappe, *Angew. Chem. Int. Ed.* **2015**, *54*, 6688.

- [16] X. Creary, T. E. Aldridge, J. Org. Chem. 1988, 53, 3890.
- [17] J. N. Ennis, Mechanisms and applications of dioxirane chemistry, Thesis, Loughborough University, 1998.
- [18] See Supporting Information for experimental details.
- [19] a) L. Zou, R. S. Paton, A. Eschenmoser, T. R. Newhouse, P. S. Baran, K. N. Houk, *J. Org. Chem.* **2013**, *78*, 4037; b) X. Du, K. N. Houk, *J. Org. Chem.* **1998**, *63*, 6480.
- [20] Z. Liu, J. Xiang, Org. Process Res. Dev. 2006, 10, 285.

# COMMUNICATION

#### Entry for the Table of Contents (Please choose one layout)

Layout 1:

TFDO.

## COMMUNICATION

