



Synthesis of acyl derivatives of salicin, salirepin, and arbutin



Elena V. Stepanova*, Maxim L. Belyanin, Victor D. Filimonov

Department of Biotechnology and Organic Chemistry, National Research Tomsk Polytechnic University, 30 Lenin Avenue, Tomsk 634050, Russia

ARTICLE INFO

Article history:

Received 15 October 2013

Received in revised form 9 February 2014

Accepted 10 February 2014

Available online 20 February 2014

Keywords:

Phenolic acyl glycosides

Populoside

Salirepin derivatives

Salicin derivatives

Benzoylarbutin

ABSTRACT

The total synthesis of two natural phenolglycosides of the family *Salicaceae*, namely: populoside and 2-(β -D-glucopyranosyloxy)-5-hydroxy benzyl (3-methoxy-4-hydroxy) cinnamate and nine not found yet in plants acyl derivatives of phenoglycosides: 2-(β -D-glucopyranosyloxy)-benzylcinnamate, 2-(β -D-glucopyranosyloxy)-benzyl (4-hydroxy) benzoate, 2-(β -D-glucopyranosyloxy)-benzyl (3-methoxy-4-hydroxy) benzoate, 2-(β -D-glucopyranosyloxy)-5-hydroxy benzyl (3,4-dihydroxy) cinnamate, 2-(β -D-glucopyranosyloxy)-5-hydroxy benzylcinnamate, 2-(β -D-glucopyranosyloxy)-5-hydroxy benzyl (4-hydroxy) benzoate, 2-(β -D-glucopyranosyloxy)-5-hydroxy benzyl (3-methoxy-4-hydroxy) benzoate, 2-(β -D-glucopyranosyloxy)-5-benzoyloxy benzylbenzoate and 4-(β -D-glucopyranosyloxy)-phenylbenzoate, starting from readily available phenols and glucose was developed for the first time.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction

The existence of phenolic glycosides in plants has been known for many years. The first, Salicin, was isolated in 1828 from willow bark.¹ Since then, many glycosides of different structures were isolated and identified.² These substances are of great interest for medicinal use because of their diverse biological activities. Plants of the family *Salicaceae* with large phenolglycoside content are widely used in traditional medicines all over the world and helpful for cure of pox, variola, pulmonary disease,³ arthritis,⁴ and other diseases. Most of phenolglycosides have anti-inflammatory activity⁵ and antioxidant activity.^{6,7}

The structure of *Salicaceae* phenolglycosides includes salicin (glucoside of salicylic alcohol) or salirepin (glucoside of gentisinic alcohol) acylated by different benzoic or cinnamic acids. It is not absolutely clear yet, but biological effects of phenolglycosides are more likely caused by the aglycon and especially by the nature of phenolic acid.

The presence of several phenolic acids in hydrolyzed methanol extracts from different species of *Salix* was established,⁸ namely, *p*-hydroxybenzoic, vanillic, cinnamic, *p*-coumaric, ferulic, and caffeic acids. Obviously, such acids exist as free or as parts of glycosides.⁹

Indeed, it is known, that caffeic acid has anti-inflammatory and analgesic activity.¹⁰ It has been found to be pharmacologically active as an antioxidant, antimutagenic, anticarcinogenic agent, and as a lipoxygenase inhibitor¹¹ and forms transitional metal

ion-caffeic acid complexes.¹² Ferulic acid has antioxidant, antimicrobial, anti-inflammatory, anti-thrombosis, and anti-cancer activities.¹³ *para*-Hydroxybenzoic acid, basic component of parabenes, shows antimicrobial activity,¹⁴ antifungal, antimutagenic, antiseptic, and estrogenic activity.¹⁵ Vanillic acid has hepatoprotective effect^{16,17} and exerts protective effects on cardiotoxic organisms.¹⁸

Thus, the structure of glycosides containing different combinations of salicin or salirepin moiety and one or more phenolic acids is the most predictable (Fig. 1). For confirmation of this, new phenolglycoside, consisting of salirepin and caffeic acid (**19**) was recently isolated.⁶

We previously reported the synthesis of some natural phenolglycosides of the family *Salicaceae*, containing cinnamic acids, namely, populosides A, B, and C.¹⁹ We suggested a simple synthetic pathway for phenolglycosides of different structures. Thus, it would be reasonable to obtain glycosides of expected structure if they were not isolated from natural sources yet to make easier further phytochemical or biological investigations.

2. Results and discussion

The initial glycoside **1** was easily obtained by bromination of tetra-*O*-acetyl-*o*-cresylglycoside, which, in turn, was obtained by glycosylation of *o*-cresole.²⁰ Glycosides **2**, **3**, and **4** were obtained according to the method described previously.¹⁹ Acylation of **1** was carried out by phenolic acids with sodium hydrocarbonate in DMF²¹ medium (Scheme 1). Acyl-glycosides **5–6** were obtained in good yields.

Acylation of compound **2**, **3**, and **4** was carried out by acyl chloride of proper acid with pyridine to produce glycosides **7–14**.

* Corresponding author. Tel.: +7 8 (3822) 56 38 61; fax: +7 8 (3822) 56 36 37.

E-mail address: eline_m@mail.ru (E.V. Stepanova).






Figure 1. Structure of synthesized phenolglycosides **15–24**, **27**.



Scheme 1. Synthesis of 2-acyl phenolglycosides. Reagents and conditions: (a) acyl chloride of proper acid (3,4-acetoxycinnamoyl chloride, cinnamoyl chloride, 4-acetoxy-3-methoxy cinnamoyl chloride, 4-acetoxycinnamoyl chloride, 3-methoxy-4-acetoxycinnamoyl chloride, benzoylchloride), pyridine, CHCl₃, 24 h, 20 °C; (b) proper acid (3,4-acetoxycinnamic acid, cinnamic acid, 4-acetoxycinnamic acid, 3-methoxy-4-acetoxycinnamic acid), NaHCO₃, DMF, RT, 24 h; (c) 36% HCl, CHCl₃, EtOH (1:1:3), 8–13 h, 30 °C.



Scheme 2. Synthesis of benzoyl-arbutin. Reagents and conditions: (d) acetobromoglucose, quinoline, Ag₂O, RT, 1 h; (c) as in the Scheme 1.

For the selective cleavage of acetyl groups in the presence of other acyl groups we applied the system, suggested previously: HCl–EtOH (96%)–chloroform in a molar ratio of HCl–glycoside 54:1. The reaction at 30 °C for 8–13 h resulted in successful acetyl group cleavage without significant cleavage of labile phenacyl groups and without breaking the glycosidic bond. All resulted glycosides **15–24** had β -configuration of anomeric center and *trans*-configuration of double bond esters of cinnamic acids ($J = 15.9$ – 16.2 Hz). Physicochemical data for populoside **15**, and glycoside **19** matched those reported.^{6,22,23}

Glycoside **24**, 5-benzoylsalireposide, is isomeric to natural compound, 3'-benzoylsalireposide, found in *Symplocos racemosa*.²⁴ Natural compound showed strong inhibitory activity against snake venom phosphodiesterase I. Thus, substance **24** has potentially the same biological activity.

We also obtained benzoyl-arbutin **27** (Scheme 2), from acetate **26**. Substance **26** is well-known intermediate product for production of arbutin²⁵ and forms by glycosylation of monobenzoylhydroquinone **25**. Arbutin is natural phenolglycoside that nowadays

is used in cosmetic industry.²⁶ Several attempts of 6'-O-benzoylarbutin synthesis were taken both by chemical^{27,28} and enzymatic²⁹ means, but obtaining arbutin with aglycon benzoyl group **26** has not been previously reported.

In conclusion, we performed the first total synthesis of two known natural phenolglycosides, namely, populoside **15** and caffeoylsalirepin **19**. The resulted substances are identical to natural samples according to their physico-chemical properties. We also synthesized nine phenolglycosides of expected structure **15–17**, **19–23**, **26** that could be found in *Salicaceae* and have biological activity.

3. Experimental

3.1. General experimental procedures

Melting points, which are uncorrected, were determined using MP50 Melting point system (Mettler toledo). UV spectroscopic data were obtained with SF-102 spectrophotometer. IR spectra were recorded with IR Fourier spectrophotometer Spectrum BX II using

KBr disks. The ^1H and ^{13}C NMR spectra were recorded on Bruker-300 spectrometer at 300 and 75.5 MHz, respectively, in CDCl_3 , $\text{DMSO}-d_6$, and $\text{MeOD}-d_4$ with TMS as an internal standard and $\text{Cr}(\text{acac})_3$ as a relaxant. The chemical shifts are given in δ (ppm) and the spin–spin coupling constants (J) in hertz. Elemental analysis was performed on a EuroEA-3000 CHNS-O elemental analyzer. GC–MS analysis was performed using Agilent 7890A/5975C GC/MSD instrument, electron energy 70 eV. The ion source temperature was 230 °C, with the quadrupole temperature 150 °C and evaporator temperature 315 °C, employing a $30.000 \times 0.25 \text{ mm} \times 0.25 \mu\text{m}$ HP–5MS fused-silica capillary column. Helium was used as carrier gas at a constant flow of 1 mL/min and an inlet temperature of 315 °C. The column temperature was initially held at 150 °C for 2 min and then the temperature was raised to 315 °C at a rate of 20 °C/min, followed by isothermal period of 25 min. The total run time was 35.25 min. TLC was performed using plates Silica gel 60 F254 Merck and Sorbfil-UV 254 using benzene–ethanol 9:1 (method A) or chloroform–methanol 4:1 (B) mixtures as eluents. HPLC analysis was carried out with the liquid chromatographer Agilent Compact LC with column 150×4.6 Exlips Plus C-18 ($5 \mu\text{m}$). Analysis was performed using 0.1% trifluoroacetic acid in H_2O – CH_3CN as mobile phase, at gradient elution (from 0% to 100% CH_3CN in 20 min) at a flow rate of 1 mL/min. Probe volume was 20 μL . UV detection was performed at 220 nm. Accurate mass measurement was performed on an Agilent 1200 series LC system coupled with an Agilent 6210 TOF mass spectrometer. Silica gel MN Kieselgel 60 0.04–0.063 mm was used for column chromatography. Commercially available solvents were used after drying with CaCl_2 .

3.2. 2-(2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosyloxy)-benzyl bromide (1)

The compound **1** was obtained according to²⁰ by glycosylation of *o*-cresole and further radical bromination of 2-(2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosyloxy)-toluene. Total yield 16% from *o*-cresole, mp 150–151 °C. UV λ_{max} (EtOH)/nm: 277. IR (KBr, $\nu_{\text{max}}/\text{cm}^{-1}$): 2960, 1750, 1603, 1490, 1380, 1240, 1210, 1040, 108, 755. ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ : 1.98, 2.01, 2.04 (s, $4 \times 3\text{H}$, Ac), 4.07 (m, 1H, H-5'), 4.21–4.30 (m, 2H, H-6'b, H-7b), 4.49–4.58 (m, 2H, C-6'a, C-7a), 4.99 (t, 1H, $J = 9.6 \text{ Hz}$, H-4'), 5.12 (m, 1H, H-3'), 5.41 (m, 1H, H-2'), 5.56 (d, 1H, $J = 7.8 \text{ Hz}$, H-1'), 7.06 (m, 2H, H-2, H-4), 7.32 (m, 1H, H-3), 7.42 (d, 1H, $J = 7.8 \text{ Hz}$, H-5). ^{13}C NMR ($\text{DMSO}-d_6$, 75.5 MHz) δ : 20.4 ($4 \times \text{CH}_3$, Ac), 28.9 (CH_2 , C-7), 61.8 (CH_2 , C-6'), 68.0 (CH, C-4'), 70.2 (CH, C-2'), 70.9 (CH, C-3'), 71.8 (CH, C-5'), 97.0 (CH, C-1'), 114.9 (CH, C-2), 122.9 (CH, C-4), 126.5 (C, C-3), 130.2 (CH, C-6), 131.2 (CH, C-5), 154.1 (C, C-1), 169.0, 169.2, 169.4, 169.9 ($4 \times \text{C}$, Ac).

3.3. 2-(2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosyloxy)-5-acetyl-oxy benzyl alcohol (2), 2-(2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosyloxy)-benzyl alcohol (3) and 2-(2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosyloxy)-5-benzoyloxy benzyl alcohol (4)

The compounds **2**, **3**, and **4** were obtained according to the method described earlier.¹⁹

3.4. Alkylation of glycoside 1. General method

A mixture of 0.150 g (0.29 mmol) of glycoside **1**, 0.35 mmol of proper acid, 0.35 mmol of sodium bicarbonate, and 1 mL of DMF was stirred at room temperature for 24–48 h. After reaction was complete (HPLC control), the reaction mixture was poured into 5 mL of water and stirred until obtained glycoside precipitated, filtered and recrystallized from ethanol. If glycoside did not precipitate, it was extracted twice with CHCl_3 . CHCl_3 layer was washed with satd Na_2CO_3 , water, dried over Na_2SO_4 , and evaporated. The residue was recrystallized from ethanol.

3.4.1. 2-(2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosyloxy)-benzyl (3,4-diacetoxy) cinnamoate (populoside hexaacetate) (5)

The compound **5** was obtained from 3,4-diacetoxy cinnamic acid. Yield 78%, mp 93–94 °C, lit.²² 93–95 °C. UV λ_{max} (EtOH)/nm: 279. IR (KBr, $\nu_{\text{max}}/\text{cm}^{-1}$): 2960, 1760, 1640, 1380, 1240, 1040, 907, 760. ^1H NMR (CDCl_3 , 300 MHz) δ : 2.04, 2.06, 2.09, 2.12, 2.30 (s, $6 \times 3\text{H}$, Ac), 3.88 (m, 1H, H-5'), 4.16 (d, 1H, $J = 10.8 \text{ Hz}$, H-6'b), 4.25 (dd, 1H, $J = 5.0$, 12.3 Hz, H-6'a), 5.08 (m, 1H, H-1'), 5.15–5.22 (m, 2H, H-4', H-3'), 5.26–5.32 (m, 3H, H-2', 2H-7), 6.35 (d, 1H, $J = 15.9 \text{ Hz}$, H-9), 7.08 (m, 2H, H-2, H-4), 7.20 (d, 1H, $J = 8.4 \text{ Hz}$, H-15), 7.26 (m, 2H, H-3, H-5), 7.36 (m, 2H, H-12, H-16), 7.62 (d, 1H, $J = 16.2 \text{ Hz}$, H-10). ^{13}C NMR (CDCl_3 , 75.5 MHz) δ : 20.7 ($6 \times \text{CH}_3$, Ac), 61.2 (CH_2 , C-6'), 61.9 (CH_2 , C-7), 68.3 (CH, C-4'), 71.0 (CH, C-2'), 72.0 (CH, C-3'), 72.6 (CH, C-5'), 99.3 (CH, C-1'), 115.9 (CH, C-2), 119.0 (CH, H-9), 122.8 (CH, C-15), 123.6 (CH, C-4), 124.0 (CH, C-12), 126.2 (CH, C-6), 126.5 (CH, C-16), 129.5 ($2 \times \text{CH}$, C-3, C-5), 133.2 (C, C-11), 142.4 (C, C-13), 143.2 (CH, C-10), 143.5 (C, C-14), 154.5 (C, C-1), 166.3 (C=O, C-8), 168.1, 169.4, 170.3 ($6 \times \text{C=O}$, Ac).

3.4.2. 2-(2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosyloxy)-benzylcinnamoate (6)

The compound **6** was obtained from cinnamic acid. Yield 65%, mp 116–118 °C. UV λ_{max} (EtOH)/nm: 278. IR (KBr, $\nu_{\text{max}}/\text{cm}^{-1}$): 1750, 1640, 1500, 1370, 1210, 1030, 907. ^1H NMR (CDCl_3 , 300 MHz) δ : 2.03, 2.04, 2.06, 2.09 (s, $4 \times 3\text{H}$, Ac), 3.85 (m, 1H, H-5'), 4.16 (d, 1H, $J = 12.3 \text{ Hz}$, H-6'b), 4.25 (dd, 1H, $J = 5.0$, 12.3 Hz, H-6'a), 5.09 (d, 1H, $J = 7.2 \text{ Hz}$, H-1'), 5.14–5.20 (m, 2H, H-4', H-3'), 5.27–5.30 (m, 3H, H-2', 2H-7), 6.45 (d, 1H, $J = 15.3 \text{ Hz}$, H-9), 6.94 (m, 2H, H-2, H-4), 7.29 (m, 1H, H-5), 7.36 (m, 4H, H-3, H-13, H-14, H-15), 7.50 (m, 2H, H-12, H-16), 7.69 (d, 1H, $J = 15.9 \text{ Hz}$, H-10). ^{13}C NMR (CDCl_3 , 75.5 MHz) δ : 20.3 ($4 \times \text{CH}_3$, Ac), 60.7 (CH_2 , C-6'), 61.6 (CH_2 , C-7), 68.0 (CH, C-4'), 70.7 (CH, C-2'), 71.7 (CH, C-3'), 72.3 (CH, C-5'), 99.1 (CH, C-1'), 115.6 (CH, C-2), 117.6 (CH, H-9), 123.3 (CH, C-4), 126.1 (C, C-6), 127.8 ($2 \times \text{CH}$, C-12, C-16), 128.6 ($2 \times \text{CH}$, C-13, C-15), 129.2 (CH, C-3), 129.8 (CH, C-5), 130.1 (CH, C-14), 134.0 (C, C-11), 144.8 (CH, C-10), 154.2 (C, C-1), 166.3 (C=O, C-8), 169.0, 169.9, 170.2 ($4 \times \text{C=O}$, Ac). Anal. Calcd for $\text{C}_{30}\text{H}_{32}\text{O}_{12}$: C, 61.64; H, 5.52. Found: C, 61.48; H, 5.50.

3.5. Acylation of glycoside 2, 3, and 4. General method

To a solution of glycoside **2**, **3**, or **4** (0.2 mmol) in 1 mL CHCl_3 , 0.22 mmol of acyl chloride and 0.26 mmol of pyridine were added. The reaction mixture was kept at room temperature for 24 h and diluted with 20 mL CHCl_3 . The solution was washed with 0.1 M H_2SO_4 , satd Na_2CO_3 , water, dried over Na_2SO_4 and evaporated. The residue was recrystallized from ethanol.

3.5.1. 2-(2,3,4,6-Tetra-*O*-acetyl- β -D-glucopyranosyloxy)-benzyl (4-acetoxy) benzoate (7)

The compound **7** was obtained from 4-acetoxy benzoic acid and glycoside **3**. Yield 53%, mp 135–137 °C. UV λ_{max} (EtOH)/nm: 235. IR (KBr, $\nu_{\text{max}}/\text{cm}^{-1}$): 1740, 1730, 1600, 1370, 1210, 1040, 910, 750. ^1H NMR (CDCl_3 , 300 MHz) δ : 2.03, 2.04, 2.05, 2.07, 2.31 (s, $5 \times 3\text{H}$, Ac), 3.84–3.88 (m, 1H, H-5'), 4.16 (dd, 1H, $J = 2.1$, 12.3 Hz, H-6'b), 4.25 (dd, 1H, $J = 5.1$, 12.3 Hz, H-6'a), 5.11 (d, 1H, $J = 7.2 \text{ Hz}$, H-1'), 5.15 (m, 1H, H-4'), 5.25–5.44 (m, 2H, H-2', H-3'), 5.30 (m, 2H, H-7), 7.11 (m, 2H, H-2, H-4), 7.16 (d, 2H, $J = 8.4 \text{ Hz}$, H-11, H-13), 7.29 (m, 1H, H-3), 7.40 (d, 1H, $J = 7.2 \text{ Hz}$, H-5), 8.09 (d, 2H, $J = 8.4 \text{ Hz}$, H-10, H-14). ^{13}C NMR (CDCl_3 , 75.5 MHz) δ : 20.6, 21.2 ($5 \times \text{CH}_3$, Ac), 61.5 (CH_2 , C-6'), 61.9 (CH_2 , C-7), 68.4 (CH, C-4'), 71.1 (CH, C-2'), 72.1 (CH, C-3'), 72.7 (CH, C-5'), 99.4 (CH, C-1'), 116.0 (CH, C-2), 121.7 ($2 \times \text{CH}$, C-11, C-13), 123.6 (CH, C-4), 126.4 (C, C-6), 127.7 (C, C-9), 129.3 (C, C-3), 129.4 (C, C-5), 131.3 ($2 \times \text{CH}$, C-10, C-14), 154.4 (C, C-1), 154.5 (C, C-12), 165.4 (C=O, C-8), 168.8,

169.4, 170.2, 170.5 ($5 \times \text{C}=\text{O}$, Ac). Anal. Calcd for $\text{C}_{30}\text{H}_{32}\text{O}_{14}$: C, 58.44; H, 5.23. Found: C, 58.31; H, 5.39.

3.5.2. 2-(2,3,4,6-Tetra-O-acetyl- β -D-glucopyranosyloxy)-benzyl (3-methoxy-4-acetoxy) benzoate (8)

The compound **8** was obtained from 3-methoxy-4-acetoxy benzoic acid and glycoside **3**. Yield 66%, mp 122–124 °C. UV λ_{max} (EtOH)/nm: 242, 291. IR (KBr, $\nu_{\text{max}}/\text{cm}^{-1}$): 2950, 1740, 1600, 1370, 1240, 1050, 1040, 910, 750. ^1H NMR (CDCl_3 , 300 MHz) δ : 2.03, 2.05, 2.07, 2.34 (s, $5 \times 3\text{H}$, Ac), 3.88 (m, 1H, H-5'), 3.88 (s, 3H, OCH_3), 4.16 (dd, 1H, $J = 2.1$, 12.0 Hz, H-6'b), 4.26 (dd, 1H, $J = 5.1$, 12.3 Hz, H-6'a), 5.09 (d, 1H, $J = 7.5$ Hz H-1'), 5.14 (m, 1H, H-4'), 5.25–5.44 (m, 2H, H-2', H-3'), 5.30 (m, 2H, H-7), 7.08 (m, 3H, H-2, H-4, H-13), 7.26 (m, 1H, H-3), 7.39 (d, 1H, $J = 6.9$ Hz, H-5), 7.68 (s, 1H, H-10), 7.68 (dd, 1H, $J = 1.2$, 9.0 Hz, H-14). ^{13}C NMR (CDCl_3 , 75.5 MHz) δ : 20.6 ($5 \times \text{CH}_3$, Ac), 56.2 (OCH_3), 61.6 (CH_2 , C6'), 61.9 (CH_2 , C-7), 68.3 (CH, C-4'), 71.0 (CH, C-2'), 72.1 (CH, C-3'), 72.7 (CH, C-5'), 99.5 (CH, C-1'), 113.6 (CH, C-10), 115.9 (CH, C-2), 122.7 (CH, C-4), 122.8 (CH, C-13), 123.7 (CH, C-14), 126.4 (CH, C-6), 128.9 (C, C-9), 129.3 (C, C-5), 129.4 (CH, C-3), 143.7 (C, C-12), 151.1 (C, C-11), 154.5 (C, C-1), 165.6 ($\text{C}=\text{O}$, C-8), 168.5, 169.3, 169.4, 170.2, 170.6 ($5 \times \text{C}=\text{O}$, Ac). Anal. Calcd for $\text{C}_{31}\text{H}_{34}\text{O}_{15}$: C, 57.58; H, 5.30. Found: C, 57.40; H, 5.55.

3.5.3. 2-(2,3,4,6-Tetra-O-acetyl- β -D-glucopyranosyloxy)-5-acetoxy benzyl (3,4-diacetoxy) cinnamoate (9)

The compound **9** was obtained from 3,4-diacetoxy cinnamoyl chloride and glycoside **4**. Yield 85%, mp 131–132 °C. UV λ_{max} (EtOH)/nm: 280. IR (KBr, $\nu_{\text{max}}/\text{cm}^{-1}$): 1750, 1640, 1500, 1370, 1210, 1030, 905. ^1H NMR (CDCl_3 , 300 MHz) δ : 2.02, 2.03, 2.06, 2.10, 2.27, 2.28, 2.29 (s, $7 \times 3\text{H}$, Ac), 3.82 (m, 1H, H-5'), 4.15 (dd, 1H, $J = 9.9$, 1.7 Hz, H-6'a), 4.24 (dd, 1H, $J = 5.3$, 12.0 Hz, H-6'b), 5.02 (m, 1H, H-1'), 5.12 (m, 1H, H-4'), 5.17 (m, 2H, H-7), 5.27 (m, 2H, H-2', H-3'), 6.39 (d, 1H, $J = 15.9$ Hz, H-9), 6.97 (dd, 1H, $J = 2.5$, 8.7 Hz, H-3), 7.09 (d, 1H, $J = 7.2$ Hz, H-2), 7.10 (s, 1H, H-5), 7.19 (d, 1H, $J = 8.4$ Hz, H-15), 7.36 (s, 1H, H-12), 7.38 (d, 1H, $J = 8.4$ Hz, H-16), 7.62 (d, 1H, $J = 16.2$ Hz, H-10). ^{13}C NMR (CDCl_3 , 75.5 MHz) δ : 20.6 ($6 \times \text{CH}_3$, Ac), 21.0 (CH_3 , Ac), 60.6 (CH_2 , C-7), 61.8 (CH_2 , C-6'), 68.1 (CH, C-4'), 70.9 (CH, C-2'), 72.0 (CH, C-3'), 72.5 (CH, C-5'), 99.7 (CH, C-1'), 117.1 (CH=CH, C-9), 118.7 (CH, C-2), 122.1 ($2 \times \text{CH}$, C-3, C-5), 122.7 (CH, C-15), 123.9 (CH, C-12), 126.5 (CH, C-16), 127.8 (C, C-6), 133.1 (C, C-11), 142.4 (C, C-13), 143.4 (C, C-14), 145.5 (CH=CH, C-10), 146.3 (C, C-4), 151.8 (C, C-1), 166.0 ($\text{C}=\text{O}$, C-8), 168.0, 169.3, 169.6, 170.2, 170.5 ($7 \times \text{C}$, Ac). HRESIMS Calcd for $[\text{M}+\text{Na}]^+$ 781.19504. Found $[\text{M}+\text{Na}]^+$ 781.19502.

3.5.4. 2-(2,3,4,6-Tetra-O-acetyl- β -D-glucopyranosyloxy)-5-acetoxy benzylcinnamoate (10)

The compound **10** was obtained from cinnamoyl chloride and glycoside **4**. Yield 40%, mp 74–76 °C. UV λ_{max} (EtOH)/nm: 278. IR (KBr, $\nu_{\text{max}}/\text{cm}^{-1}$): 1750, 1640, 1500, 1370, 1210, 1190, 1030, 905. ^1H NMR (CDCl_3 , 300 MHz) δ : 2.035, 2.04, 2.07, 2.12, 2.28 (s, $5 \times 3\text{H}$, Ac), 3.84 (m, 1H, H-5'), 4.16 (dd, 1H, $J = 2.4$, 12.3 Hz, H-6'b), 4.25 (dd, 1H, $J = 5.4$, 12.3 Hz, H-6'a), 5.01 (d, 1H, $J = 7.2$ Hz, H-1'), 5.15–5.21 (m, 2H, H-4', H-3'), 5.25 (m, 1H, H-2'), 5.29 (m, 2H, CH_2 , H-7), 6.45 (d, 1H, $J = 15.9$ Hz, H-9), 6.99 (dd, 1H, $J = 2.7$, 8.7 Hz, H-3), 7.10 (s, 1H, H-5), 7.13 (m, 1H, H-2), 7.37 (m, 3H, H-13, H-14, H-15), 7.52 (m, 2H, H-12, H-16), 7.70 (d, 1H, $J = 16.2$ Hz, H-10). ^{13}C NMR (CDCl_3 , 75.5 MHz) δ : 20.4, 20.9 ($5 \times \text{CH}_3$, Ac), 60.3 (CH_2 , C6'), 61.6 (CH_2 , C-7), 68.0 (CH, C-4'), 70.7 (CH, C-2'), 71.8 (CH, C-3'), 72.4 (CH, C-5'), 99.6 (CH, C-1'), 117.0 (CH, C-2), 117.4 (CH, H-9), 121.9 ($2 \times \text{CH}$, C-3, C-5), 127.7 (C, C-6), 127.9 ($2 \times \text{CH}$, C-12, C-16), 128.7 ($2 \times \text{CH}$, C-13, C-15), 130.2 (CH, C-14), 134.0 (CH, C-11), 145.1 (C, C-4), 146.1 (CH, C-10), 151.6 (CH, C-1), 166.2 ($\text{C}=\text{O}$, C-8), 169.1, 169.3, 169.9, 170.3 ($5 \times \text{C}=\text{O}$, Ac). Anal. Calcd for $\text{C}_{32}\text{H}_{34}\text{O}_{14}$: C, 59.81; H, 5.33. Found: C, 59.76; H, 5.47.

3.5.5. 2-(2,3,4,6-Tetra-O-acetyl- β -D-glucopyranosyloxy)-5-acetoxy benzyl (4-acetoxy) benzoate (11)

The compound **11** was obtained from 4-acetoxy benzoyl chloride and glycoside **4**. Yield 52%, mp 93–94 °C. UV λ_{max} (EtOH)/nm: 232, 274. IR (KBr, $\nu_{\text{max}}/\text{cm}^{-1}$): 1750, 1640, 1600, 1500, 1370, 1210, 1190, 1030, 906. ^1H NMR (CDCl_3 , 300 MHz) δ : 2.03, 2.04, 2.07, 2.10, 2.28, 2.32 (s, $6 \times 3\text{H}$, Ac), 3.81–3.86 (m, 1H, H-5'), 4.16 (dd, 1H, $J = 2.1$, 12.3 Hz, H-6'b), 4.25 (dd, 1H, $J = 5.1$, 12.3 Hz, H-6'a), 5.02 (d, 1H, $J = 7.2$ Hz H-1'), 5.14 (m, 1H, H-4'), 5.20–5.41 (m, 2H, H-2', H-3'), 5.28 (m, 2H, H-7), 7.02 (dd, 1H, $J = 2.7$, 8.7 Hz, H-3), 7.08. ^{13}C NMR (CDCl_3 , 75.5 MHz) δ : 20.5, 21.1 ($6 \times \text{CH}_3$, Ac), 60.9 (CH_2 , C6'), 61.7 (CH_2 , C-7), 68.1 (CH, C-4'), 70.9 (CH, C-2'), 72.0 (CH, C-3'), 72.5 (CH, C-5'), 99.7 (CH, C-1'), 117.1 (CH, C-2), 121.6 ($2 \times \text{CH}$, C-11, C-13), 121.8 (CH, C-3), 122.1 (CH, C-5), 127.5 (C, C-9), 127.8 (C, C-6), 131.2 ($2 \times \text{CH}$, C-10, C-14), 146.2 (C, C-4), 151.5 (C, C-1), 154.3 (C, C-12), 165.0 ($\text{C}=\text{O}$, C-8), 168.7, 169.2, 169.5, 170.1, 170.5 ($6 \times \text{C}=\text{O}$, Ac). Anal. Calcd for $\text{C}_{32}\text{H}_{33}\text{O}_{16}$: C, 56.97; H, 5.08. Found: C, 56.81; H, 5.01.

3.5.6. 2-(2,3,4,6-Tetra-O-acetyl- β -D-glucopyranosyloxy)-5-acetoxy benzyl (3-methoxy-4-acetoxy) benzoate (12)

The compound **12** was obtained from 3-methoxy-4-acetoxy benzoyl chloride and glycoside **4**. Yield 70%, mp 156–159 °C. UV λ_{max} (EtOH)/nm: 241, 285. IR (KBr, $\nu_{\text{max}}/\text{cm}^{-1}$): 1760, 1740, 1610, 1380, 1220, 1050, 910, 760. ^1H NMR (CDCl_3 , 300 MHz) δ : 2.03, 2.05, 2.08, 2.28, 2.33 (s, $6 \times 3\text{H}$, Ac), 3.89 (m, 1H, H-5'), 3.89 (s, 3H, OCH_3), 4.16 (dd, 1H, $J = 2.1$, 12.3 Hz, H-6'b), 4.25 (dd, 1H, $J = 5.1$, 12.3 Hz, H-6'a), 5.04 (d, 1H, $J = 7.2$ Hz H-1'), 5.14 (m, 1H, H-4'), 5.24–5.42 (m, 2H, H-2', H-3'), 5.29 (m, 2H, H-7), 6.99 (dd, 1H, $J = 2.7$, 8.7 Hz, H-3), 7.10 (m, 3H, H-2, H-5, H-13), 7.68 (m, 2H, H-10, H-14). ^{13}C NMR (CDCl_3 , 75.5 MHz) δ : 20.5, 21.0 ($6 \times \text{CH}_3$, Ac), 56.1 (OCH_3), 61.0 (CH_2 , C6'), 61.8 (CH_2 , C-7), 68.1 (CH, C-4'), 70.9 (CH, C-2'), 72.0 (CH, C-3'), 72.5 (CH, C-5'), 99.7 (CH, C-1'), 113.5 (CH, C-10), 117.0 (CH, C-2), 121.8 (CH, C-3), 122.0 (CH, C-5), 122.6 (CH, C-13), 122.8 (CH, C-14), 127.8 (C, C-9), 128.5 (C, C-6), 144.7 (CH, C-12), 145.2 (C, C-4), 151.0 (C, C-11), 151.8 (C, C-1), 165.30 ($\text{C}=\text{O}$, C-8), 168.4, 169.1, 169.2, 169.5, 170.0, 170.4 ($6 \times \text{C}=\text{O}$, Ac). Anal. Calcd for $\text{C}_{33}\text{H}_{36}\text{O}_{17}$: C, 56.25; H, 5.15. Found: C, 56.12; H, 5.34.

3.5.7. 2-(2,3,4,6-Tetra-O-acetyl- β -D-glucopyranosyloxy)-5-acetoxy benzyl (3-methoxy-4-acetoxy) cinnamoate (13)

The compound **13** was obtained from 3-methoxy-4-acetoxy cinnamoyl chloride and glycoside **4**. Yield 60%, mp 157–158 °C. UV λ_{max} (EtOH)/nm: 323, 304. IR (KBr, $\nu_{\text{max}}/\text{cm}^{-1}$): 3970, 1760, 1640, 1605, 1380, 1220, 1190, 1175, 180, 1045, 980, 908, 850, 825, 600. ^1H NMR (CDCl_3 , 300 MHz) δ : 2.02, 2.03, 2.06, 2.10, 2.27, 2.30 (s, $6 \times 3\text{H}$, Ac), 3.84 (m, 1H, H-5'), 3.84 (s, 3H, OCH_3), 4.15 (dd, 1H, $J = 10.2$, 1.7 Hz, H-6'a), 4.24 (dd, 1H, $J = 5.1$, 12.3 Hz, H-6'b), 5.02 (d, 1H, $J = 7.2$ Hz, H-1'), 5.12–5.34 (m, 5H, H-2', H-3', H-4', H-7), 6.39 (d, 1H, $J = 16.2$ Hz, H-9), 6.97–7.05 (m, 2H, H-2, H-15), 7.09 (m, 4H, H-3, H-5, H-12, H-16), 7.64 (d, 1H, $J = 15.9$ Hz, H-10). ^{13}C NMR (CDCl_3 , 75.5 MHz) δ : 20.6 ($5 \times \text{CH}_3$, Ac), 21.0 (CH_3 , Ac), 55.8 (CH_3 , OCH_3), 60.5 (CH_2 , C-7), 61.8 (CH_2 , C-6'), 68.1 (CH, C-4'), 70.9 (CH, C-2'), 72.0 (CH, C-3'), 72.5 (CH, C-5'), 99.7 (CH, C-1'), 111.2 (CH, C-12), 117.1 (CH=CH, C-9), 117.7 (CH, C-2), 121.3 (CH, C-16), 122.1 ($2 \times \text{CH}$, C-3, C-5), 123.1 (CH, C-15), 127.8 (C, C-6), 133.1 (C, C-11), 141.4 (C, C-14), 144.6 (CH=CH, C-10), 146.2 (C, C-4), 151.3 (C, C-1), 151.8 (C, C-13), 166.2 ($\text{C}=\text{O}$, C-8), 168.6, 169.3, 169.5, 170.1 ($6 \times \text{C}$, Ac). HRESIMS Calcd for $[\text{M}+\text{Na}]^+$ 753.20012, Found $[\text{M}+\text{Na}]^+$ 753.20007.

3.5.8. 2-(2,3,4,6-Tetra-O-acetyl- β -D-glucopyranosyloxy)-5-benzoyloxy benzylbenzoate (14)

The compound **14** was obtained from benzoyl chloride and glycoside **5**. Yield 77%, mp 134–135 °C. UV λ_{max} (EtOH)/nm: 230, 276,

IR (KBr, $\nu_{\max}/\text{cm}^{-1}$): 2940, 1720, 1505, 1500, 1380, 1260, 1250, 1195, 1080, 1060, 1000, 915, 900, 810, 700. ^1H NMR (CDCl_3 , 300 MHz) δ : 2.04, 2.05, 2.08, 2.09 (s, $4 \times 3\text{H}$, Ac), 3.84 (m, 1H, H-5'), 4.17 (d, 1H, $J = 10.8$ Hz, H-6'a), 4.27 (dd, 1H, $J = 5.1$, 12.0 Hz, H-6'b), 5.09 (d, 1H, $J = 7.2$ Hz, H-1'), 5.16 (m, 1H, H-4'), 5.28–5.45 (m, 4H, H-2', H-3', H-7), 7.12–7.21 (m, 2H, H-3, H-5), 7.29 (m, 1H, H-2), 7.41–7.58 (m, 5H, H-11, H-12, H-13, H-18, H-20), 7.60–7.65 (m, 1H, H-19), 8.06 (d, 2H, $J = 7.5$ Hz, H-10, H-14), 8.16 (d, 2H, $J = 7.8$ Hz, H-17, H-21). ^{13}C NMR ($\text{DMSO}-d_6$, 75 MHz) δ : 20.5 ($4 \times \text{CH}_3$, Ac), 60.9 (CH_2 , C-6'), 61.8 (CH_2 , C-7), 68.2 (CH, C-4'), 70.9 (CH, C-2'), 72.1 (CH, C-3'), 72.6 (CH, C-5'), 99.9 (CH, C-1'), 117.4 (CH, C-2), 122.0 (CH, C-5), 122.3 (CH, C-3), 128.1 (C, C-6), 128.4 ($2 \times \text{CH}$, C-11, C-13), 128.6 ($2 \times \text{CH}$, C-18, C-20), 129.2 (CH, C-9), 129.7 ($2 \times \text{CH}$, C-17, C-21), 129.9 (CH, C-10), 130.1 ($2 \times \text{CH}$, C-14, C-16), 133.1 (CH, C-12), 133.7 (C, C-19), 146.6 (C, C-4), 151.9 (C, C-1), 165.2 (C=O, C-15), 166.0 (C=O, C-8), 169.3, 170.1, 170.5 ($4 \times \text{C}=\text{O}$, Ac). Anal. Calcd for $\text{C}_{35}\text{H}_{34}\text{O}_{14}$: C, 61.94; H, 5.05. Found: C, 61.75; H, 5.13.

3.6. Selective acetyl group cleavage. General method

To glycoside (**3–12**) (0.15 mmol) in a mixture of 96%-ethanol and CHCl_3 in proportion 1.5–0.5 mL was added 0.5 mL 36% HCl. The reaction mixture was stirred and then kept at 30 °C temperature for 8–13 h. After appropriate time (HPLC control), all solvents were evaporated at low temperature using vacuum and residue was subjected to column chromatography applying gradient elution with chloroform and chloroform–ethanol mixture (from ratio 15:1 to 4:1). Analytical samples were purified by recrystallization from ethanol or water.

3.6.1. 2-(β -D-Glucopyranosyloxy)-benzyl (3,4-dihydroxy) cinnamate (populoid) (15)

Crystallization from acetone gave pale yellow crystals, yield 50%, mp 178–179 °C, lit.²² 186–188 °C, lit.⁷ 169–170 °C, lit.^{6,17} 168–169 °C. UV $\lambda_{\max}$ (EtOH)/nm: 311. IR (KBr, $\nu_{\max}/\text{cm}^{-1}$): 3450, 2920, 1680, 1600, 1370, 1280, 1160, 1090, 1050, 910, 760. ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ : 3.17 (m, 1H, H-4'), 3.26–3.47 (DMSO) (m, 3H, H-2', H-3', H-5'), 4.48 (m, 1H, H-6'b), 3.68 (d, 1H, $J = 11.4$ Hz, H-6'a), 4.84 (m, 1H, H-1'), 5.22 (m, 2H, H-7), 6.32 (d, 1H, $J = 15.9$ Hz, H-9), 6.75 (d, 1H, $J = 8.1$ Hz, H-15), 7.00 (m, 3H, H-2, H-4, H-12), 7.14 (d, 1H, $J = 7.8$ Hz, H-16), 7.28 (m, 2H, H-3, H-5), 7.50 (d, 1H, $J = 15.9$ Hz, H-10). ^{13}C NMR ($\text{DMSO}-d_6$, 75.5 MHz) δ : 60.9 ($2 \times \text{CH}_2$, C6', C-7), 69.8 (CH, C-4'), 73.2 (CH, C-2'), 76.5 (CH, C-3'), 77.1 (CH, C-5'), 101.1 (CH, C-1'), 114.0 (CH, C-12), 114.9 (CH, C-2), 115.0 (CH, C-9), 115.9 (CH, C-15), 121.5 (CH, C-4), 122.0 (CH, C-16), 125.1 (CH, C-6), 125.7 (CH, C-11), 128.2 (CH, C-3), 129.0 (CH, C-5), 145.5 ($2 \times \text{C}$, C-10, C-13), 148.2 (C, C-14), 155.1 (C, C-1), 166.4 (C=O, C-8). The data are in agreement with literature.^{6,7}

3.6.2. 2-(β -D-Glucopyranosyloxy)-benzylcinnamate (16)

Crystallization from water gave colorless crystals, yield 40%, mp 110–112 °C. UV $\lambda_{\max}$ (EtOH)/nm: 278. IR (KBr, $\nu_{\max}/\text{cm}^{-1}$): 3380, 2920, 1700, 1640, 1500, 1240, 1080, 1050, 770, 750. ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ : 3.17 (m, 1H, H-4'), 3.26–3.36 (m, 2H, H-2', H-3'), 3.43–3.50 (m, 2H, H-6'b, H-5'), 3.69 (d, 1H, $J = 10.8$ Hz, H-6'a), 4.83 (m, 1H, H-1'), 5.30 (m, 2H, H-7), 6.69 (d, 1H, $J = 16.2$ Hz, H-9), 7.01 (t, 1H, $J = 7.5$ Hz, H-4), 7.16 (d, 1H, $J = 7.8$ Hz, H-2), 7.29 (d, 1H, $J = 7.2$ Hz, H-5), 7.36 (m, 1H, H-3), 7.42 (m, 3H, H-13, H-14, H-15), 7.68 (d, 1H, $J = 16.2$ Hz, H-10), 7.74 (m, 2H, H-12, H-16). ^{13}C NMR ($\text{DMSO}-d_6$, 75.5 MHz) δ : 60.5 (CH_2 , C6'), 60.9 (CH_2 , C-7), 69.5 (CH, C-4'), 73.0 (CH, C-2'), 76.3 (CH, C-3'), 76.9 (CH, C-5'), 100.9 (CH, C-1'), 114.8 (CH, C-2), 117.8 (CH, H-9), 121.7 (CH, C-4), 124.9 (C, C-6), 128.3 ($2 \times \text{CH}$, C-12, C-16), 128.8 ($3 \times \text{CH}$, C-3, C-13, C-15), 129.2 (CH, C-5), 130.5 (CH,

C-14), 134.9 (C, C-11), 144.5 (CH, C-10), 155.0 (C, C-1), 166.0 (C=O, C-8). Anal. Calcd for $\text{C}_{22}\text{H}_{24}\text{O}_8$: C, 63.45; H, 5.81. Found: C, 63.37; H, 5.95.

3.6.3. 2-(β -D-Glucopyranosyloxy)-benzyl (4-hydroxy) benzoate (17)

Crystallization from water gave colorless crystals, yield 80%, mp 165–170 °C. UV $\lambda_{\max}$ (EtOH)/nm: 259. IR (KBr, $\nu_{\max}/\text{cm}^{-1}$): 3420, 1700, 1610, 1460, 1380, 1280, 1240, 1100, 770, 750. ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ : 3.18 (m, 1H, H-4'), 3.27–3.48 (DMSO) (m, 4H, H-2', H-3', H-5', H-6'a), 3.68 (m, 1H, H-6'b), 4.86 (m, 1H, H-1'), 5.35 (m, 2H, H-7), 6.84 (d, 2H, $J = 8.4$ Hz, H-11, H-13), 7.01 (t, 1H, $J = 7.2$ Hz, H-4), 7.16 (d, 1H, $J = 8.1$ Hz, H-2), 7.28 (m, 1H, H-3), 7.35 (d, 1H, $J = 7.5$ Hz, H-5), 7.85 (d, 2H, $J = 8.1$ Hz, H-10, H-14). ^{13}C NMR ($\text{DMSO}-d_6$, 75.5 MHz) δ : 60.7 (CH_2 , C6'), 61.0 (CH_2 , C-7), 69.7 (CH, C-4'), 73.4 (CH, C-2'), 76.5 (CH, C-3'), 77.1 (CH, C-5'), 101.0 (CH, C-1'), 115.0 ($2 \times \text{CH}$, C-11, C-13), 121.7 (CH, C-2), 120.3 (C, C-9), 121.9 (CH, C-4), 125.4 (C, C-6), 128.3 (CH, C-3), 129.1 (CH, C-5), 131.6 ($2 \times \text{CH}$, C-10, C-14), 155.0 (C, C-1), 162.0 (C, C-12), 166.5 (C=O, C-8). Anal. Calcd for $\text{C}_{20}\text{H}_{22}\text{O}_9$: C, 59.11; H, 5.46. Found: C, 58.95; H, 5.54.

3.6.4. 2-(β -D-Glucopyranosyloxy)-benzyl 3-methoxy (4-hydroxy) benzoate (18)

Crystallization from water afforded colorless crystals, yield 35%, mp 173–177 °C. UV $\lambda_{\max}$ (EtOH)/nm: 265, 292. IR (KBr, $\nu_{\max}/\text{cm}^{-1}$): 3470, 3380, 2930, 1700, 1690, 1610, 1510, 1300, 1230, 1090, 760, 750. ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ : 3.16 (m, 1H, H-4'), 3.27–3.38 (DMSO) (m, 2H, H-2', H-3'), 3.43 (m, 2H, H-5', H-6'a), 3.68 (d, 1H, $J = 11.4$ Hz, H-6'b), 3.82 (s, 3H, OCH_3), 4.86 (m, 1H, H-1'), 5.37 (m, 2H, CH_2 , H-7), 6.86 (d, 1H, $J = 8.1$ Hz, H-13), 7.01 (t, 1H, $J = 7.5$ Hz, H-4), 7.16 (d, 1H, $J = 8.1$ Hz, H-2), 7.29 (m, 1H, H-3), 7.34 (d, 1H, $J = 7.5$ Hz, H-5), 7.48 (s, 1H, H-10), 7.51 (dd, 1H, $J = 1.5$, 8.4 Hz, H-14). ^{13}C NMR ($\text{DMSO}-d_6$, 75.5 MHz) δ : 55.7 (OCH_3), 60.8 (CH_2 , C6'), 61.1 (CH_2 , C-7), 69.7 (CH, C-4'), 73.3 (CH, C-2'), 76.5 (CH, C-3'), 77.2 (CH, C-5'), 101.0 (CH, C-1'), 112.5 (CH, C-10), 115.0 (CH, C-13), 115.1 (CH, C-2), 120.5 (C, C-9), 121.9 (CH, C-4), 123.5 (CH, C-14), 125.2 (CH, C-6), 129.4 (C, C-5), 130.3 (CH, C-3), 145.4 (C, C-11), 151.5 (C, C-12), 155.2 (C, C-1), 165.4 (C=O, C-8). Anal. Calcd for $\text{C}_{21}\text{H}_{24}\text{O}_{10}$: C, 57.80; H, 5.54. Found: C, 57.71; H, 5.70.

3.6.5. 2-(β -D-Glucopyranosyloxy)-5-hydroxy benzyl (3,4-dihydroxy) cinnamate (19)

Crystallization from acetone gave yellowish crystals, yield 30%, mp 156–158 °C, lit.⁶ 157–158 °C. UV $\lambda_{\max}$ (EtOH)/nm: 248, 300, 331. IR (KBr, $\nu_{\max}/\text{cm}^{-1}$): 3272, 1680, 1610, 1560, 1520, 1460, 1380, 1280, 1210, 1080, 1050, 990, 810. ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ : 3.17 (m, 1H, H-4'), 3.26–3.47 (DMSO) (m, 4H, H-2', H-3', H-5', H-6'), 3.66 (m, 1H, H-6'), 4.62 (d, 1H, $J = 6.0$ Hz, H-1'), 5.22 (m, 2H, H-7), 6.31 (d, 1H, $J = 16.2$ Hz, H-9), 6.63 (dd, 1H, $J = 9.0$, 1.5 Hz, H-3), 6.70 (d, 1H, $J = 2.7$ Hz, H-5), 6.73 (d, 1H, $J = 8.1$ Hz, H-15), 7.00–7.03 (m, 2H, H-12, H-16), 7.06 (m, 1H, H-2), 7.50 (d, 1H, $J = 15.6$ Hz, H-10). ^{13}C NMR ($\text{DMSO}-d_6$, 75.5 MHz) δ : 60.8 (CH_2 , C-7), 61.0 (CH_2 , C-6'), 69.8 (CH, C-4'), 73.4 (CH, C-2'), 76.5 (CH, C-3'), 77.1 (CH, C-5'), 102.8 (CH, C-1'), 113.8 (CH, C-12), 114.0 (CH=CH, C-9), 114.9 (CH, C-3), 115.2 (CH, C-5), 115.9 (CH, C-15), 117.7 (CH, C-2), 121.7 (CH, C-16), 126.5 (CH, C-11), 127.1 (C, C-6), 145.6 (C, C-1, CH=CH, C-10), 147.9 (CH, C-13), 148.8 (C, C-14), 152.8 (C, C-4), 166.5 (C=O, C-8). The data agree well with those given in Ref. 6.

3.6.6. 2-(β -D-Glucopyranosyloxy)-5-hydroxy benzylcinnamate (20)

Crystallization from water gave colorless crystals, yield 34%, mp 135–139 °C. UV $\lambda_{\max}$ (EtOH)/nm: 282. IR (KBr, $\nu_{\max}/\text{cm}^{-1}$): 3425,

2920, 1700, 1635, 1500, 1210, 1080, 770. ^1H NMR (DMSO- d_6 , 300 MHz) δ : 3.23 (m, 1H, H-4'), 3.30–3.48 (DMSO) (m, 4H, H-2', H-3', H-5', H-6'a), 3.68 (m, 1H, H-6'b), 4.62 (d, 1H, J = 6.3, Hz, H-1'), 5.26 (m, 2H, CH₂, H-7), 6.62–6.75 (m, 3H, H-3, H-5, H-9), 7.00 (d, 1H, J = 9.0 Hz, H-2), 7.43 (m, 3H, H-13, H-14, H-15), 7.68 (d, 1H, J = 16.2 Hz, H-10), 7.74 (m, 2H, H-12, H-16). ^{13}C NMR (DMSO- d_6 , 75.5 MHz) δ : 60.7 (CH₂, C6'), 60.9 (CH₂, C-7), 69.8 (CH, C-4'), 73.2 (CH, C-2'), 76.4 (CH, C-3'), 77.0 (CH, C-5'), 102.6 (CH, C-1'), 114.8 (CH, C-3), 115.3 (CH, C-5), 117.4 (CH, C-2), 118.0 (CH, H-9), 126.7 (C, C-6), 128.3 (2 $\times$ CH, C-12, C-16), 128.9 (2 $\times$ CH, C-13, C-15), 130.5 (CH, C-14), 134.0 (CH, C-11), 144.7 (C, C-1), 147.9 (CH, C-10), 152.1 (CH, C-4), 166.0 (C=O, C-8). Anal. Calcd for C₂₂H₂₄O₉: C, 61.11; H, 5.59. Found: C, 61.00; H, 5.65.

3.6.7. 2-(β -D-Glucopyranosyloxy)-5-hydroxy benzyl (4-hydroxy) benzoate (21)

Crystallization from water gave colorless crystals, yield 55 %, mp 140 °C. UV λ_{max} (EtOH)/nm: 257. IR (KBr, ν_{max} /cm⁻¹): 3360, 2930, 1680, 1610, 1495, 1280, 1210, 1070, 770. ^1H NMR (DMSO- d_6 , 300 MHz) δ : 3.21 (m, 1H, H-4'), 3.30–3.44 (DMSO) (m, 4H, H-2', H-3', H-5', H-6'a), 3.67 (m, 1H, H-6'b), 4.63 (m, 1H, H-1'), 5.32 (m, 2H, CH₂, H-7), 6.64 (dd, 1H, J = 2.7, 9.3 Hz, H-3), 6.75 (d, 1H, J = 2.4, Hz, H-5), 6.85 (d, 2H, J = 8.4, Hz, H-11, H-13), 7.00 (d, 1H, J = 8.7 Hz, H-2), 7.85 (d, 2H, J = 8.4 Hz, H-10, H-14). ^{13}C NMR (CDCl₃, 75.5 MHz) δ : 61.8 (CH₂, C6'), 61.9 (CH₂, C-7), 69.8 (CH, C-4'), 74.2 (CH, C-2'), 76.5 (CH, C-3'), 77.0 (CH, C-5'), 102.6 (CH, C-1'), 114.2 (CH, C-5), 115.0 (CH, C-3), 115.5 (2 $\times$ CH, C-11, C-13), 117.8 (CH, C-2), 120.2 (C, C-9), 127.0 (C, C-6), 131.4 (2 $\times$ CH, C-10, C-14), 147.7 (C, C-1), 152.2 (C, C-4), 163.0 (C, C-12), 165.3 (C=O, C-8). Anal. Calcd for C₂₀H₂₂O₁₀: C, 56.87; H, 5.25. Found: C, 56.80; H, 5.29.

3.6.8. 2-(β -D-Glucopyranosyloxy)-5-hydroxy benzyl (3-methoxy-4-hydroxy) benzoate (22)

Crystallization from water gave colorless crystals, yield 36 %, mp 128–136 °C. UV λ_{max} (EtOH)/nm: 264, 292. IR (KBr, ν_{max} /cm⁻¹): 2970, 1695, 1520, 1290, 1215, 1080, 1240, 990, 760. ^1H NMR (DMSO- d_6 , 300 MHz) δ : 3.17 (m, 1H, H-4'), 3.22–3.44 (m, 4H, H-2', H-3', H-5', H-6'a), 3.67 (d, 1H, J = 11.4 Hz, H-6'b), 3.82 (s, 3H, OCH₃), 4.63 (m, 1H, H-1'), 5.33 (m, 2H, CH₂, H-7), 6.63 (dd, 1H, J = 2.7, 8.7 Hz, H-3), 6.74 (d, 1H, J = 3.0 Hz, H-5), 6.87 (d, 1H, J = 8.1 Hz, H-13), 7.00 (d, 1H, J = 8.7 Hz, H-2), 7.49 (s, 1H, H-10), 7.54 (dd, 1H, J = 3.0, 8.4 Hz, H-14). ^{13}C NMR (DMSO- d_6 , 75.5 MHz) δ : 55.8 (OCH₃), 60.9 (CH₂, C6'), 61.0 (CH₂, C-7), 69.8 (CH, C-4'), 73.4 (CH, C-2'), 76.6 (CH, C-3'), 77.1 (CH, C-5'), 102.6 (CH, C-1'), 112.7 (CH, C-10), 114.2 (CH, C-13), 115.0 (CH, C-5), 115.3 (CH, C-3), 117.7 (CH, C-2), 120.5 (C, C-9), 123.6 (CH, C-14), 127.1 (C, C-6), 147.3 (CH, C-11), 147.8 (C, C-1), 151.7 (C, C-12), 152.4 (C, C-4), 165.5 (C=O, C-8). Anal. Calcd for C₂₁H₂₄O₁₁: C, 55.75; H, 5.35. Found: C, 55.71; H, 5.37.

3.6.9. 2-(β -D-Glucopyranosyloxy)-5-hydroxy benzyl (3-methoxy-4-hydroxy) cinnamate (23)

Crystallization from water gave colorless crystals, yield 48%, mp 99–100 °C. UV λ_{max} (EtOH)/nm: 296, 325. IR (KBr, ν_{max} /cm⁻¹): 3405, 2930, 1690, 1600, 1500, 1270, 1210, 1175, 1075, 1035, 810. ^1H NMR (MeOD, 300 MHz) δ : 3.82 (m, 1H, H-5'), 4.15 (dd, 1H, J = 9.9, 1.7 Hz, H-6'a), 4.24 (dd, 1H, J = 5.3, 12.0 Hz, H-6'b), 5.02 (m, 1H, H-1'), 5.12 (m, 1H, H-4'), 5.17 (m, 2H, H-7), 5.27 (m, 2H, H-2', H-3'), 6.39 (d, 1H, J = 15.9 Hz, H-9), 6.97 (dd, 1H, J = 2.5, 8.7 Hz, H-3), 7.09 (d, 1H, J = 7.2 Hz, H-2), 7.10 (s, 1H, H-5), 7.19 (d, 1H, J = 8.4 Hz, H-15), 7.36 (s, 1H, H-12), 7.38 (d, 1H, J = 8.4 Hz, H-16), 7.62 (d, 1H, J = 16.2 Hz, H-10). ^{13}C NMR (MeOD, 75.5 MHz) δ : 60.6 (CH₂, C-7), 61.8 (CH₂, C-6'), 68.1 (CH, C-4'), 70.9 (CH, C-2'), 72.0 (CH, C-3'), 72.5 (CH, C-5'), 99.7 (CH, C-1'), 117.1 (CH=CH,

C-9), 118.7 (CH, C-2), 122.1 (2 $\times$ CH, C-3, C-5), 122.7 (CH, C-15), 123.9 (CH, C-12), 126.5 (CH, C-16), 127.8 (C, C-6), 133.1 (C, C-11), 142.4 (C, C-13), 143.4 (C, C-14), 145.5 (CH=CH, C-10), 146.3 (C, C-1), 151.8 (C, C-4), 166.0 (C=O, C-8). Anal. Calcd for C₂₃H₂₆O₁₁: C, 57.74; H, 5.48. Found: C, 57.67; H, 5.53.

3.6.10. 2-(β -D-Glucopyranosyloxy)-5-benzoyloxy benzylbenzoate (24)

Crystallization from water gave colorless crystals, yield 60%, mp 190–194 °C. UV λ_{max} (EtOH)/nm: 230, 276. IR (KBr, ν_{max} /cm⁻¹): 2950, 1730, 1715, 1500, 1390, 1255, 1200, 1080, 1060, 1040, 890, 795 703. ^1H NMR (DMSO, 300 MHz) δ : 3.17 (m, 1H, H-4'), 3.30–3.38 (DMSO) (m, 3H, H-2', H-3', H-5'), 3.45–3.52 (m, 1H, H-6'a), 3.69–3.75 (m, 1H, H-6'b), 4.91 (d, 1H, J = 6.6, Hz, H-1'), 5.39 (dd, 2H, J = 13.5, 3.6, Hz, H-7), 7.27 (m, 2H, H-3, H-5), 7.33 (m, 1H, H-2), 7.50–7.68 (m, 5H, H-11, H-12, H-13, H-18, H-20), 7.72–7.77 (m, 1H, H-19), 8.01 (d, 2H, J = 7.2 Hz, H-10, H-14), 8.11 (d, 2H, J = 6.9, Hz, H-17, H-21). ^{13}C NMR (DMSO, 75.5 MHz) δ : 60.9 (CH₂, C-6'), 61.4 (CH₂, C-7), 69.8 (CH, C-4'), 73.1 (CH, C-2'), 76.5 (CH, C-3'), 76.9 (CH, C-5'), 100.1 (CH, C-1'), 117.4 (CH, C-2), 121.9 (CH, C-5), 122.8 (CH, C-3), 126.1 (C, C, C-6), 129.0 (5 $\times$ CH, C-9, C-11, C-13, C-18, C-20), 129.2 (2 $\times$ CH, C-17, C-21), 129.5 (CH, C-10), 129.8 (2 $\times$ CH, C-14, C-16), 133.2 (CH, C-12), 134.0 (C, C-19), 145.0 (C, C-4), 152.9 (C, C-1), 164.9 (C=O, C-15), 165.8 (C=O, C-8). Anal. Calcd for C₂₇H₂₆O₁₀: C, 63.53; H, 5.13. Found: C, 63.60; H, 5.25.

3.7. 4-(2,3,4,6-Tetra-O-acetyl- β -D-glucopyranosyloxy)-phenylbenzoate (26)

The compound **26** was obtained by glycosylation of (**25**), yield, mp 150–151 °C, lit.²⁵ 136–138 °C, lit.³⁰ 154–155 °C. UV λ_{max} (EtOH)/nm: 283. IR (KBr, ν_{max} /cm⁻¹): ν /cm⁻¹: 1740, 1505, 1370 1230, 1200, 1065, 700. ^1H NMR (CDCl₃, 300 MHz) δ : 2.04, 2.05, 2.08 (s, 4 $\times$ 3H, Ac), 3.85 (m, 1H, H-5'), 4.15 (d, 1H, J = 12.3 Hz, H-6'b), 4.27 (dd, 1H, J = 5.4, 12.3 Hz, H-6'a), 5.05 (d, 1H, J = 6.9 Hz, H-1'), 5.15 (m, 1H, H-4'), 5.25–5.34 (m, 2H, H-2', H-3'), 7.03 (d, 2H, J = 9.0 Hz, H-2, H-6), 7.13 (d, 2H, J = 9.0 Hz, H-3, H-5), 7.48 (m, 2H, H-18, H-20), 7.61 (t, 1H, J = 7.2 Hz, H-19), 8.17 (d, 2H, J = 7.5 Hz, H-17, H-21). ^{13}C NMR (CDCl₃, 75.5 MHz) δ : 20.7 (4 $\times$ CH₃, COCH₃), 62.1 (CH₂, C6'), 68.4 (CH, C-4'), 71.3 (CH, C-2'), 72.2 (CH, C-5'), 72.8 (CH, C-3'), 99.7 (CH, C-1'), 118.2 (2 $\times$ CH, C-2, C-6), 122.8 (2 $\times$ CH, C-5, C-3), 128.7 (2 $\times$ CH, C-18, C-20), 129.5 (C, C-16), 130.3 (2 $\times$ CH, C-17, C-21), 133.8 (CH, C-19), 146.6 (C, C-4), 154.7 (C, C-1), 165.0 (C=O, C-15), 169.5, 170.3, 170.7 (4 $\times$ C, Ac). The data are in agreement with literature.²⁵

3.8. 4-(β -D-Glucopyranosyloxy)-phenylbenzoate (27)

The compound **27** was obtained according to procedure for **14–23** from glycoside **25** and was crystallized from water. Yield 36%, mp 189–196 °C. UV λ_{max} (EtOH)/nm: 230, 269. IR (KBr, ν_{max} /cm⁻¹): ν /cm⁻¹: 3540, 3390, 2890, 1712, 1600, 1510, 1455, 1400, 1275, 1240, 1200, 1080, 1040, 820, 710. ^1H NMR (DMSO, 300 MHz) δ : 3.17–3.21 (m, 1H, H-4'), 3.25–3.35 (m, 3H, H-2', H-3, H-5'), 3.44–3.53 (1H, m, H-6'a), 3.69 (dd, 1H, J = 4.5, 11.1, Hz, H-6'b), 4.87 (d, 1H, J = 6.9, Hz, H-1'), 7.09 (d, 2H, J = 8.7 Hz, H-2, H-6), 7.19 (d, 2H, J = 8.7 Hz, H-3, H-5), 7.58 (m, 2H, H-10, H-12), 7.72 (t, 1H, J = 7.2 Hz, H-11), 8.11 (d, 2H, J = 7.8 Hz, H-9, H-13). ^{13}C NMR (DMSO, 75.5 MHz) δ : 60.6 (CH₂, C6'), 69.6 (CH, C-4'), 73.2 (CH, C-2'), 76.5 (CH, C-5'), 77.0 (CH, C-3'), 100.6 (CH, C-1'), 116.9 (2 $\times$ CH, C-2, C-6), 122.6 (2 $\times$ CH, C-5, C-3), 129.0 (2 $\times$ CH, C-16, C-18, C-20), 129.7 (2 $\times$ CH, C-17, C-21), 134.0 (CH, C-19), 144.8 (C, C-4), 155.1 (C, C-1), 165.0 (C=O, C-8), 164.8. Anal. Calcd for C₁₉H₂₀O₈: C, 60.63; H, 5.36. Found: C, 60.50; H, 5.52.

Acknowledgements

This work was supported by the scientific program ‘Nauka’ No. 2387. We would like to thank the Department of Chemical Analysis of Tomsk State University for assistance in obtaining elemental analysis data.

Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.carres.2014.02.014>.

References

- Hedner, T.; Everts, B. *Clin. Rheumatol.* **1998**, *17*, 17–25.
- Boeckler, G. A.; Gershenzon, J.; Unsicker, S. B. *Phytochemistry* **2011**, *72*, 1497–1509.
- Chen, P. D. *Chin. Tradit. Herb. Drugs* **2006**, *37*, 1607–1608.
- Choudhary, M. I.; Fatima, N.; Abbasi, M. A.; Jalil, S.; Ahmad, V. U.; Atta-ur-Rahman *Bioorg. Med. Chem.* **2004**, *12*, 5793–5798.
- Lee, K. H.; Yang, M. C.; Kim, K. H.; Kwon, H. C.; Choi, S. U.; Lee, K. R. *Molecules* **2008**, *13*, 41–45.
- Si, Ch. L.; Li, Sh. M.; Liu, Z.; Kim, J. K.; Bae, Y. S. *Nat. Prod. Res.* **2011**, *25*, 1396–1401.
- Zhang, X. F.; Thuong, P. T.; Min, B. S. *J. Nat. Prod.* **2006**, *69*, 1370.
- Poblocka-Olech, L.; Krauze-Baranowska, M.; Glod, D.; Kawiak, A.; Lojkowska, E. *Phytochem. Anal.* **2010**, *21*, 463–469.
- Lim, E.; Doucet, C. J.; Li, Y.; Elias, L.; Worrall, D.; Spencer, S. P., et al *J. Biol. Chem.* **2002**, *277*, 586–592.
- Gülçin, I. *Toxicology* **2006**, *217*, 213–220.
- Pérez-Alvarez1, V.; Fernández-Martínez, E.; Morales-Ríos, M. S.; Bobadilla, R. A.; Muriel, P. *Proc. West. Pharmacol. Soc.* **2003**, *46*, 136–138.
- Lodovici, M.; Guglielmi, F.; Meoni, M.; Dolara, P. *Food Chem. Toxicol.* **2001**, *39*, 1205–1210.
- Ou, Sh.; Kwok, K. Ch. *J. Sci. Food Agric.* **2004**, *84*, 1261–1269.
- Chong, K. P.; Rossall, S.; Atong, M. *J. Agric. Sci.* **2009**, *1*, 15–20.
- Khadem, Sh.; Marles, R. J. *Molecules* **2010**, *15*, 7985–8005.
- Itoh, A.; Isoda, K.; Kondoh, M.; Kawase, M.; Watari, A.; Kobayashi, M., et al *Biol. Pharm. Bull.* **2010**, *33*, 983–987.
- Kumar, S.; Prahalathan, P.; Raja, B. *Redox Rep.* **2011**, *16*, 208–215.
- Prince, P. S. M.; Rajakumar, S.; Dhanasekar, K. *Eur. J. Pharmacol.* **2011**, *668*, 233–240.
- Belyanin, M. L.; Stepanova, E. V.; Ogorodnikov, V. D. *Carbohydr. Res.* **2012**, *363*, 66–72.
- Briggs, J. C.; Haines, A. H.; Taylor, R. J. *J. Chem. Soc., Chem. Commun.* **1992**, 1039–1041.
- Zemplén, G.; Bogner, R.; Pongor, G. *Acta Chim. Hung. Tomus.* **1959**, *19*, 285–293.
- Erickson, R. L.; Pearl, I. A.; Darling, S. F. *Phytochemistry* **1970**, *9*, 857–863.
- Lee, Y. S.; Cui, Ch. B.; Kim, J. K.; Bae, Y. S.; Lee, J. Y.; Kang, I. J., et al *Korean Soc Appl. Biol. Chem.* **2010**, *53*, 729–733.
- Choudhary, M. I.; Fatima, N.; Abbasi, M. A.; Jalil, S.; Ahmad, V. U.; Rahman, A. *Bioorg. Med. Chem.* **2004**, *12*, 5793–5798.
- Lee, Y. S. U.S. Patent 6,388,103 B2, 2002.
- Seo, D. H.; Jung, J. H.; Lee, J. E.; Jeon, E. J.; Kim, W.; Park, C. S. *Appl. Microbiol. Biotechnol.* **2012**, *95*, 1417–1425.
- Varma, M.; Varma, R. S.; Parthasarathy, M. R. *Monatsh. Chem.* **1980**, *111*, 469–47.
- Perold, G. W.; Rosenberg, M. E. K.; Howard, A. S.; Huddle, P. A. *J. Chem. Soc., Perkin Trans. 1* **1979**, 239–243.
- Yang, R. L.; Li, N.; Ye, M.; Zong, M. H. *J. Mol. Catal. B Enzym.* **2010**, *67*, 41–44.
- Robertson, A.; Waters, R. B. *J. Chem. Soc.* **1930**, 2729–2733.