



Differential effects of UTP and ATP on ion transport in porcine tracheal epithelium

*¹S.K. Inglis, ¹R.E. Olver & ¹S.M. Wilson

¹Lung Membrane Transport Group, Tayside Institute of Child Health, Ninewells Hospital and Medical School, University of Dundee, Dundee DD1 9SY

1 Isolated segments of porcine tracheal epithelium were mounted in Ussing chambers, current required to maintain transepithelial potential difference at 0 mV (short circuit current, I_{SC}) was monitored and effects of nucleotides upon I_{SC} were studied.

2 Mucosal UTP (100 μ M) evoked a transient rise in I_{SC} that was followed by a sustained fall below basal I_{SC} maintained for 30 min. Mucosal ATP (100 μ M) also stimulated a transient rise in I_{SC} but in contrast to UTP did not inhibit basal I_{SC} . Submucosal UTP and ATP both transiently increased I_{SC} .

3 UTP-prestimulated epithelia were refractory to ATP but prestimulation with ATP did not abolish the response to UTP. The epithelia thus appear to express two populations of apical receptors allowing nucleotides to modulate I_{SC} .

4 The UTP-induced rise was reduced by pretreatment with either bumetanide (100 μ M), diphenylamin-2-carboxylic acid (DPC, 1 mM), or Cl^- and HCO_3^- -free solution whilst the fall was abolished by amiloride pretreatment.

5 Thapsigargin (0.3 μ M) abolished the UTP-induced increase in I_{SC} but not the subsequent decrease. Staurosporine (0.1 μ M) inhibited basal I_{SC} and blocked UTP-induced inhibition of I_{SC} . Inhibitors of either protein kinase C (PKC) (D-erythro sphingosine) or PKA (H89) had no effect.

6 This study suggests that UTP stimulates Cl^- secretion and inhibits basal Na^+ absorption. ATP has a similar stimulatory effect, which may be mediated by activation of $P2Y_2$ receptors and an increase in $[Ca^{2+}]_{in}$, but no inhibitory effect, which is likely mediated by activation of a pyrimidine receptor and possible inhibition of a protein kinase other than PKC or PKA.

British Journal of Pharmacology (2000) **130**, 367–374

Keywords: Airway; ATP; cystic fibrosis; epithelium; ion transport; $P2Y$ receptor; pyrimidine receptor; UTP

Abbreviations: CF, cystic fibrosis; CFTR, cystic fibrosis transmembrane conductance regulator; DPC, diphenylamin-2-carboxylic acid; I_{SC} , short-circuit current; PD, potential difference; PKA, protein kinase A; PKC, protein kinase C; PMA, phorbol-12-myristate-13-acetate; R_t , transepithelial resistance

Introduction

CF is caused by mutations in the cystic fibrosis transmembrane conductance regulator (CFTR), a protein that normally forms a cyclic AMP-dependent anion channel (Anderson *et al.*, 1991) and exerts inhibitory control over epithelial Na^+ channels (Stutts *et al.*, 1995; Ismailov *et al.*, 1996). As a consequence, the airway epithelia of patients with CF are characterized by reduced anion secretion and enhanced Na^+ absorption (Boucher *et al.*, 1986). These defects disturb the volume and/or ionic composition of the liquid lining the epithelial surface and, although the mechanism has not been determined (see Boucher, 1999), this precipitates chronic chest disease that can progressively destroy the lungs.

Luminal nucleotides stimulate anion secretion across airway epithelia by acting on $P2Y$ receptors present in the apical membrane. The response is due to the activation of anion channels other than CFTR, and so it persists in CF cells (Mason *et al.*, 1991; Stutts *et al.*, 1992; 1994). Moreover, it has recently become clear that these receptors also mediate sustained inhibition of Na^+ absorption (Iwase *et al.*, 1997; Devor & Pilewski, 1999; Inglis *et al.*, 1999; Ramminger *et al.*, 1999). $P2Y$ receptor agonists can thus oppose the defects in ion transport that cause chest disease in CF patients and this has led to the suggestion that such drugs may become useful in the

treatment of this disease (Knowles *et al.*, 1996). Such responses to apical nucleotides are often attributed to the activation of receptors belonging to the $P2Y_2$ subclass (Parr *et al.*, 1994; Knowles *et al.*, 1996), which are equally sensitive to ATP and UTP. However, other $P2Y$ receptor subtypes are present in the apical membranes of at least some cultured epithelia. These receptors are activated by pyrimidine nucleotides but essentially insensitive to ATP (Inoue *et al.*, 1997; Lazarowski *et al.*, 1997; Wilson *et al.*, 1998; Ko *et al.*, 1999), and could become important therapeutic targets if they were also expressed by native tissues. In the present study, we have therefore sought to establish whether such selective 'pyrimidinoceptors' are present in the apical membranes of acutely isolated airway epithelia.

Methods

Measurement of transepithelial ion transport

Cotswold pigs (~15 kg) were obtained from a local supplier. The animals were sedated with inhaled Halothane and then killed by an intravenous overdose of sodium pentobarbitone. The trachea was removed and divided into three approximately equal segments, each of which was opened longitudinally and pinned onto a Sylgard support with the mucosa

*Author for correspondence; E-mail: s.k.inglis@dundee.ac.uk

uppermost. Epithelial sheets were then carefully dissected from each segment and mounted in Ussing chambers (area of exposed tissue 1.77 cm^2) where they were bathed with Krebs–Ringer solution (37°C) that was continually circulated by gas lifts (5% CO_2 in O_2). All preparations were first maintained under open circuit conditions until the transepithelial potential difference (PD) had stabilized (30–90 min). The transepithelial PD was then clamped at 0 mV, and the current required to maintain this PD (short circuit current, I_{SC}) continually monitored and displayed using standard techniques. Data were recorded directly to disc using a MacLab computer interface (ADInstruments Ltd., Hastings, U.K.).

Data analysis

Values of PD are reported with respect to the submucosal compartment. Positive currents are defined as those that would be carried by anions moving from the basolateral to the apical compartments and are shown as upward deflections of the traces. Experimentally-induced changes in I_{SC} were quantified by subtracting the current flowing at the peak of a response from basal I_{SC} , where basal I_{SC} is the current flowing immediately prior to addition of agonist. In some instances, responses were further characterized by integrating the current records with respect to time using commercially available software (AreaCalc 1.0, Cyber Solutions, Newport-on Tay, Fife, U.K.). The basal I_{SC} was assigned a value of zero so that all responses reflect deviations from the charge transfer occurring under unstimulated conditions. The dose response relationship was fitted using Grafit (Erithacus Software, Staines, U.K.).

Data are presented as mean $\pm$ standard error (s.e.mean) and values of n refer to the number of experiments undertaken using tissues from different animals. The statistical significance of any difference between mean values was assessed using either Student's paired t -test or analysis of variance (ANOVA) followed by Dunnett's test.

Solutions and chemicals

Krebs–Ringer solution (in mM): NaCl 112; KCl 4.7; CaCl_2 2.5; MgSO_4 2.4; KH_2PO_4 1.2; NaHCO_3 25; D-glucose 11.6. Solution was continuously bubbled with 5% CO_2 in O_2 to maintain pH 7.4. Cl^- and HCO_3^- -free solution was similar to KRB but NaHCO_3^- , NaCl and KCl were replaced with equimolar *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulphonic acid (HEPES), Na gluconate and K gluconate respectively. CaCl_2 was replaced with 11 mM Ca gluconate₂ to counteract the chelation effect of gluconate (Christoffersen & Skibsted, 1975). Solution was titrated to pH 7.4 with HCl and gassed with 100% O_2 . All drugs and nucleotides were obtained from either Sigma Chemical Co. (Poole, Dorset, U.K.) or Calbiochem (Nottingham, U.K.). Stock solutions of nucleotides and amiloride were prepared in distilled water. Stock solutions of bumetanide, diphenylamine-2-carboxylic acid (DPC), KN-93, staurosporine, phorbol-12-myristate-13-acetate (PMA), D-erythrosphingosine, H89 and forskolin were in dimethylsulphoxide whilst thapsigargin was in ethanol. Appropriate experiments showed that the solvent vehicles did not affect the electrical properties of the experimental preparations.

Results

Under unstimulated conditions the bioelectrical properties of the tracheal segments used in this study were: transepithelial

resistance (R_t) $168 \pm 6 \Omega \text{ cm}^2$; PD $-23.3 \pm 1.1 \text{ mV}$; and I_{SC} $132 \pm 3 \mu\text{A cm}^{-2}$ ($n = 190$).

The effects of mucosal UTP and ATP

Both ATP and UTP (both $100 \mu\text{M}$ mucosal) evoked transient increases in I_{SC} that reached clearly defined peaks of a similar magnitude after ~ 1 min (Figure 1A, Table 1). Thereafter, the responses to ATP and UTP differed markedly. In ATP-stimulated tissues I_{SC} decayed back towards its basal level until, after 10 min, it did not differ significantly from basal I_{SC} (basal $I_{\text{SC}} = 121 \pm 5 \mu\text{A cm}^{-2}$, 10 min after ATP addition $I_{\text{SC}} = 120 \pm 6 \mu\text{A cm}^{-2}$, $P > 0.05$, Figure 1A). In contrast, I_{SC} fell much more sharply in response to UTP. Indeed, I_{SC} remained above the basal level for only 3 min and, after 10 min, had fallen to a level significantly below the basal I_{SC} (basal: $134 \pm 8 \mu\text{A cm}^{-2}$, after 10 min with UTP: $108 \pm 6 \mu\text{A cm}^{-2}$, $P < 0.05$, Figure 1A). Although some recovery occurred I_{SC} remained depressed for at least 30 min. For the purpose of clarity, we termed the period during which I_{SC} is elevated the stimulatory phase and the period during which I_{SC} is depressed the inhibitory phase of the response. To further quantify the difference in the responses to ATP and UTP, the transepithelial charge movement occurring during the two phases of the response was quantified. During the stimulatory phase, ATP-evoked charge transfer exceeded that seen during stimulation with UTP, while the reverse was true during the inhibitory phase (Figure 1B and Table 1). Indeed, over a 16 min period, ATP caused a net stimulation of charge transfer whilst the overall action of UTP was inhibitory (Figure 1B and Table 1).

To establish the extent to which the responses were concentration dependent, epithelial segments were stimulated with one of six concentrations of UTP (0.1 – $500 \mu\text{M}$, all mucosal). Each response was compared with that evoked by $100 \mu\text{M}$ UTP in tissues isolated from the same animal. Analysis showed that both the peak and inhibitory phases of the response to UTP were dose dependent (Figure 2) and that the stimulatory phase had a slightly lower EC_{50} ($\sim 5 \mu\text{M}$) than the inhibitory phase ($\sim 20 \mu\text{M}$). Analogous experiments ($n = 3$) confirmed that $100 \mu\text{M}$ ATP elicits a monophasic response but also showed that this nucleotide did not evoke a discernible change in I_{SC} at $1 \mu\text{M}$ (Basal $I_{\text{SC}} = 97.4 \pm 12.4 \mu\text{A cm}^{-2}$, after $1 \mu\text{M}$ ATP $= 98.6 \pm 11.7 \mu\text{A cm}^{-2}$). At $10 \mu\text{M}$, however, ATP evoked a clear increase in I_{SC} ($14.6 \pm 3 \mu\text{A cm}^{-2}$) that was significantly smaller than that seen during stimulation with $100 \mu\text{M}$ ATP ($32.9 \pm 8.3 \mu\text{A cm}^{-2}$, $P < 0.05$).

Cross desensitization experiments

To test whether mucosal ATP and UTP act upon a common receptor population, epithelia were treated with UTP ($100 \mu\text{M}$ mucosal) and a second aliquot of this nucleotide (also $100 \mu\text{M}$) added once I_{SC} had stabilized. The initial stimulus elicited a response essentially identical to that described above, whilst the second simply evoked a small decrease in I_{SC} (Figure 3A, Table 2). The response to UTP is thus subject to essentially complete autologous desensitization and analogous experiments showed that this was also true of the response to ATP (Table 2).

Cross desensitization experiments showed that ATP-stimulated tissues subsequently responded to UTP (Figure 3B, Table 2). Although it was smaller than the control response, this response to UTP was clearly biphasic. Moreover, both of its phases were significantly greater than those induced by UTP in tissues pretreated with UTP ($P < 0.05$,




Figure 1 Effect of ATP and UTP on I_{sc} across tracheal epithelium. Paired tissues were treated with either UTP (100 μ M mucosal) or ATP (100 μ M mucosal). (A) Shows mean $I_{sc} \pm$ s.e. mean of 11 pairs. (B) Shows charge transfer, calculated as area between UTP-induced I_{sc} and basal I_{sc} (dotted line in A). Positive charge transfer represents increased mucosal negativity. *Indicates significant difference from effect of UTP (Paired Student's t -test, $P < 0.05$).

Table 1 Contrasting effects of mucosal ATP and UTP

	Change in I_{sc} (μ A cm ⁻²)			Time above basal I_{sc} (min)	Charge transfer (mC cm ⁻²)		
	Basal I_{sc} (μ A cm ⁻²)	Stimulatory phase	Inhibitory phase		Stimulatory phase	Inhibitory phase	Net
UTP	134 $\pm$ 8	30.6 $\pm$ 3.3	-26.8 $\pm$ 2.6	3.0 $\pm$ 0.3	3.3 $\pm$ 0.5	-16.8 $\pm$ 1.8	-13.4 $\pm$ 2.5
ATP	121 $\pm$ 5	37.8 $\pm$ 4.3	-0.5 $\pm$ 2.2*	9.2 $\pm$ 1.1*	8.1 $\pm$ 2.0*	-1.9 $\pm$ 0.6*	6.2 $\pm$ 2.4*

Table compares effects of UTP and ATP (both 100 μ M mucosal) on I_{sc} and charge transfer measured across paired tracheal segments. I_{sc} before addition of first nucleotide (Basal I_{sc}), peak increase above basal (Stimulatory phase), change in I_{sc} after 10 min (Inhibitory phase); negative value represents fall below basal I_{sc} , and time above basal I_{sc} are shown. Also charge transfer during stimulatory and inhibitory phases, and net charge transferred in 16 min are shown. Data are mean $\pm$ s.e. mean *Indicates significant difference from response to UTP (Student's paired t -test, $P < 0.05$, $n = 11$).

$n=8$) (Table 2). Further experiments revealed that the response of UTP-pretreated tissues to ATP was not significantly different from the response of tissues pretreated with ATP (Table 2). These results suggest that the apical membrane contains at least two P2Y receptor subtypes, one desensitized by both ATP and UTP and one desensitized by UTP but not by ATP.

The effects of submucosal UTP and ATP on ion transport

To test whether the biphasic effect of UTP was mediated by receptors present on the basolateral as well as the apical membrane, UTP (500 μM) was applied to the submucosal side of the epithelia. This high dose was chosen since we anticipated that the basolateral receptors may be less accessible than the apical ones. Submucosal UTP evoked a transient increase in I_{SC} ($8.1 \pm 1.5 \mu\text{A cm}^{-2}$) that was significantly smaller than the response to mucosal UTP seen in paired tissues (18.4 ± 2.2 , $n=5$, $P<0.05$). Submucosal UTP caused no inhibition of basal I_{SC} (Figure 4). Net charge transfer was positive ($5.4 \pm 1.1 \text{ mC cm}^{-2}$), and significantly different from charge transfer induced by mucosal UTP ($-14.1 \pm 4.4 \text{ mC cm}^{-2}$, $P<0.05$). Addition of mucosal UTP after submucosal UTP

induced a response (peak increase $29.8 \pm 5.6 \mu\text{A cm}^{-2}$, inhibition after 10 min $-24.4 \pm 1.5 \mu\text{A cm}^{-2}$, net charge transfer $-10.1 \pm 0.3 \text{ mC cm}^{-2}$) indistinguishable from that induced by mucosal UTP alone (peak increase $18.4 \pm 2.2 \mu\text{A cm}^{-2}$, inhibition after 10 min $-28.2 \pm 6.4 \mu\text{A cm}^{-2}$, net charge transfer $-14.1 \pm 4.4 \text{ mC cm}^{-2}$) (Figure 4). In contrast to the difference in the responses induced by mucosal UTP and ATP, the response to submucosal ATP (peak increase $11.2 \pm 2.4 \mu\text{A cm}^{-2}$, I_{SC} after 10 min $8.8 \pm 3.4 \mu\text{A cm}^{-2}$ above basal, net charge transfer $7.2 \pm 2.1 \text{ mC cm}^{-2}$) was indistinguishable from the response to submucosal UTP (Figure 4).

Ionic basis of the UTP-evoked changes in I_{SC}

Treatment with bumetanide (100 μM submucosal) to inhibit Cl^- secretion, blocked $\sim 33\%$ of basal I_{SC} (Table 3). Addition of UTP to these trachea induced an initial increase that was significantly smaller than the increase in paired control tissues ($P<0.05$, Figure 5, Table 4). The subsequent inhibition of I_{SC}




Figure 2 Dose dependency of the increase (Peak) and decrease below basal after 10 min (after 10 min) induced by UTP. Trachea were treated with either 10^{-7} , 10^{-6} , 10^{-5} , 5×10^{-5} , 10^{-4} or $5 \times 10^{-4} \text{ M}$ mucosal UTP. Responses are represented as a fraction of the response of paired tissues to $100 \mu\text{M}$ UTP. Data are mean $\pm$ s.e. mean ($n=6$ (10^{-6} , 10^{-5}), 7 (10^{-7} , 5×10^{-5}) or 3 ($5 \times 10^{-4} \text{ M}$)).




Figure 3 Cross desensitization of ATP and UTP. Paired tissues were pretreated with either UTP (100 μM mucosal; (A), or ATP (100 μM mucosal; (B), before addition of UTP (100 μM mucosal). Data are mean $\pm$ s.e. mean ($n=5$).

Table 2 Cross-desensitization of UTP and ATP

Order of Application	n	Basal I_{SC} ($\mu\text{A cm}^{-2}$)	First nucleotide ΔI_{SC} ($\mu\text{A cm}^{-2}$)		Second nucleotide ΔI_{SC} ($\mu\text{A cm}^{-2}$)	
			Stimulatory phase	Inhibitory phase	Stimulatory phase	Inhibitory phase
UTP-UTP	8	126 ± 12	32 ± 2	-25 ± 4	-5 ± 1	-5 ± 1
ATP-UTP	8	124 ± 7	39 ± 6	0 ± 3	$9 \pm 2^*$	$-10 \pm 1^*$
ATP-ATP	3	114 ± 1	34 ± 6	-1 ± 5	5 ± 1	-3 ± 1
UTP-ATP	3	158 ± 10	26 ± 10	-32 ± 5	4 ± 1	-4 ± 2

Table shows cross-desensitization experiments using UTP (100 μM mucosal) and ATP (100 μM mucosal). Tissues were treated with two nucleotides (as shown in Order of Application), the second nucleotide was added in the presence of the first nucleotide, once the I_{SC} had stabilized. I_{SC} before addition of first nucleotide (Basal I_{SC}), peak increase above basal (Stimulatory phase) and change in I_{SC} after 10 min (Inhibitory phase); negative value represent fall below basal I_{SC} in response to first and second nucleotides are shown. Data are mean $\pm$ s.e. mean, n indicates number of trachea. *Indicates significant difference from response to UTP after UTP pretreatment (Student's t -test, $P<0.05$).




Figure 4 Effects of submucosal ATP and UTP. Paired tissues were treated with either mucosal or submucosal UTP (A) or ATP (B) before addition of the same nucleotide to the contralateral side. Data are mean $\pm$ s.e.mean ($n=5$).

by UTP was not significantly different than control tissues. DPC (1 mM mucosal), a Cl^- channel inhibitor, had a similar effect on both basal I_{SC} (Table 3) and the response to UTP (Table 4). Replacement of Cl^- and HCO_3^- with the impermeant anion gluconate inhibited I_{SC} to 25% of control. The initial increase stimulated in I_{SC} by UTP was significantly smaller than in control tissues (Table 4). The subsequent inhibition also tended to be attenuated compared with control, but this was not significant.

Amiloride (10 μM mucosal), a blocker of epithelial Na^+ channels, inhibited $\sim 65\%$ of basal I_{SC} (Table 3). In the presence of amiloride the initial, stimulatory component of the response to UTP was augmented and the subsequent, inhibitory phase of this response was completely abolished (Figure 5, Table 4). In a separate series of experiments, amiloride inhibited significantly less I_{SC} after UTP treatment ($42 \pm 6 \mu\text{A cm}^{-2}$) than in paired trachea not treated with UTP ($78 \pm 6 \mu\text{A cm}^{-2}$), confirming that UTP inhibits basal amiloride-sensitive I_{SC} .

Effect of thapsigargin and Ca^{2+} /calmodulin-dependent kinase inhibitor on response to UTP

To investigate the role of changes in $[\text{Ca}^{2+}]_{\text{in}}$ in the response to UTP, epithelia were pretreated with thapsigargin, to inhibit the uptake of Ca^{2+} into intracellular stores and thereby clamp $[\text{Ca}^{2+}]_{\text{in}}$ at a high level. Thapsigargin alone (0.3 μM , mucosal and submucosal) induced a transient rise in I_{SC} (from $108 \pm 15 \mu\text{A cm}^{-2}$ to $125 \pm 11 \mu\text{A cm}^{-2}$, $n=4$) followed by a gradual decline. The initial UTP-induced stimulation of I_{SC} was abolished in the presence of thapsigargin. The subsequent UTP-induced inhibition of I_{SC} was not significantly different

Table 3 Effect of various agents on basal I_{SC}

Drug	n	I_{SC} ($\mu\text{A cm}^{-2}$)	
		Basal	+ Drug
Bumetanide	6	161 ± 14	$100 \pm 7^*$
DPC	5	115 ± 4	$73 \pm 9^*$
Amiloride	6	119 ± 8	$42 \pm 7^*$
KN93	4	115 ± 15	119 ± 6
Staurosporine	3	133 ± 14	$91 \pm 10^*$
PMA	3	141 ± 13	$121 \pm 13^*$
D-erythro sphingosine	3	128 ± 10	129 ± 14
PMA + staurosporine	4	155 ± 14	$80 \pm 4^*$
H89	6	154 ± 16	$103 \pm 18^*$

Table shows I_{SC} before (Basal) and after (+ Drug) addition of various drugs. Data are mean $\pm$ s.e.mean, n indicates number of trachea. *Indicates significant difference from basal I_{SC} (Paired Student's t -test, $P < 0.05$).

from control tissues (Table 5). Pretreatment with the Ca^{2+} /calmodulin-dependent kinase inhibitor, KN-93 (30 μM mucosal) significantly attenuated both the initial increase and secondary decrease in I_{SC} (Table 5). KN-93 itself had no effect on basal I_{SC} (Table 3).

Effect of protein kinase inhibitors on basal I_{SC} and response to UTP

Staurosporine, a relatively non-specific kinase inhibitor (0.1 μM , mucosal and submucosal) inhibited $\sim 30\%$ of basal I_{SC} (Table 3). This effect was abolished by pretreatment with amiloride ($67 \pm 16 \mu\text{A cm}^{-2}$ after amiloride alone; $61 \pm 15 \mu\text{A cm}^{-2}$ after subsequent treatment with staurosporine).

Table 4 Effect of ion transport inhibitors and ion replacement on response to UTP

Inhibitor/Ion replacement	n	Basal I_{SC} ($\mu A\ cm^{-2}$)		Stimulatory phase ΔI_{SC} ($\mu A\ cm^{-2}$)		Inhibitory phase ΔI_{SC} ($\mu A\ cm^{-2}$)	
		Control	+ Inhib/Replace	Control	+ Inhib/Replace	Control	+ Inhib/Replace
Bumetanide	6	156 $\pm$ 8	100 $\pm$ 8	31 $\pm$ 5	12 $\pm$ 4*	-32 $\pm$ 7	-30 $\pm$ 8
DPC	5	122 $\pm$ 13	73 $\pm$ 9	24 $\pm$ 4	9 $\pm$ 2*	-26 $\pm$ 6	-12 $\pm$ 2
Cl ⁻ and HCO ₃ ⁻ free	6	120 $\pm$ 6	24 $\pm$ 7	31 $\pm$ 5	13 $\pm$ 3*	-24 $\pm$ 6	-7 $\pm$ 4
Amiloride	6	121 $\pm$ 12	41 $\pm$ 6	27 $\pm$ 6	51 $\pm$ 6*	-28 $\pm$ 4	19 $\pm$ 3*

Table shows effect of UTP (100 μM mucosal) on paired tissues either in the absence (Control) or presence ($\pm$ Inhib) of an ion transport inhibitor (either bumetanide (100 μM submucosal), DPC (1 mM mucosal) or amiloride (10 μM mucosal)); or after ion replacement (Replace). I_{SC} before addition of UTP (Basal I_{SC}), peak increase above basal (Stimulatory phase) and change in I_{SC} after 10 min (Inhibitory phase); negative value represents fall below basal I_{SC} are shown. Data are mean $\pm$ s.e.mean, n indicates number of trachea.

*Indicates significant difference from paired control response (Paired Student's t -test, $P < 0.05$).




Figure 5 Effect of ion transport blockers on response to UTP (100 μM mucosal). (A) Shows effect on paired tissues either pretreated with bumetanide (100 μM submucosal; bumetanide) or left without pretreatment (control) ($n=6$). (B) Shows effect on paired tissues either pretreated with amiloride (10 μM mucosal; amiloride) or left without pretreatment (control) ($n=6$). Data are mean $\pm$ s.e.mean.

ine for 20 min, $n=3$). In the presence of staurosporine the UTP-induced increase in I_{SC} was not significantly different from the response of paired control tissues ($P > 0.05$, $n=4$), whilst the inhibitory effect of UTP was abolished (Table 5).

To investigate the role of PKC in regulation of basal Na^+ absorption tissues were treated with PMA, which activates PKC. This substance (16.2 nM, mucosal and submucosal) inhibited $\sim 14\%$ of basal I_{SC} ($P < 0.05$, $n=3$, Table 3) whilst D-erythro sphingosine, a relatively specific PKC inhibitor, (10 μM , mucosal and submucosal) had no effect on basal I_{SC} ($P > 0.05$, $n=3$, Table 3). The subsequent response to UTP was unaffected by pretreatment with either PMA or D-erythro sphingosine (Table 5). To investigate whether the effect of staurosporine was mediated by inhibition of PKC, epithelia

were treated with both staurosporine and PMA because under these conditions any effects of staurosporine on PKC are likely to be opposed by PMA and residual effects of staurosporine are thus likely to be mediated by mechanisms other than inhibition of PKC. This pretreatment significantly inhibited $\sim 46\%$ of basal I_{SC} ($P < 0.05$, $n=4$, Table 3) and the subsequent response to UTP resembled the response of epithelia pretreated with staurosporine (Table 5).

To investigate the role of PKA in regulation of basal I_{SC} , tissues were treated with the PKA inhibitor, H89 (20 μM , mucosal). H89 inhibited basal I_{SC} ($P < 0.05$, $n=6$, Table 3) but had no effect on the subsequent response to UTP (Table 5).

Discussion

The results of this study indicate that mucosal UTP has a biphasic effect on I_{SC} in porcine tracheal epithelium and similar effects have been noted in a number of other airway epithelia (Mason *et al.*, 1991; Devor & Pilewski, 1999; Inglis *et al.*, 1999; Ramminger *et al.*, 1999). We now show, however, that mucosal ATP simply evokes a monophasic response in porcine tracheal epithelium. To our knowledge, this is the first demonstration that ATP and UTP have different effects upon airway ion transport, and it suggests that the effects of nucleotides upon this tissue may be mediated by at least two distinct populations of P2Y receptors. Since UTP and ATP induce initial increases in I_{SC} of similar magnitudes, it is likely that this early response is mediated, at least in part, by P2Y₂ receptors, which are equally sensitive to these nucleotides and have been widely implicated in nucleotide-induced Cl⁻ secretion in airway epithelia (Knowles *et al.*, 1991; Mason *et al.*, 1991; Van Scott *et al.*, 1995; Yamaya *et al.*, 1996). After this initial peak, however, an inhibitory phase of the response to UTP becomes apparent. This, together with the cross desensitization experiments suggests that additional P2Y receptors sensitive to UTP but not to ATP are present in this tissue. Interestingly, submucosal UTP and ATP simply elicited a monophasic response. The inhibitory phase of the response to UTP thus appears to be mediated by ATP-insensitive receptors that are essentially confined to the apical membrane.

At least two such receptor subtypes have been described in airway epithelia. Lazarowski *et al.* (1997) showed that P2Y₆ receptors, which are activated primarily by UDP, are expressed in cultured human nasal epithelia. Since UDP has no discernible effect on either porcine trachea or bronchi (Inglis *et al.*, 1999) it is unlikely that these receptors regulate ion transport in these tissues. This may indicate that human and porcine airways express different P2Y receptor subtypes. Alternatively, it is possible that P2Y₆ receptors are not normally expressed in native airways, and that their expression

Table 5 Effects of agents that modulate signal transduction on response to UTP

Drug	n	Basal I_{SC} ($\mu A\ cm^{-2}$)		Stimulatory phase ΔI_{SC} ($\mu A\ cm^{-2}$)		Inhibitory phase ΔI_{SC} ($\mu A\ cm^{-2}$)	
		Control	+ Drug	Control	+ Drug	Control	+ Drug
Thapsigargin	4	136 ± 20	93 ± 7	34 ± 4	4 ± 1*	-32 ± 14	-12 ± 2
KN93	4	160 ± 11	106 ± 9	20 ± 3	11 ± 1*	-36 ± 2	-15 ± 5*
Staurosporine	3	137 ± 8	91 ± 10	28 ± 3	31 ± 4	-18 ± 5	-1 ± 3*
PMA	3	138 ± 12	122 ± 12	26 ± 4	32 ± 5	-17 ± 7	-22 ± 6
D-Erythro sphingosine	3	148 ± 6	130 ± 14	32 ± 4	46 ± 8	-18 ± 2	-13 ± 3
Staurosporine + PMA	4	144 ± 6	79 ± 3	32 ± 3	21 ± 3	-20 ± 3	-4 ± 3*
H89	6	122 ± 10	103 ± 18	35 ± 5	23 ± 8	-17 ± 2	-13 ± 4

Table shows effect of UTP (100 μM mucosal) on paired tissues either in the absence (Control) or presence (+Drug) of a drug that modulates signal transduction. I_{SC} before addition of UTP (Basal I_{SC}), peak increase above basal (Stimulatory phase) and change in I_{SC} after 10 min (Inhibitory phase); negative value represents fall below basal I_{SC} are shown. Data are mean $\pm$ s.e.mean, indicates number of trachea. *Indicates significant difference from paired control response (Paired student's *t*-test, $P < 0.05$).

may have been induced during the isolation and culture of human nasal epithelia. It is clear that culture conditions can acutely alter the functional expression of P2Y receptors, possibly by controlling gene expression (Clunes *et al.*, 1998; Wilson *et al.*, 1998). A second pyrimidine receptor, P2Y₄, which is sensitive to UTP with UDP acting as a weak partial agonist, has been reported in airway epithelia (Merten *et al.*, 1998; Communi *et al.*, 1999), and this may well be the receptor subtype that mediates UTP-induced inhibition of basal I_{SC} in porcine trachea. We cannot, however, rule out the possibility that another, as yet unidentified receptor mediates these responses.

Experiments with ion transport blockers and ion substitution suggest that UTP initially stimulates anion secretion before inhibiting basal Na⁺ absorption. Nucleotide-induced stimulation of Cl⁻ secretion has been described both *in vivo* (Knowles *et al.*, 1991) and *in vitro*, in native (Iwase *et al.*, 1997; Inglis *et al.*, 1999) and in cultured (e.g. Mason *et al.*, 1991; Van Scott *et al.*, 1995; Yamaya *et al.*, 1996) airways. Inhibition of Na⁺ absorption has also been reported in a number of airway epithelia (Mason *et al.*, 1991; Devor & Pilewski, 1999; Inglis *et al.*, 1999; Ramminger *et al.*, 1999). The increased peak response seen in the presence of amiloride is likely to reflect an increased Cl⁻ secretion, driven by amiloride-induced depolarization of the apical membrane. The maximal dose of UTP inhibited 40% of amiloride-sensitive I_{SC} , indicating that a substantial portion of amiloride-sensitive I_{SC} is not sensitive to inhibition by UTP. Similarly, in cultured human bronchial epithelia, 25% of amiloride-sensitive Na⁺ absorption remains after UTP-evoked inhibition (Devor & Pilewski, 1999).

Since P2Y₂ and P2Y₄ receptors allow external nucleotides to increase $[Ca^{2+}]_{in}$, we anticipated that UTP-induced regulation of Cl⁻ secretion and Na⁺ absorption would be mediated by changes in $[Ca^{2+}]_{in}$. Certainly the stimulation of Cl⁻ secretion appears to be almost completely dependent on $[Ca^{2+}]_{in}$. This is in contrast to our recent studies of porcine distal bronchi showing that their Cl⁻ secretory response to UTP is $[Ca^{2+}]_{in}$ -independent (Inglis *et al.*, 1999), and it suggests that the mechanisms that regulate Cl⁻ secretion are different in different regions of the airway. The mechanisms are clearly complex, since both Ca²⁺-dependent and Ca²⁺-independent components of nucleotide-induced Cl⁻ secretion have been reported (Stutts *et al.*, 1992; 1994; Hwang *et al.*, 1996). Although we cannot rule out the involvement of Ca²⁺-independent mechanisms, our data suggest that in porcine trachea UTP-induced Cl⁻ secretion is mediated primarily by changes in $[Ca^{2+}]_{in}$.

It is well known that increases in $[Ca^{2+}]_{in}$ can inhibit epithelial Na⁺ channels (Ishikawa *et al.*, 1998) and transepithelial Na⁺ absorption (e.g. Graham *et al.*, 1992; Koster *et al.*, 1996) so we may have expected this to be the main mechanism involved in the inhibitory phase of the response to UTP. However, it was clear that the effect of UTP upon Na⁺ absorption is not entirely dependent on $[Ca^{2+}]_{in}$, suggesting that the pyrimidine receptor expressed by this tissue is coupled to additional intracellular mechanisms that allow inhibition of basal I_{SC} . Studies using inhibitors of different kinases suggested that basal I_{SC} was reduced both by increased activity of PKC (as described in sheep trachea Graham *et al.*, 1992) and by inhibition of PKA. These results suggest that the basal rate of ion transport is under complex control, and may be set by the relative activities of PKA and PKC within the cell. Surprisingly, however, the inhibitory effect of UTP on basal Na⁺ absorption does not seem to be mediated by either PKA or PKC, since neither PMA, H89, nor the PKC inhibitor D-erythro sphingosine had any effect on this response. This was unexpected since P2Y receptor-induced activation of phospholipase C is likely to activate PKC. However, a similar lack of involvement of PKC was found in the response of bronchial epithelia to UTP (Devor & Pilewski, 1999). The inhibitory effect of UTP was however blocked by staurosporine, a non specific protein kinase inhibitor. This suggests that another, as yet unidentified protein kinase is involved in this effect of UTP. Another possible mechanism by which UTP may inhibit Na⁺ absorption is through inhibition of basolateral K⁺ channels, as described in human bronchial epithelia (Devor & Pilewski, 1999). This mechanism is also involved in inhibition of Na⁺ absorption by other Ca²⁺-mobilizing agonists, e.g. acetylcholine (Venglarik & Dawson, 1986; Inglis *et al.*, 1992).

In summary, since P2Y receptor agonists can both stimulate Cl⁻ secretion in CF (Mason *et al.*, 1991), and inhibit basal Na⁺ absorption, they provide a potential mechanism by which the ion transport defects in CF may be bypassed therapeutically. The results of this study suggest that the effects on Cl⁻ secretion and Na⁺ absorption may be mediated by distinct P2Y receptor subtypes, which may potentially be activated separately by different therapeutic drugs.

The authors thank Elaine A. Waller for technical assistance. This work was supported by Wellcome (Project Grant 049986; Programme Grant 039124; Career Development Fellowship (S.K. Inglis)), Anonymous and Tenovus Trusts.

References

- ANDERSON, M.P., GREGORY, R.J., THOMPSON, S., SOUZA, D.W., PAUL, S., MULLIGAN, R.C., SMITH, A.E. & WELSH, M.J. (1991). Demonstration that CFTR is a chloride channel by alteration of its anion selectivity. *Science*, **253**, 202–207.
- BOUCHER, R.C. (1999). Molecular insights into the physiology of the 'thin film' of airway surface liquid. *J. Physiol. Lond.*, **516.3**, 631–638.
- BOUCHER, R.C., STUTTS, M.J., KNOWLES, M.R. & GATZY, J.T. (1986). Na⁺ transport in cystic fibrosis respiratory epithelia. Abnormal basal rate and response to adenylate cyclase activation. *J. Clin. Inv.*, **78**, 1245–1252.
- CHRISTOFFERSEN, G.R.J. & SKIBSTED, L.H. (1975). Calcium ion activity in physiological salt solutions: influence of anions substituted for chloride. *Comp. Biochem. Physiol.*, **A52**, 317–322.
- CLUNES, M.T., COLLETT, A., BAINES, D.L., BOVELL, D.L., MURPHIE, H., INGLIS, S.K., MCALROY, H.L., OLVER, R.E. & WILSON, S. (1998). Culture substrate-specific expression of P2Y₂ receptors in distal lung epithelial cells isolated from foetal rats. *Br. J. Pharm.*, **124**, 845–847.
- COMMUNI, D., PAINDAVOINE, P., PLACE, G.A., PARMENTIER, M. & BOEYNAEMS, J.-M. (1999). Expression of P2Y receptors in cell lines derived from the human lung. *Br. J. Pharm.*, **127**, 562–568.
- DEVOR, D.C. & PILEWSKI, J.M. (1999). UTP inhibits Na absorption in wild-type and deltaF508 CFTR-expressing human bronchial epithelia. *Am. J. Physiol.*, **276**, C827–C837.
- GRAHAM, A., STEEL, D.M., ALTON, E.W.F.W. & GEDDES, D.M. (1992). Second-messenger regulation of sodium transport in mammalian airway epithelia. *J. Physiol. Lond.*, **453**, 475–491.
- HWANG, T.-H., SCHWIEBERT, E.M. & GUGGINO, W.B. (1996). Apical and basolateral ATP stimulates tracheal epithelial chloride secretion via multiple purinergic receptors. *Am. J. Physiol.*, **270**, C1611–C1623.
- INGLIS, S.K., COLLETT, A., MCALROY, H.L., WILSON, S.M. & OLVER, R.E. (1999). Effect of luminal nucleotides on Cl[−] secretion and Na⁺ absorption in distal bronchi. *Pflügers Arch.*, **438**, 621–627.
- INGLIS, S.K., WARD, M.R. & ACEVEDO, M. (1992). Changes in the basolateral potassium conductance induced by acetylcholine (ACh) in isolated sheep tracheal epithelium. *J. Physiol. Lond.*, **452**, 144P.
- INOUE, C.N., WOO, J.S., SCHWIEBERT, E.M., MORITA, T., HANAOKA, K., GUGGINO, S.E. & GUGGINO, W.B. (1997). Role of purinergic receptors in chloride secretion in Caco-2 cells. *Am. J. Physiol.*, **272**, C1862–C1870.
- ISHIKAWA, T., MARUNAKA, Y. & ROTIN, D. (1998). Electrophysiological characterisation of the rat epithelial Na⁺ channel (rENaC) expressed in MDCK cells. *J. Gen. Physiol.*, **111**, 825–846.
- ISMAILOV, I.I., AWAYDA, M.S., JOVOV, B., BERDIEV, B.K., FULLER, C.M., DEDMAN, J.R., KAETZEL, M.A. & BENOS, D.J. (1996). Regulation of epithelial sodium channels by the cystic fibrosis transmembrane conductance regulator. *J. Biol. Chem.*, **271**, 4725–4732.
- IWASE, N., SASAKI, T., SHIMURA, S., YAMAMOTO, M., SUZUKI, S. & SHIRATO, K. (1997). ATP-induced Cl secretion with suppressed Na⁺ absorption in rabbit tracheal epithelium. *Resp. Physiol.*, **107**, 173–180.
- KNOWLES, M.R., CLARKE, L.L. & BOUCHER, R.C. (1991). Activation by extracellular nucleotides of chloride secretion in the airway epithelia of patients with cystic fibrosis. *N. Engl. J. Med.*, **325**, 533–538.
- KNOWLES, M.R., OLIVIER, K., NOONE, P. & BOUCHER, R.C. (1996). Pharmacological modulation of salt and water transport in the airway epithelium in cystic fibrosis. *Am. J. Resp. Crit. Care Med.*, **151**, S65–S69.
- KO, W.H., LAW, W.Y.K., WONG, C.Y. & WILSON, S.M. (1999). The simultaneous measurement of I_{SC} and intracellular free Ca²⁺ in equine cultured equine sweat gland secretory epithelia. *J. Membrane Biol.*, **170**, 205–211.
- KOSTER, H.P.G., HARTOG, A., VAN OS, C.H. & BINDELS, R.J.M. (1996). Inhibition of Na⁺ and Ca²⁺ reabsorption by P2u purinoceptors requires PKC but not Ca²⁺ signalling. *Am. J. Physiol.*, **270**, F53–F60.
- LAZAROWSKI, E.R., PARADISO, A.M., WATT, W.C., HARDEN, K. & BOUCHER, R.C. (1997). UDP activates a mucosal-restricted receptor on human nasal epithelial cells that is distinct from the P2Y₂ receptor. *Proc. Natl. Acad. Sci.*, **94**, 2599–2603.
- MASON, S.J., PARADISO, A.M. & BOUCHER, R.C. (1991). Regulation of transepithelial ion transport and intracellular calcium by extracellular ATP in human and cystic fibrosis airway epithelium. *Br. J. Pharm.*, **103**, 1649–1656.
- MERTEN, M.D., SALEH, A., KAMMOUNI, W., MARCHAND, S. & FIGARELLA, C. (1998). Characterisation of two distinct P2Y receptors in human tracheal gland cells. *Eur. J. Biochem.*, **251**, 19–24.
- PARR, C.E., SULLIVAN, D.M., PARADISO, A.M., LAZAROWSKI, E.R., BIRCH, L.H., OLSEN, J.C., ERB, L., WEISMAN, G.A., BOUCHER, R.C. & TURNER, J.T. (1994). Cloning and expression of a human P2U nucleotide receptor, a target for cystic fibrosis pharmacotherapy. *Proc. Natl. Acad. Sci. U.S.A.*, **91**, 3275–3279.
- RAMMINGER, S.J., COLLETT, A., BAINES, D.L., MURPHIE, H., MCALROY, H.L., OLVER, R.E., INGLIS, S.K. & WILSON, S.M. (1999). P2Y₂ receptor-mediated inhibition of ion transport in distal lung epithelial cells. *Br. J. Pharm.*, **128**, 293–300.
- STUTTS, M.J., CANESSA, C.M., OLSEN, J.C., HAMRICK, M., COHN, J.A., ROSSIER, B.C. & BOUCHER, R.C. (1995). CFTR as a cAMP-dependent regulator of sodium channels. *Science*, **269**, 847–850.
- STUTTS, M.J., CHINET, T.C., MASON, S.J., FULLTON, J.M., CLARKE, L.L. & BOUCHER, R.C. (1992). Regulation of Cl[−] channels in normal and cystic fibrosis airway epithelial cells by extracellular ATP. *Proc. Natl. Acad. Sci. U.S.A.*, **89**, 1621–1625.
- STUTTS, M.J., FITZ, J.G., PARADISO, A.M. & BOUCHER, R.C. (1994). Multiple modes of regulation of airway epithelial chloride secretion by extracellular ATP. *Am. J. Physiol.*, **267**, C1442–C1451.
- VAN SCOTT, M.R., CHINET, T.C., BURNETTE, A.D. & PARADISO, A.M. (1995). Purinergic regulation of ion transport across nonciliated bronchiolar epithelial (Clara) cells. *Am. J. Physiol.*, **269**, L30–L37.
- VENGLARIK, C.J. & DAWSON, D.C. (1986). Cholinergic regulation of Na absorption by turtle colon: role of basolateral K conductance. *Am. J. Physiol.*, **251**, C563–C570.
- WILSON, S.M., LAW, W.Y.Y., PEDIANI, J.D., ALLEN, E.A., WILSON, G., KHAN, Z. & KO, W.H. (1998). Nucleotide-evoked calcium signals and anion secretion in equine cultured epithelia that express P2Y₂ receptors and pyrimidine nucleotide receptors. *Br. J. Pharm.*, **124**, 832–838.
- YAMAYA, M., SEKIZAWA, K., KAKUTA, Y., OHROI, T., SAWAI, T. & SASAKI, H. (1996). P2u-purinoceptor regulation of chloride secretion in cultured human tracheal submucosal glands. *Am. J. Physiol.*, **270**, L979–L984.

(Received January 18, 2000)

Accepted February 29, 2000)