

Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

A Practical Synthesis of 2'-Fluoro- 4'-hydroxyacetophenone

Xiao-Bo Zhang ^b, Qun Lu ^a & Wei-Ming Xu ^a

^a College of Material Chemistry and Chemical Engineering , Hangzhou Normal University (Xiasha Campus) , Hangzhou, P. R. China

^b Department of Chemistry , Zhejiang University (Xixi Campus) , Hangzhou, P. R. China

Published online: 08 Sep 2009.

To cite this article: Xiao-Bo Zhang , Qun Lu & Wei-Ming Xu (2009) A Practical Synthesis of 2'-Fluoro-4'-hydroxyacetophenone, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 41:5, 419-421, DOI: [10.1080/00304940903163723](https://doi.org/10.1080/00304940903163723)

To link to this article: <http://dx.doi.org/10.1080/00304940903163723>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

OPPI BRIEFS

A Practical Synthesis of 2'-Fluoro-4'-hydroxyacetophenone

Xiao-Bo Zhang,² Qun Lu,¹ and Wei-Ming Xu¹

¹College of Material Chemistry and Chemical Engineering, Hangzhou Normal University (Xiasa Campus) Hangzhou, P. R. China

²Department of Chemistry, Zhejiang University (Xixi Campus) Hangzhou, P. R. China

2'-Fluoro-4'-hydroxyacetophenone (**2**) is an important building block for many pharmaceutically active agents,¹ used to treat bacterial and parasitic infections.² A review of the literature indicated that its synthesis has rarely been reported.³

Kees and Musser *et al.* developed the only method to obtain 2'-fluoro-4'-hydroxyacetophenone (**2**) from 3-fluorophenol (**1**) through Friedel-Crafts reaction (*Scheme 1*), but the yield was not reported.³ Numerous repetitions of the procedure showed that the yield never exceeded 40% and at least 18% of 4'-fluoro-2'-hydroxyacetophenone accompanied the desired product. Furthermore, the starting compound, 3-fluorophenol (**1**) is expensive to purchase and tedious to prepare.^{4,5}




Scheme 1

Based on the report of Gryko *et al.*⁶ and as a continuation of our interest in the study of medicinal compounds,⁷ we have developed a practical pilot-scale method for the preparation of 2'-fluoro-4'-hydroxyacetophenone (**2**) from 2',4'-difluoroacetophenone (**3**) (*Scheme 2*); the latter was easily prepared by acetylation of 1,3-difluorobenzene.⁸

Submitted April 17, 2009.

Address correspondence to Wei-Ming Xu, College of Material Chemistry and Chemical Engineering, Hangzhou Normal University (Xiasa Campus), Hangzhou 310036, P. R. China. E-mail: wmxu@zju.edu.cn



Scheme 2

It is noteworthy that the use of a mixture of sodium hydroxide and calcium hydroxide is necessary in this reaction in order to ensure an adequate reaction rate and an acceptable yield of the product.

Experimental Section

Mps and bps are uncorrected. The purity of products was established on an Agilent 1100 HPLC. ^1H NMR spectra were recorded in CDCl_3 on a Bruker 400 (400 MHz) instrument with TMS as internal standard. All chemicals were reagent grade and available commercially. The elemental analysis was performed on a Flash EA1112 instrument.

2'-Fluoro-4'-hydroxyacetophenone

In a 1 L round-bottomed flask fitted with a mechanical stirrer was placed 60.0 g (0.38 mol) of 2',4'-difluoroacetophenone (**3**), 14.1 g calcium hydroxide, 16.0 g sodium hydroxide and 500 mL water. The mixture was stirred under reflux for 18 h. Then the mixture was cooled to room temperature and filtered. The filtrate was extracted with 150 mL CH_2Cl_2 to recover 11.2 g 2',4'-difluoroacetophenone (HPLC > 96%), and acidified with 100 mL of 4M hydrochloric acid followed by steam distillation to remove a small quantity of 4'-fluoro-2'-hydroxyacetophenone (about 1.1 g). The water phase was cooled to 5°C , and the precipitated crystals collected and dried *in vacuo* to afford 38.1 g (79%) of the crude product (**2**) as a white solid (HPLC > 98.5%). ^1H NMR ($\text{DMSO}-d_6$): δ 2.46 (3 H, d, $J = 4.8$ Hz), 6.65 (1 H, d, $J = 8.8$ Hz), 6.74 (1 H, d, $J = 8.8$ Hz), 7.75 (1 H, t, $J = 8.8$ Hz), 10.85 (1 H, b). ^{13}C NMR ($\text{DMSO}-d_6$): δ 193.7, 164.1 ($J = 12.1$ Hz), 163.6 ($J = 252.5$ Hz), 132.4 ($J = 4.7$ Hz), 117.1 ($J = 12.0$ Hz), 112.5 ($J = 7.8$ Hz), 103.3 ($J = 25.4$ Hz), 31.0 ($J = 7.0$ Hz). An analytical sample was prepared by recrystallization from methanol and water (*v/v* = 30/70), mp. 120–121 $^\circ\text{C}$.⁹

Anal. Calcd. for $\text{C}_8\text{H}_7\text{FO}_2$: C, 62.34; H, 4.58. Found: C, 62.44; H, 4.66.

Acknowledgment

We thank the National Natural Science Foundation of China (Grant No. 20602029) and the Science and Technology Foundation of Zhejiang Province (Grant No. 2007C11014) for financial support.

References

1. J. M. Salvino, G. C. Morton, H. J. Mason and R. F. Labaudiniere, *WO9967192* **1999**; *Chem. Abstr.*, **132**, 63782 (1999).

2. S. F. Nielsen, K. Schonning, H. Kromann, T. Boesen and M. Larsen, *WO2003097576* **2003**; *Chem. Abstr.*, **140**, 4852 (2003).
3. K. F. Kees, J. H. Musser, J. Chang, M. Skowronek and A. J. Lewis, *J. Med. Chem.*, **29**, 2329 (1986).
4. K. Momota, T. Yonezawa and Y. Inoue, *JP03232829* **1991**; *Chem. Abstr.*, **116**, 105801 (1991).
5. N. Takechi, Y. Fukai and W. Oka, *JP08188545* **1996**; *Chem. Abstr.*, **125**, 247382 (1996).
6. D. T. Gryko, D. Wyrostek, A. Nowak-Krol, K. Abramczyk and M. K. Rogacki, *Synthesis*, 4028 (**2008**).
7. H. Gu and W. M. Xu, *Org. Prep. Proced. Int.*, **40**, 465 (2008).
8. Y. Liu, S. Wang, C. Zheng, S. Song and L. Kou, *Huaxue Shijie*, **43**, 432 (2002); *Chem. Abstr.*, **139**, 133284 (2002).
9. None of the previous references has reported a melting point for this compound.