

Synthesis of (–)-pericosine B, the antipode of the cytotoxic marine natural product†

Yoshihide Usami,* Kentaro Suzuki, Koji Mizuki, Hayato Ichikawa and Masao Arimoto

Received 29th July 2008, Accepted 15th October 2008

First published as an Advance Article on the web 13th November 2008

DOI: 10.1039/b813072h

The stereoselective synthesis of (–)-pericosine B, which is the antipode of the cytotoxic metabolite of the fungus *Periconia byssoides* OUPS-N133 separated from the sea hare, was accomplished in 9 steps in 12% total yield from (–)-quinic acid, together with the synthesis of its epimer. Every crucial step of this total synthesis, including ring opening of a β -epoxide and NaBH_4 reduction of an unstable β,γ -unsaturated enone, proceeded with excellent stereoselectivity.

Introduction

The isolation and structure determination of highly functionalized C-7 cyclohexenoid natural products pericosines A–E **1–5** (Fig. 1), which are cytotoxic metabolites of the fungus *Periconia byssoides* OUPS-N133 originally separated from the sea hare *Aplysia kurodai*, were reported in 1997 and 2007 by Numata and co-workers.^{1,2} The absolute configuration of pericosines A–D **1–4** was elucidated by total syntheses.^{3–9} Compound **1** was reported to exhibit significant inhibitory activity against protein kinase EGFR and human topoisomerase II, but similar biological tests on **2–4** were not reported. We were therefore interested in the biological activity of **2** against human cancer cell lines. In addition to their significant bioactivity, it is noteworthy that pericosines C **3** and E **5** exist as a mixture of enantiomers. X-ray analysis by Numata and co-workers² established the relative stereochemistry of pericosine E **5** to originate from two monomeric pericosines **1** and **2** having different chirality. As we pointed out previously, their finding meant that the presence of the antipode of monomeric **1** or **2** in nature is possible.¹⁰ Therefore, synthesis of the antipode of **1–3** is significant. We have already reported the synthesis of the antipode of **1**^{6,7} and **3**⁸ but not that of **2**.

The total synthesis of (+)-**2**, a naturally occurring enantiomer, was reported only once in 1998 by Donohoe and co-workers⁹ in

spite of extensive effort by other groups including ours to date.^{11–13} However, the only successful synthesis had problems in terms of the use of expensive starting materials and a stoichiometric amount of toxic osmium tetroxide. Recently, we reported the determination of the absolute configuration of pericosine D by a synthetic approach.⁸ Unfortunately, the total yield of desired product **4** was quite low when relatively inexpensive (–)-quinic acid was used. Nevertheless, that work gave us a hint for a short synthesis of **2**. Following several years of failed attempts at synthesizing **2**,¹¹ we describe herein a short synthesis of the antipode of **2** and its epimer.

Results and discussion

From the results of our previous work that dealt with the determination of pericosine D **4**, we suspected that the introduction of a 6 α -methoxy group into the pericosine core 6-membered ring is possible when MeOH is used as solvent in the stereoselective ring opening of intermediate β -epoxide **8**.⁸

The synthesis of (–)-pericosine B is summarized in Scheme 1 followed by Scheme 2. Methyl quinate derivative **6** was prepared from commercially available (–)-quinic acid in 78% according to the literature.¹⁴ Compound **6** was converted into unstable diene **7** in 2 steps. Then, without purification, crude **7** was oxidized with mCPBA at 40 °C to afford an inseparable mixture of epoxides **8** and **9** in 40% yield in 3 steps.

Ring opening of a mixture of **8** and **9** with a catalytic amount of HCl in MeOH gave the desired 6 α -methoxypericosine derivative **10** in 54% yield, with small amount of **11**⁸ (1%) and 1-methoxylated alcohol **12** (1%) (whose configuration at C-1 could not be determined), with the recovery of **9** (32%). The relative

Osaka University of Pharmaceutical Sciences, 4-20-1 Nasahara, Takatsuki, Osaka, 569-1094, Japan. E-mail: usami@gly.oups.ac.jp; Fax: +81 72 690 1005; Tel: +81 72 690 1083

† Electronic supplementary information (ESI) available: ¹H- and ¹³C-NMR spectra of synthetic (–)-pericosine B, compounds **10**, **14**, natural pericosine B, and its acetone. NOESY spectrum of compound **10**. See DOI: 10.1039/b813072h








Fig. 1 Structures of naturally occurring pericosines.



Scheme 1 Stereoselective synthesis of 6 α -methoxyalcohol **10**.



Scheme 2 Completion of total synthesis of (-)-pericosine **B 2**.

stereochemistry of major product **10** was confirmed by NOESY analysis as shown in Scheme 1. Cross-peaks of H-4/H-6 and H-5/6-methoxy group were observed.

In next step, it was difficult to promote the S_N2 -type Walden inversion by Mitsunobu reaction at C-5 in **10**. Close inspection of the ^1H -NMR spectra of **10**, which had relatively large coupling constants of $J_{4,5} = 7.3$ Hz and $J_{5,6} = 6.6$ Hz, suggested a half-chair

conformation, as illustrated in Fig. 2, that inhibits S_N2 -type attack of the nucleophile from inside the pericosine core 6-membered ring, which is in a fixed conformation due to the cyclohexylidene bridge.




Fig. 2 Plausible conformation of **10**.

Then, the inversion of stereochemistry of C-5 in **10** was attempted by means of Dess–Martin oxidation followed by stereoselective reduction with NaBH_4 . Methoxy alcohol **10** was oxidized with Dess–Martin periodinane, albeit very slowly, to give crude β,γ -unsaturated enone **13**. Without purification, **13**¹⁵ was reduced with NaBH_4 at 0°C to give the desired diastereomer **14** as the sole product in 86% yield in 2 steps.

This total synthesis was completed by deprotection of the cyclohexylidene moiety in **14** with TFA in MeOH to afford (-)-pericosine **B 2** in 82% yield. All spectral data except specific rotation agreed with those of reported natural pericosine **B**. The specific rotation ($[\alpha]_D^{25} -32.6$) of synthesized compound showed almost the same value as previously synthesized (+)-**2** ($[\alpha]_D^{21} +30.6$ (c 0.8 in EtOH)) but with the opposite sign.




Fig. 3 Structures of undesired compounds.¹⁵

The overall yield of this total synthesis of **2** was 12% in 9 steps starting from (-)-quinic acid. Similarly, epimer **15** was prepared in 57% yield from **10**.

Conclusions

We have accomplished the stereoselective total synthesis of (-)-pericosine **B 2**, which has opposite chirality to the natural product, in 9 steps in 12% total yield. Its epimer **15** was also prepared. The second synthesis of pericosine **B** described herein is a toxic-reagent-free method and is also applicable to the synthesis of (+)-pericosine **B**, which was obtained as a minor component in nature and has significant biological activity, since either enantiomer of unstable diene **7**¹⁶ could be prepared from (-)-quinic acid. This synthetic route toward **2** is also a divergent one, as the common intermediate yielded pericosine **D 4**.⁸

Experimental section

General information

IR spectra were obtained with a JEOL FT/IR-680 Plus spectrometer. HRMS was determined with a JEOL JMS-700 (2) mass spectrometer. NMR spectra were recorded at 27 °C on Varian UNITY INOVA-500 and Mercury-300 spectrometers in CDCl₃ with tetramethylsilane (TMS) as internal standard. Melting points were determined on a Yanagimoto micromelting point apparatus and are uncorrected. Specific rotations were measured on a JASCO DIP-1000 polarimeter and $[\alpha]_D$ values are given in 10⁻¹ deg cm² g⁻¹. Liquid column chromatography was conducted over silica gel (Nacalai, silica gel 60, mesh 70–230 or 230–400). Analytical TLC was performed on precoated Merck glass plates (silica gel 60 F₂₅₄), and compounds were detected by dipping an ethanol solution of phosphomolybdic acid, followed by heating. Dry THF was distilled over sodium benzophenone ketyl under argon atmosphere.

Synthesis of a mixture of epoxides **8** and **9** from **6**

To a solution of diol **6** (552 mg, 1.93 mmol) in CH₂Cl₂ (25 mL) were added pyridine (780.5 μL, 9.65 mmol) and catalytic DMAP (30.0 mg). A solution of Tf₂O (729 μL, 4.25 mmol) in CH₂Cl₂ (25 mL) was added dropwise to the mixture at 0 °C with stirring. After stirring overnight at rt, the reaction mixture was treated with aqueous NaHCO₃ and then extracted with CH₂Cl₂. The organic layer was dried over MgSO₄, filtered, and evaporated to give a crude residue containing triflates and diene **7**. The mixture was dissolved in DMF (3 mL), and CsOAc (1.99 mmol) was added to the solution with stirring at rt. After 3 hr, the reaction mixture was extracted with *t*-butylmethylether and H₂O. The organic layer was washed with brine twice, dried over MgSO₄, filtered, and evaporated to afford crude diene **7**. To a solution of crude diene **7** in CH₂Cl₂ (10 mL), mCPBA (518.8 mg, max 77%, calcd *ca.* 2.3 mmol) was added and the reaction mixture was kept at 40 °C overnight. The reaction mixture was treated with aqueous NaHCO₃ and then extracted with CH₂Cl₂. The organic layer was separated, dried over MgSO₄, filtered, and evaporated to afford a crude residue that was purified by column chromatography (eluent: hexane:EtOAc = 4:1) to give a mixture of **8** and **9** (204.9 mg, 40% from **6**, ratio: **8**:**9** = 3:2 from ¹H-NMR spectrum).

Methyl (3*R*,4*R*,5*S*,6*S*)-3,4-*O*-cyclohexylidene-3,4,5-trihydroxy-6-methoxy-1-cyclohexene-1-carboxylate **10**

Methyl (4*S*,5*R*,6*R*)-4,5-*O*-cyclohexylidene-4,5,6-trihydroxy-1-methoxy-2-cyclohexene-1-carboxylate **12**

To a mixture of **8** and **9** (*ca.* 3:2) (73.9 mg, combined amount) in MeOH (5 mL) was added 1 drop of 1.0 M HCl in Et₂O with a microsyringe. After stirring overnight at rt, the reaction mixture was condensed under reduced pressure to afford a crude residue that was purified by column chromatography (eluent: hexane:EtOAc = 5:3) to give **10** (44.9 mg, 54%), **11** (0.8 mg, 1%), and **12** (0.9 mg, 1%) with recovery of **9** (23.6 mg, 32%). **10**: Colorless crystals (CH₂Cl₂); mp 132–135 °C; $[\alpha]_D^{25}$ –55.2 (*c* 0.165 in CHCl₃); IR ν_{\max} (KBr)/cm⁻¹ 3395 (OH), 1719 (C=O), 1660 (C=C); ¹H-NMR (500 MHz; CDCl₃; Me₄Si) δ 1.25–1.70

(10H, m), 3.60 (3H, s, 6-OMe), 3.80 (3H, s, COOMe), 3.94 (1H, dd, *J* = 7.3, 6.6 Hz, H-5), 4.00 (1H, dt, *J* = 6.6, 1.4 Hz, H-6), 4.17 (1H, dd, *J* = 7.3, 6.4 Hz, H-4), 4.66 (1H, ddd, *J* = 6.4, 3.9, 1.4 Hz, H-3), 6.68 (1H, dd, *J* = 3.9, 1.4 Hz, H-2); ¹³C-NMR (125.6 MHz; CDCl₃; Me₄Si) δ 23.6 (t), 24.0 (t), 25.0 (t), 35.3 (t), 37.8 (t), 52.0 (q), 60.5 (q), 70.7 (d), 72.6 (d), 75.9 (d), 78.0 (d), 111.7 (s), 132.3 (d), 134.7 (s), 166.4 (s); EIMS *m/z* 298 (M⁺, 76%); HREIMS *m/z* calcd for C₁₅H₂₂O₆ (M)⁺ 298.1416, found 298.1415.

12: Colorless oil; $[\alpha]_D^{25}$ +146.8 (*c* 0.5 in CHCl₃); IR ν_{\max} (liquid film)/cm⁻¹ 3553 (OH), 1724 (C=O), 1660 (C=C); ¹H-NMR (500 MHz; CDCl₃; Me₄Si) δ 1.35–1.77 (10H, m), 2.63 (1H, d, *J* = 3.4 Hz, 6-OH), 3.45 (3H, s, –OMe), 3.78 (3H, s, COOMe), 3.91 (1H, dd, *J* = 8.4, 3.4 Hz, H-4), 4.45 (1H, dd, *J* = 8.4, 7.1 Hz, H-5), 4.78 (1H, ddd, *J* = 7.1, 3.4, 1.1 Hz, H-6), 5.86 (1H, dd, *J* = 10.1, 1.1 Hz, H-1), 6.17 (1H, dd, *J* = 10.1, 3.4 Hz, H-2); ¹³C-NMR (125.6 MHz; CDCl₃; Me₄Si) δ 23.5 (t), 24.0 (t), 25.1 (t), 34.5 (t), 37.6 (t), 52.2 (q), 52.8 (q), 71.9 (d), 73.1 (d), 76.5 (d), 83.3 (s), 111.2 (s), 127.7 (d), 129.9 (d), 170.0 (s); EIMS *m/z* 298 (M⁺, 85%); HREIMS *m/z* calcd for C₁₅H₂₂O₆ (M)⁺ 298.1416, found 298.1416.

Methyl (3*R*,4*R*,5*R*,6*S*)-3,4-*O*-cyclohexylidene-3,4,5-trihydroxy-6-methoxy-1-cyclohexene-1-carboxylate **14**

To a solution of **10** (13.8 mg, 0.046 mmol) in CH₂Cl₂ (2 mL) was added Dess–Martin periodinane (99.1 mg, 0.23 mmol) at rt with stirring. After 24 hr, the reaction mixture was diluted with *tert*-butylmethylether and treated with aq. Na₂S₂O₄ and aq. NaHCO₃. The organic layer was separated, dried over MgSO₄, and filtered and the solvent was removed under reduced pressure to afford crude ketone **13**. Data of crude **13**: IR ν_{\max} (liquid film)/cm⁻¹ 1726 (C=O), 1612 (C=C); ¹H-NMR (300 MHz; CDCl₃; Me₄Si) δ 1.25–1.70 (10H, m), 3.60 (3H, s, 6-OMe), 3.80 (3H, s, COOMe), 3.94 (1H, dd, *J* = 7.3, 6.6 Hz, H-5), 4.00 (1H, dt, *J* = 6.6, 1.4 Hz, H-6), 4.17 (1H, dd, *J* = 7.3, 6.4 Hz, H-4), 4.66 (1H, ddd, *J* = 6.4, 3.9, 1.4 Hz, H-3), 6.68 (1H, dd, *J* = 3.9, 1.4 Hz, H-2); ¹³C-NMR (75.5 MHz; CDCl₃; Me₄Si) δ 24.1 (t), 25.2 (t), 30.0 (t), 35.8 (t), 37.2 (t), 52.6 (q), 59.6 (q), 74.5 (d), 77.1 (d), 113.3 (s), 133.0 (d), 134.3 (s), 164.6 (s), 200.9 (s); EIMS *m/z* 296 (M⁺, 94%); HREIMS *m/z* calcd for C₁₅H₂₀O₆ (M)⁺ 296.1260, found 296.1262.

To a suspension of NaBH₄ in MeOH (0.5 mL) was added crude ketone **13** dissolved in CH₂Cl₂ (2 mL) at 0 °C with stirring. After 1 hr, the reaction mixture was treated with aq. NH₄Cl and extracted with CH₂Cl₂. The organic layer was dried over MgSO₄ and filtered. The solvent was removed under reduced pressure to afford a crude residue that was purified by column chromatography (eluent: EtOAc: hexane = 1: 1) to give **14** (11.9 mg, 86% in 2 steps). **14**: Colorless oil; $[\alpha]_D^{25}$ +25.1 (*c* 0.12 in CHCl₃); IR ν_{\max} (liquid film)/cm⁻¹ 3497 (OH), 1720 (C=O), 1656 (C=C); ¹H-NMR (300 MHz; CDCl₃; Me₄Si) δ 1.25–1.70 (10H, m), 3.17 (1H, d, *J* = 11.5 Hz, 5-OH), 3.58 (3H, s, 6-OMe), 3.81 (3H, s, COOMe), 3.83 (1H, m, H-5), 4.28 (1H, dd, *J* = 4.9, 0.6 Hz, H-6), 4.48 (1H, ddd, *J* = 5.7, 3.4, 0.7 Hz, H-3), 4.66 (1H, dd, *J* = 5.7, 3.2 Hz, H-4), 6.82 (1H, br d, *J* = 3.4 Hz, H-2); ¹³C-NMR (75.5 MHz; CDCl₃; Me₄Si) δ 24.1 (t), 24.3 (t), 25.3 (t), 36.0 (t), 37.5 (t), 52.3 (q), 61.3 (q), 68.1 (d), 72.1 (d), 73.0 (d), 74.2 (d), 111.5 (s), 129.7 (s), 137.3 (d), 166.0 (s); EIMS *m/z* 298 (M⁺, 69%); HREIMS *m/z* calcd for C₁₅H₂₂O₆ (M)⁺ 298.1416, found 298.1419.

(–)-Pericosine B: Methyl (3*R*,4*R*,5*R*,6*R*)-6-methoxy-3,4,5-trihydroxy-1-cyclohexene-1-carboxylate 2

To a solution of alcohol **14** (13.2 mg) in MeOH (0.5 mL) was added TFA (0.5 mL, excess) at 0 °C and the reaction mixture was stirred for 1 hr. After stirring for another 4 hr at rt, the reaction mixture was condensed under reduced pressure to afford a crude residue that was purified by silica gel chromatography (eluent: 3–5% MeOH in CH₂Cl₂) to give (–)-**2** (7.9 mg, 82%). (–)-**2**: Colorless crystals (hexane-EtOAc); mp 69–71 °C; $[\alpha]_D^{25}$ –32.6 (*c* 0.35 in EtOH); IR ν_{\max} (liquid film)/cm^{–1} 3433 (OH), 1713 (C=O), 1651 (C=C); ¹H-NMR (500 MHz; acetone-*d*₆; Me₄Si) δ 3.60 (3H, s, 6-OMe), 3.78 (3H, s, COOMe), 3.85 (1H, dd, *J* = 4.1, 2.0 Hz, H-5), 3.98 (1H, m, H-4), 4.20 (1H, m, H-3), 4.26 (1H, ddd, *J* = 4.1, 1.1, 0.9 Hz, H-6), 6.74 (1H, dd, *J* = 2.5, 1.1 Hz, H-2); ¹³C-NMR (125.6 MHz; acetone-*d*₆; Me₄Si) δ 52.2 (q), 61.5 (q), 69.5 (d), 70.0 (d), 72.8 (d), 77.0 (d), 130.5 (s), 141.9 (d), 166.9 (s); EIMS *m/z* 219 (M⁺, 0.8%), 186 (M⁺–MeOH, 8%); HREIMS *m/z* calcd for C₉H₁₅O₆ (M + H)⁺ 219.0868, found 219.0860.

Methyl (3*R*,4*R*,5*S*,6*S*)-6-methoxy-3,4,5-trihydroxy-1-cyclohexene-1-carboxylate 15

Alcohol **10** (11.4 mg) was converted to (–)-**15** (4.7 mg, 57%) by the same process as above. (–)-**15**: Colorless crystals (hexane-EtOAc); mp 94–97 °C; $[\alpha]_D^{25}$ –75.6 (*c* 0.23 in EtOH); IR ν_{\max} (KBr)/cm^{–1} 3418 (OH), 1716 (C=O), 1651 (C=C); ¹H-NMR (500 MHz; acetone-*d*₆; Me₄Si) δ 3.51 (3H, s, 6-OMe), 3.676 (1H, dd, *J* = 7.3, 4.1 Hz, H-4), 3.76 (3H, s, COOMe), 3.98 (1H, ddd, *J* = 4.8, 0.9, 0.7 Hz, H-6), 4.11 (1H, dd, *J* = 7.3, 4.8 Hz, H-5), 4.32 (1H, br dd, *J* = 4.1, 3.9 Hz, H-3), 6.69 (1H, ddd, *J* = 3.9, 0.9, 0.5 Hz, H-2); ¹³C-NMR (125.6 MHz; acetone-*d*₆; Me₄Si) δ 52.0 (q), 59.7 (q), 66.7 (d), 70.5 (d), 71.2 (d), 79.3 (d), 132.3 (s), 139.2 (d), 167.2 (s); EIMS *m/z* 219 (M⁺, 0.4%), 186 (M⁺–MeOH, 6%); HREIMS *m/z* calcd for C₉H₁₅O₆ (M + H)⁺ 219.0868, found 219.0861.

Acknowledgements

We are grateful to Dr. K. Minoura and Ms. M. Fujitake of this University for NMR and MS measurements, respectively. We also thank Dr. T. Yamada of this University for providing copies of ¹H- and ¹³C-NMR spectra of natural pericosine B and its acetone, which appear in the ESI[†]. This work was supported in part by a Grant-in-Aid from “Dousoukai” of Osaka University of Pharmaceutical Sciences awarded to Y. Usami.

References

- 1 A. Numata, M. Iritani, T. Yamada, K. Minoura, E. Matsumura, T. Yamori and T. Tsuruo, *Tetrahedron Lett.*, 1997, **38**, 8215–8218.
- 2 T. Yamada, M. Iritani, H. Ohishi, K. Tanaka, M. Doi, K. Minoura and A. Numata, *Org. Biomol. Chem.*, 2007, **5**, 3979–3986.
- 3 Y. Usami, C. Hatsuno, H. Yamamoto, M. Tanabe and A. Numata, *Chem. Pharm. Bull.*, 2004, **52**, 1130–1133 (Erratum: *Chem. Pharm. Bull.* 2005, **53**, 271.).
- 4 Y. Usami and Y. Ueda, *Chem. Lett.*, 2005, **34**, 1062–1063.
- 5 Y. Usami and Y. Ueda, *Synthesis*, 2007, 3219–3225.
- 6 Y. Usami, Y. Horibe, I. Takaoka, H. Ichikawa and M. Arimoto, *Synlett*, 2006, 1598–1600.
- 7 Y. Usami, I. Takaoka, H. Ichikawa, Y. Horibe, T. Tomiyama, M. Otsuka, Y. Imanishi and M. Arimoto, *J. Org. Chem.*, 2007, **68**, 6127–6134.
- 8 Y. Usami, K. Mizuki, H. Ichikawa and A. Arimoto, *Tetrahedron: Asymmetry*, 2008, **19**, 1460–1463.
- 9 T. J. Donohoe, K. Blades, M. Helliwell, M. J. Warning and N. J. Newcombe, *Tetrahedron Lett.*, 1998, **39**, 8755–8758.
- 10 Y. Usami, H. Ichikawa and M. Arimoto, *Int. J. Mol. Sci.*, 2008, **9**, 401–421.
- 11 Y. Usami and A. Numata, *Chem. Pharm. Bull.*, 2004, **52**, 1125–1129.
- 12 H. Okamura, Y. Nakamura, K. Morishige, R. Ohura, H. Shimizu, T. Iwakawa, M. Nakatani, *Proceedings of the 40th Tennen Yuki Kagobutsu Toronkai Koen Yoshisyu*, 1988, Fukuoka, Japan, pp. 187–192.
- 13 J. Garcia, Ruano, J. Lopez-Cantarero, A. M. Martin, Castro, H. Adams and J. H. Rodriguez, Ramos, *Phosphorus, Sulfur and Silicon and the Related Elements*, 2005, **180**, 1493–1494.
- 14 G. Ulibarri, W. Nadler, T. Skrydstrup, H. Audrain, A. Chianori, C. Riche and A. A. Grierson, *J. Org. Chem.*, 1995, **60**, 2753–2761.
- 15 β,γ -Unsaturated enone **13** was so unstable that purification by silica gel chromatography afforded a complex mixture including more stable **16** shown in Fig. 3 formed by double bond migration. Therefore, other oxidizing agents could not be used in the preparation of **13**. Dess–Martin oxidation at 40 °C to accelerate the reaction resulted in the formation of **16** and aromatized product **17**. **Methyl (4*R*,5*R*)-4,5-cyclohexylidene-2-methoxy-3-oxo-1-cyclohexene-1-carboxylate 16**: Colorless crystals (CH₂Cl₂); mp 60–62 °C; $[\alpha]_D^{25}$ –1.8 (*c* 0.085 in CHCl₃); IR ν_{\max} (KBr)/cm^{–1} 1732 (C=O), 1693 (C=O), 1617 (C=C); ¹H-NMR (500 MHz; CDCl₃; Me₄Si) δ 1.35–1.70 (10H, m), 2.91 (1H, dd, *J* = 9.2, 4.8 Hz, H-6_A), 3.09 (1H, dd, *J* = 9.2, 2.1 Hz, H-6_B), 3.79 (3H, s, -OMe), 3.85 (3H, s, COOMe), 4.36 (1H, d, *J* = 5.3 Hz, H-4), 4.62 (1H, ddd, *J* = 5.3, 4.8, 2.1 Hz, H-5); ¹³C-NMR (125.6 MHz; CDCl₃; Me₄Si) δ 23.7 (t), 23.8 (t), 24.9 (t), 26.4 (t), 35.2 (t), 37.1 (t), 52.4 (q), 60.6 (q), 71.4 (d), 76.3 (d), 76.5 (d), 110.5 (s), 127.4 (s), 151.2 (s), 166.4 (s), 193.1 (s); EIMS *m/z* 296 (M⁺, 76%); HREIMS *m/z* calcd for C₁₅H₂₀O₆ (M)⁺ 296.1260, found 296.1259. **Methyl 3,4-dihydroxy-2-methoxybenzoate 17**: Yellow oil; IR ν_{\max} (liquid film)/cm^{–1} 3538 (OH), 1714 (C=O), 1604 (C=C); ¹H-NMR (CDCl₃) δ 3.89 (3H, s, -OMe), 3.93 (3H, s, COOMe), 5.87 (2H, br s, -OH), 6.74 (1H, d, *J* = 8.9 Hz), 7.46 (1H, d, *J* = 8.9 Hz); ¹³C-NMR (CDCl₃) δ 51.9 (q), 62.3 (q), 110.9 (d), 115.2 (s), 123.9 (d), 136.7 (s), 148.1 (s), 148.6 (s), 165.4 (s); EIMS *m/z* 198 (M⁺, 76%), 166 (M⁺–MeOH, 88%); HREIMS *m/z* calcd for C₉H₁₀O₅ (M)⁺ 198.0520, found 198.0524.
- 16 Y. Usami, K. Mizuki, K. Suzuki, E. Sakamoto, H. Ichikawa, M. Arimoto, *Proceedings of the 50th Tennen Yuki Kagobutsu Toronkai Koen Yoshisyu*, 2008, Fukuoka, Japan, pp. 547–552.