Journal of Medicinal Chemistry

Orally available soluble epoxide hydrolase/ phosphodiesterase 4 dual inhibitor treats inflammatory pain

Rene Blöcher, Karen M Wagner, Raghavender R Gopireddy, Todd R Harris, Hao Wu, Bogdan Barnych, Sung Hee Hwang, Yang K Xiang, Ewgenij Proschak, Christophe Morisseau, and Bruce D. Hammock *J. Med. Chem.*, Just Accepted Manuscript • DOI: 10.1021/acs.jmedchem.7b01804 • Publication Date (Web): 03 Apr 2018 Downloaded from http://pubs.acs.org on April 3, 2018

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Orally available soluble epoxide hydrolase/phosphodiesterase 4 dual inhibitor treats inflammatory pain

René Blöcher¹, Karen M. Wagner¹, Raghavender R. Gopireddy², Todd R. Harris¹, Hao Wu¹, Bogdan Barnych¹, Sung Hee Hwang¹, Yang K. Xiang², Ewgenij Proschak³, Christophe Morisseau¹ and Bruce D. Hammock^{1*}

¹Department of Entomology and Nematology and UC Davis Comprehensive Cancer Center, University of California Davis, One Shields Avenue, CA 95616, Davis, U.S.A. ²Department of Pharmacology, University of California Davis, One Shields Avenue, CA 95616, Davis, U.S.A., and VA Northern California Health Care System, CA 95655 Mather, U.S.A.

³Institute of Pharmaceutical Chemistry, Goethe-University Frankfurt, Max-von-Laue-Strasse 9, D-60438 Frankfurt am Main, Germany

Abstract

Inspired by previously discovered enhanced analgesic efficacy between soluble epoxide hydrolase (sEH) and phosphodiesterase 4 (PDE4) inhibitors, we designed, synthesized and characterized 21 novel sEH/PDE4 dual inhibitors. The best of these displayed good efficacy in in vitro assays. Further pharmacokinetic studies of a subset of 4 selected compounds led to the identification of a bioavailable dual inhibitor *N*-(4-methoxy-2-(trifluoromethyl)benzyl)-1-propionylpiperidine-4-carboxamide (**MPPA**). In a lipopolysaccharide induced inflammatory pain rat model, **MPPA** rapidly increased in the blood ($T_{max} = 30$ min; $C_{max} = 460$ nM) after oral administration of 3 mg/kg and reduced inflammatory pain with rapid onset of action correlating with blood levels over a time course of 4 hours. Additionally, **MPPA** does not alter self-motivated exploration of rats with inflammatory pain or the withdrawal latency in control rats.

ACS Paragon Plus Environment

Introduction

Multi-target ligands are designed to improve treatment of complex diseases¹. Target combinations are selected to improve the therapeutic impact on one or several physiological disorders. For example, phosphodiesterase 4 (PDE4) is the major enzyme degrading cAMP into 5-AMP and thus, PDE4 inhibition increases cAMP levels which subsequently leads to a down-regulation of multiple inflammatory mediators.² PDE4 is therefore a valuable target in the treatment of inflammatory diseases such as psoriasis, psoriatic arthritis (PsA),² chronic obstructive pulmonary diseases (COPD)³ and asthma.⁴ Additionally, cAMP levels in neurons are shown to be correlated with symptoms of neuronal disorders such as depression,⁵ schizophrenia⁶ and Alzheimer's disease.⁷ Nevertheless, the historical challenge of targeting PDE4 for therapy is the limited therapeutic index of PDE4 inhibitors due to vasculitis and severe dose limiting side effects such as headache, gastric hyper secretion emesis, and nausea.^{3,8–10} Separately from PDE4, soluble epoxide hydrolase (sEH) inhibition has beneficial physiological effects in animal models of inflammation,¹¹ pain,¹² hypertension¹³ and depression.¹⁴ The sEH metabolizes endogenous cytochrome P450s (CYP450) derived epoxyfatty acids (EpFAs) such as epoxyeicosatrienoic acids (EETs) from arachidonic acid.¹³ Inhibition of sEH elevates EpFA levels, which have anti-hypertensive, anti-inflammatory, and analgesic properties.⁶

In a previous study sEH inhibitors and PDE4 inhibitors were shown separately to increase plasma levels of natural anti-inflammatory analgesic epoxy fatty acids. The same study showed that co-administration of PDE4 and sEH inhibitors resulted in an enhanced analgesic effect compared to the individual treatments.¹⁵ This enhanced analgesia of the combination motivated us to develop a bioavailable sEH/PDE4 dual inhibitor. The development of polypharmacological agents requires the combination of at least two pharmacophores in one molecule. To achieve this goal we followed a medicinal chemistry driven approach,^{16–18}

where the essential structural moieties of known ligands were identified and combined in one novel molecular design, which subsequently was evaluated and re-designed in an iterative process. The pharmacokinetics (PK) and biological properties of the best compounds were evaluated in an animal model. In the future these tools will enable us to investigate the benefits of the dual ligand strategy in analgesia. ^{3,8–10}

Results

Dual ligand design and structural development

The design of dual modulators requires a combination of the pharmacophores from two inhibitors into one novel molecule. The choice of template inhibitors is essential, and was based on potency, water solubility, bioavailability and accessibility of space surrounding the inhibitors within the target binding-pocket. The initial dual inhibitor design was inspired by the sEH inhibitor GSK 2256294 and the PDE4 inhibitor Rolipram (Scheme 1). GSK 2256294 was chosen due to its subnanomolar potency on recombinant human and rat sEH and its successful phase I clinical trial.^{19,20} Rolipram was selected based on its high *in vivo* efficacy against PDE4, and its previous usage in combination with a sEH inhibitor against pain.¹⁵ The efficacy of PDE4 inhibitors, such as in Roflumilast, that have more bulky functional groups in place of a pyrrolidone, as well as cocrystal structures of PDE4 with the dialkoxyphenyl family of inhibitors²¹ suggest that the PDE4 active site is spacious enough to accommodate larger substituents on the dialkoxy phenyl ring of Rolipram. Therefore, the 2-, 4-substituted benzyl amide moiety was abstracted from GSK 2256294 and combined with the 3-(cyclopentyloxy)-4-methoxybenzene moiety from Rolipram, leading to the 1st dual inhibitor design (Scheme 1). Within this initial dual inhibitor design, the 2-trifluoromethyl-4-methoxy substitution of the benzyl amide (R1, R2 in Scheme 1) was evaluated by replacement. Interestingly, the initial design has a core structure similar to Apremilast, an FDA approved PDE4 inhibitor (PDE4I) (in red, Scheme 1). This similarity inspired the second dual inhibitor design, in which the 3(cyclopentyloxy)-4-methoxy substitution of the central benzene moiety was evaluated by replacement. After discarding the 3-(cyclopentyloxy)-4-methoxy benzene substitution, the derivatives maintained PDE4 inhibition. This finding led to the assumption that the 4-methoxy benzyl amide moiety might be sufficient as a PDE4I pharmacophore. Therefore, structural fragments were introduced in the design improving the pharmacokinetic properties of the dual inhibitors. TPPU is an in house potent sEHI with good bioavailability.²² To increase structural diversity and improve bioavailability the 1-(piperidin-1-yl)propan-1-one fragment of TPPU was combined with the 2-trifluoro-4-methoxy benzyl amide moiety of the piperidine fragment were evaluated within this structural class.

Synthesis

Scheme 2 shows the synthetic pathways toward sEH/PDE4 dual inhibitors. A central method was the amide synthesis. In general a benzoic acid derivative or a piperidine carboxylic acid derivative was activated with 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide and a catalytic amount of 4-dimethylaminopyridine under dry conditions, followed by the addition of a benzylamine derivative to produce compounds 1-11, 14 and 19-21.¹⁶ Benzoic acids used for the synthesis of compounds 14 and 19-20 were prepared as described below. To introduce the 2-diflurormethoxy substitution, methyl 4-hydroxy benzoate and sodium chloro difluoro acetate were reacted under basic conditions and gave, after decarboxylation, the 4-(difluoromethoxy)benzoic acid methyl ester (12).²³ After ester hydrolysis the substituted benzoic acid (13) was used for the final amide synthesis step. For the preparation of the 3-(cyclopentyloxy)benzoic acid (17) and 3-(cyclopentyloxy)-4-fluorobenzoic acid (18), 4-fluoro substituted and non-substituted 3-hydroxy methyl benzoates were reacted with cyclopentyl bromide under basic conditions in presence of a catalytic amount of potassium iodide to generate the cyclopentyloxy substituted intermediates 15 and 16.⁴ Again after ester hydrolysis

the benzoic acid derivatives **17** and **18** were used for the final amide coupling step. The synthesis of the piperidine containing inhibitors started with preparation of the intermediate **21**. Deprotection of the amine was performed with trifluoroacetic acid, delivering **22**.²⁴ Two methods were used for the following final synthetic step, each depending on the required substitution. To introduce the amide substitution, appropriate acid chlorides were purchased or generated with oxalylchloride and further reacted with **22** under dry basic conditions to produce **23-25**.²⁵ To synthesize the ethyl and propyl substituted piperidine derivatives (**26**, **27**), **22** was reacted with ethyl iodide or propyl iodide under basic conditions.

Inhibition of sEH and PDE4 activities

The *in vitro* profile of the initial dual inhibitor **1** showed an IC₅₀ on the human sEH at a subnanomolar concentration ($0.6 \pm 0.1 \text{ nM}$), a cAMP increase of more than 200% compared to Rolipram at 1 µM and a moderate water solubility of 37.5 µM. Replacing the 2-trifluoromethyl-4-methoxy benzyl substitution by simpler 2-trifluoromethyl and 2-methyl benzyl substituents resulted in a significant drop in potency on both targets, while the moderate water solubility stayed constant.

Within the second structural class (Group 2) of dual sEH/PDE4 inhibitors the substitution pattern at the central benzene moiety was investigated. The first compound of this class, **4**, contained a non-substituted central benzene fragment, resulting in a slight decrease of potency on both targets. Nevertheless, the sEH IC₅₀ was in the low one digit nanomolar range and the cAMP increase was $160 \pm 12\%$ of Rolipram at 1 μ M, while the water solubility improved up to 100 μ M. Introduction of the fluorine substitution at position 4 of the benzene ring (**5**) only affected the PDE4 potency by limiting the cAMP increase to the level of Rolipram at 1 μ M. Introduction of the *N*-acetamide substitution motif of the PDE4 inhibitor Apremilast in the *meta* position of the benzene moiety did not improve the potency of the compound (**6**) compared to the non-substituted inhibitor **4**. Shifting the acetamide substitution in the *para* position (10), elongating the substituent to propionamide (7) or introducing an additional methyl group in *para* position (8) decreased inhibition on both targets as well as water solubility. Thus, the Apremilast inspired design did not improve the dual inhibitors. A para methoxy benzene substitution (11) also did not improve target inhibition compared to the nonsubstituted inhibitor (4), while the para diffuoromethoxy benzene substituted inhibitor 14 even decreased water solubility to 10 μ M. The *meta* cyclopentoxy benzene substitution in dual inhibitor 19 improved the sEH inhibition compared to the non-substituted inhibitor 4, but water solubility decreased by one order of magnitude and cAMP increase relative to Rolipram at 1 µM slightly decreased too. For inhibitor 20, the meta cyclopentyloxy benzene substitution was combined with a *para* fluoro benzene substituent in order to increase metabolic stability. The additional *para* fluoro substituent did not change sEH inhibition (IC_{50} 0.4 nM), improved the cAMP increase relative to Rolipram at 1 μ M and restored the water solubility to 100 μ M. The third structural class (Group 3) of dual sEH/PDE4 inhibitors enfolds in different Nsubstitution of the piperidine moiety. The dual inhibitor 22 contains a non-substituted piperidine, which led to a significant loss in sEH inhibitory potency (IC₅₀ 370 ± 170 nM), a higher cAMP increase than Rolipram at 1 µM and an improved water solubility of 1 mM. Ethyl and propyl substituted modulators (26, 27) similarly had low sEH inhibitory potencies and high water solubility, but only 60% cAMP increase compared to Rolipram at 1 μ M. Dual inhibitor 23/MPPA with a propionamide piperidine substitution showed a low, one digit nanomolar sEH IC₅₀, 200% cAMP increase compared to Rolipram at 1 μ M and 100 μ M water solubility. Increasing the alkyl chain of the N-substitution to butyramide (24) and secbutylamide (25) decreased sEH inhibition and even more dramatically the water solubility. The *in vitro* screening process shown in **Table 1**, revealed 4 compounds with both adequate

potency on both targets and suitable water solubility. To better characterize those 4 compounds (1, 19, 20 and 23/MPPA), the melting point and IC_{50} value for human and rat recombinant enzymes were measured and compared to the single target model ligands (TPPU

and Rolipram). Previous studies have shown a difference in potency of sEH inhibition comparing human and rat sEH.²⁶ **Table 2** shows a drop in potency on the rsEH compared to hsEH for compound 1, 19 and 20 by 3 fold or more. 23/MPPA demonstrated a comparatively much lower potency (75-fold) on the rsEH than on the hsEH.

More than 20 PDE4 isoforms are known today^{27,30}, we decided therefore to use a cell based activity assay for our screening. This provides the benefit of a broad PDE4 evaluation and a better estimate towards the later *in vivo* effect of our compound. We additionally tried to evaluate one of our most potent compounds (**23**) on recombinant PDE4 enzyme and were therefore forced to pick one out of 20 isoforms. We chose PDE4B1, but did sadly not find inhibition up 10 μ M. Out of commercial reasons we did not screen the remaining 19 PDE4 isoforms.

Pharmacokinetics

To identify the optimal candidate for use *in vivo*, the pharmacokinetics of 4 dual compounds were determined with oral administration by cassette dosing. Compounds **1**, **19**, **20** and **23/MPPA** were administered at 0.3 mg/kg by oral gavage to 4 rats. Compound **1** was detected at 0.5 h with a 3 ± 2.2 nM blood concentration, while compound **20** was not observed above the limit of detection in the blood (**Table S2**). Compound **19** did not increase over single digit nM blood concentration over 12 h. **23/MPPA** reached the highest blood concentration of 60 ± 13 nM and was observed in blood for several hours. Therefore, **23/MPPA** having the best observed pharmacokinetics of the *in vivo* cassette compounds was chosen for further *in vivo* evaluated for PK parameters (**Table S3**). A 3 mg/kg dose that demonstrated *in vivo* efficacy resulted in a 460 \pm 16 nM blood concentration (n=3) at 0.5 h and remained in the blood several hours. The blood concentration of the 3 mg/kg dose is 3-fold above the rat sEH IC₅₀ and stays above the inhibitory concentration for about 2 h. Single administration of 30 and

100 mg/kg (each n=1) dosed to rats resulted in concentrations reaching 1,060 nM and 1,730 nM respectively. Thus, **MPPA** is absorbed quickly after oral administration and therefore has potential as a rapid acting pharmaceutical agent.

In vivo Efficacy

The efficacy of **MPPA** was evaluated in an LPS induced inflammatory pain model. For this assay the compound or vehicle control were administered by oral gavage followed immediately by an intraplantar injection of lipopolysaccharide (LPS) to one hindpaw and measured over a time course for their thermal withdrawal latency (Figure 1). Baseline scores were assessed before the begin of the assay and normalized to 100%. The LPS results in lower than baseline scores that continue to decrease over the first several hours indicating a painful state. Oral administration of 3 mg/kg MPPA resulted in significantly increased thermal withdrawal latencies (interpreted as pain relief) compared to the vehicle control. The rapid onset of anti-inflammatory analgesia at 30 min correlates directly with the increase of MPPA in the blood ($T_{max} = 30$ min) and decreases with the clearance from the blood after 4 hours. Interestingly, though hypoalgesia is apparent at the initial time point in the LPS assay, MPPA did not alter nociceptive thresholds in naïve animals. The dual inhibitor MPPA had no effect on withdrawal latency in naïve animals (Figure S4) while the PDE4 inhibitor Rolipram at a similar dose (3 mg/kg) significantly increased withdrawal latency. This effect of Rolipram in the absence of enhanced pain perception suggests a central nervous system activity. In addition to efficacy testing, we also observed spontaneous locomotion in the treated animals. Alteration of a normal pain response and sedation can be serious side effects. In LPS treated rats, MPPA administration did not decrease exploration in an open field assay compared to the vehicle. This contrasted with Rolipram which altered locomotion in the open field assay (Figure S5). This may indicate that the sEH/PDE4 dual inhibitor MPPA does not have effects similar to Rolipram. However, future investigations of **MPPA** in causing known side effects of classical PDE4Is such as vasculitis, headache, emesis and nausea will be required.

Discussion

In this study, 21 dual sEH/PDE4 inhibitors were synthesized and their in vitro potencies were assessed. The dual ligands with the most potential from each Group (1-3) were selected based on potency on both targets and water solubility. The pharmacokinetic properties of these selected compounds (1, 19, 20, MPPA) revealed good blood concentrations of 1-(piperidin-1yl)propan-1-one fragment, which was anticipated in the design process. In comparison, the cyclopentyloxy substituted benzyl fragments resulted in lower blood concentrations. Surprisingly, the addition of fluorine on the cyclopentyloxy benzyl fragment diminished the lowered these even further. MPPA, containing the 1-(piperidin-1-yl)propan-1-one fragment, was chosen out of the PK cassette and further investigated. The dose escalation revealed a fast absorption of MPPA combined with a rapid clearance. At a 3 mg/kg dose, the blood concentration of MPPA in rats reached levels above the IC₅₀ of both targets but deceased substantially after 4 hours. The efficacy evaluation showed a direct correlation between compound blood concentration and analgesic efficacy. Therefore, a future goal of this study is to identify labile positions in the structure of MPPA and substitute those with more metabolically stable moieties while maintaining potency on the targets. Finally, MPPA did not alter nociceptive responses in naïve rats compared to Rolipram. Nevertheless the dual inhibitor developed will need to be further investigated towards major potential side effects known from PDE inhibition to ensure the safety of the compound.

Conclusion

In conclusion, this study presents a proof of concept for the sEH/PDE4 dual inhibitor approach on analgesia and therefore adds to the concept of previous poly-pharmacological

tools targeting sEH in combination with other enzymes.^{16,18,31} Nevertheless, several experiments will have to be performed in the future to clarify the therapeutic improvements of the dual inhibitor. Further *in vivo* experiments, including ferret or other emetic models are necessary to define the side effects caused by the dual inhibitor in comparison to classical PDEIs.

Despite this, the dual ligand **MPPA** opens the path to future investigations of sEH/PDE4 target interactions and the crosstalk between cAMP and EpFA¹⁵. In addition, **MPPA** has the potential to impact further pathological conditions in addition to inflammatory pain, such as depression and COPD.^{3,5,14,20} The efficacy was limited by the rapid elimination of the compound suggesting that the next generation design of the dual inhibitor should focus on improving metabolic stability, while maintaining the rapid absorption and activity. Still, **MPPA** represents a new potential approach for pain management as well as a tool for investigating the biology of pain.

EXPERIMENTAL SECTION

Chemistry

General. All reagents and solvents were purchased from commercial suppliers and were used directly without further purification. All reactions were carried out at room temperature unless otherwise specified. Reactions were monitored by thin-layer chromatography (TLC) on Merck F_{254} silica gel 60 aluminum sheets, and spots were revealed with UV light (254 mm), potassium permanganate or Ninhydrin stains. ¹H NMR spectra were recorded on a 400 MHz Bruker Avance III HD Nanobay Spectrometer with deuterated chloroform (CDCl₃; δ = 7.24 ppm) or deuterated dimethyl sulfoxide (DMSO-*d*₆) containing TMS as internal standard. ¹³C NMR spectra were recorded on a Bruker Avance III HD Nanobay spectrometer at 100 MHz. The percent purity of the inhibitors reported in this manuscript were determined by HPLC-UV using Agilent 1200 series HPLC system equipped with Phenomenex Luna2 C18 reverse phase

column (C18, 4.6 mm \times 150 mm, 5 μ m) coupled with Agilent G1314 UV–vis detector (detection at 200, 210, 254 and 360) with solvent gradient acetonitrile/water 20 to 95% over 15 min and they are expressed as percent total OD at 254 and 360 nm. The purity of all final compounds was above 95%. HRMS spectra were recorded on Thermo Electron LTQ-Orbitrap XL Hybrid MS in ESI mode.

General synthetic procedure A for compounds 1-11, 14 and 19-21 shown on the example 3-(cyclopentyloxy)-4-methoxy-N-(4-methoxy-2-(trifluoromethyl)benzyl) (1). 0.2 g (0.85 mmol) of 3-(cyclopentyloxy)-4-methoxybenzoic acid, 160 mg (1 mmol) of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide and 20 mg (0.17 mmol) 4-(dimethylamino)pyridine were dissolved in 10 ml dichloromethane and stirred under a nitrogen atmosphere for 1 h. Subsequently, 160 μ l (0.93 mmol) (4-methoxy-2-(trifluoromethyl)phenyl)methanamine was added to the mixture. The reaction was further stirred for 12 h. After completion of the reaction, dichloromethane was removed under reduced pressure. The residue was dissolved in 20 ml of ethyl acetate and washed with aqueous sodium hydroxide solution (2 M, 3 x 20 mL), aqueous hydrochloric acid solution (2 M, 3 x 20 mL) and brine (20 mL). The organic solvent was dried over magnesium sulfate and removed under reduced pressure. The crude product was purified by recrystallization from ethyl acetate and hexane mixture. The pure product remained as a white solid.

General synthetic method B to produce compounds 13, 17 and 18 shown with the example 4-(difluoromethoxy)benzoic acid (13). 0.47 g (2.3 mmol) methyl 4- (difluoromethoxy)benzoate (12) and 0.47 g (12 mmol) sodium hydroxide were stirred in a mixture of 10 ml of tetrahydrofuran, 10 ml of methanol and 5 ml of water under a nitrogen atmosphere at 40 °C for 12 h. The organic solvents were removed under reduced pressure and the remaining aqueous phase was acidified through dropwise addition of concentrated hydrochloric acid solution. The precipitated product was filtered, dried and used in the next step without further purification.

General synthetic method C for compounds 15 and 16, shown with the synthesis of methyl 3- (cyclopentyloxy)benzoate (15). 1 g (6.6 mmol) of methyl 3-hydroxybenzoate, 22 mg (0.13 mmol) of potassium iodide and 1.4 g (9.9 mmol) of potassium carbonate are stirred in 6.5 ml of dimethylformamide at 65 °C. Subsequently 0.92 ml (8.6 mmol) of bromocyclopentane was added dropwise. The reaction was further stirred for 21 h, cooled down to rt and diluted with 50 ml dichloromethane, washed with aqueous sodium hydroxide solution (2 M, 3 x 50 mL) and brine (50 mL). The organic solvent was dried over magnesium sulfate and evaporated under reduced pressure. The product was used in the next step without further purification.

General synthetic method D to produce compounds 23/MPPA and 24, shown on the example *N*-(4-methoxy-2-(trifluoromethyl)benzyl)-1-propionylpiperidine-4-carboxamide 23/MPPA. 28 μ L (0.38 mmol) of propionic acid and 40 μ L (0.47 mmol) of oxalyl chloride were stirred in 1 mL of dry dichloromethane under a nitrogen atmosphere for 1 h, to generate propionyl chloride. In a separate flask 0.1 g (0.32 mmol) of *N*-(4-methoxy-2-(trifluoromethyl)benzyl)piperidine-4-carboxamide 22 and 80 μ L (0.5 mmol) triethylamine were stirred in 2 ml of dry dichloromethane at 0 °C under a nitrogen atmosphere. The produced propionyl chloride solution was now added slowly to the amine containing reaction flask and the whole mixture was stirred for 4 h. The reaction was diluted with 7 mL of dichloromethane, washed with hydrochloric acid aqueous solution (2 M, 3 x 10 mL), sodium hydroxide aqueous solution (2 M, 3 x 10 mL) and brine (10 mL). The organic layer was dried over magnesium sulfate and evaporated under reduced pressure. The remaining crude material was purified by flash column chromatography (acetone/ethyl acetate = 50:50).

General synthetic method E to produce compounds 26 and 27, illustrated with the synthesis of 1-ethyl-N-(4-methoxy-2-(trifluoromethyl)benzyl)piperidine-4-carboxamide 26. 0.15 g (0.48 mmol) of N-(4-methoxy-2-(trifluoromethyl)benzyl)piperidine-4-carboxamide 22, 0.13 g (0.95 mmol) of potassium carbonate and 38 μ L (0.48 mmol) ethyl iodide were

Page 13 of 36

stirred in 3.2 mL of dimethylformamide for 24 h. After the reaction was completed, the pure product was precipitated with the addition of 2 M sodium hydroxide aqueous solution.

3-(Cyclopentyloxy)-4-methoxy-*N***-(4-methoxy-2-(trifluoromethyl)benzyl) (1).** Yield: 0.18 g (50%). ¹H NMR (400 MHz, Chloroform-*d*) δ 7.86 – 6.25 (m, 6H), 4.89 – 4.81 (m, 1H), 4.74 (d, *J* = 6.1, 1.3 Hz, 2H), 3.89 (s, 3H), 3.85 (s, 3H), 2.33 – 1.20 (m, 8H); ¹³C-NMR (DMSO-*d*₆) δ 166.7, 157.1, 151.3, 147, 128.2, 128, 127.8, 126.9, 126.4, 124.1, 121.3, 118.2, 115.1, 112.6, 109.3, 80.1, 55.2, 54.1, 42.7, 31.7, 30.6, 24.1, 23.7; HRMS: measured [M+Cl]⁻ 374.1523 (calculated: 374.1521).

3-(Cyclopentyloxy)-4-methoxy-*N***-(2-(trifluoromethyl)benzyl)benzamide (2)**. Yield: 0.16 (49%). ¹H NMR (400 MHz, Chloroform-*d*) δ 7.90 – 6.47 (m, 7H), 4.88 – 4.84 (m, 1H), 4.82 (d, *J* = 6.4 Hz, 2H), 3.89 (s, 3H), 2.08 – 1.50 (m, 8H); ¹³C-NMR (DMSO-*d*₆) δ 168.3, 156.1, 148.3, 147.5, 127.2, 127, 126.9, 126.7, 126.5, 123.4, 120.3, 119.2, 116.3, 112.1, 108.3, 79.1, 53.1, 41.7, 33.7, 31.4, 25.1, 23.1; HRMS: measured [M+Cl]⁻ 428.1239 (calculated: 428,1237). **3-(Cyclopentyloxy)-4-methoxy-***N***-(2-methylbenzyl)benzamide (3)**. Yield: 0.15 (62%). ¹H NMR (400 MHz, Chloroform-*d*) δ 7.63 – 6.07 (m, 7H), 4.93 – 4.81 (m, 1H), 4.65 (d, *J* = 5.4 Hz, 2H), 3.89 (s, 3H), 2.40 (s, 3H), 2.08 – 1.54 (m, 8H); ¹³C-NMR (DMSO-*d*₆) δ 167.3, 155.1, 147.3, 146.5, 128.1, 128, 127.9, 125.7, 120.8, 119.1, 117.3, 113.1, 109.3, 79.5, 52.1, 42.7, 31.7, 30.4, 24.1, 22.1, 19.1; HRMS: measured [M+Cl]⁻ 458.1344 (calculated: 458.1342).

N-(4-Methoxy-2-(trifluoromethyl)benzyl)benzamide (4). Yield 0.11 g (34%). ¹H NMR (400 MHz, Chloroform-*d*) δ 7.36 (m, 8H), 6.40 (s, 1H), 3.7 (s, 3H); ¹³C-NMR (DMSO-*d*₆) δ 166.1, 158.1, 150.2, 146, 127.8, 127.6, 127.3, 126.9, 126.1, 123.1, 120.3, 119.2, 116.1, 112.6, 109.1, 53.1, 40.7, HRMS: measured [M+HCOO]⁻ 354.0950 (calculated: 354.0953).

4-Fluoro-*N***-(4-methoxy-2-(trifluoromethyl)benzyl)benzamide (5)**. Yield: 0.21 g (66%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.05 (t, *J* = 5.8 Hz, 1H), 8.26 – 6.91 (m, 7H), 4.58 (d, *J* = 5.7 Hz, 2H), 3.82 (s, 3H); ¹³C-NMR (DMSO-*d*₆) δ 168.1, 155.1, 148.2, 147, 130.8, 129.6,

129.7,128.9, 128.1, 126.1, 125.1, 120.1, 118.2, 117.1, 113.6, 107.1, 51.1, 40.5, HRMS: measured [M+HCOO]⁻ 372.0856 (calculated: 372.0854).

3-Acetamido-*N***-(4-methoxy-2-(trifluoromethyl)benzyl)benzamide (6)**. Yield: 0.23 mg (56%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.05 (s, 1H), 8.94 (t, *J* = 5.8 Hz, 1H), 8.24 – 6.44 (m, 7H), 4.52 (d, *J* = 5.6 Hz, 2H), 3.77 (s, 3H), 2.00 (s, 3H); ¹³C-NMR (DMSO-*d*₆) δ 166.9, 159.1, 151.2, 145, 141.3, 128.6, 127.4, 127.3, 127.1, 126. 8, 122.1, 121.3, 118.3, 117.1, 112.6, 109.7, 55.8, 42.6; HRMS: measured [M-H]⁻ 365.1114 (calculated: 365.1113).

N-(4-Methoxy-2-(trifluoromethyl)benzyl)-3-propionamidobenzamide (7). Yield: 0.14 (41%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.08 (s, 1H), 9.04 (t, *J* = 5.9 Hz, 1H), 8.13 – 7.01 (m, 7H), 4.63 (d, *J* = 5.6 Hz, 2H), 3.88 (s, 3H), 2.40 (q, *J* = 7.5 Hz, 2H), 1.15 (t, *J* = 7.5 Hz, 3H); ¹³C-NMR (DMSO-*d*₆) δ 167.9, 155.1, 150.2, 146, 142.3, 129.6, 128.5, 128.3, 128.2, 128, 121.1, 119.3, 118.1, 116.1, 110.6, 109.7, 56.8, 41.6, 10.1; HRMS: measured [M+HCOO]⁻ 425.1320 (calculated: 425.1321).

3-Acetamido-*N*-(**4-methoxy-2-(trifluoromethyl)benzyl)-4-methylbenzamide** (**8**). Yield: 0.1 g (29%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.43 (s, 1H), 8.95 (t, *J* = 5.8 Hz, 1H), 8.32 – 6.80 (m, 6H), 4.57 (d, *J* = 5.6 Hz, 2H), 3.82 (s, 3H), 2.25 (s, 3H), 2.08 (s, *J* = 2.7 Hz, 3H). ¹³C-NMR (DMSO-*d*₆) δ 166, 158.1, 152.2, 146, 142.3, 128.3, 128.1, 128, 127.8, 121.1, 120.3, 119.3, 118.1, 112.7, 109.9, 56.8, 40.6, 17.3; HRMS: measured [M+HCOO]⁻ 435.1321 (calculated: 435.1321).

N-(4-Methoxy-2-(trifluoromethyl)benzyl)-2-oxoindoline-6-carboxamide (9). Yield: 0.14 (44%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.56 (s, 1H), 9.00 (t, *J* = 5.8 Hz, 1H), 7.79 – 6.95 (m, 6H), 4.57 (d, *J* = 5.6 Hz, 2H), 3.82 (s, 3H), 3.55 (s, 2H); ¹³C-NMR (DMSO-*d*₆) δ 176.7, 167.8, 155.1, 147.2, 141, 127, 126.9, 126.8, 126.6, 126.1, 124, 123.5, 120.1, 119.7, 118.1, 113.6, 109.6, 52.1, 42.7, 36.5; HRMS: measured [M+HCOO]⁻ 409.1008 (calculated: 409.1007).

4-Acetamido-N-(4-methoxy-2-(trifluoromethyl)benzyl)benzamide (10). Yield: 0.16 (50%).
¹H NMR (400 MHz, DMSO-d₆) δ 10.23 (s, 1H), 8.95 (t, J = 5.8 Hz, 1H), 8.14 – 7.11 (m, 7H),
4.63 (d, J = 5.6 Hz, 2H), 3.87 (s, 3H), 2.14 (s, 3H); ¹³C-NMR (DMSO-d₆) δ 166.3, 157.1,
152.2, 147, 142.3, 128.6, 128.5, 128.3, 127.9, 123.1, 122.3, 119.3, 118.1, 112.8, 109.3, 56.8,
41.6; HRMS: measured [M-H]⁻ 365.1111 (calculated: 365.1113).

4-Methoxy-*N***-(4-methoxy-2-(trifluoromethyl)benzyl)benzamide (11)**. Yield: 0.16 g (48%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.88 (t, *J* = 5.8 Hz, 1H), 8.17 – 7.73 (m, 2H), 7.57 – 7.33 (m, 1H), 7.32 – 7.14 (m, 2H), 7.09 – 6.92 (m, 2H), 4.57 (d, *J* = 5.7 Hz, 2H), 3.83 (s, 3H), 3.82 (s, 3H); ¹³C-NMR (DMSO-*d*₆) δ 167.7, 156.1, 150.3, 148, 129.2, 128, 127.9, 127.7, 127.4, 123.1, 120.3, 119.2, 116.1, 113.6, 109.1, 56.2, 53.1, 41.7; HRMS: measured [M+HCOO]⁻ 384.1055 (calculated: 384.1057).

Methyl 4-(difluoromethoxy)benzoate (12). 1 g (6.6 mmol) methyl 4-hydroxybenzoate, 4.3 g (13 mmol) cesium carbonate and 2.5 g (16 mmol) sodium chloro difluoro acetate were dissolved in 48 ml dimethylformamide and 7 ml water under a nitrogen atmosphere. The mixture was first stirred for 15 min at rt and subsequently heated to 100 °C for 2 h. After completion of the reaction, the mixture was diluted with 50 mL of water and the product extracted with 100 mL of dichloromethane. The organic phase was dried over magnesium sulfate and evaporated under reduced pressure. The product was used in the next step without further purification. Yield: 0.47 g (35%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.02 (d, *J* = 8.9 Hz, 2H), 7.40 (s, 1H), 7.30 (d, *J* = 8.8 Hz, 2H), 3.85 (s, 3H).

4-(Difluoromethoxy)benzoic acid (13). Yield: 0.18 (41%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.02 (s, 1H), 8.00 (d, *J* = 8.8 Hz, 2H), 7.39 (s, 1H), 7.32 – 7.25 (m, 2H).

4-(Difluoromethoxy)-*N***-(4-methoxy-2-(trifluoromethyl)benzyl)benzamide** (14). Yield: 0.11 g (37%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.05 (t, *J* = 5.7 Hz, 1H), 8.05 – 7.92 (m, 2H), 7.47 – 7.43 (m, 1H), 7.36 (s, 1H), 7.32 – 7.26 (m, 2H), 7.26 – 7.20 (m, 2H), 4.58 (d, *J* = 5.6 Hz, 2H), 3.82 (s, 3H); ¹³C-NMR (DMSO-*d*₆) δ 167.8, 167.3, 167.1, 166.9, 157.1, 152.3,

148.3, 128.2, 128.1, 128, 127.9, 127.6, 123.2, 121.3, 119.1, 117.1, 112.6, 109.1, 50.1, 42.7; HRMS: measured [M-H]⁻ 374.0813 (calculated: 374.0811).

Methyl 3- (cyclopentyloxy)benzoate (15) Yield: 1.1 g (76%). ¹H NMR (400 MHz, DMSO d_6) δ 7.85 – 7.03 (m, 4H), 4.87 (tt, J = 5.9, 2.5 Hz, 1H), 3.84 (s, 3H), 2.16 – 1.42 (m, 8H).

Methyl 3-(cyclopentyloxy)-4-fluorobenzoate (16). Yield: 1 g (70%). ¹H NMR (400 MHz,

DMSO-*d*₆) δ 8.00 – 6.91 (m, 3H), 5.18 – 4.68 (m, 1H), 3.85 (s, 3H), 2.20 – 1.37 (m, 8H).

3-(Cyclopentyloxy)benzoic acid (17). Yield: 0.4 g (85%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.95 (s, 1H), 7.92 – 6.80 (m, 4H), 5.02 – 4.59 (m, 1H), 2.12 – 1.48 (m, 8H).

3-(Cyclopentyloxy)-4-fluorobenzoic acid (18). Yield: 0.8 g (71%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.06 (s, 1H), 7.95 – 6.87 (m, 3H), 4.99 – 4.90 (m, 1H), 2.02 – 1.54 (m, 8H).

3-(Cyclopentyloxy)-*N***-(4-methoxy-2-(trifluoromethyl)benzyl)benzamide (19)**. Yield: 0.1 g (30%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.00 (t, *J* = 5.8 Hz, 1H), 7.79 – 6.89 (m, 7H), 4.96 – 4.81 (m, 1H), 4.57 (d, *J* = 5.7 Hz, 2H), 3.82 (s, 3H), 2.04 – 1.47 (m, 8H); ¹³C-NMR (DMSO-*d*₆) δ 167.1, 156.2, 151.1, 148, 128.4, 128.2, 128.1, 127.9, 127.2, 126.5, 123.9, 120.3, 119.2, 116.1, 113.6, 109.1, 80.2, 53.1, 41.7, 30.7, 30.6, 24.1, 23.7; HRMS: measured [M+HCOO]⁻ 438.1527 (calculated: 438.1528).

3-(Cyclopentyloxy)-4-fluoro-*N*-(**4**-methoxy-**2**-(trifluoromethyl)benzyl)benzamide (**20**). Yield: 0.1 (33%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.04 (t, *J* = 5.8 Hz, 1H), 7.82 – 7.09 (m, 7H), 4.95 (tt, *J* = 5.9, 2.4 Hz, 1H), 4.59 (d, *J* = 5.6 Hz, 2H), 3.82 (s, 3H), 2.12 – 1.50 (m, 8H);); ¹³C-NMR (DMSO-*d*₆) δ 166.9, 166.4; 155.2, 155.3, 152.1, 147, 128.5, 128.3, 128.2, 128, 127.5, 122.9, 120.1, 119.2, 117.1, 112.6, 109.7, 80.6, 52.7, 40.7, 30.6, 30, 24.3, 23.9; HRMS: measured [M-H]⁻ 410.1379 (calculated: 410.1377).

tert-Butyl 4-((4-methoxy-2-(trifluoromethyl)benzyl)carbamoyl)piperidine-1-carboxylate (21). Yield: 1.1 g (42%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.34 (t, *J* = 5.8 Hz, 1H), 7.69 – 6.86 (m, 3H), 4.35 (d, *J* = 5.6 Hz, 2H), 3.95 (d, *J* = 13.2 Hz, 2H), 3.82 (s, 3H), 2.71 (d, *J* = 24.0 Hz, 2H), 2.42 – 2.31 (m, 1H), 1.70 (d, *J* = 13.2 Hz, 2H), 1.45 (d, *J* = 4.2 Hz, 2H), 1.40 (s,

N-(4-Methoxy-2-(trifluoromethyl)benzyl)piperidine-4-carboxamide (22). 1.3 g (3 mmol) *tert*-Butyl 4-((4-methoxy-2-(trifluoromethyl)benzyl)carbamoyl)piperidine-1-carboxylate **21** and 2.8 mL (36 mmol) of trifluoroacetic acid were stirred in 16 mL of dichloromethane at 0 °C for 2 h. The organic phase was diluted with 4 mL dichloromethane and extracted with hydrochloric acid aqueous solution (10 %, 3 x 20 mL). The aqueous layer was basified through the addition of solid sodium hydroxide pellets. From the basic aqueous layer the product was extracted with ethyl acetate (3 x 60 mL). The organic phase was dried over magnesium sulfate and evaporated. The remaining product was used in the next step without further purification. Yield: 0.73 g (77%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.27 (t, *J* = 5.8 Hz, 1H), 7.81 – 6.84 (m, 3H), 4.34 (d, *J* = 5.7 Hz, 2H), 3.82 (s, 3H), 2.99 (dt, *J* = 12.3, 3.4 Hz, 2H), 2.66 – 2.45 (m, 3H), 1.76 – 1.32 (m, 4H); ¹³C-NMR (DMSO-*d*₆) δ 172.8, 158.7, 157.8, 128.4, 128.2, 128.1, 127.8, 125.3, 124.7, 123.9, 117.9, 109.1, 56.8, 44.7, 43.2, 41.7, 39.9; 29.4, 29.3; HRMS: measured [M+H]⁺ 317.1493 (calculated: 317.1492).

N-(4-Methoxy-2-(trifluoromethyl)benzyl)-1-propionylpiperidine-4-carboxamide

23/MPPA. Yield: 70 mg (63%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.34 (t, *J* = 5.8 Hz, 1H), 7.63 – 6.99 (m, 3H), 4.35 (d, *J* = 6.0 Hz, 2H), 3.82 (s, 3H), 3.10 – 2.88 (m, 2H), 2.73 – 2.63 (m, 3H), 2.39 – 2.23 (m, 2H), 1.88 – 1.64 (m, 2H), 1.34 – 1.19 (m, 2H), 0.98 (t, *J* = 7.4 Hz, 3H); ¹³C-NMR (DMSO-*d*₆) δ 176.9, 173.8, 156.5, 128.5, 128.2, 127.9, 127.8, 125.1, 124.8, 123.4, 118.7, 109.9, 56.9, 43.7, 42.2, 40.9, 38.9; 28.9, 28.8, 26.1, 10.2; HRMS: measured [M+FA]⁻ 417.1633 (calculated: 417.1633).

1-Butyryl-*N***-(4-methoxy-2-(trifluoromethyl)benzyl)piperidine-4-carboxamide** (24). Yield: 0.05 g (37%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.35 (t, *J* = 5.8 Hz, 1H), 7.53 – 7.04 (m, 3H), 4.35 (d, *J* = 5.9 Hz, 2H), 3.94 – 3.85 (m, 1H), 3.82 (s, 3H), 3.01 (t, *J* = 12.7 Hz, 2H),

2.71 – 2.55 (m, 1H), 2.28 (td, J = 7.3, 3.1 Hz, 2H), 1.73 (t, 2H), 1.59 – 1.44 (m, 3H), 1.42 – 1.27 (m, 2H), 0.89 (t, J = 7.4 Hz, 3H); ¹³C-NMR (DMSO- d_6) δ 176.7, 175.8, 155.5, 128.4, 128.2, 128, 127.9, 124.1, 124, 122.4, 119.7, 109.3, 54.9, 42.7, 41.2, 40.7, 37.9; 29.9, 28.8, 27.1, 19.2, 13.2; HRMS: measured [M-H]⁻ 385.0268 (calculated: 385.0267).

N-(4-Methoxy-2-(trifluoromethyl)benzyl)-1-(2-methylbutanoyl)piperidine-4-

carboxamide (25). 0.1 g (0.32 mmol) of *N*-(4-methoxy-2-(trifluoromethyl)benzyl)piperidine-4-carboxamide (**22**) and 44 μ L (0.32 mmol) triethylamine were stirred in 5 mL dry dichloromethane at 0 °C under a nitrogen atmosphere. Subsequently, 39 μ L (0.32 mmol) of 2methylbutanoyl chloride was added and the reaction mixture was stirred for 4 h. The reaction was diluted with 5 mL of dichloromethane and washed with hydrochloric acid aqueous solution (2 M, 3 x 10 mL), sodium hydroxide aqueous solution (2 M, 3 x 10 mL) and once with 10 mL brine. The organic phase was dried over magnesium sulfate and removed under reduced pressure. The crude material was purified by flash column chromatography (acetone/ethyl acetate = 50:50). Yield: 82 mg (65%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.37 (t, *J* = 5.8 Hz, 1H), 7.49 – 7.09 (m, 3H), 4.39 – 4.48 (m, 1H), 4.37 – 4.34 (m, 2H), 4.07 – 3.95 (m, 1H), 3.82 (s, 3H), 3.1 – 2.96 (m, 1H), 2.71 (h, *J* = 6.7 Hz, 1H), 2.58 (t, *J* = 11.9 Hz, 1H), 2.47 (dd, *J* = 11.5, 3.9 Hz, 1H), 1.83 – 1.7 (m, 2H), 1.62 – 1.21 (m, 2H), 0.97 (t, *J* = 7.7 Hz, 3H), 0.81 (q, *J* = 7.1 Hz, 3H); ¹³C-NMR (DMSO-*d*₆) δ 176.9, 175.7, 155.4, 128.3, 128.1, 128, 127.8, 125.2, 125.1, 123.4, 119.1, 109.2, 53.9, 42.5, 41, 40.2, 39.2, 37.9; 29.7, 27.5, 17.3, 10.2; HRMS: measured [M+HCOO] 445.1947 (calculated: 445.1949).

1-Ethyl-*N***-(4-methoxy-2-(trifluoromethyl)benzyl)piperidine-4-carboxamide (26).** Yield: 40 mg (24%). %). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.13 (t, *J* = 5.7 Hz, 1H), 7.66 – 6.43 (m, 3H), 4.12 (d, *J* = 5.7 Hz, 2H), 3.72 (s, 3H), 3.31 (dt, *J* = 12.1, 3.3 Hz, 2H), 2.67 – 2.48 (m, 3H), 1.77 – 1.42 (m, 4H), 3.02 – 2.98 (m, 2H), 0.95 – 1.16 (m, 3H); ¹³C-NMR (DMSO-*d*₆) δ 172.8, 158.7, 157.8, 128.2, 128.1, 127.8, 127.7, 125.3, 124.7, 123.9, 117.9, 109.1, 56.8, 49.9,

44.7, 43.2, 41.7, 39.9; 29.4, 29.3, 13.3; HRMS: measured [M+H]⁺ 345.1814 (calculated: 345.1815).

N-(4-Methoxy-2-(trifluoromethyl)benzyl)-1-propylpiperidine-4-carboxamide (27). Yield: 0.06 g (35%). ¹H NMR (400 MHz, DMSO- d_6) δ 8.9 (t, J = 5.7 Hz, 1H), 7.55 – 6.41 (m, 3H), 4.3 (d, J = 5.8 Hz, 2H), 3.75 (s, 3H), 3.21 (dt, J = 12.2, 2.9 Hz, 2H), 2.77 – 2.38 (m, 3H), 1.76 – 1.41 (m, 4H), 2.51 – 2.39 (m, 2H), 1.46 – 1.31 (m, 2H), 0.93 – 0.79 (m, 3H) ; ¹³C-NMR (DMSO- d_6) δ 173.8, 157.7, 156.8, 129.2, 128.6, 128.5, 128.3, 128.1, 125.7, 123.8, 117.6, 109.4, 59.3, 55.8, 47.9, 43.7, 42.2, 41.9, 39.7; 29.1, 21.2, 12.3; HRMS: measured [M+H]⁺ 359.1970 (calculated: 359.1971).

Biochemistry

sEH activity assay. The assay was performed as previously described.³² All hsEH and rsEH IC₅₀ values were determined by a fluorescence-based assay system in a 96-well format. Non-fluorescent MNPC cyano(6-methoxy-naphthalen-2-yl)methyl trans-[(3-phenyloxiran-2-yl)methyl] carbonate was used as the assay substrate at a concentration of 5 μ M. This substrate is hydrolyzed by the sEH to the fluorescent 6-methoxynaphthaldehyde.³² The formation of the product was measured ($\lambda_{em} = 330$ nm, $\lambda_{ex} = 465$ nm) by a Molecular Device M-2 plate reader. All measurements were performed in triplicate.

PDE4 activity cell assay.

HEK293 cell culture and transfection.

HEK293 cells were cultured in DMEM (Gibco life technologies) containing 10% FBS, 100 units/ml penicillin-streptomycin, and 2 mM L-glutamine. Cell transfections were performed with the PIE Transfection reagent (Qiagen) according to manufacturer's instructions.

Förster Resonance Energy Transfer (FRET) Measurements to track PKA activity. HEK293 cells were transfected with PM-AKAR3 plasmid DNA (a plasma membranetargeted PKA activity reporter) according to methods described previously.²⁸ Images were acquired using a Zeiss AXIO inverted fluorescence microscope (Carl Zeiss micrscopy, LLC, Thornwood, NY) with a 40X oil-emersion objective lens and a charge-coupled device camera controlled by Metafluor software (Molecular Devices, Sunnyvale, CA). FRET was recorded by exciting the donor fluorophore at 430-455 nm and measuring emission fluorescence with two filters (475DF40 for cyan and 535DF25 for yellow). Images were subjected to background subtraction, and were acquired every 30 seconds with exposure time of 200 ms. The donor/acceptor FRET ratio was calculated and normalized to the ratio value of baseline. The binding of cAMP to AKAR3 increases YFP/CFP FRET ratio with increasing concentration.^{27,33}

PDE4B1 activity assay. PDE4B1 IC₅₀ for rat and human were conducted by BPS Bioscience according to standard procedures. MPPA in a range of $0.001 - 10 \mu$ M and Rolipram, as a reference compound, were assessed for IC50.

Water solubility approximation. Solutions of the compound under investigation were prepared in 0.1 M PBS buffer at pH 7.4 and 1% DMSO and placed in a 96-well transparent flat bottom microtiter plate. Precipitation of the compounds turbidity was measured at 650 nm using a microplate reader (Infinite M200).

Pharmacology and Behavior

Pharmacokinetic study in rat. All the animal experiments were performed according to the protocols approved by the Animal Use and Care Committee of University of California Davis. Male Sprague-Dawley rats (n=4, 8 weeks old, 250-300 g) were used in the pharmacokinetic study of sEH/PDE4 dual inhibitors. Inhibitors were dissolved in 100% polyethylene glycol 300 to form a clear solution. A cassette of four inhibitors (inhibitor 1, 19, 20 and 23; 0.3 mg/kg of each inhibitor, 0.9-1.2 mL per body weight) in solution was given by oral gavage. Whole blood (10 μ L) was collected with a pipette from the tail vein punctured by a lancet at 0, 0.25, 0.5, 1, 2, 4, 8, 12, 24, 48 and 72 hours after oral dosing with the inhibitor. Each blood

Page 21 of 36

Journal of Medicinal Chemistry

sample was immediately transferred to a tube containing 100 μ L of water with 0.1% EDTA and mixed and stored at -80°C until analysis. The blood samples were processed and sEH/PDE4 dual inhibitor concentrations determined according to the previously reported method by Liu et al.³⁴

Evaluation of analgesic efficacy in rat pain model. A thermal withdrawal latency (Hargreaves's) assay was performed as previously described.¹⁵ Thermal withdrawal thresholds (TWL) were determined before dosing to determine a baseline score. After baseline determination, 500 μ L of 100% PEG300 vehicle or 3 mg/kg **MPPA** were oral gavaged. Immediately post oral gavage, 50 μ l of a 0.2 μ g/ml solution of lipopolysaccharide (LPS) from *Escherichia coli* (0111:B4; Sigma Aldrich L2630) in saline was injected intraplantar in one hind paw (ipsilateral paw). The rats were then assessed for TWL over a time course of 4 hours (0.5, 1, 2 and 4 hrs). The ipsilateral TWLs were measured 5 times per rat per time point and scores are reported as an average of a group of rats (n=6-9).

Open field assay. Each rat was placed in an open top acrylic box with a floor area size of 40 cm x 40 cm and walls 30 cm high marked with a 16 square grid. The exploration of the box was observed over 2 min. The score represents the sum of the number of squares the rat passed with both of its hind paws plus the number of vertical rears. Individual rats are compared to themselves (pre- and post-treatment) to calculate the percent of baseline (pre-treatment) scores and were then averaged.

Statistics

Statistical analysis for the PDE4 cell assay was performed using GraphPad Prism 6 software (La Jolla, CA). Results are shown as mean \pm SEM and were analyzed by one-way ANOVA with post hoc Bonferroni's multiple compression test. Behavioral nociceptive assay data were analyzed using SigmaPlot 11.0 for Windows (Systat Software Inc., San Jose, CA). The applied statistical methods are reported in the figure legend with p values \leq 0.05 considered significant.

ACS Paragon Plus Environment

Supporting Information

The results for PDE4 screening value tables and figures, Pharmacokinetics table, Naïve control TWLs and open field assay are given in supporting information as well as the molecular formula strings. This material is available free of charge via the internet at http://pubs.acs.org.

Corresponding author: Bruce D. Hammock

Tel: 530-751-7519

e-mail: bdhammock@ucdavis.edu

Acknowledgements

This work was supported, in part, by National Institutes of Health grants: NIEHS R01 ES002710 and NIEHS/Superfund Research Program P42 ES004699 and Grants NIEHS T32ES007059, NIH 5T32DC008072-05 and 4T32HL086350-09 (to K.W.). A personal thanks is directed at the Services for International Students and Scholars (SISS) and within this organization to Simone Kueltz. EP thanks the German Research Foundation (DFG; Sachbeihilfe PR1405/2-2; Heisenberg-Professur PR1405/4-1; SFB 1039 Teilprojekt A07) for financial support.

Abbreviations

% of Bl – percent of baseline; 5-AMP – 5 adenosine monophosphate; cAMP – cyclic adenosine monophosphate; Cmax – the maximum (or peak) serum concentration that a drug achieves in a specified compartment; COPD – chronic obstructive pulmonary diseases; Ctrl – control; CYP450 – cytochrome P450; DHETs – dihydroxy eicosatrienoic acid; DMSO – Dimethylsulfoxide;; EETs – epoxyeicosatrienoic acids; ESI – Electrospray ionization; FBS –

Fetal bovine serum; FDA – Food and Drug Administration; FRET – Förster Resonance Energy Transfer; GSK – Glaxosmithkline; HPLC – High-performance liquid chromatography; HRMS – High-resolution mass spectrometry; hsEH – human soluble epoxide hydrolase; IC50 – the concentration of an inhibitor where the response (or binding) is reduced by half; LPS – lipopolysaccharide; mg/kg – mg compound per kg bodyweight; PD – pharmacodynamics; PDE4 – phosphodiesterase 4; PDE4I – phosphodiesterase 4 inhibitor; PEG300 – Polyethylene Glycol 300; PK – pharmacokinetic; PKA – protein kinase A; PO – per oral; PsA – psoriatic arthritis; rsEH – rat soluble epoxide hydrolase; rt – room temperature; sEH – soluble epoxide hydrolase; sEHI – soluble epoxide hydrolase inhibitor; Tmax – the amount of time that a drug is present at the maximum concentration in blood; TWL – Thermal withdrawal threshold; ws – water solubility.

References

- Reddy, A. S.; Zhang, S. Polypharmacology: Drug Discovery for the Future. *Expert Rev. Clin. Pharmacol.* 2013, 6 (1), 41–47.
- Perez-Aso, M.; Montesinos, M. C.; Mediero, A.; Wilder, T.; Schafer, P. H.; Cronstein,
 B. Apremilast, a Novel Phosphodiesterase 4 (PDE4) Inhibitor, Regulates Inflammation
 through Multiple cAMP Downstream Effectors. *Arthritis Res. Ther.* 2015, *17*, 249.
- Brown, W. M. Treating COPD with PDE 4 Inhibitors. *Int J Chron Obstruct Pulmon Dis.* 2007, 2 (4), 517-533.
- (4) Ashton, M. J.; Cook, D. C.; Fenton, G.; Karlsson, J. A.; Palfreyman, M. N.; Raeburn, D.; Ratcliffe, A. J.; Souness, J. E.; Thurairatnam, S.; Vicker, N. Selective Type IV Phosphodiesterase Inhibitors as Antiasthmatic Agents. The Syntheses and Biological Activities of 3-(Cyclopentyloxy)-4- Methoxybenzamides and Analogues. *J. Med. Chem.* 1994, *37* (11), 1696–1703.
- (5) Fujita, M.; Richards, E. M.; Niciu, M. J.; Hines, C. S.; Pike, V. W.; Jr, C. A. Z. cAMP

Signaling in Brain Is Decreased in Unmedicated Depressed Patients and Increased by Treatment with a Selective Serotonin Reuptake Inhibitor. *Mol Psychiatry*. **2017**, *22* (5), 754-759.

- Brandon, N. J. Uncovering the Function of Disrupted in Schizophrenia 1 through Interactions with the cAMP Phosphodiesterase PDE4: Contributions of the Houslay Lab to Molecular Psychiatry. *Cell. Signal.* 2016, 28 (7), 749–752.
- Gurney, M. E.; D'Amato, E. C.; Burgin, A. B. Phosphodiesterase-4 (PDE4) Molecular
 Pharmacology and Alzheimer's Disease. *Neurotherapeutics* 2015, *12* (1), 49–56.
- (8) Dal Piaz, V.; Giovannoni, M. P. Phosphodiesterase 4 Inhibitors, Structurally Unrelated to Rolipram, as Promising Agents for the Treatment of Asthma and Other Pathologies. *Eur. J. Med. Chem.* 2000, *35* (5), 463–480.
- Rock, E. M.; Benzaquen, J.; Limebeer, C. L.; Parker, L. A. Potential of the Rat Model of Conditioned Gaping to Detect Nausea Produced by Rolipram, a Phosphodiesterase-4 (PDE4) Inhibitor. *Pharmacol. Biochem. Behav.* 2009, *91* (4), 537–541.
- (10) Prickaerts, J., Heckman, P.R.A., Blokland, A. Investigational Phosphodiesterase Inhibitors in Phase I and Phase II Clinical Trials for Alzheimer's Disease. *Expert Opin Investig Drugs.* 2017, 26 (9), 1033-1048.
- Mclellan, G. J.; Aktas, Z.; Hennes-beean, E.; Kolb, A. W.; Larsen, V.; Schmitz, E. J.;
 Clausius, H. R.; Yang, J.; Hwang, S. H.; Morisseau, C.; Inceoglu, B.; Hammock, B. D.;
 Brandt, C. R. Induced Uveitis in the Rabbit. *J. Ocul. Biol.* 2017, *4* (1), 1–17.
- Wagner, K.; Inceoglu, B.; Hammock, B. D. Soluble Epoxide Hydrolase Inhibition,
 Epoxygenated Fatty Acids and Nociception. *Prostaglandins Other Lipid Mediat*. 2011,
 96 (1–4), 76–83.
- Morisseau, C.; Hammock, B. D. Impact of Soluble Epoxide Hydrolase and Epoxyeicosanoids on Human Health. *Annu. Rev. Pharmacol. Toxicol.* 2013, *53*, 37–58.
- (14) Ren, Q.; Ma, M.; Ishima, T.; Morisseau, C.; Yang, J.; Wagner, K. M.; Zhang, J. Gene

| 2 | |
|----------|--|
| 3 4 | |
| 5 6 | |
| 7 8 | |
| 9 10 | |
| 11 | |
| 13 | |
| 14 | |
| 16 17 | |
| 18 19 | |
| 20 21 | |
| 22 23 | |
| 24 25 | |
| 26 27 | |
| 28 29 | |
| 30 31 | |
| 32 | |
| 34 | |
| 36 | |
| 37 | |
| 39 40 | |
| 41 42 | |
| 43 44 | |
| 45 46 | |
| 47 48 | |
| 49 50 | |
| 51 52 | |
| 53 54 | |
| 55 56 | |
| 57 | |
| 58 59 | |
| 60 | |

Deficiency and Pharmacological Inhibition of Soluble Epoxide Hydrolase Confers Resilience to Repeated Social Defeat Stress. *Proc Natl Acad Sci U S A*. **2016**, *113* (13), E1944-E1952.

- (15) Inceoglu, B.; Wagner, K.; Schebb, N. H.; Morisseau, C.; Jinks, S. L.; Ulu, A.;
 Hegedus, C.; Rose, T.; Brosnan, R.; Hammock, B. D. Analgesia Mediated by Soluble
 Epoxide Hydrolase Inhibitors Is Dependent on cAMP. *Proc. Natl. Acad. Sci. USA*.
 2011, *108* (12), 5093–5097.
- (16) Blöcher, R.; Lamers, C.; Wittmann, S. K.; Merk, D.; Hartmann, M.; Weizel, L.; Diehl, O.; Brüggerhoff, A.; Boß, M.; Kaiser, A.; Schader, T.; Göbel, T.; Grundmann, M.; Angioni, C.; Heering, J.; Geisslinger, G.; Wurglics, M.; Kostenis, E.; Brüne, B.; Steinhilber, D.; Schubert-Zsilavecz, M.; Kahnt, A. S.; Proschak, E. N-Benzylbenzamides: A Novel Merged Scaffold for Orally Available Dual Soluble Epoxide Hydrolase/Peroxisome Proliferator-Activated Receptor Gamma Modulators. *J. Med. Chem.* 2016, *59* (1), 61–81.
- (17) la Buscató, E.; Blöcher, R.; Lamers, C.; Klingler, F.-M.; Hahn, S.; Steinhilber, D.;
 Schubert-Zsilavecz, M.; Proschak, E. Design and Synthesis of Dual Modulators of
 Soluble Epoxide Hydrolase and Peroxisome Proliferator-Activated Receptors. *J. Med. Chem.* 2012, *55* (23), 10771–10775.
- Meirer, K.; Glatzel, D.; Kretschmer, S.; Wittmann, S.; Hartmann, M.; Blöcher, R.;
 Angioni, C.; Geisslinger, G.; Steinhilber, D.; Hofmann, B.; Fürst, R.; Proschak, E.
 Design, Synthesis and Cellular Characterization of a Dual Inhibitor of 5-Lipoxygenase and Soluble Epoxide Hydrolase. *Molecules* 2016, *22* (1), E45.
- Podolin, P. L.; Bolognese, B. J.; Foley, J. F.; Long, E.; Peck, B.; Umbrecht, S.; Zhang, X.; Zhu, P.; Schwartz, B.; Xie, W.; Quinn, C.; Qi, H.; Sweitzer, S.; Chen, S.; Galop, M.; Ding, Y.; Belyanskaya, S. L.; Israel, D. I.; Morgan, B. A.; Behm, D. J.; Marino, J. P.; Kurali, E.; Barnette, M. S.; Mayer, R. J.; Booth-Genthe, C. L.; Callahan, J. F. In

Vitro and in Vivo Characterization of a Novel Soluble Epoxide Hydrolase Inhibitor. *Prostaglandins Other Lipid Mediat*. **2013**, *104–105*, 25–31.

- (20) Lazaar, A. L.; Yang, L.; Boardley, R. L.; Goyal, N. S.; Robertson, J.; Baldwin, S. J.; Newby, D. E.; Wilkinson, I. B.; Tal-Singer, R.; Mayer, R. J.; Cheriyan, J.
 Pharmacokinetics, Pharmacodynamics and Adverse Event Profile of GSK2256294, a Novel Soluble Epoxide Hydrolase Inhibitor. *Br. J. Clin. Pharmacol.* 2016, *81* (5), 971– 979.
- (21) Card, G. L.; England, B. P.; Suzuki, Y.; Fong, D.; Powell, B.; Lee, B.; Luu, C.;
 Tabrizizad, M.; Gillette, S.; Ibrahim, P. N.; Artis, D. R.; Bollag, G.; Milburn, M. V.;
 Kim, S. H.; Schlessinger, J.; Zhang, K. Y. J. Structural Basis for the Activity of Drugs
 That Inhibit Phosphodiesterases. *Structure* 2004, *12* (12), 2233–2247.
- (22) Lee, K. S. S.; Liu, J. Y.; Wagner, K. M.; Pakhomova, S.; Dong, H.; Morisseau, C.; Fu, S. H.; Yang, J.; Wang, P.; Ulu, A.; Mate, C. A.; Nguyen, L. V.; Hwang, S. H.; Edin, M. L.; Mara, A. A.; Wulff, H.; Newcomer, M. E.; Zeldin, D. C.; Hammock, B. D. Optimized Inhibitors of Soluble Epoxide Hydrolase Improve in Vitro Target Residence Time and in Vivo Efficacy. *J. Med. Chem.* 2014, *57* (16), 7016–7030.
- (23) Brown, A.; Rawson, D.; Storer, R.; Swain, A. Chemical Compounds, U.S. Patent 007868 A2; 2012.
- (24) Anandan, S. K.; Webb, H. K.; Chen, D.; Wang, Y. X.; Aavula, B. R.; Cases, S.; Cheng, Y.; Do, Z. N.; Mehra, U.; Tran, V.; Vincelette, J.; Waszczuk, J.; White, K.; Wong, K. R.; Zhang, L. N.; Jones, P. D.; Hammock, B. D.; Patel, D. V.; Whitcomb, R.; MacIntyre, D. E.; Sabry, J.; Gless, R. 1-(1-Acetyl-Piperidin-4-Yl)-3-Adamantan-1-Yl-Urea (AR9281) as a Potent, Selective, and Orally Available Soluble Epoxide Hydrolase Inhibitor with Efficacy in Rodent Models of Hypertension and Dysglycemia. *Bioorganic Med. Chem. Lett.* 2011, *21* (3), 983–988.
- (25) Zhao, Z.; Pissarnitski, D. A.; Josien, H. B.; Bara, T. A.; Clader, J. W.; Li, H.; McBriar,

| 1 | |
|-----|--|
| 1 | |
| 2 | |
| З | |
| | |
| 4 | |
| 5 | |
| 5 | |
| 6 | |
| 7 | |
| 0 | |
| ð | |
| 9 | |
| 10 | |
| 10 | |
| 11 | |
| 12 | |
| 12 | |
| 13 | |
| 14 | |
| 1 5 | |
| 15 | |
| 16 | |
| 17 | |
| 17 | |
| 18 | |
| 10 | |
| 19 | |
| 20 | |
| 21 | |
| 21 | |
| 22 | |
| 23 | |
| 24 | |
| 24 | |
| 25 | |
| 26 | |
| 20 | |
| 27 | |
| 28 | |
| 20 | |
| 29 | |
| 30 | |
| 21 | |
| 31 | |
| 32 | |
| 22 | |
| 22 | |
| 34 | |
| 35 | |
| 55 | |
| 36 | |
| 37 | |
| 20 | |
| 38 | |
| 39 | |
| 40 | |
| 40 | |
| 41 | |
| 42 | |
| 42 | |
| 43 | |
| 44 | |
| 45 | |
| 45 | |
| 46 | |
| 17 | |
| 47 | |
| 48 | |
| 49 | |
| -72 | |
| 50 | |
| 51 | |
| 5. | |
| 52 | |
| 53 | |
| E 4 | |
| 54 | |
| 55 | |
| 56 | |
| 50 | |
| 57 | |
| 58 | |
| 50 | |
| 59 | |

M. D.; Rajagopalan, M.; Xu, R.; Terracina, G.; Hyde, L.; Song, L.; Zhang, L.; Parker,

E. M.; Osterman, R.; Buevich, A. V. Substituted 4-Morpholine N-Arylsulfonamides as Gamma-Secretase Inhibitors. *Eur. J. Med. Chem.* **2016**, *124*, 36–48.

- (26) Rose, T. E.; Morisseau, C.; Liu, J. Y.; Inceoglu, B.; Jones, P. D.; Sanborn, J. R.;
 Hammock, B. D. 1-Aryl-3-(1-Acylpiperidin-4-Yl)urea Inhibitors of Human and Murine
 Soluble Epoxide Hydrolase: Structure-Activity Relationships, Pharmacokinetics, and
 Reduction of Inflammatory Pain. *J. Med. Chem.* 2010, *53* (19), 7067–7075.
- (27) Soto, D.; De Arcangelis, V.; Zhang, J.; Xiang, Y. K.; Liu, S.; Li, Y.; Kim, S.; Fu, Q.;
 Parikh, D.; Sridhar, B.; Shi, Q.; Zhang, X.; Guan, Y.; Chen, X.; Xiang, Y. K.
 Phosphodiesterases Coordinate cAMP Propagation Induced by Two Stimulatory G
 Protein-Coupled Receptors in Hearts. *Circ. Res.* 2009, *109* (17), 6578–6583.
- Bian, H.; Zhang, J.; Wu, P.; Varty, L. A.; Jia, Y.; Mayhood, T.; Hey, J. A.; Wang, P. Differential Type 4 cAMP-Specific Phosphodiesterase (PDE4) Expression and Functional Sensitivity to PDE4 Inhibitors among Rats, Monkeys and Humans. *Biochem Pharmacol.* 2004, 68 (11), 2229-2236.
- (29) Halpin, D. M. G. ABCD of the Phosphodiesterase Family: Interaction and Differential Activity in COPD. *Int J Chron Obstruct Pulmon Dis.* 2008, 3 (4), 543–561.
- (30) Richter, W., Xie, M., Scheitrum, C., Krall, J., Movsesian, M.A., Conti, M. Conserved Expression and Functions of PDE4 in Rodent and Human Heart. *Basic Res Cardiol.* 2011, *106*, 249-262.
- (31) Hwang, S. H.; Wagner, K. M.; Morisseau, C.; Liu, J.-Y.; Dong, H.; Wecksler, A. T.;
 Hammock, B. D. Synthesis and Structure-Activity Relationship Studies of Urea-Containing Pyrazoles as Dual Inhibitors of Cyclooxygenase-2 and Soluble Epoxide Hydrolase. *J. Med. Chem.* 2011, *54* (8), 3037–3050.
- (32) Jones, P. D.; Wolf, N. M.; Morisseau, C.; Whetstone, P.; Hock, B.; Hammock, B. D.Fluorescent Substrates for Soluble Epoxide Hydrolase and Application to Inhibition

Studies. Anal. Biochem. 2005, 343 (1), 66-75.

- Barbagallo, F.; Xu, B.; Reddy, G. R.; West, T.; Wang, Q.; Fu, Q.; Li, M.; Shi, Q.;
 Ginsburg, K. S.; Ferrier, W.; Isidori, A. M.; Naro, F.; Patel, H. H.; Bossuyt, J.; Bers,
 D.; Xiang, Y. K. Genetically Encoded Biosensors Reveal PKA Hyperphosphorylation
 on the Myofilaments in Rabbit Heart Failure. *Circ. Res.* 2016, *119* (8), 931–943.
- (34) Liu, J. Y.; Lin, Y. P.; Qiu, H.; Morisseau, C.; Rose, T. E.; Hwang, S. H.;

Chiamvimonvat, N.; Hammock, B. D. Substituted Phenyl Groups Improve the
Pharmacokinetic Profile and Anti-Inflammatory Effect of Urea-Based Soluble Epoxide
Hydrolase Inhibitors in Murine Models. *Eur. J. Pharm. Sci.* 2013, 48 (4–5), 619–627.



Scheme 1. Development of sEH/PDE4 dual inhibitor design.



a) EDC, DMAP, DCM, 12 h; b) 1. Sodium chloro difluoro acetate, Cs_2CO_3 , DMF, H_2O , 2 h 2. 100 °C, 12 h; c) NaOH, H_2O /THF/MeOH, 40 °C, 12 h; d) Cyclopentyl bromide, K_2CO_3 , KI, DMF, 65 °C, 21h; e) TFA, DCM, 12 h; Propionic acid or butanoic acid or 2-methylbutyric acid, (COCl)₂, DCM, TEA, 4 h; g) MeI or EtI, K_2CO_3 , DMF, 12 h.

Scheme 2. Synthetic routes for the production of sEH/PDE4 dual inhibitors

Table 1. Evaluation of dual sEH/PDE4 inhibitors in vitro activity



ACS Paragon Plus Environment

| | | | | | | _ |
|-------------------|-------------------|--|---------------|----------------------------|-------------------|---|
| 11 | OMe | - | 1 ± 0.2 | 160 ± 17 | 100 | - |
| 14 | OCHF ₂ | - | 1.5 ± 0.4 | 150 ± 14 | 10 | |
| 10 | _ | cyclopenty | 0.4 ± 0.1 | 147 + 9 | 10 | |
| 17 | _ | $\begin{array}{c} - & 0.4 \pm 0.1 \\ \\ \text{loxy} \end{array}$ | 0.7 ± 0.1 | 14/ 4 / | 10 | |
| 20 | F | cyclopenty | 0.4 ± 0 | 222 + 8 | 100 | |
| -0 | - | loxy | 0.1 – 0 | | 100 | |
| F F F | NH N.R5 | | | | | - |
| | | | hsEH | cAMP increase ^b | w.s. ^c | |
| ID | R ₅ | | IC_{50}^{a} | (% of Rolipram) | [µM] | |
| | | | [nM] | | | |
| TPPU ^D | - | | 3.7 | 40 ± 10 | 200 | |
| 21 | BOC | | 1.7 ± 0.7 | 100 ± 12 | 100 | |
| 22 | - | | 370 ± 170 | 138 ± 5 | 1000 | |
| 23/MPPA | COEt | | 2.1 ± 0.5 | 200 ± 19 | 100 | |
| 24 | COPr | | 4 ± 2 | 110 ± 16 | 50 | |
| 25 | s-Bu | | 7 ± 3 | 160 ± 11 | 0.5 | |
| | | | | | | |
| 26 | Et | | 109 ± 16 | 60 ± 10 | 1000 | |

In vitro evaluation of sEH/PDE4 dual inhibitors Group 1: 1-3, Group 2: 4-11, 14, 19-20 and Group 3: 21-27. ^aThe sEH IC₅₀ values are recorded on recombinant human sEH protein (hsEH) using MNPC as substrate at 5 μ M concentration³²; ^bPDE4 inhibition is evaluated by cAMP increase induced by 1 μ M concentration of the dual inhibitors in live human embryonic kidney (HEK) cells, transfected with a PKA biosensor and visualized relative to Rolipram at 1 μ M; ^cwater solubility (w.s.) is measured in sodium phosphate buffer (0.1 M,

| 2 | |
|------------|--|
| 2 | |
| 3 | |
| 4 | |
| 5 | |
| 6 | |
| 7 | |
| / | |
| 8 | |
| 9 | |
| 10 | |
| 11 | |
| 10 | |
| 12 | |
| 13 | |
| 14 | |
| 15 | |
| 16 | |
| 10 | |
| 17 | |
| 18 | |
| 19 | |
| 20 | |
| 20 | |
| 21 | |
| 22 | |
| 23 | |
| 24 | |
| 25 | |
| 26 | |
| 20 | |
| 27 | |
| 28 | |
| 29 | |
| 30 | |
| 31 | |
| 27 | |
| 22 | |
| 33 | |
| 34 | |
| 35 | |
| 36 | |
| 37 | |
| 20 | |
| 20 | |
| 39 | |
| 40 | |
| 41 | |
| 42 | |
| <u>4</u> २ | |
| ر⊤- ۸۸ | |
| 44 | |
| 45 | |
| 46 | |
| 47 | |
| 48 | |
| 10 | |
| 49 | |
| 50 | |
| 51 | |

pH 7.4) with 1% DMSO by precipitation. All experiments were performed in triplicates.

Results are average \pm standard deviation of at least three measurements.

| ID | $\begin{array}{c} IC_{50}(hsEH)^{a}\\ [nM] \end{array}$ | IC ₅₀ (rsEH) ^b [nM] | EC ₅₀ (cAMP increase) ^c [nM] | Melting point ^d (°C) |
|-------------------|---|--|--|---------------------------------------|
| TPPU ^d | 3.7 | 2.8 | - | 226 |
| Rolipram | >100 000 | - | 340 ± 0.06 | 127 |
| 1 | 0.6 ± 0.1 | 14 ± 1 | 1.9 ± 0.03 | 144 |
| 19 | 0.4 ± 0.1 | 1.3 ± 0.2 | 128 ± 0.05 | 106 |
| 20 | 0.4 ± 0 | 1.7 ± 0.2 | 3.9 ± 0.06 | 99 |
| 23/MPPA | 2.1 ± 0.5 | 150 ± 16 | 8.1 ± 0.05 | 133 |

Table 2. In vitro evaluation for laboratory species

IC₅₀ evaluation of dual inhibitors. ^aThe sEH IC₅₀ on recombinant human sEH (hsEH); ^bThe sEH IC₅₀ on recombinant rat sEH (rsEH); ^cIC₅₀ evaluated on cAMP increase in live HEK cells, transfected with a PKA biosensor; All experiments were performed in triplicates; ^dMelting points were measured on an automated melting point system from Stanford Research Systems (OptiMelt-MPA100);^dLiterature values.²⁶ Results are average \pm standard deviation of at least three measurements.



Figure 1. **MPPA** is efficacious against LPS induced inflammatory pain. Baseline thermal withdrawal latencies were determined for each rat before administration of compounds. After baseline determination, **MPPA** and the vehicle control were administered by oral gavage immediately before a 50 µl intraplantar injection of LPS in saline (arrow) to induce inflammatory pain in male rats. The efficacy of **MPPA** peaked at 30 min post LPS injection and was largely dissipated by 4 hours. The efficacy at 3 mg/kg was statistically significant compared to vehicle control. Scores are the mean \pm SEM reported as percent of baseline (baseline score normalized to 100%) calculated as the score *100/baseline score (Mann-Whitney Rank Sum Test, T = 448.000 n(small)= 24 n(big)= 36 *p≤0.001, n=6-9/group).

ACS Paragon Plus Environment

Table of contents graphic

sEH/PDE4 dual inhibitor



MPPA

| rsEH IC ₅₀ – 150 ± 16 nM |
|---------------------------------------|
| PDE4 IC ₅₀ – 8.1 ± 0.05 nM |
| in vivo at 3 mg/kg PO in rat |
| T _ 30 min |

hsEH IC₅₀ – 2.1 ± 0.5 nM

in vitro

 $T_{max} - 30 \text{ min}$ $C_{max} - 460 \pm 16 \text{ nM}$ cleared from plasma after 4h

