


Potassium Fluoride-induced 1,4-Elimination of *o*-[(Trimethylsilyl)methyl]benzyl Acetates: A Versatile Generation of *o*-Quinodimethanes

Ryoichi Kuwano* and Takenori Shige

Department of Chemistry, Graduate School of Sciences, Kyushu University, 6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8581

(Received February 24, 2005; CL-050238)


o-Quinodimethane intermediate was generated from *o*-[(trimethylsilyl)methyl]benzyl acetate with potassium fluoride. The *o*-quinodimethane in hand reacted with electron-deficient olefins, affording [4+2] cycloadducts, tetrahydronaphthalenes.



Scheme 2.

Cycloaddition of olefins to *o*-quinodimethane¹ intermediates has offered a versatile and straightforward access to 1,2,3,4-tetrahydronaphthalene skeletons.² A variety of methods for in situ generation of *o*-quinodimethane have been developed,^{3–7} and have been utilized for the syntheses of steroids⁸ and other natural products to date.⁹ In particular, fluoride-induced 1,4-elimination from α' -heterosubstituted α -silyl-*o*-xylenes is an important and versatile method for the preparation of *o*-quinodimethane.^{10–13} The 1,4-elimination occurred under mild conditions compared with other methods, e.g. thermal ring-opening of benzocyclobutenes.¹⁴ The generation of *o*-quinodimethane from the α -silyl-*o*-xylenes, however, has been conducted in the presence of either tetrabutylammonium fluoride or cesium fluoride, which are expensive and hard to handle in air because of their hygroscopicity.

Herein, we report a new protocol for the generation of *o*-quinodimethanes via 1,4-elimination of *o*-(silylmethyl)benzylic acetates (**1**).^{12,15} Potassium fluoride, which is handled with ease in air and is an inexpensive fluoride source,¹⁶ was employed for the induction of 1,4-elimination. The *o*-quinodimethanes reacted with electron-deficient olefins to give the corresponding tetrahydronaphthalenes **2** (Scheme 1).




Scheme 1.

A mixture of *o*-[(trimethylsilyl)methyl]benzyl acetate (**1a**) with methyl acrylate in DMF was heated at 100 °C in the presence of KF. The resulting mixture afforded methyl 1,2,3,4-tetrahydronaphthalene-2-carboxylate (**2a**) in 88% yield (Scheme 2).¹⁷ When tetrabutylammonium fluoride was used as a fluoride source, **2a** was obtained in trace amount (2%), and 2-methylbenzyl acetate resulting from protodesilylation was generated in 50% yield.¹⁸ The desired product **2a** was formed by use of KF even at room temperature (26% at 48 h), while the reaction afforded 2-methylbenzyl acetate in 40% yield.¹⁹ This observation may indicate that the KF-induced activation of silyl group occurs even at room temperature but the following 1,4-elimination of the acetoxy group seems to compete with the protodesilylation. No consumption of **1a** was observed in any solvents other than

DMF. Lewis basicity of DMF may assist the activation of C–Si bond by KF.²⁰

When **1a** was treated with KF in the absence of any dienophile, a mixture of spiro-di-*o*-xylene **3** and [2,2]orthocyclophane **4** was obtained as seen in Scheme 3. Although the formation of **4** might result from thermally forbidden [4,4] cycloaddition of *o*-quinodimethane, the observation is consistent with Errede's results.²¹













Scheme 3.

The *o*-quinodimethane intermediate generated from **1a** reacted with acrylonitrile as well as with methyl acrylate, and the cycloaddition product **2b** was obtained in high yield (Table 1, Entry 1). Unexpectedly, acrolein was not so reactive toward the *o*-quinodimethane intermediate (Entry 2). The reactions of electron-deficient trans-olefins, *trans*-benzalacetone, and dimethyl fumarate, afforded [4+2] cycloadducts **2d** and **2e** in good yields, respectively (Entries 3 and 4). In contrast, no cycloaddition of *o*-quinodimethane was observed when cis-olefins such as maleic anhydride and dimethyl maleate were employed as a dienophile.²² Use of acetylenedicarboxylate resulted in a complex mixture.²² The reaction of 4-methyl-*o*-quinodimethane precursor **1b** with methyl acrylate gave a mixture of two regioisomeric [4+2] cycloadducts with no selectivity (Entry 5).


The *o*-quinodimethane precursor **1a** was prepared with acetylation of benzylic alcohol **5**, which was readily synthesized from *o*-methylbenzyl alcohol.^{11b} We attempted the one-pot generation of *o*-quinodimethane from **5** (Scheme 4). After **5** was treated with acetic anhydride, methyl acrylate and KF were added to the resulting solution containing **1a**. Heating the mixture at 100 °C for 1 h produced **2a** in 66% yield.²³

In summary, we have developed a new method for generating *o*-quinodimethane intermediates from **1**. Potassium fluoride, which is an inexpensive and tractable fluoride source, was employed to activate the silyl group and induced 1,4-elimination providing the reactive intermediate. The *o*-quinodimethane in

Table 1. Cycloaddition of *o*-quinodimethane with dienophiles^a

Entry	1	Dienophile	Product (yield/%) ^b
1			 2b (81)
2 ^c			 2c (26)
3			 2d (63)
4			 2e (77)
5 ^c			 2f (68) ^d

^aReactions were conducted on a 1 mmol scale in DMF (20 mL) at 100 °C for 3 h. 1/dienophile/KF was 1/1.25/1.5. ^bIsolated yield. ^cThe reactions were conducted for 1 h. ^dYield of the mixture of methyl 5- and 6-methyl-1,2,3,4-tetrahydronaphthalene-2-carboxylates (ca. 1:1).

**Scheme 4.** One-pot synthesis of tetrahydronaphthalene from **3**.

hand reacted with electron-deficient dienophiles to give tetrahydronaphthalenes.

This work was partly supported by the Sumitomo Foundation and a Grant-in-Aid for Young Scientist (A) (no. 16685011) from MEXT.

References and Notes

- M. P. Cava and D. R. Napier, *J. Am. Chem. Soc.*, **79**, 1701 (1957).
- Reviews: a) J. L. Charlton and M. M. Alauddin, *Tetrahedron*, **43**, 2873 (1987). b) J. L. Segura and N. Martin, *Chem. Rev.*, **99**, 3199 (1999).
- M. P. Cava and A. A. Deana, *J. Am. Chem. Soc.*, **81**, 4266 (1959).
- K. Alder and M. Fremery, *Tetrahedron*, **14**, 190 (1961).
- P. G. Sammes and S. M. Mellows, *J. Chem. Soc., Chem. Commun.*, **1971**, 21.
- F. Jung, M. Molin, R. Van der Elzen, and T. Durst, *J. Am. Chem. Soc.*, **96**, 935 (1974).
- T. Tuschka, K. Naito, and B. Rickborn, *J. Org. Chem.*, **48**, 70 (1983).
- Review: H. Nemoto and K. Fukumoto, *Tetrahedron*, **54**, 5425 (1998).
- Recent examples: a) Y. Matsuya, K. Sasaki, M. Nagaoka, H. Kakuda, N. Toyooka, N. Imanishi, H. Ochiai, and H. Nemoto, *J. Org. Chem.*, **69**, 7989 (2004). b) K. C. Nicolaou and D. L. F. Gray, *J. Am. Chem. Soc.*, **126**, 607 (2004).
- Generation from α -silyl- α' -ammonium-*o*-xylenes: a) Y. Ito, M. Nakatsuka, and T. Saegusa, *J. Am. Chem. Soc.*, **102**, 863 (1980). b) Y. Ito, M. Nakatsuka, and T. Saegusa, *J. Am. Chem. Soc.*, **104**, 7609 (1982). c) Y. Ito, Y. Amino, M. Nakatsuka, and T. Saegusa, *J. Am. Chem. Soc.*, **105**, 1586 (1983).
- Generation from α -silyl- α' -sulfonyl-*o*-xylenes: a) B. D. Lenihan and H. Shechter, *Tetrahedron Lett.*, **35**, 7505 (1994). b) B. D. Lenihan and H. Shechter, *J. Org. Chem.*, **63**, 2072 (1998).
- SiO₂-induced generation of *o*-quinodimethane from α -silyl- α' -acetoxy-*o*-xylenes: A. R. Beard, S. J. Hazell, J. Mann, and C. Palmer, *J. Chem. Soc., Perkin Trans. 1*, **1993**, 1235.
- Acid-induced generation of *o*-quinodimethane from α -stannyl- α' -hydroxy-*o*-xylenes: H. Sano, H. Ohtsuka, and T. Migita, *J. Am. Chem. Soc.*, **110**, 2014 (1988).
- Examples: a) W. Oppolzer, *J. Am. Chem. Soc.*, **93**, 3833 (1971). b) T. Kametani, M. Tsubuki, Y. Shiratori, Y. Kato, H. Nemoto, M. Ihara, K. Fukumoto, F. Satoh, and H. Inoue, *J. Org. Chem.*, **42**, 2672 (1977).
- I. Fleming, I. T. Morgan, and A. K. Sarkar, *J. Chem. Soc., Perkin Trans. 1*, **1998**, 2749.
- The costs of fluoride sources are as follows: TBAF (1.0 M THF solution), ¥37,400/mol; CsF, ¥25,519/mol; KF, ¥581/mol (from Aldrich).
- Procedure of the reaction of **1a** with methyl acrylate is as follows: Under nitrogen atmosphere, **1a** (241 mg, 1.02 mmol) and methyl acrylate (109 mg, 1.26 mol) were added to a suspension of KF (88 mg, 1.51 mmol) in dry DMF (20 mL). The mixture was stirred at 100 °C for 3 h. After cooling, the mixture was diluted with water and was extracted with hexane. The organic layer was washed with brine, was dried with MgSO₄, and was evaporated under reduced pressure. The residue was purified with a flash column chromatography on silica gel (EtOAc/hexane = 1/5), giving **2a** (170 mg, 88%): ¹H NMR (400 MHz, CDCl₃, TMS) δ 1.86 (dddd, *J* = 6.6, 10.8, 13.0, 23.9 Hz, 1H), 2.17–2.25 (m, 1H), 2.70–2.79 (m, 1H), 2.79–2.93 (m, 2H), 2.95–3.06 (m, 2H), 3.73 (s, 3H), 7.06–7.13 (m, 4H); ¹³C {¹H} NMR (100 MHz, CDCl₃) δ 25.8, 28.5, 31.6, 39.9, 51.7, 125.7, 125.8, 128.8, 129.0, 134.8, 135.6, 175.8. lit. S.-i. Inaba, R. M. Wehmeyer, M. W. Forkner, and R. D. Rieke, *J. Org. Chem.*, **53**, 339 (1988).
- The reaction was conducted in dry DMF (5 mL) at 80 °C. The rest of **1a** was converted into **3** and **4**. The water contained in TBAF solution might cause the protodesilylation. The reaction employing KF under the identical conditions gave **2a** in 70% yield.
- The remaining **1a** was recovered.
- S. Kobayashi and K. Nishio, *J. Org. Chem.*, **59**, 6620 (1994).
- The pathway of the [4+4] cycloaddition might involve the formation of a biradical intermediate: L. A. Errede, *J. Am. Chem. Soc.*, **83**, 949 (1961).
- Many examples of reactions of *o*-quinodimethanes with maleic anhydride and acetylenedicarboxylates have been reported.^{10,13}
- Procedure of the one-pot transformation of **5** into **2a** is as follows: Under nitrogen atmosphere, pyridine (95 mg, 1.20 mmol) and acetic anhydride (114 mg, 1.12 mmol) were added to a solution of **5** (193 mg, 0.99 mmol) and 4-(dimethylamino)pyridine (1.5 mg, 12 μ mol) in dry DMF (2 mL). The mixture was stirred at room temperature for 3 h. After DMF (18 mL), KF (87 mg, 1.50 mmol), and methyl acrylate (118 mg, 1.37 mmol) were added, the mixture was stirred at 100 °C for 1 h. The resulting mixture was purified according to the protocol described in note 17, giving **2a** (125 mg, 66%).