

Synthesis of a Capillarisin Sulfur-Analogue Possessing Aldose Reductase Inhibitory Activity by Selective Isopropylation

Yasushi IGARASHI,*^a Hiroaki KUMAZAWA,^a Toshihoro OHSHIMA,^a Hisanori SATOMI,^a
Susumu TERABAYASHI,^a Shuichi TAKEDA,^a Masaki ABURADA,^{a,b} and Ken-ichi MIYAMOTO^c

^a Research Division, Tsumura & Co.; 3586 Yoshiwara, Ami-machi, Inashiki-gun, Ibaraki 300–1192, Japan; ^b Faculty of Pharmacy, Musashino University; 1–1–20 Shinmachi, Nishitokyo, Tokyo 202–8585, Japan; and ^c Department of Hospital Pharmacy, School of Medicine, Kanazawa University; Kanazawa 920–8641, Japan.

Received February 9, 2005; accepted June 12, 2005

We describe the synthesis of 2-[(4-hydroxyphenyl)thio]-7-isopropoxy-5,6-dimethoxy-4*H*-chromen-4-one **2** from 3,4,5-trimethoxyphenol **6** via the key intermediate, 3-iodo-7-isopropoxy-5,6-dimethoxy-4*H*-chromen-4-one **3**. An important feature of this synthetic scheme involves selective alkylation, which can be achieved by two different routes. One route involves the selective isopropylation of a triacetate derivative **4** under basic conditions. The second route employs the selective demethylation of a trimethoxy derivative **5** under acidic conditions followed by isopropylation. The product of these alternative routes, compound **3**, is then converted to a capillarisin sulfur analogue **2** in a one-pot reaction via the imidazolyl intermediate **22**.

Key words capillarisin; selective isopropylation; substitution reaction; aldose reductase inhibitor; 3,4,5-trimethoxyphenol

Capillarisin **1**,¹⁾ a bioactive constituent of *Artemisiae Capillaris Herba*, is an antioxidant,²⁾ as well as a choleric³⁾ and antitumor agent⁴⁾ and an inhibitor of aldose reductase (AR).⁵⁾ We have synthesized a number of capillarisin analogues and tested their ability to inhibit AR.⁶⁾ Among them, the sulfur analogue **2**, which was prepared from capillarisin **1** by a semi-synthetic method (Chart 1), showed excellent *in vitro* activity with an IC₅₀ of 3 × 10^{−8} mol/l against rat lens AR. Therefore compound **2** is an excellent lead structure for an AR inhibitor. 6-Demethoxycapillarisin has been synthesized by two groups,^{7,8)} but total synthesis of **1** has not been reported. During the development of a drug to treat diabetic complications, we devised a synthetic route to the sulfur analogue **2** from a commercially available compound.

A key feature of this synthesis is the selective construction of the A-ring substituted with trialkoxy groups and introduction of the phenylthio group into the 2-position in its structure. The retrosynthetic analysis of **2** is outlined in Fig. 1. A substitution reaction using 3-iodochromone **3** with 4-hydroxybenzenethiol affords **2**. To selectively arrange the 7-isopropoxy-5,6-dimethoxy groups on the A-ring of **3**, two synthetic routes were investigated. The first route involves the selective isopropylation using a triacetate derivative **4** (route a), and the second route involves selective demethylation of the trimethoxy derivative **5** (route b). Compounds **4** and **5** can be prepared from 3,4,5-trimethoxyphenol **6**. In this paper, we describe the total synthesis of the novel capillarisin derivative **2** by these two methods.

Synthesis of 3 via 4 (Route a) The starting material **7**, is commercially available or can easily be prepared from 3,4,5-trimethoxyphenol **6**.⁹⁾ Condensation of **7** with *N,N*-dimethylformamide dimethyl acetal gave **8** in 91% yield. Cycliza-

tion¹⁰⁾ of **8** using iodine afforded 3-iodochromone **9** in 84% yield (Chart 2).

To obtain the desired compounds, **14** or **15**, demethylation of compound **9** was carried out under various conditions (Table 1). Demethylation proceeded in the order of the position 5, 6 and 7 using Lewis acids (runs 1–4). Interestingly, the reaction of **9** with sulfuric acid (run 5) occurred selec-




Fig. 1. Retrosynthetic Analysis of **2**




Chart 2. Synthesis of **9**

Table 1. Demethylation of **9**

Run	Reagent	Solv.	Temp.	Time	Products (Yield)
1	BCl ₃	CH ₂ Cl ₂	rt.	20 h	10 (82%), 11 (18%)
2	BCl ₃	CH ₂ Cl ₂	−78 °C	10 min	10 (90%)
3	TiCl ₄	Tol., CH ₂ Cl ₂	rt.	20 h	10 (46%), 11 (21%)
4	AlCl ₃	Tol.	100 °C	6 h	12 (94%)
5	H ₂ SO ₄		80 °C	2 h	13 (96%)
6	LiI	Dioxane	Reflux	20 h	10 (100%)

10: R¹=Me, R²=Me, R³=H
11: R¹=Me, R²=H, R³=H
12: R¹=H, R²=H, R³=H
13: R¹=Me, R²=H, R³=Me
14: R¹=H, R²=Me, R³=Me
15: R¹=H, R²=Me, R³=H







Chart 1. Conversion of Capillarisin **1** to Sulfur-Analogue **2**

* To whom correspondence should be addressed. e-mail: igarashi_yasushi@mail.tsumura.co.jp


Chart 3. Isopropylation of **12**Chart 4. Synthesis of Key Intermediate **3** via **4**Chart 5. Alternative Synthetic Method of Key Intermediate **3**

tively at the 6-position to afford **13** in 96% yield. Additionally, compound **10** was exclusively obtained after treatment with lithium iodide (run 6). However, the desired compounds, **14** and **15**, could not be obtained by this method.

We then tried selective isopropylation at the 7-position using the trihydroxy derivative **12** under various conditions (base; K_2CO_3 , Cs_2CO_3 or Li_2CO_3 ; solvent; DMF, MEK, acetone or EtOH). However, the yield (maximum 43%) of desired compound **16** was unsatisfactory (Chart 3). Presumably the reactivity of the 7-hydroxy group was equal to that of the 6-hydroxy group, and the substrate **12** is unstable under basic conditions.

We judged that direct isopropylation of **12** is difficult, and therefore adopted a selective alkylation methodology using methyl gallate.¹¹ Compound **12** was converted to the triacetate **4** in 83% yield, and then reacted with 2-iodopropane in the presence of potassium carbonate. As expected, the 7-acetoxy group *para* to the carbonyl group reacted selectively to give the desired compound **17** in 86% yield. Hydrolysis of **17** followed by methylation afforded the key intermediate **3** in 86% yield (Chart 4).

Synthesis of **3 via **5** (Route b)** Since selective demethylation of **9** failed (Table 1), conversion of **9** to **3** was achieved

Chart 6. One-Pot Conversion of **3** to **2**

in 5 steps (route a). Thus we planned an alternative route involving the selective demethylation at the 7-position *via* intermediate **5**, possessing a carbonyl function at the 8-position.

Friedel–Crafts reaction of 3,4,5-trimethoxyphenol **6** using acetic anhydride and aluminum chloride in nitromethane afforded the diacetyl derivative **18** in 74% yield. Condensation of **18** with *N,N*-dimethylformamide dimethyl acetal (1 eq) followed by treatment with iodine gave **5**. Demethylation of **5** using titanium tetrachloride as Lewis acid proceeded selectively to afford the 5,7-dihydroxy-6-methoxy derivative **20** as expected. The acetyl group of **20** was then removed by reaction with trifluoromethanesulfonic acid¹²) to afford **14** in 89% yield. Isopropylation at the 7-position followed by methylation at the 5-position of **14** gave compound **3** in 62% yield (Chart 5).

Preparation of **2 from **3**** The conversion of **3** to **2** is outlined in Chart 6. A direct substitution reaction of the 3-iodo-derivative **3** with 4-hydroxybenzenethiol gave only the reduced product **21** in 88% yield. It was reported that 3-iodochromone reacted with imidazole to give 2-(1-imidazolyl)chromone.¹³ Treatment of **3** with imidazole in the presence of potassium carbonate in DMF gave the imidazolyl derivative **22**, which without isolation then underwent a substitution reaction with 4-hydroxybenzenethiol to give **2** in 86% yield. These two steps could proceed either in a one-pot reaction or step-by-step.

Conclusion

We have developed two synthetic routes for the preparation of the key intermediate **3** in the synthesis of a caparrisin-analogue **2**. Both reaction schemes use readily available starting reagents. One route employs the selective isopropylation of **4**, whereas the second route involves the selective demethylation of **5**. Compound **2** was prepared from **3** in a one-pot reaction *via* the imidazolyl derivative **22**.

Experimental

Melting points were measured by the use of a Yanaco micro melting point apparatus and are uncorrected. ¹H-NMR spectra were recorded on a JEOL FX-200 (200 MHz) using tetramethylsilane as an internal standard. Infrared spectra were recorded using a Hitachi 270-30 spectrophotometer. Mass spectra were recorded on a JEOL DX-300 spectrometer. Elemental analyses were performed on a Heraeus CHN-O-Rapid.

(*E*)-3-(Dimethylamino)-6'-hydroxy-2',3',4'-trimethoxyacrylophenone (8**)** A mixture of 6'-hydroxy-2',3',4'-trimethoxyacetophenone (607 mg, 2.69 mmol) and *N,N*-dimethylformamide dimethyl acetal (0.534 ml, 4.03 mmol) was stirred at 100 °C for 2 h. The reaction mixture was purified by flash chromatography on silica gel (hexane/EtOAc = 1/1) to give the title

compound (691.1 mg, 91%). Recrystallization from EtOAc–hexane afforded yellow crystals: mp 113–114 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 2.96 (3H, br s), 3.18 (3H, br s), 3.81 (3H, s), 3.86 (3H, s), 3.87 (3H, s), 6.25 (1H, s), 6.33 (1H, d, $J=12.4$ Hz), 7.95 (1H, d, $J=12.4$ Hz), 14.84 (1H, s). IR (KBr) cm^{-1} : 1616, 1528. EI-MS m/z : 281 (M^+). *Anal.* Calcd for $\text{C}_{14}\text{H}_{19}\text{NO}_5$: C, 59.78; H, 6.81; N, 4.98. Found: C, 59.71; H, 6.96; N, 5.09.

3-Iodo-5,6,7-trimethoxy-4H-chromen-4-one (9) A mixture of **8** (1.897 g, 6.75 mmol) and iodine (3.40 g, 13.4 mmol) in toluene (19 ml) was stirred at room temperature for 1.5 h. The reaction mixture was quenched with $\text{Na}_2\text{S}_2\text{O}_3$ aq and extracted with EtOAc. The organic phase was dried over Na_2SO_4 , concentrated and then recrystallized from EtOAc–hexane to give the title compound (1.871 g, 77%). The filtrate of the recrystallization was purified by flash chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{EtOAc}=10/1$) to give the title compound (177 mg, 7%): mp 183–184 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 3.90 (3H, s), 3.95 (3H, s), 3.97 (3H, s), 6.68 (1H, s), 8.13 (1H, s). IR (KBr) cm^{-1} : 1628, 1612. EI-MS m/z : 362 (M^+). *Anal.* Calcd for $\text{C}_{12}\text{H}_{11}\text{IO}_5$: C, 39.80; H, 3.06. Found: C, 39.95; H, 3.19.

5,6,7-Trihydroxy-3-iodo-4H-chromen-4-one (12) To a suspension of **9** (497.9 mg, 1.38 mmol) in toluene (10 ml) was added AlCl_3 (915 mg, 6.88 mmol) at room temperature. The mixture was then heated and stirred at 100 °C for 6 h. After cooling, the reaction mixture was poured into 3N-HCl, and stirred at room temperature for 1 h. The reaction mixture was extracted with EtOAc, dried over Na_2SO_4 and concentrated. The residue was washed with CH_2Cl_2 to give the title compound (415.6 mg, 94%): mp 227–229 °C. $^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ : 6.50 (1H, s), 8.65 (1H, s), 12.11 (1H, s). IR (KBr) cm^{-1} : 3472, 3420, 1650, 1612. EI-MS m/z : 320 (M^+). *Anal.* Calcd for $\text{C}_9\text{H}_5\text{IO}_5$: C, 33.78; H, 1.57. Found: C, 33.75; H, 1.72.

5,6,7-Triacetox-3-iodo-4H-chromen-4-one (4) To a solution of **12** (1.02 g, 3.19 mmol) in pyridine (5 ml) was added acetic anhydride (1.51 ml, 15.95 mmol), and the mixture was stirred at room temperature for 4 h. The reaction mixture was poured into 3N-HCl, and the resulting solid was filtered, washed with H_2O , and dried *in vacuo* to give the title compound (1.18 g, 83%). Recrystallization from EtOAc–hexane afforded colorless crystals: mp 200–202 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 2.33 (3H, s), 2.34 (3H, s), 2.45 (3H, s), 7.40 (1H, s), 8.21 (1H, s). IR (KBr) cm^{-1} : 1774, 1646, 1626. EI-MS m/z : 446 (M^+). HR-EI-MS m/z : 445.9482 (Calcd for $\text{C}_{15}\text{H}_{11}\text{IO}_8$: 445.9499).

5,6-Diacetox-3-iodo-7-isopropoxy-4H-chromen-4-one (17) A mixture of **4** (928 mg, 2.08 mmol), 2-iodopropane (0.416 ml, 4.16 mmol), and K_2CO_3 (861 mg, 6.24 mmol) in DMF (4.6 ml) was stirred at 40 °C for 6 h. H_2O was added to the mixture, and the mixture was extracted with EtOAc, dried over Na_2SO_4 and concentrated. The residue was purified by flash chromatography on silica gel (hexane/EtOAc=2/1) to give the title compound (795.5 mg, 86%). Recrystallization from EtOAc–hexane afforded colorless crystals: mp 138–139 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 1.39 (6H, d, $J=6.1$ Hz), 2.32 (3H, s), 2.44 (3H, s), 4.44 (1H, septet, $J=6.1$ Hz), 6.79 (1H, s), 8.14 (1H, s). IR (KBr) cm^{-1} : 1762, 1626. EI-MS m/z : 446 (M^+). *Anal.* Calcd for $\text{C}_{16}\text{H}_{13}\text{IO}_7$: C, 43.07; H, 3.39. Found: C, 43.08; H, 3.51.

3-Iodo-7-isopropoxy-5,6-dimethoxy-4H-chromen-4-one (3) 1) To a solution of **17** (793.8 mg, 1.78 mmol) in acetone (8 ml) was added 6N-HCl, and the mixture was stirred at 70 °C for 5 h. After adding H_2O the mixture was extracted with EtOAc, dried over Na_2SO_4 and concentrated to give 5,6-dihydroxy-3-iodo-7-isopropoxy-4H-chromen-4-one (554 mg, 86%). Recrystallization from EtOAc–hexane afforded colorless crystals: mp 151–152 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 1.45 (6H, d, $J=6.1$ Hz), 4.70 (1H, septet, $J=6.1$ Hz), 5.43 (1H, s), 6.50 (1H, s), 8.17 (1H, s), 11.96 (1H, s). IR (KBr) cm^{-1} : 3508, 1660, 1596. EI-MS m/z : 362 (M^+). *Anal.* Calcd for $\text{C}_{12}\text{H}_{11}\text{IO}_5$: C, 39.80; H, 3.06. Found: C, 39.97; H, 3.24.

2) A mixture of the above compound (554 mg, 1.53 mmol), K_2CO_3 (1.47 g, 10.68 mmol) and iodomethane (0.665 ml, 10.68 mmol) in DMF (8 ml) was stirred at 40 °C for 5 h. After adding H_2O the mixture was extracted with EtOAc, dried over Na_2SO_4 and concentrated. The residue was purified by flash chromatography on silica gel (hexane/EtOAc=4/1) to give the title compound (597.8 mg, 100%). Recrystallization from EtOAc–hexane afforded colorless crystals: mp 99–100 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 1.45 (6H, d, $J=6.1$ Hz), 3.87 (3H, s), 3.96 (3H, s), 4.65 (1H, septet, $J=6.1$ Hz), 6.65 (1H, s), 8.11 (1H, s). IR (KBr) cm^{-1} : 1638, 1608, 1478. EI-MS m/z : 390 (M^+). *Anal.* Calcd for $\text{C}_{14}\text{H}_{15}\text{IO}_5$: C, 43.10; H, 3.88. Found: C, 43.06; H, 3.82.

2,6-Diacetyl-3,4,5-trimethoxyphenol (18) To a solution of 3,4,5-trimethoxyphenol (3.21 g, 16.96 mmol) and acetic anhydride (6.41 ml, 67.84 mmol) in nitromethane (31 ml) was added AlCl_3 (6.77 g, 50.88 mmol) at 0 °C. The mixture was then stirred at room temperature for 20 h. 3N-HCl was added to the reaction mixture, and the mixture was stirred at room tem-

perature for 1 h, extracted with EtOAc, dried over Na_2SO_4 and concentrated. The residue was purified by flash chromatography on silica gel (hexane/EtOAc=6/1) to give the title compound (3.35 g, 74%). Recrystallization from hexane afforded colorless crystals: mp 85–86 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 2.59 (6H, s), 3.79 (3H, s), 4.00 (6H, s), 13.31 (1H, s). EI-MS m/z : 268 (M^+). *Anal.* Calcd for $\text{C}_{13}\text{H}_{16}\text{O}_6$: C, 58.20; H, 6.01. Found: C, 58.18; H, 6.04.

(E)-3'-Acetyl-3-(dimethylamino)-2'-hydroxy-4',5',6'-trimethoxyacrylophenone (19) A mixture of **18** (3.33 g, 12.4 mmol) and *N,N*-dimethylformamide dimethyl acetal (1.65 ml, 12.4 mmol) in DMF (3.3 ml) was stirred at 100 °C for 6 h. The reaction mixture was purified by flash chromatography on silica gel (hexane/EtOAc=1/4) to give the title compound (2.70 g, 67%). Recrystallization from EtOAc–hexane afforded yellow crystals: mp 113–114 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 2.60 (3H, s), 2.91 (3H, s), 3.12 (3H, s), 3.81 (3H, s), 3.94 (3H, s), 3.97 (3H, s), 5.78 (1H, br d, $J=12.4$ Hz), 7.65 (1H, br s), 13.96 (1H, br s). EI-MS m/z : 323 (M^+). *Anal.* Calcd for $\text{C}_{16}\text{H}_{21}\text{NO}_6$: C, 59.43; H, 6.55; N, 4.33. Found: C, 59.21; H, 6.53; N, 4.31.

8-Acetyl-3-iodo-5,6,7-trimethoxy-4H-chromen-4-one (5) A mixture of **19** (1.23 g, 3.81 mmol) and iodine (1.94 g, 7.62 mmol) in CH_2Cl_2 (12 ml) was stirred at room temperature for 2 h. The reaction mixture was quenched with $\text{Na}_2\text{S}_2\text{O}_3$ aq and extracted with EtOAc. The organic phase was dried over Na_2SO_4 , concentrated, and purified by flash chromatography on silica gel (hexane/EtOAc=8/3) to give the title compound (1.27 g, 83%). Recrystallization from EtOAc–hexane afforded colorless crystals: mp 139–140 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 2.56 (3H, s), 3.93 (3H, s), 3.99 (3H, s), 4.05 (3H, s), 8.11 (1H, s). EI-MS m/z : 404 (M^+). *Anal.* Calcd for $\text{C}_{14}\text{H}_{13}\text{IO}_6$: C, 41.61; H, 3.24. Found: C, 41.35; H, 3.16.

8-Acetyl-5,7-dihydroxy-3-iodo-6-methoxy-4H-chromen-4-one (20) To a solution of **5** (100 mg, 0.248 mmol) in toluene (2 ml) was added TiCl_4 (0.74 ml, 1 M solution in toluene) at room temperature, and the mixture was then stirred at 100 °C for 1 h. After cooling, the reaction mixture was poured into 3N-HCl and stirred at 80 °C for 0.5 h. The reaction mixture was extracted with EtOAc, dried over Na_2SO_4 and concentrated. The residue was purified by flash chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{EtOAc}=30/1$) to give the title compound (68 mg, 73%). Recrystallization from EtOAc afforded colorless crystals: mp 224–226 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 2.76 (3H, s), 3.93 (3H, s), 8.31 (1H, s), 13.26 (1H, s), 14.54 (1H, s). EI-MS m/z : 376 (M^+). *Anal.* Calcd for $\text{C}_{12}\text{H}_9\text{IO}_6$: C, 38.32; H, 2.41. Found: C, 38.11; H, 2.47.

5,7-Dihydroxy-3-iodo-6-methoxy-4H-chromen-4-one (14) To a solution of **20** (222 mg, 0.59 mmol) in 1,2-dichloroethane (4 ml) was added trifluoromethanesulfonic acid (0.2 ml) and H_2O (0.2 ml) at room temperature. The mixture was then refluxed for 20 h. The reaction mixture was extracted with CH_2Cl_2 , dried over Na_2SO_4 and concentrated. The residue was purified by flash chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{EtOAc}=30/1$) to give the title compound (174.8 mg, 89%). Recrystallization from EtOAc–hexane afforded colorless crystals: mp 175–176 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 4.03 (3H, s), 6.52 (1H, s), 6.57 (1H, s), 8.14 (1H, s), 12.45 (1H, s). EI-MS m/z : 334 (M^+). *Anal.* Calcd for $\text{C}_{10}\text{H}_7\text{IO}_5$: C, 35.95; H, 2.11. Found: C, 35.91; H, 2.03.

2-[(4-Hydroxyphenyl)thio]-7-isopropoxy-5,6-dimethoxy-4H-chromen-4-one (2) A mixture of **3** (225.2 mg, 0.578 mmol), imidazole (79 mg, 1.16 mmol) and K_2CO_3 (798 mg, 5.78 mmol) in DMF (2.3 ml) was stirred at 80 °C for 2 h. After cooling, a solution of 4-hydroxybenzenethiol (146 mg, 1.16 mmol) in acetone (1 ml) was added to the reaction mix, and the mixture was stirred at room temperature for 2 h. The reaction mixture was poured into 3N-HCl, extracted with CH_2Cl_2 , dried over Na_2SO_4 and concentrated. The residue was purified by flash chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{acetone}=10/1$) to give the title compound (193 mg, 86%). Recrystallization from EtOH afforded colorless crystals: mp 213–215 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 1.45 (6H, d, $J=6.1$ Hz), 3.86 (3H, s), 3.92 (3H, s), 4.65 (1H, septet, $J=6.1$ Hz), 5.64 (1H, s), 6.70 (1H, s), 6.85 (2H, d, $J=8.8$ Hz), 7.36 (2H, d, $J=8.8$ Hz), 9.82 (1H, s). IR (KBr) cm^{-1} : 3230, 1612, 1578. EI-MS m/z : 388 (M^+), 373, 331.

Reference and Notes

- 1) Komiya T., Tsukui M., Oshio H., *Chem. Pharm. Bull.*, **23**, 1387–1388 (1975).
- 2) Chu C., Tseng T., Hwang J., Chou F., Wang C., *Toxicology*, **73**, 263–268 (1999).
- 3) Nakata K., Sakaguchi H., *Wakan Iyaku Gakkaishi*, **1**, 222–229 (1984).
- 4) Xu Q., Mori H., Sakamoto O., Koda A., Nishioka I., Ogawa Y., Hosaka K., *Wakan Iyaku Gakkaishi*, **6**, 1–7 (1989).
- 5) Yamaguchi T., Sato S., Chin M., Ageta T., Nakajima K., Mitsuhashi

- H., *Wakan Iyaku Gakkaishi*, **5**, 374—375 (1988).
- 6) Igarashi Y., Yamaguchi T., Ogawa Y., Tomita M., Hayashi H., Sato T., Hosaka K., WO 9209594 (1992).
- 7) Takeno H., Hashimoto M., *J. Chem. Soc., Chem. Commun.*, **1981**, 474—475 (1981).
- 8) Okutani T., Kawakita K., Aki O., Morita K., *Heterocycles*, **6**, 1581—1586 (1977).
- 9) Although the acetophenone **7** was a commercially available material, it was synthesized by the following method. Friedel–Crafts reaction of 3,4,5-trimethoxyphenol and acetic anhydride using zinc chloride in nitromethane followed by hydrolysis using sodium hydrogencarbonate to give **7** in 91% yield.
- 10) Gammil R. B., *Synthesis*, **1979**, 901—903 (1979).
- 11) Zhu J., Chastanet J., Beugelmans R., *Synth. Commun.*, **26**, 2479—2486 (1996).
- 12) Keumi T., Morita T., Ozawa Y., Kitajima H., *Bull. Chem. Soc. Jpn.*, **62**, 599—601 (1989).
- 13) Sugita Y., Yokoe I., *Heterocycles*, **43**, 2503—2511 (1996).