

Green Synthesis of Schiff Bases Derived from 4,5-Diazafluorene-9-one in Aqueous Medium Under Microwave Irradiation

HUI CANG*, YU LU, WENYAN SHI, JINGLING SHAO and QI XU

Chemical and Biological Engineering College, Yancheng Institute of Technology, Yancheng 224051, Jiangsu Province, P.R. China

*Corresponding author: Fax: +86 515 88298618; Tel: +86 515 88298611; E-mail: canghui@ycit.edu.cn

(Received: 4 April 2012;

Accepted: 24 December 2012)

AJC-12595

A series of Schiff bases were successfully synthesized with satisfactory yield from various amines and 4,5-diazafluorene-9-one in water by using polystyrene sulfonic acid as a catalyst under microwave irradiation.

Key Words: Schiff base, 4,5-Diazafluorene-9-one, Microwave irradiation, Aqueous medium.


INTRODUCTION

4,5-Diazafluorene-9-one and their derivatives are also strong bidentate chelating and bridging agents¹, which have been widely explored due to their easy accessibility, easy modification and relative air stability². Synthesis of Schiff bases containing a bipyridine type metal receptor³, has attracted considerable attention due to their many applications in the fields of catalyst⁴, conducting and photoresponsive materials⁵ and anion sensor⁶ *etc.* But, higher temperatures, increased reaction times, plenty of organic solvent and the use of a catalyst were generally required in previous synthesis of Schiff bases from 4,5-diazafluorene-9-one^{7,8}.

In recent years, microwave irradiation was demonstrated to be a valuable tool in organic synthesis resulting in faster and cleaner reactions that sometimes exhibit different reactivities due to specific microwave absorption⁹. From an environmental view point, we report herein, for the first time, the reaction (**Scheme-I**) can be successfully accomplished with satisfactory yield in water by using polystyrene sulfonic acid (PSSA) as a catalyst under microwave irradiation (MW), which is a rapid and environmentally friendly preparation of schiff bases from 4,5-diazafluorene-9-one compared to conventional heating method.

EXPERIMENTAL

All reagents and solvents were purchased commercially as AC grade and were used without further purification unless noted. The molecular weight of polystyrene sulfonic acid is 70,000. Organic solvents were dried over 4 Å activated molecular sieves prior to use. 4,5-diazafluorene-9-one was obtained as light yellow needle by oxidation of 1,10-phenanthroline



Scheme-I: Synthesis of schiff bases derived from 4,5-diazafluorene-9-one

with KMnO₄ in a KOH solution following a literature method¹⁰. The microwave reaction was carried out in the WF-4000M closed type microwave synthesis system with the refluxing equipment. Melting points were determined in open capillaries and uncorrected. ¹H NMR spectra were obtained using Bruker DRX300 spectrometer with TMS. The IR spectra were determined as potassium bromide pellet on a Bruker Equinox55 FT-IR spectrophotometer. UV-VIS spectra were performed with a UV-2501PC spectrophotometer. Mass spectra and element analysis were done with LC-MSD-Trap-SL and Elementar Vario EL III.

General procedure for preparation of L₁-L₁₁: All the Schiff bases derived from 4,5-diazafluorene-9-one were synthesized by modification of a literature procedure¹¹. A mixture of 4,5-diazafluorene-9-one (455 mg, 2.5 mmol) and amines (3 mmol) were dissolved in 20 % polystyrene sulfonic acid solution in water (5 mL) and then were exposed to microwave irradiation (700 W, 100 °C) for 7 min. The desired

precipitate was filtered after cooling, washed with ethanol 3 times and dried in vacuum oven at 50 °C. The results of yield, melt point and UV-visible spectra were listed in Table-1.

9-Imine-4,5-diazafluorene (L₁): ¹H NMR (CDCl₃, 300 MHz): δ 8.82 (dd, 2H, *J*₁ = 5.1, *J*₂ = 1.5 Hz, pyridyl H *ortho* to N), 8.05 (dd, 2H, *J*₁ = 5.1, *J*₂ = 1.5 Hz, pyridyl H *para* to N), 7.38 (t, 2H, *J*₁ = 7.6, *J*₂ = 5.1 Hz, pyridyl H *meta* to N), 5.45 (s, 1H, =NH). IR (KBr, $\nu_{\max}$, cm⁻¹): 3314, 3058, 1655, 1590, 1555, 1420, 1200, 801, 726. Elemental analysis calcd. (%) for C₁₁H₇N₃: C, 72.92; H, 3.89; N, 23.19. Found (%): C, 72.89; H, 3.92; N, 23.15. MS: *m/z* = 181.01 (Calcd.: 181.06).

9-Hydrazone-4,5-diazafluorene (L₂): ¹H NMR (CDCl₃, 300 MHz): δ 8.75 (dd, 1H, *J*₁ = 5.1, *J*₂ = 1.6 Hz, pyridyl H *ortho* to N), 8.65 (dd, 1H, *J*₁ = 5.0, *J*₂ = 1.6 Hz, pyridyl H *ortho* to N), 8.27 (dd, 1H, *J*₁ = 7.7, *J*₂ = 1.6 Hz, pyridyl H *para* to N), 8.16 (dd, 1H, *J*₁ = 7.8, *J*₂ = 1.6 Hz, pyridyl H *para* to N), 7.50 (t, 1H, *J*₁ = 7.7, *J*₂ = 5.1 Hz, pyridyl H *meta* to N), 7.39 (t, 1H, *J*₁ = 7.7, *J*₂ = 5.0 Hz, pyridyl H *meta* to N), 6.65 (s, 2H, -NH₂). IR (KBr, $\nu_{\max}$, cm⁻¹): 3376, 3320, 3058, 1646, 1593, 1561, 1206, 799, 723. Elemental analysis calcd. (%) for C₁₂H₉N₃: C, 67.34; H, 4.11; N, 28.55. Found (%): C, 67.30; H, 4.10; N, 28.52. MS: *m/z* = 196.01 (calcd.: 196.06).

9-Oxime-4,5-diazafluorene (L₃): Sodium carbonate (0.016 g, 1.5 mmol) was used as a neutralizer in the reaction. ¹H NMR (CDCl₃, 300 MHz): δ 13.14 (s, 1H, -OH), 8.73 (dd, 1H, *J*₁ = 5.1, *J*₂ = 1.6 Hz, pyridyl H *ortho* to N), 8.67 (dd, 1H, *J*₁ = 5.0, *J*₂ = 1.6 Hz, pyridyl H *ortho* to N), 8.27 (dd, 1H, *J*₁ = 7.7, *J*₂ = 1.6 Hz, pyridyl H *para* to N), 8.17 (dd, 1H, *J*₁ = 7.8, *J*₂ = 1.6 Hz, pyridyl H *para* to N), 7.53 (t, 1H, *J*₁ = 7.7, *J*₂ = 5.1 Hz, pyridyl H *meta* to N), 7.46 (t, 1H, *J*₁ = 7.7, *J*₂ = 5.0 Hz, pyridyl H *meta* to N). IR (KBr, $\nu_{\max}$, cm⁻¹): 3141, 3036, 1627, 1593, 1564, 1495, 1399, 1008, 949, 748. Elemental analysis calcd. (%) for C₁₁H₇N₃O: C, 67.00; H, 3.58; N, 21.31. Found (%): C, 69.92; H, 3.56; N, 21.36. MS: *m/z* = 197.10 (calcd.: 197.06).

9-Anilino-4,5-diazafluorene (L₄): ¹H NMR (CDCl₃, 300 MHz): δ 8.81 (dd, 1H, *J*₁ = 4.9, *J*₂ = 1.5 Hz, pyridyl H *ortho* to N), 8.65 (dd, 1H, *J*₁ = 4.9, *J*₂ = 1.6 Hz, pyridyl H *ortho* to N), 8.26 (dd, 1H, *J*₁ = 7.8, *J*₂ = 1.6 Hz, pyridyl H *para* to N), 7.43 (m, 3H, pyridyl H *meta* to N and Ar-H), 7.28 (t, 1H, *J*₁ = 7.7, *J*₂ = 5.1 Hz, pyridyl H *meta* to N), 7.02 (m, 3H, Ar-H), 6.87 (dd, 1H, *J*₁ = 7.8, *J*₂ = 1.5 Hz, pyridyl H *para* to N). IR (KBr, $\nu_{\max}$, cm⁻¹): 3045, 1656, 1592, 1561, 1401, 748, 710. Elemental analysis calcd. (%) for C₁₇H₁₁N₃: C, 79.36; H, 4.31; N, 16.33.

Found (%): C, 79.30; H, 4.35; N, 16.36. MS: *m/z* = 257.13 (calcd.: 257.10).

9-(4-Methylanilino)-4,5-diazafluorene (L₅): ¹H NMR (CDCl₃, 300 MHz): δ 8.80 (dd, 1H, *J*₁ = 5.0, *J*₂ = 1.6 Hz, pyridyl H *ortho* to N), 8.64 (dd, 1H, *J*₁ = 4.9, *J*₂ = 1.6 Hz, pyridyl H *ortho* to N), 8.25 (dd, 1H, *J*₁ = 7.6, *J*₂ = 1.5 Hz, pyridyl H *para* to N), 7.39 (t, 1H, *J*₁ = 7.6, *J*₂ = 5.1 Hz, pyridyl H *meta* to N), 7.25 (m, 2H, pyridyl H *meta* to N and Ar-H), 6.98 (m, 2H, pyridyl H *para* to N and Ar-H), 6.90 (d, 2H, *J* = 8.2 Hz, Ar-H), 2.43 (s, 3H, -CH₃). IR (KBr, $\nu_{\max}$, cm⁻¹): 3046, 2954, 2869, 1642, 1590, 1557, 1399, 752, 729. Elemental analysis calcd. (%) for C₁₈H₁₃N₃: C, 79.68; H, 4.83; N, 15.49. Found (%): C, 79.62; H, 4.85; N, 15.46. MS: *m/z* = 271.13 (calcd.: 271.11).

9-(4-Methoxyanilino)-4,5-diazafluorene (L₆): ¹H NMR (CDCl₃, 300 MHz): δ 8.80 (dd, 1H, *J*₁ = 5.0, *J*₂ = 1.6 Hz, pyridyl H *ortho* to N), 8.65 (dd, 1H, *J*₁ = 5.0, *J*₂ = 1.6 Hz, pyridyl H *ortho* to N), 8.23 (dd, 1H, *J*₁ = 7.7, *J*₂ = 1.5 Hz, pyridyl H *para* to N), 7.38 (t, 1H, *J*₁ = 7.7, *J*₂ = 5.0 Hz, pyridyl H *meta* to N), 7.09 (dd, 1H, *J*₁ = 7.8, *J*₂ = 1.7 Hz, pyridyl H *para* to N), 6.98 (m, 5H, pyridyl H *meta* to N and Ar-H), 3.92 (s, 3H, -OCH₃). IR (KBr, $\nu_{\max}$, cm⁻¹): 3043, 2999, 2837, 1634, 1590, 1559, 1497, 1400, 1234, 1023, 845, 757. Elemental analysis calcd. (%) for C₁₈H₁₃N₃O: C, 75.25; H, 4.56; N, 13.95. Found (%): C, 75.20; H, 4.58; N, 13.97. MS: *m/z* = 287.07 (calcd.: 287.11).

9-(4-Hydroxyanilino)-4,5-diazafluorene (L₇): ¹H NMR (CDCl₃, 300 MHz): δ 9.53 (s, 1H, -OH), 8.79 (dd, 1H, *J*₁ = 4.9, *J*₂ = 1.5 Hz, pyridyl H *ortho* to N), 8.67 (dd, 1H, *J*₁ = 4.9, *J*₂ = 1.5 Hz, pyridyl H *ortho* to N), 8.27 (dd, 1H, *J*₁ = 7.6, *J*₂ = 1.5 Hz, pyridyl H *para* to N), 7.54 (t, 1H, *J*₁ = 7.6, *J*₂ = 4.9 Hz, pyridyl H *meta* to N), 7.27 (t, 1H, *J*₁ = 7.6, *J*₂ = 4.9 Hz, pyridyl H *meta* to N), 7.13 (dd, 1H, *J*₁ = 7.6, *J*₂ = 1.5 Hz, pyridyl H *para* to N), 6.90 (m, 4H, Ar-H). IR (KBr, $\nu_{\max}$, cm⁻¹): 3144, 3037, 1638, 1592, 1561, 1504, 1401, 1276, 1230, 1165, 758, 708. Elemental analysis calcd. (%) for C₁₇H₁₁N₃O: C, 74.71; H, 4.06; N, 15.38. Found (%): C, 74.65; H, 4.00; N, 15.42. MS: *m/z* = 272.85 (calcd.: 273.09).

9-(4-Carboxyanilino)-4,5-diazafluorene (L₈): ¹H NMR (CDCl₃, 300 MHz): δ 12.93 (s, 1H, -COOH), 8.82 (dd, 1H, *J*₁ = 4.9, *J*₂ = 1.5 Hz, pyridyl H *ortho* to N), 8.68 (dd, 1H, *J*₁ = 4.9, *J*₂ = 1.5 Hz, pyridyl H *ortho* to N), 8.29 (dd, 1H, *J*₁ = 7.7, *J*₂ = 1.5 Hz, pyridyl H *para* to N), 8.02 (d, 2H, *J* = 8.6 Hz, Ar-H), 7.56 (t, 1H, *J*₁ = 7.7, *J*₂ = 4.9 Hz, pyridyl H *meta* to N),

TABLE-1
RESULTS OF YIELD, MELTING POINT AND UV-VISIBLE SPECTRA OF L₁₋₁₁

No.	Yield (%)	m.p. (°C)	UV-VIS $\lambda_{\max}$ (> 235 nm, $\epsilon \times 10^4$) ^a
L ₁	78	195-196	249 (2.40), 304 (0.99), 317 (1.01).
L ₂	82	208-210	245 (2.45), 303 (0.98), 313 (1.13).
L ₃	80	>300	242 (2.41), 308 (1.00), 315 (0.96).
L ₄	87	188-189	240 (2.40), 302 (0.95), 315 (1.02).
L ₅	89	225-226	244 (2.56), 302 (0.98), 314 (1.13), 416 (0.20).
L ₆	85	202-203.	252 (3.80), 300 (1.44), 314 (1.04), 435 (0.29).
L ₇	90	>300	244 (2.69), 302 (1.15), 314 (0.97), 433 (0.26)
L ₈	81	>300	245 (2.73), 265 (2.51), 302 (1.01), 314 (0.91), 394 (0.16)
L ₉	93	216-217	246 (3.24), 302 (1.15), 314 (1.01), 405 (0.20)
L ₁₀	85	188-189	242 (2.03), 303 (1.22), 314 (1.18), 396 (2.50)
L ₁₁	83	>300	CH ₃ CN: 303 (0.58), 385 (1.30), 406 (1.27), 562 (0.32)

^aL₁-L₁₀ in CHCl₃ and L₁₁ in CH₃CN.

7.24 (t, 1H, $J_1 = 7.7$, $J_2 = 4.9$ Hz, pyridyl H *meta* to N), 7.17 (d, 2H, $J = 8.6$ Hz, Ar-H), 6.83 (dd, 1H, $J_1 = 7.7$, $J_2 = 1.5$ Hz, pyridyl H *para* to N). IR (KBr, $\nu_{\max}$, cm^{-1}): 3098, 3032, 1702, 1702, 1663, 1598, 1566, 1398, 1259, 1164, 752, 723. Elemental analysis: calcd. (%) for $\text{C}_{18}\text{H}_{11}\text{N}_3\text{O}_2$: C, 71.75; H, 3.68; N, 13.95. Found (%): C, 71.70; H, 3.63; N, 13.89. MS: $m/z = 300.99$ (calcd.: 301.09).

9-(4-Chloroanilino)-4,5-diazafluorene (L_9): ^1H NMR (CDCl_3 , 300 MHz): δ 8.83 (dd, 1H, $J_1 = 5.0$, $J_2 = 1.5$ Hz, pyridyl H *ortho* to N), 8.70 (dd, 1H, $J_1 = 5.0$, $J_2 = 1.5$ Hz, pyridyl H *ortho* to N), 8.26 (dd, 1H, $J_1 = 7.6$, $J_2 = 1.5$ Hz, pyridyl H *para* to N), 7.43 (m, 3H, pyridyl H *meta* to N and Ar-H), 7.07 (t, 1H, $J_1 = 7.7$, $J_2 = 5.0$ Hz, pyridyl H *meta* to N), 7.03 (dd, 1H, $J_1 = 7.8$, $J_2 = 1.5$ Hz, pyridyl H *para* to N), 6.95 (d, 2H, $J = 8.2$ Hz, Ar-H). IR (KBr, $\nu_{\max}$, cm^{-1}): 3046, 1656, 1592, 1561, 1400, 1087, 839, 749, 730. Elemental analysis calcd. (%) for $\text{C}_{18}\text{H}_{10}\text{N}_3\text{Cl}$: C, 69.99; H, 3.45; N, 14.40. Found (%): C, 69.93; H, 3.46; N, 14.45. MS: $m/z = 291.01$ (calcd.: 291.06).

9-Phenylhydrazone-4,5-diazafluorene (L_{10}): ^1H NMR (DMSO , 300 MHz): δ 10.72 (s, 1H, -NH), 8.90 (d, 1H, $J = 7.7$ Hz, pyridyl H *para* to N), 8.70 (dd, 2H, $J = 4.6$ Hz, pyridyl H *ortho* to N), 8.29 (dd, 1H, $J = 7.7$ Hz, pyridyl H *para* to N), 7.59 (m, 3H, pyridyl H *meta* to N and Ar-H), 7.45 (m, 3H, pyridyl H *meta* to N and Ar-H), 7.10 (m, 1H, Ar-H). IR (KBr, $\nu_{\max}$, cm^{-1}): 3385, 3047, 1652, 1591, 1560, 1404, 750, 713. Elemental analysis: calcd. (%) for $\text{C}_{17}\text{H}_{12}\text{N}_4$: C, 74.98; H, 4.44; N, 20.58. Found (%): C, 74.92; H, 4.41; N, 20.64. MS: $m/z = 272.14$ (calcd.: 272.11).

9-(2,4-Dinitrophenylhydrazone)-4,5-diazafluorene (L_{11}): ^1H NMR (DMSO , 300 MHz): δ 12.01 (s, 1H, -NH), 8.96 (s, 1H, Ar-H), 8.84 (d, 1H, $J = 5.0$ Hz, pyridyl H *ortho* to N), 8.76 (d, 1H, $J = 4.9$ Hz, pyridyl H *ortho* to N), 8.59 (d, 1H, $J = 7.8$ Hz, pyridyl H *para* to N), 8.53 (d, 1H, $J = 7.8$ Hz, pyridyl H *para* to N), 8.42 (m, 2H, Ar-H), 7.73 (t, 1H, $J_1 = 7.7$, $J_2 = 5.0$ Hz, pyridyl H *meta* to N), 7.56 (t, 1H, $J_1 = 7.8$, $J_2 = 5.0$ Hz, pyridyl H *meta* to N). IR (KBr, $\nu_{\max}$, cm^{-1}): 3414,

3100, 1616, 1594, 1557, 1496, 1404, 1334, 1095, 754, 740, $\lambda_{\max}$ (> 235 nm, ϵ): Elemental analysis: calcd. (%) for $\text{C}_{17}\text{H}_{10}\text{N}_6\text{O}_4$: C, 56.36; H, 2.78; N, 23.20. Found (%): C, 56.32; H, 2.74; N, 23.25. MS: $m/z = 362.14$ (calcd.: 362.08).

Conclusion

This work has demonstrated an efficient and general microwave irradiation protocol for the synthesis of schiff bases derived from 4,5-diazafluorene-9-one under microwave irradiation using polystyrene sulfonic acid as a catalyst in aqueous medium. The simple product isolation *via* filtration precludes the use of an organic solvent thus culminating in an environmentally benign aqueous protocol for the synthesis of schiff bases derived from 4,5-diazafluorene-9-one.

ACKNOWLEDGEMENTS

This work was funded by the Applied Basic Research Project of Yancheng Institute of Technology of Chian, Jiangsu (Grant No. XKR2011005).

REFERENCES

1. D.A.V. Griend, D.K. Bediako, M.J. Devries, A.D. Nathan and P.H. Lee, *Inorg. Chem.*, **47**, 656 (2008).
2. Y.X. Wang, W. Perez, G.Y. Zheng, D.P. Rillema and C.L. Huber, *Inorg. Chem.*, **37**, 2227 (1998).
3. A.L. Eckermann, K.D. Barker, M.R. Hartings, M.A. Ratner and T.J. Meade, *J. Am. Chem. Soc.*, **127**, 11880 (2005).
4. J. Hou, W.H. Sun, D. Zhang, L. Chen, W. Li, D. Zhao and H. Song, *J. Mol. Catal. A: Chem.*, **231**, 221 (2005).
5. C.Y. Duan, Z.L. Lu, X.Z. You, Z.Y. Zhou, T.C.W. Mak, Q. Luo and J.Y. Zhou, *Polyhedron*, **17**, 4131 (1998).
6. Z.H. Lin, Y.G. Zhao, C.Y. Duan, B.G. Zhang and Z.P. Bai, *J. Chem. Soc. Dalton Trans.*, 3678 (2006).
7. H. Cang, S.Q. Wang, W.Z. Yang and J.T. Wang, *J. Chem. Res.*, **2009**, 101 (2009).
8. Y. Wang and D.P. Rillema, *Tetrahedron*, **53**, 12377 (1997).
9. C.O. Kappe, *Angew. Chem. Int. Ed.*, **43**, 6250 (2004).
10. L.J. Henderson, F.R. Fronczek Jr. and W.R. Cherry, *J. Am. Chem. Soc.*, **106**, 5876 (1984).
11. V. Polshettiwar and R.S. Varma, *Tetrahedron Lett.*, **48**, 5649 (2007).