



## OpdA, a bacterial organophosphorus hydrolase, prevents lethality in rats after poisoning with highly toxic organophosphorus pesticides

Steven B. Bird<sup>a,\*</sup>, Tara D. Sutherland<sup>b</sup>, Chip Gresham<sup>c</sup>, John Oakeshott<sup>b</sup>, Colin Scott<sup>b</sup>, Michael Eddleston<sup>d</sup>

<sup>a</sup> Department of Emergency Medicine, Division of Toxicology, University of Massachusetts Medical School, 55 Lake Avenue North, Worcester, MA 01655, USA

<sup>b</sup> CSIRO Entomology, Canberra, ACT 2601, Australia

<sup>c</sup> Department of Medical Toxicology, Banner Good Samaritan Medical Center, Phoenix, AZ, USA

<sup>d</sup> Clinical Pharmacology Unit, University of Edinburgh and Scottish Poisons Information Bureau, Royal Infirmary of Edinburgh, UK

### ARTICLE INFO

#### Article history:

Received 30 January 2008

Received in revised form 13 February 2008

Accepted 14 February 2008

Available online 10 March 2008

#### Keywords:

Organophosphorus (OP)

Hydrolase

Acetylcholinesterase (AChE)

Pralidoxime (2-PAM)

### ABSTRACT

Organophosphorus (OP) pesticides poison more than 3,000,000 people every year in the developing world, mostly through intentional self-poisoning. Advances in medical therapy for OP poisoning have lagged, and current treatment is not highly effective with mortality of up to 40% in even the most advanced Western medical facilities. Administration of a broadly active bacterial OP hydrolase to patients in order to hydrolyze OPs in circulation might allow current therapies to be more effective. The objective of this work was to evaluate the efficacy of a new recombinant bacterial OP hydrolase (OpdA), cloned from *Agrobacterium radiobacter*, in rat models of two chemically distinct but highly toxic and rapidly acting OP pesticides: dichlorvos and parathion. Without OpdA treatment, median time to death in rats poisoned with  $3 \times \text{LD}_{50}$  of dichlorvos or parathion was 6 min and 25.5 min, respectively. Administration of a single dose of OpdA immediately after dichlorvos resulted in 100% survival at 24 h, with no additional antidotal therapy. After parathion poisoning, OpdA alone caused only a delay to death. However, an additional two doses of OpdA resulted in 62.5% survival at 24 h after parathion poisoning. In combination with pralidoxime therapy, a single dose of OpdA increased survival to 75% after parathion poisoning. Our results demonstrate that OpdA is able to improve survival after poisoning by two chemically distinct and highly toxic OP pesticides.

© 2008 Elsevier Ireland Ltd. All rights reserved.

### 1. Introduction

Occupational exposure and intentional self-poisoning with organophosphorus (OP) pesticides are major global health problems (Jeyaratnam, 1990; Van der Hoek et al., 1998). The World Health Organization estimates that as many as 3,000,000 people per year are poisoned by pesticides; many are due to OP pesticides, resulting in around 200,000 deaths (Jeyaratnam). Although the greatest burden is borne by the developing world (Buckley et al., 2004; Eddleston and Phillips, 2004), it is also an important cause of fatal self-poisoning in developed countries (Bruyndonckx et al., 2002). Highly toxic and widely available OPs such as parathion also pose a threat to municipal water supplies from intentional or unintentional contamination.

\* Corresponding author. Tel.: +1 508 421 1422; fax: +1 508 421 1490.

E-mail addresses: [birds@ummhc.org](mailto:birds@ummhc.org) (S.B. Bird), [tara.sutherland@csiro.au](mailto:tara.sutherland@csiro.au) (T.D. Sutherland), [chipgresham@yahoo.com](mailto:chipgresham@yahoo.com) (C. Gresham), [john@oakeshott.csiro.au](mailto:john@oakeshott.csiro.au) (J. Oakeshott), [colin.scott@csiro.au](mailto:colin.scott@csiro.au) (C. Scott), [edlestonm@yahoo.com](mailto:edlestonm@yahoo.com) (M. Eddleston).

OPs inhibit acetylcholinesterase (AChE, EC 3.1.1.7), resulting in overstimulation at cholinergic synapses. Clinical management of moderate and severe poisoning is difficult, requiring prolonged intensive care and use of large doses of atropine, oxime cholinesterase reactivators, and benzodiazepines (Eddleston et al., 2007). However, these therapies are insufficient, not always available in the developing world, and do not prevent the post-poisoning neurocognitive dysfunction that is common with severe poisonings (Dunn and Sidell, 1989). Overall mortality after OP poisoning in the developing world is as high as 25%, and in the most sophisticated Western hospitals mortality is as high as 40% (Eyer et al., 2003). This overall difference in mortality between the developed and developing world is due to the vast numbers of poisoned patients in the agricultural areas of the developing world, the majority of whom are not critically ill. However, OP pesticide poisoning is uncommon in the developed world, and patients who ingest OP pesticides for self-harm typically have substantial ingestions and are more likely to be critically ill.

OpdA is a bacterial enzyme capable of hydrolyzing a wide variety of OP pesticides *in vitro* (Fig. 1) (Yang et al., 2003). The addition of an




Fig. 1. Hydrolysis of dichlorvos (A) and parathion (B) by OpdA.

OP-degrading enzyme should improve the clinical results obtained with standard therapies by decreasing the concentration of OP pesticides in circulation. Clinical use of an enzyme with a broad range of substrates would be useful in the event of poisoning with many OPs, even when the identity of the pesticide is unknown.

We sought to determine the *in vivo* efficacy of OpdA in rat models of two chemically distinct and highly toxic OP pesticides: dichlorvos and parathion. Demonstration of OpdA's effectiveness should provide the impetus for further development of this enzyme for eventual use in humans.

## 2. Methods

All animals were acquired and cared for in accordance with the guidelines published in the *Guide for the Care and Use of Laboratory Animals* (National Institutes of Health Publications No. 85-23, Revised 1985). The Institutional Animal Care and Use Committee of the University of Massachusetts Medical School approved the study protocol.

### 2.1. Subjects

Male Wistar rats weighing  $250 \pm 50$  g were obtained from Charles River Laboratories (Wilmington, MA, USA). Animals were housed in pairs, maintained on 12-h light:12-h dark cycle and provided food and water *ad libitum* except for 2 h prior to experimentation.

### 2.2. Expression of highly active OpdA

OpdA is a metalloenzyme that is catalytically active with a variety of metal ions ( $\text{Zn}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Co}^{2+}$  or  $\text{Cd}^{2+}$ ) with highest activity toward organophosphate insecticides when  $\text{Co}^{2+}$  is present in the active site (Jackson et al., 2006; Yang et al., 2003). Therefore, highly active OpdA was prepared by expressing the enzyme in the presence of cobalt as follows. OpdA discovery and cloning are described elsewhere (Horne et al., 2002). The OpdA gene in the plasmid pCy76-opdA (Yang et al., 2003) was transformed into electrocompetent *E. coli* DH5 $\alpha$  cells. A 20 ml starter culture in Luria–Bertani medium supplemented with 50  $\mu\text{g}/\text{ml}$  ampicillin (LB Amp) was inoculated then incubated at 30 °C for 8 h. This inoculum was used to inoculate 1 l of Terrific broth medium (24 g/l of yeast extract, 12 g/l of tryptone, 100 mM  $\text{K}_2\text{HPO}_4/\text{KH}_2\text{PO}_4$ , pH 7.0, 0.4% glycerol, and 50  $\mu\text{g}/\text{ml}$  ampicillin) supplemented with 1 mM  $\text{CoCl}_2$ . After 30 h at 37 °C, cells were harvested and resuspended in 50 mM HEPES buffer (pH 8.0). Cells were lysed using a French Press, and cell debris removed by centrifugation at  $30,000 \times g$  for 30 min. The soluble fraction was passed through a DEAE Fractogel column that does not bind OpdA. Following 12-h dialysis against 50 mM HEPES (pH 7.0) the protein solution was loaded on to a Sulphopropyl-Sepharose column. Protein was using a linear gradient from 0 to 1 M NaCl, with OpdA eluting at around 150 mM NaCl. SDS/PAGE analysis of the eluted OpdA indicated a purity of greater than 95%. The protein was stored in 50 mM HEPES (pH 7.0)/150 mM NaCl at 4 °C until required.

### 2.3. Ethyl parathion and dichlorvos activity assays with purified OpdA

Ethyl parathion OpdA assays were conducted in duplicate in assay buffer (50 mM HEPES, 10% methanol, 1 mM  $\text{CoCl}_2$ , pH 7.0) with an ethyl parathion range of 0–100  $\mu\text{M}$  and 2 nM of purified OpdA. The initial rates of the reactions were determined by measuring the change in absorbance at 412 nM over time, until 10% of the substrate had been converted. The rate of conversion was calculated from a standard

curve of 0–10  $\mu\text{M}$  nitrophenol (the ethyl parathion hydrolysis product) measured at 412 nM (using a Molecular Devices SpectraMax 190). The kinetic parameters for the hydrolysis of ethyl parathion by OpdA were estimated using hyperbolic regression (using the Hyper32.exe enzyme kinetic analysis software; a freeware package). The  $k_{\text{cat}}$  of OpdA with parathion was calculated to be  $1500 \text{ s}^{-1}$ , and the  $K_m$  was estimated at 1  $\mu\text{M}$ .

Dichlorvos assays were conducted in duplicate in assay buffer with 0–1 mM dichlorvos and 19 nM OpdA. The hydrolysis products were detected by mass spectrometry (MS) using an Agilent G1969 LC/MS TOF after filtration through a liquid chromatography (LC) guard column at a flow rate of  $1 \text{ ml min}^{-1}$  and 80% acetonitrile, 0.002% formic acid. A 5  $\mu\text{l}$  sample was analyzed by MS, with the fragmentor set at 120 V. The mass spectral fragmentation patterns of authentic dichlorvos standards (Sigma–Aldrich, Castle Hill, New South Wales, Australia) were used to validate the identity of the substrates. The assay reactions were assayed at 3-min intervals for 12 min at each substrate concentration. The kinetic parameters for this reaction were estimated using hyperbolic regression (as above) giving a  $K_m$  of 183  $\mu\text{M}$  and  $k_{\text{cat}}$  of  $149 \text{ s}^{-1}$ .

### 2.4. Poisoning models

Animals were briefly anesthetized with isoflurane while a 24-gauge intravenous lateral tail vein catheter was placed. Immediately upon awakening, 3 times the oral  $\text{LD}_{50}$  for parathion ( $\text{LD}_{50} = 6 \text{ mg/kg}$ ) (Sigma–Aldrich, St. Louis, MO, USA) or dichlorvos ( $\text{LD}_{50} = 50 \text{ mg/kg}$ ) (Sigma–Aldrich) suspended in peanut oil was given by gavage feeding tube in a volume of 1.5 ml/kg. Three times the  $\text{LD}_{50}$  of the pesticides were used in order to mimic severe human poisoning and to assure that all, or nearly all, control animals would die, thereby decreasing the number of animals needed to demonstrate statistical significance. All injections were given via a 24-gauge catheter (Surflo catheter, Terumo Corporation, Somerset, NJ, USA) through a lateral tail vein in volume of 0.5 ml. Nitrile gloves and chemical safety goggles were worn when working with the OP pesticides or handling animals after poisoning. Consumable supplies were soaked in a dilute OpdA solution in order to hydrolyze any residual pesticide before disposal in a biohazards container for collection by the institutional Environmental Health and Safety department. Animal carcasses were burned in an incinerator equipped with an afterburner and scrubber.


### 2.5. Data analysis

Blinded outcomes of interest were survival to 4 and 24 h. Grouped survival data were compared using a two-tailed Fisher's exact test. For 80% power to detect a 50% reduction in mortality, assuming an alpha of 0.05, 8 animals per group were required. For all analyses, a *p* value of less than 0.05 was considered significant. All statistical analyses were performed with GraphPad Prism software version 4 for Mac (GraphPad Software, Inc., San Diego, CA, USA).

## 3. Results

### 3.1. Safety of repeated doses of OpdA in rats

Since OpdA had not previously been administered to animals, we first examined whether the hydrolase would elicit severe allergic reactions in the rat at OpdA doses likely to be needed in efficacy studies. 0.5 mg of OpdA (a 10-fold excess of OpdA estimated for effective hydrolysis of dichlorvos based on extrapolations from *in vitro* kinetic data) was injected into a tail vein of four rats once a



**Fig. 2.** Acute mortality after poisoning with  $3 \times \text{LD}_{50}$  of dichlorvos or parathion. Survival after poisoning with dichlorvos (■) or parathion (▲) alone. Rats were given 0.5 ml of saline placebo IV concomitantly with poisoning by  $3 \times \text{LD}_{50}$  dichlorvos or parathion via gavage.

week for 4 weeks. Rats were continuously observed for 4 h after each injection, then regularly for 24 h. No evidence of an allergic reaction was observed in any rat, and all exhibited normal behavior and normal weight gain over the 4-week period. With no evidence of an allergic reaction, we proceeded to test the efficacy of OpdA against dichlorvos and parathion.


### 3.2. Dichlorvos and parathion oral poisoning models

To mimic human poisoning, we developed oral poisoning models for parathion and dichlorvos in the Wistar rat. To assure severe poisoning and at least a lethality of 90% in controls, all rats received  $3 \times \text{LD}_{50}$  parathion ( $\text{LD}_{50} \sim 6 \text{ mg/kg}$ ) or dichlorvos ( $\text{LD}_{50} \sim 50 \text{ mg/kg}$ ) via gavage feeding tube. The  $\text{LD}_{50}$ s for dichlorvos and parathion were determined from exhaustive literature reviews. All rats that received dichlorvos developed signs of cholinergic toxicity (muscle fasciculations and tremors; gait disturbances; salivation; and urination) within 3 min and died by 12 min (median time to death 6 min), while all rats that received parathion developed cholinergic signs within 20 min and died by 35 min (median time to death 25.5 min) (Fig. 2). This rapid onset of poisoning is consistent with clinical experience of human poisoning with these pesticides (Eyer et al., 2003; Peng et al., 2004).

### 3.3. Efficacy of OpdA versus dichlorvos and parathion

Because of the difference in speed of poisoning onset, we gave a single dose of 0.15 mg/kg OpdA (or an equal volume of 0.9% normal saline) intravenously immediately after giving dichlorvos and immediately or 10 min after giving parathion. We chose 0.15 mg/kg of OpdA in order to give a margin of error based upon an estimated minimum dose of 0.05 mg/kg OpdA derived from *in vitro* data with dichlorvos, while remaining within the range of larger OpdA doses given in the preliminary safety dosing studies. All dichlorvos-poisoned rats that received OpdA survived to 4 h and 24 h ( $p = 0.0002$  vs. placebo). Delaying OpdA administration to 3 min after dichlorvos administration resulted in complete loss of OpdA efficacy (Fig. 3A).

A single dose of OpdA either immediately or 10 min after poisoning had no survival effect on parathion-poisoned rats, with all dead at 4 h. However, repeat doses of 0.15 mg/kg OpdA at 45 min and 90 min after poisoning improved survival: all animals survived to 4 h after this repeated treatment with OpdA ( $p = 0.0002$  vs. placebo). Twenty-four hours after poisoning 5/8 animals were alive (OpdA vs. placebo group,  $p = 0.026$ ) (Fig. 3B).



**Fig. 3.** OpdA improves survival after poisoning with  $3 \times \text{LD}_{50}$  of (A) dichlorvos, or (B) parathion. (A) 24-h survival after poisoning with dichlorvos. Rats were given 0.15 mg/kg of OpdA or the same volume of saline placebo concomitantly with poisoning by  $3 \times \text{LD}_{50}$  dichlorvos via gavage. (■) dichlorvos plus saline placebo; (▲) dichlorvos plus single dose 0.15 mg/kg OpdA. Each group contained 8 rats. (B) 24-h survival after poisoning with parathion. Rats were given OpdA 0.15 mg/kg or placebo, with or without 2-PAM, after poisoning with  $3 \times \text{LD}_{50}$  parathion via gavage. (■) Parathion plus saline placebo; (×) parathion plus single dose 0.15 mg/kg OpdA; (▲) parathion plus 3 doses 0.15 mg/kg OpdA; (▼) parathion plus 2-PAM bolus and infusion of 30 mg/kg/h for 24 h; (◆) parathion plus single dose 0.15 mg/kg OpdA plus 2-PAM bolus and infusion of 30 mg/kg/h for 24 h. Each group contained 8 rats with the exception of the 24-h 2-PAM alone group, which contained 10 rats.

OpdA demonstrates prolonged efficacy against dichlorvos and parathion. The prolonged *in vivo* enzymatic activity of an OP hydrolyase is essential if it is to be used effectively against OPs, like parathion, that partition out of the blood and into adipose tissue and other physiologic compartments. To test the duration of OpdA's effect versus parathion and dichlorvos, 250 g male Wistar rats were given 1.5 mg/kg i.v. OpdA, followed by  $3 \times \text{LD}_{50}$  oral dichlorvos 30, 60, or 90 min later. All rats survived to 24 h and no animals exhibited any signs of OP toxicity ( $p = 0.0002$  vs. placebo). When these experiments were repeated using  $3 \times \text{LD}_{50}$  parathion, again all animals survived to 24 h ( $p = 0.0002$  vs. placebo). Lastly, a cohort of 4 rats was given 1.5 mg/kg of OpdA 180 min before parathion poisoning. Again, all rats survived to 24 h. Therefore, OpdA maintains clinically relevant enzymatic activity *in vivo* for several hours.

### 3.4. OpdA improves efficacy of pralidoxime versus parathion

High OP concentrations are thought to limit the effectiveness of oxime cholinesterase reactivators in OP poisoning (Eddleston et al., 2005; Sogorb et al., 2004). To address this issue we treated parathion-poisoned rats with either pralidoxime chloride (2-PAM) alone, or 2-PAM with OpdA. Rats received a bolus dose of 30 mg/kg

**Table 1**  
Physicochemical properties of the studied pesticides

OP pesticide	Chemistry			WHO toxicity class	Rat oral LD <sub>50</sub> (mg/kg) <sup>a</sup>	Human mortality (%)	Fat solubility ( $K_{ow}$ log $p$ ) <sup>b</sup>
Dichlorvos	Aliphatic	Dimethyl	Oxon	Ib	25–100	34	Low (1.43)
Parathion	Aromatic	Diethyl	Thion	Ia	2–13	40	High (3.83)

<sup>a</sup> Values vary according to source: Crop Protection Handbook 2003, 2nd WHO Classification of Pesticide Toxicity.<sup>b</sup> National Toxicology Program, Forest Stewardship Council Pesticides Policy 2007.**Table 2**  
Enzymatic activity of OpdA vs. OPs representative of different chemical classes

Enzyme	Chemistry of OP							
	Ethyl-aromatic thion (parathion)		Methyl-aromatic thion (methyl parathion)		Methyl-aromatic oxon (methyl paraoxon)		Methyl-aliphatic (dichlorvos)	
	$K_m$	$k_{cat}$	$K_m$	$k_{cat}$	$K_m$	$k_{cat}$	$K_m$	$k_{cat}$
OpdA	110	1500	100	1200	230	2500	183	149

 $K_m$  is expressed in  $\mu\text{M}$  and  $k_{cat}$  in  $\text{s}^{-1}$ .

2-PAM 10 min after parathion administration, followed by a continuous IV infusion of 30 mg/kg/h (standard 2-PAM doses used in previous OP rat studies) for 24 h. 2-PAM treatment alone resulted in 4 of 10 animals surviving to 4 h and 1 of 10 animals surviving to 24 h (24-h 2-PAM infusion vs. placebo,  $p=1.00$ ). After a single IV dose of 0.15 mg/kg OpdA administered concomitantly with the 2-PAM bolus and infusion, 8 of 8 animals survived to 4 h, and 6 of 8 animals survived to 24 h ( $p=0.007$  compared to single dose OpdA alone;  $p=0.013$  compared to 2-PAM alone).

#### 4. Discussion

Since their first description in 1946 (Mazur, 1946), several OP degrading enzymes have been isolated. These enzymes include phosphotriesterase (PTE, aka OPH) (Lewis et al., 1988), paraoxonase (PON) (Ortigoza-Ferado et al., 1984), DFPase (Ahmad and Forgash, 1976), sarinase (Adie, 1956), and OpdA. OpdA is a particularly good candidate enzyme for clinical applications, partly because it is a very efficient hydrolase, performing at near diffusion-limited rates towards its favored substrates *in vitro*. In addition, and unlike some of the other OP-degrading enzymes, OpdA is active against a wide range of phosphotriester OP pesticides, including substrates with a phosphoryl sulfur (rather than an oxygen) and substrates with both methoxy and ethoxy groups. One consequence of the breadth of OpdA's substrate range is that treatments based on OpdA have a high probability of success against OP poisoning even when the exact identity of the OP pesticide is unknown. Additionally, phosphoryl sulfur containing OP pesticides (such as parathion) can, in principle, be hydrolyzed before they are activated by P450s, so long as they are available to the enzyme in circulation.

As a test for broad-spectrum hydrolytic activity, we sought to test the efficacy and safety of OpdA in two very different rat models of severe oral OP poisoning. We chose parathion and dichlorvos because their different chemical structures result in variable requirements for bioactivation and differences in fat solubility (Table 1). However, both pesticides are highly toxic, fast acting, widely used globally; parathion and dichlorvos are the two most commonly used OPs in Chinese suicides, where an estimated 170,000 deaths occur annually from pesticide poisoning (Eddleston and Phillips, 2004; Phillips et al., 2002), and therefore highly clinically relevant.

We found no evidence of allergic reactions to this bacterial hydrolase in rats. Clinical use of another microbial enzyme, streptokinase, has shown that repeated administration is safe and effective within 5 days or after 1 year (Fears et al., 1992). In the acute OP pesticide-poisoning scenario, it is unlikely that OpdA would be

required after the first few hours or days. Therefore, theoretical concerns regarding prolonged re-exposure to a bacterial OP hydrolase such as OpdA are likely not applicable.

OpdA was surprisingly effective against dichlorvos and parathion, two quite different OP insecticides. *In vitro* enzymatic studies had previously suggested that OpdA would be poorly effective against dimethylated oxon OPs, such as dichlorvos (Horne et al., 2002). The  $K_{cat}$  of OpdA towards dichlorvos is  $149 \pm 26 \text{ s}^{-1}$  at room temperature, compared to  $\sim 1500 \text{ s}^{-1}$  towards parathion (Table 2). The similar clinical responses observed after dichlorvos and parathion poisoning suggest that *in vitro* catalytic activities above a certain threshold may not be important in determining the clinical effectiveness of an OP hydrolase.

Despite excellent efficacy against dichlorvos, a single dose of OpdA was not effective against parathion whether OpdA was given concomitantly with poisoning or 10 min later. Differences in pesticide toxicokinetics and need for bioactivation of the two OPs explain the differences in speed of poisoning onset and the effectiveness of OpdA: dichlorvos does not require bioactivation (because of its oxon group) and is therefore immediately toxic. In addition, dichlorvos is rapidly present at high concentrations in the blood due to its low distribution into fat, and is therefore available for hydrolysis by OpdA. Parathion, in contrast, requires bioactivation from a thion to oxon and distributes rapidly to fat. This latter characteristic decreases the amount available in the circulation for OpdA hydrolysis and provides a reservoir for subsequent leaching of parathion back into circulation. Three doses of OpdA at 45-min intervals were needed to counter parathion returning from the fat into the blood.

2-PAM is considered by most researchers and clinicians to be an integral part of therapy after OP poisoning (Eddleston et al., 2007). However, a 2-PAM bolus followed by continuous 24-h infusion was ineffective as a treatment for parathion poisoning. The addition of a single bolus dose of OpdA markedly increased the effectiveness of 2-PAM. This result suggests that early use of OpdA lowers the amount of pesticide that reaches the fat, improving 2-PAM's ability to reactivate AChE. The combination of OpdA and 2-PAM may well improve outcomes for other fat soluble OPs that have a long half-life, such as fenthion (Eddleston et al., 2005), that are also associated with severe poisonings in the developing world.

With dichlorvos poisoning, OpdA was required immediately to have a beneficial effect, which will make it difficult for it be useful clinically for very fast acting pesticides. However, it is possible that OpdA may be effective at later time points in poisoning with lower dose of OP pesticides. OpdA may also be effective in poisoning with slowly activated and poorly fat-soluble OPs, such as dimethoate, for which there is more time for OpdA administration.

Given the profound human health burden caused by OP pesticide poisoning, and potential terrorist use of OP pesticides, these studies should provide the impetus for accelerated research and human trials of broad spectrum, low-cost therapeutic OP hydrolases such as OpdA.

## Acknowledgments

Support for this research was provided by NIEHS grant K08 ES012897, the Emergency Medicine Foundation, and the Orphan Medical/Jazz Pharmaceuticals research award from the American College of Medical Toxicology. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIEHS.

## References

- Adie, P.A., 1956. The purification of sarinase from bovine plasma. *Can. J. Biochem. Physiol.* 34, 1091–1094.
- Ahmad, S., Forgash, A.J., 1976. Nonoxidative enzymes in the metabolism of insecticides. *Ann. Clin. Biochem.* 13, 141–164.
- Bruyndonckx, R.B., Meulemans, A.I., Sabbe, M.B., Kumar, A.A., Delooz, H.H., 2002. Fatal intentional poisoning cases admitted to the University Hospitals of Leuven Belgium from 1993 to 1996. *Eur. J. Emerg. Med.* 9, 238–243.
- Buckley, N.A., Roberts, D., Eddleston, M., 2004. Overcoming apathy in research on organophosphate poisoning. *BMJ* 329, 1231–1233.
- Dunn, M.A., Sidell, F.R., 1989. Progress in medical defense against nerve agents. *JAMA* 262, 649–652.
- Eddleston, M., Buckley, N.A., Eyer, P., Dawson, A.H., 2008. Management of acute organophosphorus pesticide poisoning. *Lancet* 371, 597–607.
- Eddleston, M., Eyer, P., Worek, F., Mohamed, F., Senarathna, L., von Meyer, L., Juszczak, E., Hittarage, A., Azhar, S., Dissanayake, W., Sheriff, M.H., Szinicz, L., Dawson, A.H., Buckley, N.A., 2005. Differences between organophosphorus insecticides in human self-poisoning: a prospective cohort study. *Lancet* 366, 1452–1459.
- Eddleston, M., Phillips, M.R., 2004. Self poisoning with pesticides. *BMJ* 328, 42–44.
- Eyer, F., Meischner, V., Kiderlen, D., Thiermann, H., Worek, F., Haberkorn, M., Felgenhauer, N., Zilker, T., Eyer, P., 2003. Human parathion poisoning A toxicokinetic analysis. *Toxicol. Rev.* 22, 143–163.
- Fears, R., Ferres, H., Glasgow, E., Standring, R., Hogg, K.J., Gemmill, J.D., Burns, J.M., Rae, A.P., Dunn, F.G., Hillis, W.S., 1992. Monitoring of streptokinase resistance titre in acute myocardial infarction patients up to 30 months after giving streptokinase or anistreplase and related studies to measure specific antistreptokinase IgG. *Br. Heart J.* 68, 167–170.
- Horne, I., Sutherland, T.D., Harcourt, R.L., Russell, R.J., Oakeshott, J.G., 2002. Identification of an opd (organophosphate degradation) gene in an *Agrobacterium* isolate. *Appl. Environ. Microbiol.* 68, 3371–3376.
- Jackson, C.J., Carr, P.D., Kim, H.K., Liu, J.W., Herrald, P., Mitic, N., Schenk, G., Smith, C.A., Ollis, D.L., 2006. Anomalous scattering analysis of *Agrobacterium radiobacter* phosphotriesterase: the prominent role of iron in the heterobinuclear active site. *Biochem. J.* 397, 501–508.
- Jeyaratnam, J., 1990. Acute pesticide poisoning: a major global health problem. *World Health Stat. Q.* 43, 139–144.
- Lewis, V.E., Donarski, W.J., Wild, J.R., Raushel, F.M., 1988. Mechanism and stereochemical course at phosphorus of the reaction catalyzed by a bacterial phosphotriesterase. *Biochemistry* 27, 1591–1597.
- Mazur, A., 1946. An enzyme in animal tissue capable of hydrolysing the phosphorus-fluorine bond of alacyl fluorophosphates. *J. Biol. Chem.* 164, 271–289.
- Ortigoza-Ferado, J., Richter, R.J., Hornung, S.K., Motulsky, A.G., Furlong, C.E., 1984. Paraoxon hydrolysis in human serum mediated by a genetically variable arylesterase and albumin. *Am. J. Hum. Genet.* 36, 295–305.
- Peng, A., Meng, F.Q., Sun, L.F., Ji, Z.S., Li, Y.H., 2004. Therapeutic efficacy of charcoal hemoperfusion in patients with acute severe dichlorvos poisoning. *Acta Pharmacol. Sin.* 25, 15–21.
- Phillips, M.R., Li, X., Zhang, Y., 2002. Suicide rates in China 1995–99. *Lancet* 359, 835–840.
- Sogorb, M.A., Vilanova, E., Carrera, V., 2004. Future applications of phosphotriesterases in the prophylaxis and treatment of organophosphorus insecticide and nerve agent poisonings. *Toxicol. Lett.* 151, 219–233.
- Van der Hoek, W., Konradsen, F., Athukorala, K., Wanigadewa, T., 1998. Pesticide poisoning: a major health problem in Sri Lanka. *Soc. Sci. Med.* 46, 495–504.
- Yang, H., Carr, P.D., McLoughlin, S.Y., Liu, J.W., Horne, I., Qiu, X., Jeffries, C.M., Russell, R.J., Oakeshott, J.G., Ollis, D.L., 2003. Evolution of an organophosphate-degrading enzyme: a comparison of natural and directed evolution. *Protein Eng.* 16, 135–145.