

Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

A Practical Synthesis of 2-Fluoro-4-bromobiphenyl

Ya-Yu Qiu ^a, Haining Gu ^a, Peng-Fei Zhang ^b & Wei-Ming Xu ^a

^a College of Material Chemistry and Chemical Engineering, Hangzhou Normal University (Xiasha Campus), Hangzhou, 310036, P. R. China

^b Center of Analysis and Measurement, Zhejiang University, (Xixi Campus), Hangzhou, P. R. China

Published online: 20 Nov 2009.

To cite this article: Ya-Yu Qiu, Haining Gu, Peng-Fei Zhang & Wei-Ming Xu (2009): A Practical Synthesis of 2-Fluoro-4-bromobiphenyl, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 41:6, 539-541

To link to this article: <http://dx.doi.org/10.1080/00304940903328219>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

A Practical Synthesis of 2-Fluoro-4-bromobiphenyl


Ya-Yu Qiu,¹ Haining Gu,¹ Peng-Fei Zhang,² and Wei-Ming Xu²

¹College of Material Chemistry and Chemical Engineering, Hangzhou Normal University (Xiasha Campus) Hangzhou 310036, P. R. China

²Center of Analysis and Measurement, Zhejiang University (Xixi Campus), Hangzhou, P. R. China

2-Fluoro-4-bromobiphenyl (**2**) is the key intermediate for the manufacture of flurbiprofen,^{1,2} a non-steroidal anti-inflammatory and analgesic material. Though compound **2** is a useful molecule, a review of the literature, including patents, indicated the absence of an easily-performed synthesis. Lang *et al.*³ had developed a cross-coupling reaction of 2-fluoro-4-bromiodobenzene with phenylboronic acid to obtain 2-fluoro-4-bromobiphenyl in good yield (87%). However, the high cost associated with the use (and removal) of palladium and toxic phenylboronic acid have limited the more widespread use of such protocol in large-scale production. Alternatively, diazotization of 2-fluoro-4-bromoaniline with sodium nitrite and an organic⁴ or inorganic acid⁵ has commonly been used to generate the diazonium salt for coupling with benzene. However, in spite of all precautions the formation of dark decomposition products is unavoidable thus lowering the yield and making the product difficult to isolate.

As part of our research program on the study of diazotizing systems,⁶ we have developed and now report a practical pilot-scale method for the preparation of 2-fluoro-4-bromobiphenyl (*Scheme 1*) from methyl nitrite and 2-fluoro-4-bromoaniline, easily prepared by bromination of 2-fluoroaniline according to the literature procedure.⁴ Methyl nitrite is a toxic, volatile, heat-sensitive potentially explosive gas that is not commercially available; its transport is prohibited in the U.S. and the European Union. Therefore its generation previously described by us⁶ is reproduced in this article. Proper precautions and a good hood should be used.



Scheme 1

Submitted June 4, 2009.

Address correspondence to Wei-Ming Xu, College of Material Chemistry and Chemical Engineering, Hangzhou Normal University (Xiasha Campus), Hangzhou 310036, P. R. China. E-mail: wmxu@zju.edu.cn

The above process is efficient, no acid is used⁷ during the diazotization, and the reaction mixture is easy to work up because of the use of methyl nitrite.⁶ In this reaction, proper control of temperature is important to maintain an adequate reaction rate and satisfactory yield.

Experimental Section

Mps and bps are uncorrected. The purity of products was established on an Agilent 1100 HPLC. ¹H NMR spectra were recorded in CDCl₃ on a Bruker 400 (400 MHz) instrument with TMS as internal standard. Infrared spectrum was obtained as a KBr pellet on a Shimadzu IR-408 instrument. All chemicals were reagent grade and available commercially. The elemental analysis was performed on a Flash EA1112 instrument.

Methyl Nitrite (prepared just prior to use)

In a 500 mL round-bottomed flask, fitted with a pressure equalizing addition funnel filled with 125 mL 12M hydrochloric acid, and a gas tube in a well ventilated hood, was placed 86.3 g (1.25 mol) of sodium nitrite, 70 mL methanol and 125 mL water. The temperature was raised to 35°C and hydrochloric acid was added dropwise over 1.5 h. The colorless methyl nitrite evolved was passed through a water trap to remove acid and then bubbled through the reaction mixture below *via* the gas tube.

2-Fluoro-4-bromobiphenyl

In a 3 L round-bottomed flask, fitted with a mechanical stirrer, and a gas tube in a well-ventilated hood, was placed 224.0 g (1.18 mol) of 2-fluoro-4-bromoaniline, 15.0 g (0.15 mol) CuCl and 2 L benzene. The temperature was raised to 43°C and methyl nitrite (76.3 g, 1.25 mol, generated as described above) was added through the tube for about 1.5 h while the temperature was kept between 43–47°C by means of water bath. When the reaction was complete, the temperature was raised to 50°C and the mixture was stirred for another 20 min. to expel excess methyl nitrite. The mixture was then cooled to room temperature, washed with 1500 mL water and dried over MgSO₄. Benzene was then removed by distillation and recycled. The residue was evaporated under vacuum (105–107°C/0.5 mmHg) to afford 222.0 g (75%) of the product as a white solid, mp. 36–37°C (HPLC >98.5%).

¹H NMR (DMSO-*d*₆): δ 7.31–7.55 (8 H, m). ¹³C NMR (DMSO-*d*₆): δ 159.4 (*J* = 250.4 Hz), 134.7, 131.7 (*J* = 3.4 Hz), 128.8, 128.7, 128.5, 128.2 (*J* = 13.1 Hz), 128.0, 127.7, 127.6, 121.2 (*J* = 9.0 Hz), 119.7 (*J* = 26.2 Hz). An analytical sample was prepared by recrystallization from benzene and methanol (v/v 5/95), mp. 36.5–37°C, *lit.*⁴ mp. 36–37°C.

Anal. Calcd for C₁₂H₈BrF: C, 57.40; H, 3.21. Found: C, 57.24; H, 3.13.

Acknowledgment

We thank the National Natural Science Foundation of China (Grant No. 20602029) and the Science and Technology Foundation of Zhejiang Province (Grant No. 2007C11014) for financial support.

References

1. D. S. Huang, D. Dapremont, P. Berget, X. Her and D. Sanchez, WO2008144198 **2008**; *Chem. Abstr.*, **149**, 582566 (2008).
2. F. Han, S. M. Li, R. Yin, X. L. Shi and Q. Jia, *Drug Development and Industrial Pharmacy*, **34**, 453 (2008); *Chem. Abstr.*, **148**, 568629 (2008).
3. J. F. Lang and W. Groves, US5254776 **1993**; *Chem. Abstr.*, **120**, 106562 (1993).
4. Z.Y. Wang, M.H. Liang, Z. Ma and Z.R. Sheng, *Zhongguo Xiandai Yingyong Yaoxue*, **23**, 463 (2006); *Chem. Abstr.*, **148**, 144462 (2006).
5. T. A. Hylton and J. A. Walker, EP0032620 **1981**; *Chem. Abstr.*, **96**, 476256 (1981).
6. H. Gu, B. Yu, P. F. Zhang and W. M. Xu, *Org. Prep. Proced. Int.*, **41**, 162 (2009).
7. When the diazotization was performed with methyl nitrite purified by passing through a water trap and a 10% sodium carbonate trap, the same yield of product was obtained.