

## Photolysis of *cis*- and *trans*-3,5-Diphenyl-1-pyrazolines: Competing Trimethylene and Cycloreversion Pathways

By MANFRED P. SCHNEIDER,\* HERBERT BIPPI, HERMANN RAU,\* DIETER UFERMANN, and MICHAEL HÖRMANN  
(Institut für Chemie, Universität Hohenheim, D-7000 Stuttgart 70, West Germany)


**Summary** In the photolysis of the title compounds up to 30% of the reactions are proceeding *via* cycloreversions producing styrene and phenyldiazomethane.

DECOMPOSITIONS of 1-pyrazolines have frequently been interpreted in terms of various trimethylene intermediates<sup>1</sup> [(a) Scheme]. [3 + 2] Cycloreversions, yielding olefins and diazocompounds [(b) Scheme], can, in principle, compete effectively with these processes and cycloreversion products have been characterised for 1-pyrazolines with a bicyclo-[3.1.0] framework.<sup>2</sup> Cycloreversion products have also

been suggested, on the basis of product studies,<sup>3</sup> e.s.r. spectroscopy, and trapping experiments,<sup>4</sup> to be involved in the decomposition of simple monocyclic 1-pyrazolines, but no definite or quantitative evidence for these primary products has been given. To differentiate between these processes and thus to provide a concise mechanistic interpretation a quantitative determination of all primary products is crucial. This is especially important in cases, *e.g.* with the title compounds, where both pathways may ultimately lead to identical reaction products [(c) Scheme].<sup>5</sup>

Degassed 10<sup>-3</sup> molar solutions of *cis*- or *trans*-3,5-diphenyl-1-pyrazoline (**1c**) or (**1t**) in a number of solvents (Table) were irradiated at -20 °C [+ 4 °C for benzene; (**1c**) is stable only at low temperature] with monochromatic (330 ± 10 nm, Osram XBO 1600 W Xenon lamp with monochromator) u.v. light. The reactions were monitored by the decay of the *n*→*π*\* transitions of (**1c**) and (**1t**) at 330 and 332 nm, respectively. After completion, the resulting solutions containing *cis*- and *trans*-1,2-diphenylcyclopropane (**2c**) and (**2t**), styrene (**3**), and phenyldiazomethane (**4**) were treated with a slight excess of HOAc [(**4**)→benzyl acetate] and analysed by H.P.L.C. and V.P.C. The results are presented in the Table.

Alternatively, a spectroscopic low conversion (≤30%) study of the photolysis of (**1t**) at room temperature allows a direct and quantitative determination of (**1t**) at 330 nm and (**4**) (which in high concentration is actually visible as a red colour) at 308 nm, their absorption coefficients being



SCHEME

TABLE Properties of biradical and cycloreversion products (mol %) in the photolysis of (1c) and (1t) from chromatographic and spectroscopic analysis<sup>c</sup>

Compound	Solvent	Biradical <sup>a</sup>		Cycloreversion <sup>a</sup>	Cycloreversion <sup>b</sup>
		(2c)	(2t)	(3)	(4)
(1c)	Acetonitrile	55	40	4	
(1t)		7	75	8	12
(1c)	EtOH	50	40	5	
(1t)		6.5	80	10	10
(1c)	Benzene	52	35	13	
(1t)		6	80	13	22
(1c)	n-Hexane	40	33	17	
(1t)		6	63	23	30

<sup>a</sup> Chromatographic analysis    <sup>b</sup> Spectroscopic analysis    <sup>c</sup> Some decomposition of (4) cannot be excluded completely, leading to different results from the chromatographic and spectroscopic analyses

known. The proportions of (4), formed in the early stages of the reaction, are given in the Table. Careful spectroscopic and chromatographic studies at various stages of the reaction revealed that the amount of (3) and (4) passes through a maximum, and decreases again with continued irradiation. This is the reason for the higher values obtained spectroscopically.

The data indicate that cycloreversions (avoiding biradical intermediates) are responsible for up to 30% of the decomposition reactions of (1c) and (1t), an amount which has certainly been underestimated in the past. This may be due to the fact that broadband irradiation has usually been employed. On the basis of the present work, there-

fore, the direct fragmentation into carbenes, previously discussed<sup>3b</sup> seems rather unlikely.

The results suggest that for a number of 1-pyrazolines cycloreversions and trimethylene processes may have rather similar energy requirements and that mechanistic interpretation must be based on a careful and quantitative evaluation of all possible reaction pathways.

We thank the Deutsche Forschungsgemeinschaft for financial support of this research and the Fonds der Chemischen Industrie for a stipend for one of us (H.B.).

(Received, 13th June 1980, Com 649)

<sup>1</sup> For a critical review see R. G. Bergman in 'Free Radicals,' Vol 1, ed J. K. Kochi, Wiley-Interscience, New York, 1973, pp 191-237.

<sup>2</sup> By <sup>1</sup>H n.m.r. spectroscopy M. Schneider and B. Csacsko, *Angew. Chem., Int. Ed. Engl.*, 1977, **16**, 867 and references cited therein.

<sup>3</sup> (a) R. Moore, A. Mishra, and R. J. Crawford, *Can. J. Chem.*, 1968, **46**, 3305, (b) K. L. Rinehart and T. V. van Aiken, *J. Am. Chem. Soc.*, 1960, **82**, 5251, 1962, **84**, 3736.

<sup>4</sup> S. L. Buchwalter and G. L. Closs, *J. Org. Chem.*, 1975, **40**, 2549, G. Ege, K. Gilbert, and B. Hahn, *Tetrahedron Lett.*, 1979, 1571.

<sup>5</sup> Cf. earlier work with (1t) J. W. Timberlake and B. K. Bandlish, *Tetrahedron Lett.*, 1971, 1395, (1c) and (1t) M. Schneider and H. Strohacker, *Tetrahedron*, 1976, **32**, 619.