## LETTERS

# Eu(OTf)<sub>3</sub>-Catalyzed Highly Regioselective Nucleophilic Ring Opening of 2,3-Epoxy Alcohols: An Efficient Entry to 3-Substituted 1,2-Diol Derivatives

Shun-ichiro Uesugi, Tsubasa Watanabe, Takamichi Imaizumi, Masatoshi Shibuya, Naoki Kanoh, and Yoshiharu Iwabuchi\*

Department of Organic Chemistry, Graduate School of Pharmaceutical Sciences, Tohoku University, Aobayama, Sendai 980-8578, Japan

#### **(5)** Supporting Information

**ABSTRACT:** In our study of the total synthesis of (+)-irciniastatin A, we found a need to develop a method that enables a C3-selective nucleophilic ring opening of 2,3-epoxy alcohol by MeOH, by which we found that the use of combined catalytic amounts of Eu(OTf)<sub>3</sub> and 2,6-di-*tert*-butyl-4-methylpyridine (DTBMP) enables the intended trans-



formation to obtain 3-methoxy-1,2-diol efficiently. Promising features of a protocol that effects a highly regioselective nucleophilic ring opening of 2,3- and 3,4-epoxy alcohols using various nucleophiles including alcohols, thiols, and unprotected amines are described.

A strategy employing the asymmetric epoxidation of allylic alcohols<sup>1</sup> followed by the stereoselective modification of an epoxide product<sup>2</sup> provides chemists with a reliable latitude for the design and synthesis of optically active compounds with multiple stereogenic centers. The Ti(O-*i*-Pr)<sub>4</sub>-mediated nucleophilic ring opening of the glycidol (2,3-epoxyl alcohol) unit, developed by Sharpless and co-workers, provides many applications for realizing a regio- and diastereocontrolled introduction of various substituents including ethers, esters, amino groups, chalcogens, and halogens at position C3.<sup>3</sup> Some other conditions have also been reported over the past decades.<sup>4–6</sup> However, we have recently realized that there is still room for developing effective conditions for this kind of transformation. For example, in our efforts to realize the total synthesis of (+)-irciniastatin A,<sup>7</sup> we were unable to achieve a C3 epoxide opening of 2,3-epoxy alcohol by MeOH using the preexisting protocols (Figure 1). In addition,



Figure 1. Outline of our synthesis of (+)-irciniastatin A.

literature survey revealed that most of the reported methods require stoichiometric amounts of Lewis acids to achieve regioselective ring opening, and thus, an effective catalytic method is desired.<sup>8</sup>

We therefore developed an efficient and catalytic protocol to effect the C3-selective epoxide opening of 2,3-epoxy alcohol using various nucleophiles. Herein, we report a protocol for  $Eu(OTf)_3$ -catalyzed regioselective nucleophilic ring opening of diverse 2,3-epoxy alcohols and their derivatives using alcohols, thiols, and amines. This reagent system is also applicable to the C4-selective epoxide opening of 3,4-epoxy alcohols.

First, we screened Lewis acid conditions that led to the C3-selective ring opening of 2,3-epoxy alcohol 1 with MeOH (Table 1). The treatment of 1 with  $BF_3 \cdot OEt_2$  in MeOH at 0 °C yielded the desired 1,2-diol 2 and the undesired 1,3-diol 5 in a 1:1 ratio, indicating that the stereoelectronic requirements for epoxide opening at C2 and C3 are essentially identical (entry 1). The Sharpless conditions<sup>3</sup> using  $Ti(O-i-Pr)_4$  gave an inseparable 3:1 mixture of 2 and 5 in 54% yield after refluxing in MeOH for 5 days (entry 2).<sup>9</sup> The unexpectedly slow reaction would be due to the formation of  $Ti(OMe)_4$  or  $Ti(OMe)_n(O-i-Pr)_{4-n}$  in MeOH, as indicated by the formation of a white precipitate in the reaction mixture. The Crotti conditions<sup>4</sup> using LiClO<sub>4</sub> resulted in low conversion even after 3 days (entries 3 and 4). In contrast, a promising reactivity was observed in the case of using 1.2 equiv of  $Yb(OTf)_{3}$ , although the selectivity was moderate (3:1, entry 5).<sup>10</sup> We assumed that the generation of the undesired isomer (C2-Nu isomer) 5 was caused by TfOH, which was generated in small amounts from Yb(OTf)<sub>3</sub> during the reaction. Considering this hypothesis, several bases for scavenging TfOH were examined, and the addition of 1.0 equiv of 2,6-di-tert-butyl-4-methylpyridine (DTBMP) was found to improve the regioselectivity (>20:1, entry 6). We then sought for the optimal catalytic conditions. Decreasing the amount of Yb(OTf)<sub>3</sub> to 30 mol % required an

Received: June 25, 2014

Table 1. Optimization of Regioselective Ring-OpeningReaction of Epoxy Alcohol 1

но	, ō	≈	onditions MeOH		Me	но	OH Me
	1			<b>2</b> (C3-OMe	isomer)	5 (C2-O	Me isomer)
entry	Lewis acid	(equiv)	base (equiv	) temp	time	yield <sup>a</sup>	selectivity <sup>b</sup> 2 / 5
1	$BF_3{\cdot}OEt_2$	(1.5)	—	0 °C	6 h	65%	1:1
2	Ti(OiPr)₄	(1.5)	—	reflux	5 days	54%	3:1
3	LiClO <sub>4</sub>	(2.0)	—	reflux	3 days	trace	—
4	LiClO <sub>4</sub>	(10 M)	—	reflux	3 days	10%	6:1
5	Yb(OTf) <sub>3</sub>	(1.2)	—	rt	18 h	70%	3:1
6	Yb(OTf) <sub>3</sub>	(1.0)	DTBMP (1.	0) rt	8 h	64%	>20:1
7	Yb(OTf) <sub>3</sub>	(0.3)	DTBMP (1.	2) reflux	18 h	82%	8.3:1
8	Sc(OTf)3	(0.2)	DTBMP (0.	3) reflux	10 h	87%	4:1
9	La(OTf) <sub>3</sub>	(0.2)	DTBMP (0.	3) reflux	10 h	90%	17:1
10	Eu(OTf) <sub>3</sub>	(0.2)	DTBMP (0.	3) reflux	13 h	92%	18:1
11	Eu(OTf) <sub>3</sub>	(0.2)	DTBMP (0.	2) reflux	4.5 h	86%	18:1
-							1.

<sup>*a*</sup>Combined yields of the isolated C3 and C2 regioisomers. <sup>*b*</sup>All ratios were determined by <sup>1</sup>H spectroscopy.

Table 2. Regioselective Ring-Opening Reactions of 2,3-Epoxy Alcohol Derivatives with MeOH



increase in the reaction temperature to reflux for completion and reduced the regioselectivity (8.3:1, entry 7). Further screening of lanthanoid triflates eventually led to the identification of  $Eu(OTf)_3$  as the ideal catalyst for this particular substrate (entries 8–11).

RO	$\begin{array}{c} R^{2} \\ R^{1} \\ \hline \\ \mathbf{9a} \cdot \mathbf{m} \\ R = H, PMB \\ n = 0, 1 \end{array}$	X m X n M	ol % Eu(OTf) <sub>3</sub> ol % DTBMP eOH (0.2 M) temp.	R0 R <sup>1</sup> C3(C4)- (1	OM n R OH OMe isor <b>0a-m</b> )	e R <sup>2</sup> mer	RO R <sup>1</sup> C2(C3)-OM (11a	OH R <sup>3</sup> OMe Me isomer -m)
entry	epoxide		product		temp	time	yield	selectivity C3 / C2
1 <sup>a</sup>	но-	_/ 9a	оМе но ОН	10a	60 °C	3 h	88%	12:1
2 <sup>a</sup>	РМВО	/ 9b	ОМе РМВО ОН	 10b	60 °C	12 h	86%	25:1
3 <sup>a</sup>	но Но Ме		HO Ne Me OH	10c	30 °C	11 h	74%	C3 only
4 <sup>a</sup>	но о	~_ 9d	но Ме ОМе ОН	10d	30 °C	4 h	87%	C3 only
5 <sup>a</sup>	HO	TE 9e	S OMe HO	TES	60 °C	5 h	90%	4:1
6 <sup>a</sup>	но	∕_ 9f	HO HO O Me	, 9 10f	rt	18 h	92%	14:1 <sup>d</sup>
7 <sup>a</sup>	HO		OH HO ÖMe	- 10g	rt	9 h	90%	12:1 <sup>d</sup>
8 <sup>b</sup>	OH	9h	HOOH	10h	60 °C	12 h	96%	C3 only
9 <sup>c</sup>	OH O	9i		10i	30 °C	24 h	64%	C3 only
10 <sup>c</sup>	OH V	9j	HO MeO	10j	rt	1 h	97%	C3 only
11 <sup>a</sup>	OH V	9k	HO MeO <sup>(1)</sup>	10k	60 °C	6 h	85%	39:1
12 <sup>a</sup>	HO	91	HO HO MeO'''	101	60 °C	44 h	65%	43:1
13ª	HQ	9m	HO HO MeO'''	10m	60 °C	44 h	NR <sup>e</sup>	ND <sup>f</sup>

Table 3. Regioselective Ring-Opening Reactions of Epoxy

Alcohol Derivatives with MeOH

<sup>a</sup>20 mol % Eu(OTf)<sub>3</sub>/DTBMP was used. <sup>b</sup>10 mol % Eu(OTf)<sub>3</sub>/ DTBMP was used. <sup>c</sup>5 mol % Eu(OTf)<sub>3</sub>/DTBMP was used. <sup>d</sup>Selectivity of C4/C3 regioisomer. <sup>e</sup>No reaction. <sup>f</sup>Not determined.

Encouraged by the high reactivity and regioselectivity of the Eu(OTf)<sub>3</sub>-catalyzed, DTBMP-assisted epoxide ring opening reaction with MeOH, we examined the compatibility of this catalytic system using protected 2,3-epoxy-1-hexanols **6b**–f (Table 2). Fortunately, methyl-, MOM-, and PMB-protected substrates gave the corresponding C3-methoxylated products as the sole detectable products.<sup>11</sup> However, TBS- and Ac-protected substrates resulted in the detachment of the

Table 4. Ring-Opening Reactions with Various Alcohol/Thiol Nucleophiles

но		X mol % Eu(OTf) <sub>3</sub> X mol % DTBMP NuH (0.2 M) 60 °C	HO HO OH C3-Nu isomer ( <b>12a-g</b> )		OH HO Nu C2-Nu isomer ( <b>13a-g</b> )	
entry	NuH	product		time	yield	selectivity C3 / C2
1 <sup>a,b,c</sup>	MeOH	OMe HO∕↓. OH	12a	1 h (5 h) <sup>b</sup> (3.5 h) <sup>c</sup>	93% (87%) <sup>b</sup> (81%) <sup>c</sup>	C3 only (20:1) <sup>b</sup> (45:1) <sup>c</sup>
2 <sup>a,b,c</sup>	<i>i</i> PrOH	0/Pr HO∕∕ OH	12b	1 h (4 h) <sup>b</sup> (4 h) <sup>c</sup>	93% (83%) <sup>b</sup> (83%) <sup>c</sup>	25:1 (12:1) <sup>b</sup> (25:1) <sup>c</sup>
3 <sup>a,b,c</sup>	BnOH	OBn HO OH	12c	1 h (4 h) <sup>b</sup> (4 h) <sup>c</sup>	91% (88%) <sup>b</sup> (78%) <sup>c</sup>	25:1 (12:1) <sup>b</sup> (30:1) <sup>c</sup>
4 <sup>a,c</sup>	allyl alcohol	QAIIyi HO OH	12d	1 h (4 h) <sup>c</sup>	98% (83%) <sup>c</sup>	25:1 (25:1) <sup>c</sup>
5 <sup>a,c</sup>	<i>t</i> BuOH	OtBu HO OH	12e	3 h (4 h) <sup>c</sup>	48% (50%) <sup>c</sup>	13:1 (10:1) <sup>c</sup>
6 <sup>a,c</sup>	PrSH	HO OH	12f	2 h (3 h) <sup>c</sup>	91% (85%) <sup>c</sup>	C3 only (C3 only) <sup>c</sup>
7 <sup>a,c</sup>	PhSH	SPh HO OH	12g	1.5 h (3 h) <sup>c</sup>	80% (78%) <sup>c</sup>	15:1 (25:1) <sup>c</sup>

<sup>*a*</sup>20 mol %  $Eu(OTf)_3/DTBMP$  was used. <sup>*b*</sup>5 mol %  $Eu(OTf)_3/DTBMP$  was used. <sup>*c*</sup>5 equiv of NuH was added. (Toluene was used as the solvent.)

protecting group together with nucleophilic ring opening (entries 5 and 6).

Next, we explored the scope and limitations of the catalytic system using various epoxy alcohols (Table 3). cis-2,3-Epoxy alcohol 9a gave the corresponding C3 adduct 10a with sufficient regioselectivity of 12:1. Interestingly, the PMB-protected cisepoxide 9b showed higher regioselectivity (entry 2). Importantly, this method was also found to be applicable to the trisubstituted epoxides 9c and 9d to provide the corresponding C3-methoxylated products 10c and 10d, respectively (entries 3 and 4). The substrate 9e with an alkyne moiety provided moderate C3 selectivity (entry 5). Note that the catalytic system could be used for the C4-selective opening of 3,4-epoxy alcohols: 3,4-epoxy-1-alcohols 9f and 9g, respectively, gave 10f and 10g in good yields with sufficient C4 regioselectivities (entries 6 and 7). Various cyclic substrates were found to undergo a C3-selective nucleophilic ring opening with high regioselectivity (entries 9–12). Cyclic cis-epoxy alcohols gave the corresponding C3 adducts, whereas anti-epoxy alcohol 9m was recovered unreacted (entry 13).

After screening the reaction with MeOH, we became interested in the regioselective epoxide ring opening reaction using other nucleophiles. We thus explored the scope of the reaction using other solvents (Table 4). We confirmed that various alcohols attack at the C3 position with high regioselectivity (entries 1-5). In addition, thiol nucleophiles can also be used with high regioselectivity (entries 6 and 7). It should be stressed that these nucleophiles can be applied not only as solvents but also as reagents at several equivalents.

Furthermore, we found that various amines can be used as nucleophiles for the system (Table 5). Some types of aniline

### Table 5. Ring-Opening Reactions with Various Amine Nucleophiles

HO^	6a	X mol % Eu(OTf) <sub>3</sub> 2 equiv Nucleophile toluene (0.2 M) temp	HO C3-Nu is	Nu 3 OH somer ( <b>14a-g</b> )	HO C2-Nu i	OH 2 Nu somer ( <b>15a-g</b> )
entry		product	temp	time	yield	selectivity C3 / C2
1 <sup>a</sup>	HO	NHPh Vivon OH 14a	rt	7 h (7 h) <sup>b</sup>	90% (90%) <sup>b</sup>	25:1 (10:1) <sup>b</sup>
2 <sup>a,b</sup>	но́	HN OH 14b	rt	5 h (5 h) <sup>b</sup>	93% (90%) <sup>b</sup>	33:1 (18:1) <sup>b</sup>
3ª	но	HN HN OH 14c	rt	6 h	88%	20:1
4 <sup>a</sup>	НО		rt	6 h	87%	26:1
5 <sup>a,c</sup>	HO	NHBn V OH 14e	60 °C	2 h	88%	9:1
6 <sup>a,c</sup>	HO	NHPMB V OH 14f	60 °C	3 h	85%	7:1
7 <sup>a,d</sup>	НО	O N OH 14g	60 °C	40 min	81%	3.6:1

<sup>*a*</sup>20 mol %  $Eu(OTf)_3$  was used. <sup>*b*</sup>10 mol %  $Eu(OTf)_3$  was used in toluene (0.5 M). <sup>*c*</sup>The ratio was determined after Boc protection. <sup>*d*</sup>The ratio was determined after acetylation.

nucleophile were successfully employed to achieve excellent yields and regioselectivities, even at room temperature (entries 1-4). Note that the addition of DTBMP is not necessary in the case of amine nucleophiles.<sup>12</sup> Moreover, the use of some aliphatic amines as nucleophiles resulted in the formation of the corresponding 3-amino-1,2-diols with good selectivity (entries 5-7). These results strongly indicate that this method can be applied to the synthesis of nitrogen-containing natural products.

In summary, we have developed a highly stereo- and regioselective ring-opening reaction of epoxy alcohols and their derivatives using catalytic amounts of  $Eu(OTf)_3$  and DTBMP. This method has the following advantages: (1) excellent regio-selectivities are achieved with various epoxides, including unprotected and protected 2,3-epoxy alcohols, 3,4-epoxy alcohols, and cyclic epoxy alcohols; (2) various *O*-, *S*-, and *N*-nucleophiles can be used; (3) the reaction conditions are not sensitive to water or air; (4) the reaction is induced using commercially available europium salt. The present protocol can therefore be widely applied to the synthesis of complex molecules with continuous stereocenters.

ASSOCIATED CONTENT

#### **S** Supporting Information

Experimental procedures and characterization data for all new compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

#### AUTHOR INFORMATION

#### **Corresponding Author**

\*E-mail: y-iwabuchi@m.tohoku.ac.jp.

#### Notes

The authors declare no competing financial interest.

#### ACKNOWLEDGMENTS

This work was partly supported by the Ministry of Education, Culture, Sports, Science and Technology, Japan (MEXT) Platform for Drug Discovery, Informatics, and Structural Life Science, and a Grant-in-Aid for Scientific Research on Innovative Areas "Advanced Molecular Transformations by Organocatalysis."

#### REFERENCES

(1) (a) Katsuki, T.; Sharpless, K. B. J. Am. Chem. Soc. 1980, 102, 5974.
(b) Hanson, R. M.; Sharpless, K. B. J. Org. Chem. 1986, 51, 1922.
(c) Zhang, W.; Basak, A.; Kosugi, Y.; Hoshino, Y.; Yamamoto, H. Angew. Chem., Int. Ed. 2005, 44, 4389. (d) Egami, H.; Ogama, T.; Katsuki, T. J. Am. Chem. Soc. 2010, 132, 5886. (e) Olivares-Romero, J. L.; Li, Z.; Yamamoto, H. J. Am. Chem. Soc. 2013, 135, 3411. (f) Wang, C.; Yamamoto, H. J. Am. Chem. Soc. 2014, 136, 1222.

(2) For synthetic applications of chiral epoxy alcohols, see: Riera, A.; Moreno, M. *Molecules* **2010**, *15*, 1041.

(3) (a) Caron, M.; Sharpless, K. B. J. Org. Chem. 1985, 50, 1557.
(b) Chong, J. M.; Sharpless, K. B. J. Org. Chem. 1985, 50, 1560. (c) Behrens, C. H.; Ko, S. Y.; Sharpless, K. B.; Walker, F. J. J. Org. Chem. 1985, 50, 5687.
(d) Behrens, B. H.; Sharpless, K. B. J. Org. Chem. 1985, 50, 5696.

(4) Chini, M.; Crotti, P.; Flippin, L. A.; Gardelli, C.; Giovani, E.; Macchia, F.; Pineschi, M. J. Org. Chem. 1993, 58, 1221.

(5) For examples of the regioselective ring opening of 2,3-epoxy alcohols with O-, S-, and halide nucleophiles, see: (a) Onaka, M.; Sugita, K.; Takeuchi, H.; Izumi, Y. J. Chem. Soc., Chem. Commun. 1988, 117.
(b) Narender, M.; Reddy, M. S.; Nageswar, Y. V. D.; Rao, K. R. Helv. Chim. Acta 2007, 90, 1107.

(6) For examples of the regioselective ring opening of 2,3-epoxy alcohols with *N*-nucleophiles, see: (a) Maruoka, K.; Sano, H.; Yamamoto, H. Chem. Lett. **1985**, *14*, 599. (b) Onaka, M.; Sugita, K.; Izumi, Y. Chem. Lett. **1986**, *15*, 1327. (c) Caron, M.; Carlier, P. R.; Sharpless, K. B. J. Org. Chem. **1988**, *53*, 5185. (d) Canas, M.; Poch, M.; Verdaguer, X.; Moyano, A.; Pericàs, M. A.; Riera, A. Tetrahedron Lett. **1991**, *32*, 6931. (e) Benedetti, F.; Berti, F.; Norbedo, S. Tetrahedron Lett. **1998**, *39*, 7971. (f) Martín, R.; Islas, G.; Moyano, A.; Pericàs, M. A.; Riera, A. Tetrahedron **2001**, *57*, 6367. (g) Pastó, M.; Rodríguez, B.; Riera, A.; Pericás, M. A. Tetrahedron Lett. **2003**, *44*, 8369. (h) Surendra, K.; Krishnaveni, N. S.; Rao, K. R. Synlett **2004**, *36*, 506.

(7) Watanabe, T.; Imaizumi, T.; Chinen, T.; Nagumo, Y.; Shibuya, M.; Usui, T.; Kanoh, N.; Iwabuchi, Y. Org. Lett. **2010**, *12*, 1040.

(8) Very recently, Yamamoto and Wang have reported an effective catalytic system using a tungsten salt that has high regioselectivity. Wang, C.; Yamamoto, H. J. Am. Chem. Soc. **2014**, *136*, 6888.

(9) For some "unexpected" results using reported methods, see:
(a) Alegret, C.; Santacana, F.; Riera, A. J. Org. Chem. 2007, 72, 7688.
(b) Tan, L.; Ma, D. Angew. Chem., Int. Ed. 2008, 46, 3614.

(10) (a) Meguro, M.; Asao, N.; Yamamoto, Y. *Tetrahedron Lett.* **1994**, 35, 7395. (b) Meguro, M.; Asao, N.; Yamamoto, Y. *J. Chem. Soc., Chem. Commun.* **1995**, 1021.

(11) Under Sharpless conditions, epoxide opening reaction did not proceed using methyl ether 6b (see ref 3a).

(12) For details of the reaction conditions, see the Supporting Information.