


Jian-Chao Liu, Hong-Wu He*, Qing-Yun Ren and Ming-Wu Ding

Key Laboratory of Pesticide & Chemical Biology, Ministry of Education, College of Chemistry, Central China Normal University, Wuhan, 430079, P. R. China

e-mail: he1208@public.wh.hb.cn

Received July 18, 2005



The thienopyridine derivative **2**, obtained from reaction of acetoacetic ester with **1** in the presence of tin tetrachloride, was treated with triphenylphosphine in hexachloroethane and Et_3N to give iminophosphorane **3**. Iminophosphorane **3** reacted with phenyl isocyanate to give carbodiimide **4**, which was further treated with phenols or ethenol to produce 2-substituted 5,8,9-trimethyl-3-phenyl-thieno[3',2'-5,6]pyrido[4,3-*d*]pyrimidin-4(3*H*)-ones **5** in presence of catalytic amount of K_2CO_3 or EtONa . The structures of compounds **5** were confirmed by ^1H NMR, IR, MS, and elemental analysis.

J. Heterocyclic Chem., **43**, 803 (2006).

Introduction.

The derivatives of heterocycles containing pyrido-pyrimidine system are of great importances because of their remarkable biological properties [1]. For example, some pyridopyrimidine derivatives were used as a novel class of adenosine kinase inhibitors [2,3]. Thus, a large number of general methods for the preparation of fused pyridines have been reported [4-7].

Recently we have been interested in the synthesis of fused pyrimidinones *via* aza-Wittig reaction of β -ethoxycarbonyl iminophosphorane with aromatic isocyanate and subsequent reaction with various nucleophiles under mild conditions [8]. As a continuation of our research for new bioactive heterocycles [9], here we wish to report an efficient synthesis of 2-substituted thieno[3',2'-5,6]pyrido[4,3-*d*]pyrimidin-4(3*H*)-ones, a series of compounds which have not been reported before.

Results and Discussion.

The 2,3-dimethyl-4-cyano-5-aminothiophene **1** [10] was converted to thienopyridine derivative **2** *via* reaction with acetoacetic ester and tin tetrachloride under heating (Scheme 1). The iminophosphorane **3** was subsequently obtained in a satisfactory yield when **2** was treated with triphenylphosphine, hexachloroethane and Et_3N . Iminophosphorane **3** reacted with phenyl isocyanate to give carbodiimide **4**. The direct reaction of carbodiimide **4**

with phenols did not cyclize to produce 2-aryloxy-thieno[3',2'-5,6]pyrido[4,3-*d*]pyrimidin-4(3*H*)-ones **5**. However, the reaction took place to give compound **5** in good yields under the condition of heating in presence of catalytic amount of K_2CO_3 . Irrespective of the fact whether the substitutes on the phenols were electron-withdrawing or electron-releasing groups, the cyclization was carried out smoothly. Thin layer chromatography was employed to follow the progress of every above reaction. Most of the compounds **5** were readily soluble in polar organic solvents and melt within 223-300 °C. The results are listed in Table 1.

It is noteworthy that the direct reaction of carbodiimide **4** with EtOH gave a complex mixture. However, when the reaction was achieved in presence of catalytic amount of EtONa , the reaction took place completely and 2-ethoxy-thieno[3',2'-5,6]pyrido[4,3-*d*]pyrimidin-4(3*H*)-one **5r** was obtained in satisfactory yields.

The structure of compounds **5** was confirmed by their elemental analysis and spectral data. IR spectral data of compound **5** are presented in Table 2. The elemental analysis and ^1H NMR spectra are given in Table 3. The EI-mass spectra are presented in Table 4. For example, the IR spectra of compound **5a** revealed $\text{C}=\text{O}$ absorption bands at 1699 cm^{-1} and 1688 cm^{-1} and absorption bands at 3136 cm^{-1} due to Ar-H group. The ^1H NMR spectral data of **5a** show the signals of three

CH₃ at 1.98 ppm, 2.39 ppm, 3.06 ppm as singlet with the other signals appeared at 7.14~7.60 (m, 10H, Ar-H). The MS spectrum of **5a** shows an obvious molecule ion peak at *m/z* 413 with 100% abundance.

The structure of **5a** was also established on the basis of elemental analysis data. The difference between found value and calculated value of elemental analysis of all compounds was under 0.5%.

Scheme 1




Table 1

Physical Constants of Compounds **5**

Compd.	R	Formula	Color	m. p. (°C)	r. t. (h)	Yield (%)
5a	Ph-	C ₂₄ H ₁₉ N ₃ O ₂ S	colorless crystals	268~269	14	87
5b	2-MePh-	C ₂₅ H ₂₁ N ₃ O ₂ S	colorless crystals	258~260	13	60
5c	4-CH ₃ Ph-	C ₂₅ H ₂₁ N ₃ O ₂ S	colorless crystals	243~244	13	84
5d	4-ClPh-	C ₂₄ H ₁₈ ClN ₃ O ₂ S	colorless crystals	267~268	12	68
5e	2,4-diClPh-	C ₂₄ H ₁₇ Cl ₂ N ₃ O ₂ S	colorless crystals	240~241	14	69
5f	2-ClPh-	C ₂₄ H ₁₈ ClN ₃ O ₂ S	colorless crystals	260~261	13	70
5g	4-BrPh-	C ₂₄ H ₁₈ BrN ₃ O ₂ S	colorless crystals	276~277	12	67
5h	2,4-diFPh-	C ₂₄ H ₁₇ F ₂ N ₃ O ₂ S	colorless crystals	250~252	13	76
5i	3-FPh-	C ₂₄ H ₁₈ FN ₃ O ₂ S	colorless crystals	290~292	11	82
5j	2-Cl-4-FPh-	C ₂₄ H ₁₇ FCIN ₃ O ₂ S	colorless crystals	254~255	12	67
5k	3-Cl-4-FPh-	C ₂₄ H ₁₇ FCIN ₃ O ₂ S	colorless crystals	266~267	13	75
5l	2-Cl-5-CH ₃ Ph-	C ₂₅ H ₂₀ ClN ₃ O ₂ S	colorless crystals	271~273	12	40
5m	4-Cl-3-CH ₃ Ph-	C ₂₅ H ₂₀ ClN ₃ O ₂ S	colorless crystals	257~259	12	74
5n	3,5-diFPh-	C ₂₄ H ₁₇ F ₂ N ₃ O ₂ S	colorless crystals	>300	13	59
5o	4-NO ₂ Ph-	C ₂₄ H ₁₈ N ₄ O ₄ S	colorless crystals	235~237	13	60
5p	2-NO ₂ Ph-	C ₂₄ H ₁₈ N ₄ O ₄ S	colorless crystals	299~300	13	44
5q	PentaClPh-	C ₂₄ H ₁₄ Cl ₅ N ₃ O ₂ S	colorless crystals	>300	11	43
5r	CH ₃ CH ₂ -	C ₂₀ H ₁₉ N ₃ O ₂ S	colorless crystals	223~225	14	55

Table 2

IR Spectral Data of Compounds **5**

Compd.	IR (KBr, ν/ cm ⁻¹)
5a	3136(Ph-H), 2910(C-H), 1699,1688(C=O), 1617, 1561, 1511, 1490, 1401, 1373, 1266, 1205, 807, 731, 690
5b	3136(Ph-H), 2925(C-H), 1733,1697(C=O), 1617, 1593, 1562, 1514, 1491, 1400, 1372, 1319, 1266, 1240, 1157, 750, 688
5c	3125(Ph-H), 2917(C-H), 1699(C=O),1618,1561,1504,1402,1370,1266,1201,835,807,700
5d	3135(Ph-H), 2924(C-H), 1700,1687(C=O), 1616, 1560, 1510, 1488, 1402, 1263, 1208, 1012, 844, 740, 690
5e	3137(Ph-H), 2926(C-H), 1699(C=O), 1622, 1563, 1516, 1403, 1371, 1251, 1225, 1096, 1049, 834, 757, 690
5f	3137(Ph-H), 2935(C-H), 1696(C=O), 1618, 1562, 1514, 1477, 1401, 1371, 1263, 1221, 1051, 806, 745

Table 2 (continued)

Compd.	IR (KBr, ν/cm^{-1})
5g	3098(Ph-H), 2927(C-H), 1701(C=O), 1614, 1560, 1512, 1485, 1402, 1371, 1322, 1263, 1207, 1066, 1009, 842, 740, 630
5h	3124(Ph-H), 2921(C-H), 1700(C=O), 1621, 1561, 1506, 1402, 1372, 1265, 1195, 1143, 965, 830, 746, 703
5i	3136(Ph-H), 2936(C-H), 1696(C=O), 1621, 1592, 1562, 1516, 1489, 1400, 1373, 1264, 1141, 944, 810, 706
5j	3124(Ph-H), 2927(C-H), 1732, 1701(C=O), 1623, 1594, 1561, 1490, 1420, 1371, 1265, 1182, 1047, 830, 702
5k	3117(Ph-H), 2924(C-H), 1699(C=O), 1620, 1560, 1493, 1400, 1319, 1247, 1194, 1054, 873, 773, 693
5l	3136(Ph-H), 2926(C-H), 1699(C=O), 1621, 1562, 1518, 1399, 1367, 1268, 1236, 1171, 1050, 806, 703, 687
5m	3136(Ph-H), 2926(C-H), 1700(C=O), 1622, 1562, 1519, 1476, 1401, 1371, 1323, 1265, 1243, 1163, 1047, 877, 752
5n	3134(Ph-H), 2932(C-H), 1701, 1686(C=O), 1618, 1560, 1510, 1491, 1466, 1400, 1375, 1264, 1156, 1120, 1091, 992
5o	3116(Ph-H), 2928(C-H), 1700(C=O), 1621, 1562, 1520, 1490, 1419, 1403, 1342, 1322, 1261, 1214, 1160, 863, 734
5p	3147(Ph-H), 2935(C-H), 1695(C=O), 1620, 1562, 1525, 1492, 1401, 1349, 1263, 1220, 1047, 806, 733, 692
5q	3094(Ph-H), 2915(C-H), 1737, 1721(C=O), 1656, 1592, 1573, 1495, 1423, 1396, 1376, 1270, 1200, 1009, 806, 705, 662
5r	3152(Ph-H), 2927(C-H), 1689(C=O), 1604, 1589, 1560, 1515, 1493, 1419, 1340, 1310, 1263, 1174, 764, 694

Table 3
Elemental Analysis and ^1H NMR Spectral Data of Compounds **5**

Compd.	Calcd. (Found)/%			^1H NMR (δ / ppm, CDCl_3 , TMS, 400 MHz)
	C.	H.	N.	
5a	69.73(69.47)	4.60(4.63)	10.17(10.33)	1.98 (s, 3H, CH_3), 2.39(s, 3H, CH_3), 3.06 (s, 3H, CH_3), 7.14~7.60 (m, 10H, Ar-H)
5b	70.24(69.95)	4.95(4.83)	9.83(10.02)	2.06 (s, 3H, CH_3), 2.39(s, 3H, CH_3), 2.41(s, 3H, CH_3), 3.04 (s, 3H, CH_3), 6.96~7.62 (m, 9H, Ar-H)
5c	70.24(70.50)	4.95(5.02)	9.83(9.86)	2.02(s, 3H, CH_3), 2.38(s, 3H, CH_3), 2.40(s, 3H, CH_3), 3.05(s, 3H, CH_3), 7.07~7.60 (m, 9H, Ar-H)
5d	64.36(64.09)	4.02(4.03)	9.39(9.50)	2.04 (s, 3H, CH_3), 2.40(s, 3H, CH_3), 3.03 (s, 3H, CH_3), 7.10~7.62 (m, 9H, Ar-H)
5e	59.75(59.90)	3.53(3.67)	8.71(8.53)	1.96 (s, 3H, CH_3), 2.41(s, 3H, CH_3), 3.05 (s, 3H, CH_3), 7.19~7.61 (m, 8H, Ar-H)
5f	64.36(64.69)	4.02(4.18)	9.39(9.41)	1.89 (s, 3H, CH_3), 2.39(s, 3H, CH_3), 3.07 (s, 3H, CH_3), 7.23~7.61 (m, 9H, Ar-H)
5g	58.54(58.35)	3.66(3.79)	8.54(8.49)	2.04 (s, 3H, CH_3), 2.41(s, 3H, CH_3), 3.03 (s, 3H, CH_3), 7.04~7.60 (m, 9H, Ar-H)
5h	64.14(63.86)	3.79(3.84)	9.35(9.48)	2.01 (s, 3H, CH_3), 2.65(s, 3H, CH_3), 3.06 (s, 3H, CH_3), 7.26~7.61 (m, 8H, Ar-H)
5i	66.82(66.64)	4.18(4.25)	9.74(9.90)	2.05 (s, 3H, CH_3), 2.41(s, 3H, CH_3), 3.05 (s, 3H, CH_3), 6.96~7.60 (m, 9H, Ar-H)
5j	61.87(61.85)	3.65(3.72)	9.02(9.11)	1.98 (s, 3H, CH_3), 2.52(s, 3H, CH_3), 3.06 (s, 3H, CH_3), 7.26~7.56 (m, 8H, Ar-H)
5k	61.87(62.16)	3.65(3.82)	9.02(9.09)	2.10 (s, 3H, CH_3), 2.43(s, 3H, CH_3), 3.04 (s, 3H, CH_3), 7.26~7.60 (m, 8H, Ar-H)
5l	65.00(64.79)	4.36(4.45)	9.10(9.04)	1.94 (s, 3H, CH_3), 2.35(s, 3H, CH_3), 2.40(s, 3H, CH_3), 3.07 (s, 3H, CH_3), 7.06~7.60 (m, 8H, Ar-H)
5m	65.00(64.86)	4.36(4.27)	9.10(9.29)	2.06 (s, 3H, CH_3), 2.39(s, 3H, CH_3), 2.41(s, 3H, CH_3), 3.03 (s, 3H, CH_3), 6.94~7.62 (m, 8H, Ar-H)
5n	64.13(64.22)	3.81(3.92)	9.35(9.32)	2.14 (s, 3H, CH_3), 2.44(s, 3H, CH_3), 3.07 (s, 3H, CH_3), 6.78~7.61 (m, 8H, Ar-H)
5o	62.87(62.65)	3.96(3.91)	12.22(12.37)	2.00 (s, 3H, CH_3), 2.41(s, 3H, CH_3), 3.06 (s, 3H, CH_3), 7.26~8.34 (m, 9H, Ar-H)
5p	62.87(63.03)	3.96(3.99)	12.22(12.03)	1.86 (s, 3H, CH_3), 2.38(s, 3H, CH_3), 3.06 (s, 3H, CH_3), 7.26~8.24 (m, 9H, Ar-H)
5q	49.21(48.97)	2.41(2.56)	7.17(7.30)	2.53 (s, 3H, CH_3), 2.67(s, 3H, CH_3), 3.01 (s, 3H, CH_3), 7.26~7.59 (m, 5H, Ar-H)
5r	65.75(66.01)	5.21(5.28)	11.51(11.40)	1.33(d, 3H, $J=7.2\text{Hz}$, CH_3), 2.51(s, 3H, CH_3), 2.71 (s, 3H, CH_3), 3.03(s, 3H, CH_3), 4.55(m, 2H, $J=7.2\text{Hz}$, CH_2), 7.25~7.54(m, 5H, Ar-H)

Table 4
The EI-mass Spectra of Compounds **5**

Compd.	MS (EI, m/z , %)
5a	415 ($M^+ + 2$, 6), 414 ($M^+ + 1$, 21), 413 (M^+ , 100), 321 (11), 320 (66), 189 (12), 76 (24), 64 (10), 49 (9)
5b	429 ($M^+ + 2$, 8), 428 ($M^+ + 1$, 27), 427 (M^+ , 100), 320 (10), 90 (19), 76 (31), 64 (12), 50 (7)
5c	428 ($M^+ + 1$, 11), 427 (M^+ , 28), 320 (11), 252 (12), 169 (10), 166 (48), 98 (27), 95 (15), 82 (64), 67 (38), 66 (34), 55 (100), 40 (48), 38 (38)
5d	449 ($M^+ + 1$, 20), 448 (M^+ , 32), 447 (100), 336 (25), 126 (17), 110 (19), 98 (48), 76 (58), 74 (19), 72 (10), 58 (13), 49 (13)
5e	485 ($M^+ + 3$, 15), 484 ($M^+ + 2$, 20), 483 ($M^+ + 1$, 64), 482 (M^+ , 33), 481 (100), 160 (8), 134 (17), 132 (31), 76 (96), 50 (15), 44 (14), 41 (22)
5f	449 ($M^+ + 1$, 38), 448 (M^+ , 29), 447 (100), 127 (6), 110 (20), 98 (31), 76 (69), 74 (18), 50 (14)
5g	494 ($M^+ + 2$, 26), 493 ($M^+ + 1$, 90), 492 (M^+ , 31), 491 (100), 320 (9), 144 (9), 142 (10), 76 (35), 62 (10), 49 (7)
5h	451 ($M^+ + 2$, 9), 450 ($M^+ + 1$, 25), 449 (M^+ , 100), 128 (10), 118 (12), 100 (19), 76 (73), 62 (13), 50 (16), 49 (9)
5i	433 ($M^+ + 2$, 8), 432 ($M^+ + 1$, 27), 431 (M^+ , 100), 320 (8), 111 (13), 94 (53), 82 (43), 76 (85), 74 (25), 56 (12), 50 (25)
5j	467 ($M^+ + 2$, 27), 466 ($M^+ + 1$, 23), 465 (M^+ , 74), 144 (10), 128 (12), 118 (32), 116 (30), 90 (28), 76 (100), 62 (11), 50 (30)
5k	467 ($M^+ + 2$, 19), 465 (M^+ , 88), 337 (14), 319 (100), 190 (13), 188 (51), 127 (6), 76 (29), 50 (5)
5l	464 ($M^+ + 2$, 11), 463 ($M^+ + 1$, 36), 462 (M^+ , 30), 461 (100), 426 (20), 124 (12), 112 (9), 76 (72), 62 (7), 50 (15)
5m	464 ($M^+ + 3$, 24), 463 ($M^+ + 2$, 16), 461 (M^+ , 85), 432 (13), 319 (100), 307 (24), 305 (22), 189 (18), 172 (11), 112 (10), 76 (38)
5n	451 ($M^+ + 2$, 6), 450 ($M^+ + 1$, 19), 449 (M^+ , 65), 320 (10), 112 (28), 100 (21), 76 (100), 50 (24)
5o	460 ($M^+ + 2$, 8), 459 ($M^+ + 1$, 26), 458 (M^+ , 100), 427 (6), 320 (37), 76 (17), 56 (9), 54 (9), 38 (8)
5p	458 (M^+ , 100), 441 (53), 412 (35), 411 (64), 337 (20), 336 (31), 333 (15), 319 (61), 92 (19), 76 (28), 66 (21)
5q	585 (M^+ , 1), 340 (8), 339 (21), 337 (100), 292 (27), 245 (31), 218 (11), 190 (26), 189 (12), 171 (24), 119 (10)
5r	365 (M^+ , 100), 337 (13), 336 (14), 76 (5), 63 (3)

EXPERIMENTAL

Melting points were determined with a WRS-1B Digital melting point apparatus and are uncorrected. EI-MS spectra were measured on a Finnigan Trace Mass Spectrometer. IR spectra were recorded on a Shimadzu IR-408 Infrared Spectrometer. ^1H NMR spectra were taken on a Varian XL-300 Spectrometer. Elemental Analysis were recorded on a Varian EL III elemental analysis instrument. All of the solvents and materials were reagent grade and purified as required.

Thienopyridine derivative **2** was prepared according to the literature procedures [10] in 69%, Colorless crystals, mp 131~132 °C. ^1H NMR (CDCl_3 , δ): 6.65~6.73 (s, 2H, NH_2), 4.40 (m, 2H, $J=7.2\text{Hz}$, CH_2CO), 2.77 (s, 3H, CH_3), 2.57 (s, 3H, CH_3), 2.39 (s, 3H, CH_3), 1.20 (s, 3H, $J=7.2\text{Hz}$, CH_3).

Preparation of iminophosphorane **3**.

A solution of thienopyridine derivative **2** (1.06 g, 4 mmol) in CH_3CN (15mL) was added to triphenylphosphine (1.31 g, 5 mmol) and C_2Cl_6 (1.19 g, 5 mmol). The mixture was treated with triethylamine (8.0 mL), then stirred for 18~24 h at 0 °C, the solution was condensed and the residue was recrystallized from $\text{CH}_3\text{CH}_2\text{OH}$ to give iminophosphorane **3** in yield 93%, Colorless crystals, mp 174~175 °C. ^1H NMR (CDCl_3 , δ): 7.42~7.63 (m, 18H, 3Ph-H), 3.16 (m, 2H, $J=7.2\text{Hz}$, CH_2CO), 2.46 (s, 3H, CH_3), 2.35 (s, 3H, CH_3), 2.11 (s, 3H, CH_3), 1.02 (s, 3H, $J=7.2\text{Hz}$, CH_3).

Preparation of carbodiimides **4**.

To a solution of iminophosphorane **3** (0.525 g, 1 mmol) in anhyd CH_2Cl_2 (10 mL) was added aromatic isocyanate (1.1 mmol) under N_2 at r.t. After the reaction mixture was left unstirred for 5-12 h, the solvent was removed under reduced pressure and Et_2O /petroleum ether was added to precipitate triphenylphosphine oxide. Removal of the solvent gave carbodiimides **4**, which were used directly without further purification.

General Procedure for the Preparation of Compounds **5a~5q**.

To the solution of **4** prepared above in CH_3CN (15 mL) was added phenol (1.1 mmol) and catalytic K_2CO_3 . The mixture was

stirred for 11~14 h at 80 °C, the solution was concentrate and the residue was recrystallized from CH_3CN to give 2-aryloxy-5,8,9-trimethyl-3-phenylthieno[3',2'-5,6]pyrido[4,3-*d*]pyrimidin-4(3*H*)-ones **5a~5q**.

Preparation of compound **5r**.

To the solution of **4** prepared above in $\text{CH}_3\text{CH}_2\text{OH}$ (15 mL) were added several drops of $\text{CH}_3\text{CH}_2\text{ONa}$ in $\text{CH}_3\text{CH}_2\text{OH}$. After the mixture was stirred for 14 h at room temperature, the solution was condensed and the residue was recrystallized from $\text{CH}_3\text{CH}_2\text{OH}$ to give 2-ethyloxy-5,8,9-trimethyl-3-phenyl thieno[3',2'-5,6]pyrido[4,3-*d*]pyrimidin-4(3*H*)-ones **5r**.

Acknowledgement.

We gratefully acknowledge financial support of this work by the National Key Project for Basic Research and the National Natural Science Foundation of China (No. 2003CB114400 and Project No. 20372023)

REFERENCES AND NOTES

- [1] G. L. Anderson and A. D. Broom. *J. Org. Chem.*, **42**, 997 (1977).
- [2] G. Z. Zheng, C. H. Lee, J. K. Pratt and R. J. Perner. *Bioorg. Med. Chem. Lett.*, **11**, 2071 (2001).
- [3] G. Z. Zheng, Y. Mao, C. H. Lee and J. K. Pratt. *Bioorg. Med. Chem. Lett.*, **13**, 3041 (2003).
- [4] A. M. Thompson, A. J. Bridges, D. W. Fry, A. J. Kraker and W. A. Denny. *J. Med. Chem.*, **38**, 3780 (1995).
- [5] G. W. Rewcastle, B. D. Palmer, A. M. Thompson and A. J. Bridges. *J. Med. Chem.*, **39**, 1823 (1996).
- [6] H. Maruoka, M. Yamazaki and Y. Tomioka. *J. Heterocyclic Chem.*, **41**, 641 (2004).
- [7] M. M. Abdelkhalik, A. M. Eltoukhy, S. M. Agamy and M. H. Elnagdi. *J. Heterocyclic Chem.*, **41**, 431 (2004).
- [8] M. W. Ding, S. J. Yang and J. Zhu. *Synthesis*, 75 (2004).
- [9] F. L. Yang, Z. J. Liu, X. B. Huang and M. W. Ding. *J. Heterocyclic Chem.*, **41**, 77 (2004).
- [10] A. C. Veronese, R. Callegari and C. F. Morelli. *Tetrahedron*, **51**, 12277 (1995).