

S0040-4039(96)00408-X

A Concise Synthesis of *Ortho*-Condensed Oxane-Oxene, Oxepene, Oxocene and Oxonene Ring Systems


Eleuterio Alvarez, Mercedes Delgado, María Teresa Díaz, Liu Hanxing, Ricardo Pérez and Julio D. Martín*

Instituto Universitario de Bio-Orgánica "Antonio González", Universidad de La Laguna-CSIC,
 Carretera Vieja de La Esperanza, 2; 38206 La Laguna, Tenerife, Spain

Abstract: An efficient strategy for the synthesis of *trans*-fused oxabicyclic systems involving thioannulation followed by a Ramberg-Bäcklund olefination as the key step is described. Copyright © 1996 Elsevier Science Ltd

As a part of a program¹ aimed at the total synthesis of ciguatoxin,² and in an attempt to develop a general synthetic strategy for the synthesis of *trans*-fused unsaturated oxacycles, we focused on an approach in which unsaturated rings are introduced via thioannulation followed by a Ramberg-Bäcklund olefination.³


Figure 1: Ciguatoxin (partial structure)




According to this strategy, cycloalkenes would be generated from the O-linked acyclic precursor via sulfur connection to form the 1,*n*-oxathiane and successive α -halogenation and oxidation at sulfur, followed by SO₂-extrusion reactions.⁴ To assess the general applicability and scope of this method in terms of ring size, a number of *ortho*-condensed oxane:oxathiacycles were synthesized starting from the common tri-O-acetyl-D-glucal (Schemes 1-3).⁵

Compounds **10** and **18** (Scheme 1) were synthesized in 60% and 70% yield, respectively, by treatment of **9** and **17** with Na₂S/Al₂O₃, HMPA, at 100 °C.⁶

An alternative thioannulation pathway is shown in Scheme 2. The whole process involves treatment of the tosyl derivative **6** with NaH/AcSH to give **19**, which was further iodinated using (Si₂)₂BH followed by I₂/NaOH oxidation to yield iodide **20**.⁷ Thioannulation to **21** proceeded smoothly by treatment of **20** with MeONa in MeOH at -25 °C under H₂ atmosphere (87% yield).




^a **Reagents and Conditions:** (a) 1.2 equiv of Et₃SiH, 1.0 equiv of BF₃·OEt₂, CH₂Cl₂, 0 °C, 2 h, 99%; (b) 1.0 equiv of NaOMe, MeOH, 30 min, 100%; (c) H₂, Pd/C 10% cat, MeOH, 8 h, 99%; (d) 1.0 equiv of TsCl, 1.5 equiv of NEt₃, DMAP cat, CH₂Cl₂, 0 °C, 8 h, 77%; (e) 1.5 equiv of NaH, 1.5 equiv of allyl bromide, *n*-Bu₄NI cat, DMF, 0 °C, 10 h, 93%; (f) cat OsO₄, 1.1 equiv of NMO, THF:H₂O (1:1), 25 °C, 4 h, 99%; (g) 1.5 equiv of NaIO₄, MeOH:H₂O (7:1), 0 °C, 30 min, then 2.0 equiv of NaBH₄, MeOH, 0 °C, 15 min, 95%; (h) 3.0 equiv of PPh₃, 3.0 equiv of imidazole, 2.0 equiv of I₂, benzene, 0 °C, 1 h, 75%; (i) 1.1 equiv of Na₂S·Al₂O₃, HMPA (0.01 M), 100 °C, 12 h, 60%; (j) 2.4 equiv of TBSCl, 5.0 equiv of imidazole, CH₂Cl₂, 25 °C, 6 h, 100%; (k) 7.0 equiv of TFA, THF:H₂O (1:1), 0 °C, 30 min, 81%; (l) 3.0 equiv of SO₃·Py complex, 5.0 equiv of NEt₃, DMSO:CH₂Cl₂ (1:1), 0 °C, 2 h, 85%; (m) 1.3 equiv of Ph₃PCH₂Br, 1.2 equiv of *n*-BuLi, toluene, 0 °C, 12 h, 54%; (n) 2.0 equiv of *n*-Bu₄NF, THF, 0 °C, 12 h, 100%; (o) 1.5 equiv of NaH, 1.3 equiv of allyl bromide, DMF, 0 °C, 2 h, 83%; (p) 3.0 equiv of (Sia)₂BH, THF, 0 °C, 12 h; then 2.2 equiv of I₂, MeOH, 3.0 equiv of NaOH, 25 °C, 2 h, 48%; (q) 1.1 equiv of Na₂S·Al₂O₃, HMPA (0.01 M), 100 °C, 24 h, 70%.




^a **Reagents and Conditions:** (a) 2.5 equiv of AcSH, 2.0 equiv of NaH, DMF, 0 °C, 10 h, 81%; (b) 1.5 equiv of (Sia)₂BH, THF, 0 °C, 8 h, then 1.1 equiv of I₂, MeOH, 1.5 equiv of NaOH, 25 °C, 40 min, 58%; (c) 2.0 equiv of NaOMe, MeOH, 0-25 °C, 12 h, 87%.

Cyclization of **30**, under similar conditions, gave in modest yield (40%) the oxathiacycle **31**, due to the formation of dimer **32** (40%)⁸ and trimer **33** (12%)⁸ (Scheme 3). However, thioannulation of the one-carbon higher homologous **34** was successful and provided **36** as a single isomer in high yield (89%). These results,

clearly reveal that the intramolecular transition state arrangement is affected not only by ring size but also by the conformation of the bicyclic skeleton.







^a **Reagents and Conditions:** (a) 50.0 equiv of DMSO, 25.0 equiv of Ac₂O, 12.5 equiv of AcOH, 25 °C, 2 days, 80%; (b) RuCl₃ cat, 4.0 equiv of NaIO₄, H₂O:CH₃CN:CCl₄ (3:2:2), 25 °C, 15 min, 100%; (c) 2.0 equiv of CH₂=CHCH₂SiMe₃, 2.0 equiv of AlCl₃, CH₂Cl₂, -78 °C → 20 °C, 6 h, 80%; (d) 3.0 equiv of NaCN, DMSO, 50 °C, 8 h, 95%; (e) 1.5 equiv of DIBAL, Et₂O, 0 °C, 6 h, then HCl (1.0 N), then 2.0 equiv of NaBH₄, MeOH, 0 °C, 30 min, 55%; (f) 1.1 equiv of TsCl, 1.5 equiv of NEt₃, DMPA cat, CH₂Cl₂, 8 h, 90%; (g) 2.5 equiv of AcSH, 2.0 equiv of NaH, DMF, 0 °C, 95%; (h) 3.0 equiv of (Sia)₂BH, THF, 0 °C, 12 h, then H₂O:NaOH (15%):H₂O₂ (30%) (2:5:5), 70%; (i) 2.0 equiv of I₂, 3.0 equiv of PPh₃, 3.0 equiv of imidazol, benzene, 0 °C, 95%; (j) 2.0 equiv of NaOMe, MeOH, 0-25 °C, 12 h, gave **31** (40%), **32** (40%) and **33** (12%).



Unsaturated oxabicycles **36-39** were formed from their respective oxathiacyclic precursors via the Ramberg-Bäcklund olefination process³ (Table 1). Fortunately, SO₂-extrusion precludes undesired β-eliminations in entries 1 and 2. In entry 4, where the size of the unsaturated ring allows the possibility of *cis*- and *trans*-geometry of the double bond, a *cis:trans* mixture was obtained.

Table 1. Synthesis of Unsaturated Polyethers by SO_2 -extrusion Reactions.

Entry	Oxathiane ^a	Product	Yield (%)
1	10	 36	40
2	21	 37	41
3	18	 38	37
4	31	 39	48

^a Reagents and Conditions: (i) 1.5 equiv of NCS, CCl_4 , 0 °C, 4-5 h; (ii) 1.5 equiv of MCPBA, CH_2Cl_2 , 0-25 °C, 8-10 h; (iii) 1.2 equiv of ^tBuOK, THF, 0 °C, 4-5 h. Since the intermediate α -chloro sulfides and α -chloro sulfones are formed as mixtures of regio- and stereoisomers, it is convenient to use them in a crude form, and to withhold purification until the Ramberg-Bäcklund reaction itself has been completed.

Acknowledgement. Financial support from the Ministry of Education and Science, Spain (PB92-0487) and the EU (Contract CII-CT92-0049) is gratefully acknowledged.

References and Notes

- For our recent stereoselective synthesis of several ciguatoxin frameworks, see: Alvarez, E.; Díaz, M.T.; Pérez, R.; Ravelo, J.L.; Regueiro, A.; de Vera, J.A.; Zurita, D.; Martín, J.D. *J. Org. Chem.* **1994**, *59*, 2848.
- For an overview, see: Yasumoto, T.; Murata, M. *Chem. Rev.* **1993**, *93*, 1897.
- For an extensive review of the Ramberg Bäcklund reaction, see: Clough, J.M. In "Comprehensive Organic Synthesis"; Trost, B.M., Fleming, Y., Pattenden, G. Eds.; Pergamon Press: Oxford, 1991; Vol. 3, pp 861-886.
- Alvarez, E.; Díaz, M.T.; Hanxing, L.; Martín, J.D. *J. Am. Chem. Soc.* **1995**, *117*, 1437.
- All new compounds exhibited satisfactory spectral and exact mass data. Yields have not been maximized and refer to spectroscopically and chromatographically homogeneous materials.
- Kerwin, S.M. *Tetrahedron Lett.* **1994**, *35*, 1023. Tan, L.C.; Pagni, R.M.; Kabalta, G.W.; Hillmyer, M.; Woosley, J. *Tetrahedron Lett.* **1992**, *33*, 7709. Czech, B.; Quici, S.; Regen, S.L. *Synthesis* **1980**, 113.
- Brown, H.C.; Ratke, M.W.; Rogic, M.M. *J. Am. Chem. Soc.* **1968**, *90*, 5038.
- As expected, the symmetry of **32** and **33** was reflected in the ¹H and ¹³C NMR spectra, which exhibited resonances for 20 hydrogens and 11 carbons, respectively. Dimer, **32**, FABMS, m/e, 455 (M+Na)⁺; ¹H NMR (400 MHz, $CDCl_3$) δ 3.85 (br d, J = 11.0 Hz, 1H), 3.70 (ddd, J = 9.0, 5.1, 5.1 Hz, 1H), 3.30 (m, 2H), 3.20 (ddd, J = 7.6, 7.6, 3.0 Hz, 1H), 2.98 (ddd, J = 10.5, 4.4, 4.4 Hz, 1H), 2.67 (ddd, J = 14.2, 9.7, 4.7 Hz, 1H), 2.54 (m, 3H), 2.22 (br d, J = 12.5 Hz, 1H), 2.10 (m, 1H), 1.66 (m, 7H), 1.30 (ddd, J = 17.5, 12.5, 5.0 Hz, 1H); ¹³C NMR ($CDCl_3$) δ 79.5 (d), 77.3 (d), 68.1 (t), 67.8 (t), 32.4 (t), 31.0 (t), 29.3 (t), 29.2 (t), 27.4 (t), 26.7 (t), 25.4 (t). Trimer, **33**, FABMS, m/e, 671 (M+Na)⁺; ¹H NMR (400 MHz, $CDCl_3$) δ 3.86 (br d, J = 10.9 Hz, 1H), 3.63 (m, 1H), 3.31 (m, 2H), 3.18 (ddd, J = 8.8, 8.8, 2.6 Hz, 1H), 3.63 (m, 1H), 3.31 (m, 2H), 3.18 (ddd, J = 8.8, 8.8, 2.6 Hz, 1H), 2.95 (ddd, J = 10.5, 9.2, 4.5 Hz, 1H), 2.68 (ddd, J = 9.5, 9.5, 4.8 Hz, 1H), 2.54 (m, 3H), 2.20 (br d, J = 15.0 Hz, 1H), 2.14 (m, 1H), 1.66 (m, 7H), 1.30 (m, 1H); ¹³C NMR ($CDCl_3$) δ 79.6 (d), 77.7 (d), 68.1 (t), 67.7 (t), 32.6 (t), 31.5 (t), 29.3 (t), 29.2 (t), 28.0 (t), 26.3 (t), 25.4 (t).

(Received in UK 24 January 1996; revised 27 February 1996; accepted 1 March 1996)