

Convergent synthesis of the ABCDE ring framework of ciguatoxin


Kenji Maeda, Tohru Oishi, Hiroki Oguri and Masahiro Hirama*

Department of Chemistry, Graduate School of Science, Tohoku University, and CREST, Japan Science and Technology Corporation (JST), Sendai 980-8578, Japan. E-mail: hirma@ykbcs.chem.tohoku.ac.jp

Received (in Cambridge, UK) 19th April 1999, Accepted 4th May 1999

An alkylation–metathesis sequence is shown to be a powerful method to synthesize the ABCDE ring framework of ciguatoxin 1.

Ciguatoxin (CTX1B, **1**) is the principal toxin that causes ciguatera poisoning.¹ Its gigantic structure and unique agonist activity against the sodium channel have attracted considerable attention among synthetic organic chemists.² During the course of our synthetic studies directed toward **1**, we developed an efficient method of constructing *trans*-fused 6,6,6-tetracyclic ether systems ($n = 7$ –10) by combining intermolecular alkylation and ring-closing metathesis (RCM) reactions.³ We describe herein a convergent synthesis of the ABCDE ring framework of **1** starting from the AB ring **4** and E-ring fragments **5**, using the described alkylation–metathesis strategy (Scheme 1).




Scheme 1


The synthesis of the iodide **4** is shown in Scheme 2. Tri-*O*-acetyl-*D*-glucal was converted to **6** via methanolysis of the acetate and successive protection of the resulting triol as the tetraisopropylidisiloxanediyl (TIPDS) and benzyl ether. Epoxidation of **6** using Spilling's method⁴ followed by addition of $\text{CH}_2=\text{CHCH}_2\text{MgBr}$ resulted in the formation of **7** in 53% overall yield. Allylation of the secondary alcohol of **7** gave **8**, which was subjected to RCM reaction using Grubbs' catalyst, $(\text{PCy}_3)_2\text{Cl}_2\text{Ru}=\text{CHPh}$ **9**⁵ to afford the 7,6-bicyclic system **10** in 94% yield.⁶ The TIPDS group of **10** was removed using TBAF and the 1,3-diol was converted to the *p*-methoxybenzylidene acetal **11**. Reductive cleavage of the benzylidene acetal with DIBAL-H followed by conversion of the resulting primary alcohol to an iodide gave **4** regioselectively.⁷

Alkylation of the *tert*-butyl ester **5**, which was readily prepared from *D*-glucose⁸ with **4** using LDA in the presence of

1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidone (DMPU) gave a separable mixture of the desired diastereomer **12** and the epimer **13** in a 3:2 ratio in 58% combined yield (Scheme 3). When HMPA was used instead of DMPU in this alkylation



Scheme 2 Reagents and conditions: i, K_2CO_3 (cat.), MeOH; ii, $(\text{TIPDS})\text{Cl}_2$, pyridine; iii, BnBr , NaH, DMF, THF, 0 °C to room temp., 63% (3 steps); iv, NBS, $\text{THF}-\text{H}_2\text{O}$ (9:1); v, KHMDS, 18-crown-6, toluene, -75 to 0 °C, 4 h; vi, $\text{H}_2\text{C}=\text{CHCH}_2\text{MgBr}$, 0 °C to room temp., 53% (3 steps); vii, $\text{H}_2\text{C}=\text{CHCH}_2\text{Br}$, NaH, 0 °C to room temp., 93%; viii, **9** (cat.), CH_2Cl_2 , 94%; ix, TBAF, THF; x, PPTS (cat.), *p*- $\text{MeOC}_6\text{H}_4\text{CH}(\text{OMe})_2$, $(\text{CH}_2\text{Cl})_2$; xi, DIBAL-H, CH_2Cl_2 , -78 to -20 °C, 75% (3 steps); xii, I_2 , PPh_3 , imidazole, toluene, 82%.



Scheme 3 Reagents and conditions: i, LDA, DMPU, THF, -78 to 0 °C, 58% (based on recovery of **4**; 23%); ii, PPTS (cat.), MeOH; iii, TBDPSCl , imidazole, DMF, 71% (2 steps); iv, DDQ, $(\text{CH}_2\text{Cl})_2-\text{H}_2\text{O}$ (20:1), 91%; v, CSA (cat.), toluene, 80 °C, 80%; vi, imidazole, toluene, reflux, 80% (**15**:**16** = 1:1); vii, $\text{H}_2\text{C}=\text{CHMgBr}$, Et_2O , -78 to -60 °C, 91%; viii, CSA (cat.), $\text{HC}(\text{OMe})_3$, CH_2Cl_2 , 75%; ix, $\text{BF}_3\cdot\text{OEt}_2$, Et_3SiH , CH_2Cl_2 , -45 °C, 96%; x, **9**, CH_2Cl_2 , 40 °C, 93%.

reaction, the ratio became 1:1. Although further attempts to improve the diastereoselectivity and to epimerize **13** were unsuccessful at this stage, we found that the epimerization could be obtained at a later stage (*vide infra*). Acidic methanolysis of the acetonide **12** followed by protection of the resulting 1,3-diol as TBDPS ethers gave **14**. Removal of the *p*-methoxybenzyl group using DDQ followed by treatment with CSA in toluene at 80 °C gave the δ -lactone **15**. The epimeric lactone **16** was also synthesized from **13** in an analogous manner. This lactone was found to undergo epimerization by treatment with imidazole in toluene under reflux to give a separable 1:1 mixture of **15** and **16** in 80% yield, while stronger bases such as Bu^tOK and DBU caused only decomposition. Thus, the undesired epimer **13** was successfully converted into **15**. Treatment of the lactone **15** with CH₂=CHMgBr followed by conversion of the resultant hemiacetal to the corresponding methyl acetal afforded **17** as a 1:1 mixture of anomers. Reduction of the anomeric mixture **17**⁹ with Et₃SiH in the presence of BF₃·OEt₂¹⁰ gave **3** as a single isomer in 96% yield. Finally RCM reaction of **3** with Grubbs' catalyst **9** at 40 °C in CH₂Cl₂ gave the ABCDE ring **2** in 93% yield.¹¹

In conclusion, we have demonstrated that the alkylation–metathesis strategy is versatile for the convergent synthesis of the pentacyclic segment **2** of **1**. Further studies directed toward the total synthesis of **1** are currently in progress in our laboratory.

Notes and references

- 1 M. Murata, A. M. Legrand, Y. Ishibashi, M. Fukui and T. Yasumoto, *J. Am. Chem. Soc.*, 1990, **112**, 4380; M. Satake, A. Morohashi, H. Oguri, T. Oishi, M. Hirama, N. Harada and T. Yasumoto, *J. Am. Chem. Soc.*, 1997, **119**, 11325.
- 2 For recent synthetic studies, see: T. Oishi, M. Shoji, K. Maeda, N. Kumahara and M. Hirama, *Synlett*, 1996, 1165; T. Oishi, K. Maeda and M. Hirama, *Chem. Commun.*, 1997, 1289; T. Oishi, M. Shoji, N. Kumahara and M. Hirama, *Chem. Lett.*, 1997, 845; T. Oishi, Y. Nagumo and M. Hirama, *Synlett*, 1997, 307; S. Hosokawa and M. Isobe, *Synlett*, 1996, 351; T. Oka, K. Fujiwara and A. Murai, *Tetrahedron*, 1996, **37**, 1179; E. Alvarez, M. Delgado, M. T. Díaz, L. Hanxing, R. Pérez and J. D. Martín, *Tetrahedron Lett.*, 1996, **37**, 2865; J. L. Ravelo, A. Regueiro, E. Rodríguez, J. A. Vera and J. D. Martín, *Tetrahedron Lett.*, 1996, **37**, 2869; S. Hosokawa, K. Kira and M. Isobe, *Chem. Lett.*, 1996, 473; M. Inoue, M. Sasaki and K. Tachibana, *Tetrahedron Lett.*, 1997, **38**, 1611; T. Oka, K. Fujiwara and A. Murai, *Tetrahedron Lett.*, 1997, **38**, 8053; M. Inoue, M. Sasaki and K. Tachibana, *Angew. Chem., Int. Ed.*, 1998, **37**, 965; M. Sasaki, T. Noguchi and K. Tachibana, *Tetrahedron Lett.*, 1999, **40**, 1337.
- 3 T. Oishi, Y. Nagumo and M. Hirama, *Chem. Commun.*, 1998, 1041.
- 4 C. H. Marzabadi and C. D. Spilling, *J. Org. Chem.*, 1993, **58**, 3761.
- 5 P. Schwab, R. H. Grubbs and J. W. Ziller, *J. Am. Chem. Soc.*, 1996, **118**, 100.
- 6 M. Delgado and J. D. Martín, *Tetrahedron Lett.*, 1997, **38**, 6299.
- 7 R. Johansson and B. Samuelsson, *J. Chem. Soc., Chem. Commun.*, 1984, 201.
- 8 I. Kadota, A. Ohno, Y. Matsukawa and Y. Yamamoto, *Tetrahedron Lett.*, 1998, **39**, 6373.
- 9 We found that reduction of the ketal rather than the corresponding hemiacetal using Et₃SiH and BF₃·OEt₂ gave a remarkably higher yield of the reduction product; 0–70% yield in the case of hemiacetal under the same reduction conditions. Also see ref. 3.
- 10 D. L. Lewis, J. K. Cha and Y. Kishi, *J. Am. Chem. Soc.*, 1982, **104**, 4976.
- 11 The stereochemistry of **2** was unambiguously determined by ¹H NMR analysis. *Selected data for 2*: δ_{H} (500 MHz, CDCl₃) 0.98 (9H, s, TBDPS), 0.99 (9H, s, TBDPS), 1.55 (1H, q, *J* 11.4, H14_{ax}), 1.53–1.60 (1H, m, H22), 1.62–1.69 (1H, m, H22'), 1.71–1.78 (1H, m, H21), 2.05–2.14 (1H, m, H21'), 2.29 (1H, dt, *J* 11.4, 4.1, H14_{eq}), 2.31–2.38 (1H, m, H8), 2.64 (1H, ddd, *J* 16.0, 7.7, 3.5, H8'), 3.08–3.15 (2H, m, H12, H13), 3.28(0) (1H, ddd, *J* 8.4, 7.7, 3.8, H9), 3.28(5) (1H, ddd, *J* 11.4, 8.9, 4.1, H15), 3.31–3.40 (2H, m, H20, H24), 3.35 (1H, dd, *J* 9.0, 8.4, H10), 3.48 (1H, t, *J* 9.0, H11), 3.62–3.70 (3H, m, H23, H25, H25'), 3.88 (1H, dq, *J* 8.9, 2.3, H16), 4.01 (1H, dq, *J* 16.0, 2.8, H5), 4.12 (1H, dq, *J* 9.0, 2.3, H19), 4.29 (1H, dd, *J* 16.0, 5.7, H5'), 4.83 (1H, d, *J* 11.5, CH₂Ph), 4.89 (1H, d, *J* 11.5, CH₂Ph), 5.64 (1H, dt, *J* 13.0, 2.3, H18), 5.77 (1H, dddd, *J* 11.8, 5.1, 3.5, 2.8, H7), 5.83 (1H, dt, *J* 13.0, 2.3, H17), 5.87 (1H, ddt, *J* 11.8, 5.7, 2.8, H6), 7.23–7.43 (17H, m, Ph), 7.52–7.56 (4H, m, Ph), 7.59–7.66 (4H, m, Ph).

Communication 9/03063H