Synthesis, structure, and reactivity of (triphenylarsoranylidene)methylcyclohepta-2,4,6-trienone derivatives: reactions with heterocumulenes and an activated acetylene

Yuhki Mitsumoto and Makoto. Nitta*

JERKIN

Department of Chemistry, School of Science and Engineering, Waseda University, Shinjuku-ku, Tokyo 169-8555, Japan

Received (in Cambridge, UK) 10th June 2002, Accepted 12th August 2002 First published as an Advance Article on the web 9th September 2002

Stable arsonium ylide derivatives **7a,b** bearing cyclohepta-2,4,6-trienyl and electron-withdrawing CO₂Et and CN groups, respectively, have been prepared for the first time through a reaction of 2-chlorocyclohepta-2,4,6-trienone with the corresponding arsonium methylide derivatives in the presence of Bu'OK. Compounds **7a,b** are isolated as stable crystalline compounds, which do not undergo hydrolysis even in acidic conditions. The X-ray crystal analysis revealed that their As–O bond distances (2.31 Å for **7a**, 2.39 Å for **7b**) lie below the sum of the van der Waals radii (3.37 Å), and thus, there is appreciable bonding interaction between the arsenic and the oxygen elements. With a view to constructing a series of cyclohepta-annulated heterocycles and in order to gain a better understanding of a series of arsonium ylides, **7a,b** were allowed to react with heterocumulenes such as carbon disulfide, phenyl isothiocyanate, diphenylcarbodiimido, and phenyl isocyanate, in a Wittig type reaction followed by electrocyclization or a formal [8 + 2]-type cycloaddition eliminating triphenylarsine sulfide or oxide to give 2*H*-cyclohepta[*b*]furan-2-thione, its imine, 2-phenylimino-2*H*-cyclohepta[*b*]pyrrole, and 2*H*-cyclohepta[*b*]furan-2-one. On the other hand, the reactions of **7a,b** with dimethyl acetylenedicarboxylate (DMAD) give azulene derivatives.

Introduction

Whereas extensive studies on phosphonium ylides have been published, relatively little attention has been paid to arsonium ylides. Arsonium ylides may be represented as a resonance hybrid of a pentacovalent arsenic structure of **1-A** and ylide structure **1-B**; they are easily prepared, and are of particular interest as regards their reactivity (Fig. 1). They are more



C-sp2 orbitals and the larger and more diffuse 4d orbitals of arsenic, and to decreased electrostatic interaction across the ylide bond. Therefore, contribution of the "covalent" canonical form 1-A should become smaller as compared with the corresponding phosphonium ylides.1 Arsonium ylides are commonly less stable than the corresponding phosphonium ylides, but those bearing electron-withdrawing groups such as monobenzoyl² and two pentafluoropropionyl^{3,4} are relatively stable, and X-ray crystallographic analysis of isolated compounds was undertaken. Previously, Kawamoto and co-workers have reported structural studies^{5,6} of 2-(triphenylphosphoranylidenemethyl)cyclohepta-2,4,6-trienone derivatives 2a-c and their reactivities.^{7,8} The X-ray crystallographic analysis of 2a,b has revealed that there is appreciable coordinative interaction between the phosphorus and the oxygen atoms in the compounds. In this context, we have reported recently the reactions of $2a-c^9$ and their aza-analogues 3-6,^{10,11} with heterocumulenes to provide new methodology for constructing cycloheptaannulated five-membered heterocycles (heteroazulenes). In order to gain a better understanding of this class of arsonium ylides, we have embarked on the preparation of 2-[triphenylarsoranylidene(ethoxycarbonyl)methyl)cyclohepta-2,4,6-trienone 7a and 2-[triphenylarsoranylidene(cyano)methyl]cyclohepta-2,4,6-trienone 7b. We report herein the first synthesis and X-ray structure analyses of 7a,b, their reactivities toward heterocumulenes and dimethyl acetylenedicarboxylate (DMAD) resulting in the formation of heteroazulenes and an azulene derivative.

Results and discussion

(a) Synthesis

reactive than analogous phosphonium or sulfonium ylides, and they may provide alkenes or epoxides, depending on the nature of the substituents on both the ylidic carbon atom and the arsenic atom and also, to some extent, the solvent used.¹ The difference between arsonium and phosphonium ylides is commonly ascribed to less efficient $p\pi$ -d π overlap between the

Previously, the synthesis of (triphenylphosphoranylidenemethyl)cyclohepta-2,4,6-trienone derivatives 2a-c has been accomplished by the reaction of 2-chlorocyclohepta-2,4,6trienone 10 with triphenylphosphonium methylide derivatives

2268 J. Chem. Soc., Perkin Trans. 1, 2002, 2268–2274

DOI: 10.1039/b205612g

in THF.^{6,9} The mechanistic aspects of the reaction are also clarified through the reaction of 2-chloro- and 2-methoxy-3,5,7-trideuteriocyclohepta-2,4,6-trienones with triphenylphosphonium methylide.¹² This procedure is applicable to the preparation of (triphenylarsoranylidenemethyl)cyclohepta-2,4,6-trienone derivatives **7a,b**. The reactions of 2-chlorocyclohepta-2,4,6-trienone **10** with arsonium ylides **9a,b**, which are generated *in situ* through the reaction of ethoxycarbonylmethyltriphenylarsonium bromide **8a**¹³ and cyanomethyltriphenylarsonium bromide **8b**^{13,14} with Bu'OK, easily afforded **7a,b**, respectively (Scheme 1). The structures of new com-



Scheme 1 Reagents and conditions: i, Bu'OK, THF, rt.

pounds **7a,b** were confirmed from inspection of the spectroscopic data including ¹H NMR spectra, IR, UV–Vis, and mass spectral data, as well as elemental analysis, and finally X-ray crystal structure analysis. We propose the pathways for the formation of **7a,b** as outlined in Scheme 1, whereby arsonium ylides **9a,b** attack the C-7 position of **10** (abnormal substitution)^{12,15} to give intermediates **11a,b**, which undergo hydrogen migration regenerating arsoranes **12a,b** and subsequent elimination of HCl give **7a,b**, respectively. The yields of **7a,b** are rather modest, and thus, the attack of **9a,b** at the 7-position of **10** does not seem to occur selectively. This assumption was confirmed by monitoring the reaction by ¹H NMR to yield a complicated mixture of unassignable products.

(b) Structure

The assigned ¹H NMR spectra of **7a,b** by H–H COSY resemble each other, and they are similar to those of **2a–c.**⁹ ORTEP drawings of **7a,b** are shown in Fig. 2. Fig. 2 shows that the arsenic atoms of **7a,b** lie between the center of a trigonal bipyramid configuration [(O2 and C8 for **7a**; O1 and C10 for **7b**) in apical position, (C8, C10, and C16 for **7a**; C10, C16, and C22 for **7b**) in equatorial position, respectively] and the center of a tetrahedral configuration, with the bond angles shown in Fig. 3. In compound **7a**, the angles of C6–As1–C16, C16– As1–C10 and C6–As1–C10 are 118.8°, 111.5° and 120.0°, respectively, and the angles of C8–As1–O2, C6–As1–C8, C8–As1–C16, C8–As1–C10 are 171.8°, 102.2°, 94.7°, and 104.4°, respectively. These values are similar to those of compound 7b. On the other hand, the intramolecular As1-O1 distances (2.311 Å for 7a and 2.391 Å for 7b) are longer than that of the As–O covalent bond (1.74–1.90 Å) in spirobi(1,3,2- λ^5 dioxarsolane) derivative ¹⁶ and are significantly shorter than the sum of the van der Waals radii (3.37 Å).¹⁷ Thus, evidently, there is appreciable interaction between the oxygen and arsenic atoms of the compounds 7a,b as found previously for 2-acetylcyclopentadienylidenearsorane 13¹⁸ (Fig. 4). The ylidic C-As bond length (1.881 Å for 7a and 1.881 Å for 7b) falls between the sum of the covalent radii of singly-bonded and doubly-bonded carbon and arsenic (1.98 and 1.775 Å), respectively,¹⁹ signifying an appreciable amount of single-bond character, and consequently of dipolar character. Thus, the canonical structures 7a,b-A and 7a,b-B best represent the actual bonding in 7a,b (Fig. 4). The cyclohepta-2,4,6-trienone moieties in compounds 7a,b are nearly planar, and bond length alternations are clearly seen (1.37–1.49 Å for 7a; 1.37–1.47 Å for 7b); the results are also in agreement with the evidence obtained from the ¹H NMR spectra. The carbonyl absorptions appearing at v_{max} 1568 cm⁻¹ (for 7a) and 1588 cm⁻¹ (for 7b) are slightly lower than that found in cyclohepta-2,4,6-trienone $(v_{max} 1594 \text{ cm}^{-1})$,²⁰ and the C=O bond lengths (1.270 Å for 7a and 1.260 Å for 7b) do not differ appreciably from that of cyclohepta-2,4,6-trienone $(1.26 \text{ Å})^{10,21}$ The compounds **7a,b** are stable crystalline solids which can be kept in air without significant decomposition. Many arsonium ylides are easily hydrolysed in the presence of moisture or in acidic solutions to give an arsine oxide and an organic residue,¹ but both 7a and 7b were recovered quantitatively after neutralization of the solution of 7a,b in 0.5 M H_2SO_4 . These features suggest that the compounds 7a,b seem to be stabilized by not only the electron-withdrawing groups of -CO₂Et (for 7a) and -CN (for 7b) as well as the cyclohepta-2,4,6-trienyl group but also the coordinative interaction of the carbonyl oxygen to the arsenic element.

(c) Reactivity

Although reactions of phosphonium ylides with heterocumulenes have been reported in several papers,²² the reactions of the arsonium analogues with heterocumulenes are not familiar at all. In our previous study,⁹ 2-(triphenylphosphoranylidenemethyl)cyclohepta-2,4,6-trienone derivatives 2a-c were revealed to react with heterocumulenes to afford cycloheptaannulated heterocycles. In relation to the study and to clarify the relative reactivities of 7a,b toward phosphoranes 2a,b, the reactions of 7a,b with heterocumulenes 14-17 and DMAD 27 were investigated. Reaction of compound 7a with carbon disulfide 14 was accomplished to give 3-ethoxycarbonyl-2Hcyclohepta[b]furan-2-thione 20a (Scheme 2). On the other hand, the reaction of compound 7b with 14 under reflux did not proceed to give 20b, and 7b was recovered. Under similar conditions, the phosphonium ylides 2a,b did not react with 14, and the starting materials were recovered. Thus, the arsonium vlide 7a seems to be more reactive than the phosphonium ylides 2a,b, and cyano-substituted derivative 7b is less reactive as compared with 7a. The reaction conditions and the yields of the products are summarized in Table 1 (Entries 1-4). Similarly, the reactions of compounds 7a,b with phenyl isothiocyanate 15 were also carried out to give (Z)-3-ethoxycarbonyl- and (Z)-2-imino-3cyano-2H-cyclohepta[b]furan (Z)-21a,b, respectively (Table 1, Entries 5 and 6). The structures of compounds 20a, (Z)-21a and (Z)-21b were confirmed on the basis of comparison of the physical data with those of the authentic specimens.9 Compounds 7a,b undergo Wittig-type reaction with 14 and 15 to eliminate Ph₃As=S to lead to the intermediates 18a and 19a,b as outlined in Scheme 2. The intermediates 18a and 19a,b then undergo 10π -electron cyclization to give 20a and (Z)-21a,b, respectively (Scheme 2). The Wittig-type reactions of arsonium ylides 7a,b with 15 occur selectively to eliminate Ph₃As=S to



Fig. 2 ORTEP drawing of compounds **7a** and **7b** with thermal ellipsoid plot (30% for **7a** and 50% for **7b** probability). Selected bond lengths (Å). For **7a**: Asl–O2 2.311, As1–C8 1.966(3), As1–C16 1.926(3), As1–C10 1.926(3), As1–C6 1.881(3), C9–O2 1.270(4), C5–C6 1.404(4), C5–C9 1.488(4), C5–C15 1.419(4), C14–C15 1.396(6), C14–C29 1.377(6), C19–C29 1.367(6), C19–C24 1.388(5), C9–C24 1.414(4). For **7b**: Asl–O1 2.391, As1–C10 1.945(2), As1–C16 1.938(2), As1–C22 1.930(2), As1–C8 1.881(2), C1–O1 1.260(3), C2–C8 1.407(3), C1–C2 1.466(3), C2–C3 1.406(3), C3–C4 1.380(3), C4–C5 1.388(3), C5–C6 1.373(3), C6–C7 1.388(3), C1–C7 1.427(3).





give (Z)-18a,b, respectively, while in the reactions of phosphonium ylides 2a,b with 14, products 20a,b were obtained in addition to (Z)-21a,b (Table 1, Entries 7 and 8).⁹

On the other hand, the reactions of 7a,b with diphenylcarbodiimide 16, which was prepared in situ by the reaction of phenyl isocyanate in the presence of a catalytic amount of Ph₃AsO,²³ did not undergo Wittig-type reaction and afforded 3-ethoxycarbonyl- and 3-cyano-N-phenyl-2-phenylimino-2Hcyclohepta[b]imidazole 24a,b as single products, respectively (Scheme 3). The reaction conditions and the yields of the products are summarized in Table 1 (Entries 9 and 10). The structures of new compounds 24a,b were assigned on the basis of the ¹H and ¹³C NMR spectra, IR, UV–Vis spectra, mass spectral data, analytical data as well as comparison of the physical data with those of the related derivatives.^{11,24} Furthermore, it was found that neither 24a nor 24b is a mixture of (E)- and (Z)isomers. Although no evidence for the stereochemical situation for **24a**,**b** was obtained, we propose the (*E*)-isomers for **24a**,**b**, because of the steric hindrance of two phenyl groups. The





reactions of **7a**,**b** with **16** are considered to proceed *via* a formal [8 + 2]-type cycloaddition to give the intermediates **23a**,**b**, which then eliminate Ph₃AsO to yield **24a**,**b** (Scheme 3). The formation of **24a**,**b** would be chemical evidence for the appreciable coordinating interaction between the carbonyl-oxygen and arsenic element in **7a**,**b**. Furthermore, compound **7a**,**b** seems to be reactive as compared with phosphonium ylides **2a**,**b** (Table 1, Entries 11 and 12), in which **7a** undergoes Wittig-type reaction.⁹

Contrary to our expectation, the reactions of **7a,b** with phenyl isocyanate **17** afforded 3-ethoxycarbonyl- and 3-cyano-2*H*-cyclohepta[*b*]furan-2-one **26a,b** along with **24a,b**, respect-

Table 1 Results for the reactions of compounds 2a,b and 7a,b with heterocumulenes 14–17 and DMAD 27

Entry	Compd.	Cumulene or 27	Ratio of 14-17 or 27	Solvent ^a	Time t/h	Product Yield (%)
1	7a	14	Excess	PhH-CS ₂	4 days	20a (49)
2	7b	14	Excess	PhH-CS ₂	4 days	7b (90)
3	2a	14	Excess	PhH-CS ₂	4 days	2a (87)
4	2b	14	Excess	PhH-CS ₂	4 days	2b (67)
5	7a	15	5	Xylene	15.5	(Z)-21a (27)
6	7b	15	10	Xylene	11	(Z)-21b (49)
7 ^b	2a	15	5	Xylene	24	20a (15), (Z)- 21a (5)
8 ^b	2b	15	5	Xylene	140	20b (9), (Z)- 21b (59)
9	7a	16	5	Xylene	15	24a (21)
10	7b	16	10	Xylene	63	24b (71)
11 ^b	2a	16	5	Xylene	77	(Z)-21a (14)
12 ^b	2b	16	5	Anisole	24	2b (71)
13	7a	17	5	Xylene	15	24a (5), 26a (42)
14	7b	17	10	Xylene	64	24b (57), 26b (23)
15 ^b	2a	17	5	Xylene	24	(Z)-21a (19), 26a (77)
16 ^b	2b	17	5	Xylene	168	(Z)-21b (22), 26b (58)
17	7a	27	5	Toluene	3	31a (29)
18	7a	27	5	Xylene	1	31a (22)
19	7b	27	5	Xylene	36	31b (71)
20	2a	27	5	Xylene	12	2a(0)
21	2b	27	5	Xylene	40	2b (23)

^{*a*} Unless otherwise specified, the reaction was carried out under reflux. ^{*b*} Ref. 9.



Scheme 2 Conditions: i, CS₂-PhH or xylene, reflux.





ively (Scheme 4). The reaction conditions and the yields of the products are summarized in Table 1 (Entries 13 and 14). The structures of compounds 26a,b were confirmed on the basis of comparison of the physical data with those of the authentic specimens.9 We propose the pathways for the formation of 24a,b and 26a,b as outlined also in Scheme 4. The reaction of 7a,b with 17 gives the intermediates 25a,b and PhN=AsPh₃, which probably reacts with excess 17 to generate diphenylcarbodiimide 16 and Ph₃AsO. The latter compound further reacts with 17 to generate PhN=AsPh3 and CO2, and thus, 16 is generated in a catalytic process.^{11,23} The intermediates 25a,b undergo 10π -electron cyclization to yield **26a,b**, respectively, and the generated 16 reacts with 7a,b to give 24a,b (vide supra). The reactions of 7a,b with 17 do not give 19a,b (Scheme 2), while those of 2a,b with 17 give 19a,b to result in the formation of (Z)-21a,b (Table 1, Entries 15 and 16).9

Finally, we investigated the reactions of **7a,b** with dimethyl acetylenedicarboxylate (DMAD) **27**. Compound **7a,b** reacted with **27** to give dimethyl 1-(ethoxycarbonyl)azulene-2,3-dicarboxylate **31a** and dimethyl 1-cyanoazulene-2,3-dicarboxylate **31b**, respectively (Scheme 5). The reaction conditions and the yields of the products are summarized in Table 1 (Entries

J. Chem. Soc., Perkin Trans. 1, 2002, 2268–2274 2271



Scheme 5 Conditions: i, toluene or xylene, reflux.

17–19). Under similar conditions, attempted reaction of phosphonium ylides 2a,b with 27 did not proceed, and complete or partial decomposition of 2a,b was observed (Table 1, Entries 20 and 21). The structures of $31a^{25}$ and $31b^{26}$ were assigned on the basis of comparison of the physical data. We propose two reaction pathways for the formation of 31a,b as follows; Michael-type addition of the ylide carbon atom of 7a,b to 27 leads to the intermediates 28a,b and/or 29a,b. Then, the As–O bonded intermediates 28a,b (*cf.* 23a,b) eliminate Ph₃AsO to result in the formation of azulene derivatives 31a,b. An alternative pathway is a ring-opening of 29a,b to give the intermediates 30a,b and the following intramolecular Wittig-type reaction gives 31a,b.

In summary, we have demonstrated the synthesis of 2-(triphenylarsoranylidenemethyl)cyclohepta-2,4,6-trienone derivatives **7a,b** for the first time. Compounds **7a,b** were isolated as fairly stable crystalline compounds and X-ray structure analysis clarified that they are stabilized by not only the electron-withdrawing groups but also the interaction of the carbonyl oxygen of the cyclohepta-2,4,6-trienone with the arsenic element. The compounds **7a,b** seem to be more reactive than the corresponding phosphonium ylides **2a,b**, and they react with heterocumulenes and DMAD to provide a variety of cyclohepta-annulated five-membered heterocycles and an azulene derivative. Further studies concerning the synthesis of the related compounds will be continued.

Experimental

IR spectra were recorded on a Horiba FT-710 spectrometer. UV–Vis spectra were recorded on a Shimadzu UV-3101PC spectrometer. Mass spectra and high-resolution mass spectra (FAB) were run on JEOL JMS-AUTOMASS150 and JMS-SX102A spectrometers. ¹H NMR spectra and ¹³C NMR spectra were recorded on an AVANCE600 spectrometer using CDCl₃ as a solvent, and the chemical shifts are given relative to internal SiMe₄ standard; *J*-values are given in Hz. Arsonium salts **8a**¹² and **8b**^{12,13}were prepared according to the procedure reported in the literature. Mps were recorded on a Yamato MP-21 apparatus and are uncorrected. All reactions except hydrolysis were carried out under anhydrous conditions and dry nitrogen atmosphere.

Preparation of 2-[triphenylarsoranylidene(ethoxycarbonyl)methyl]cyclohepta-2,4,6-trienone 7a and 2-[triphenylarsoranylidene(cyano)methyl]cyclohepta-2,4,6-trienone 7b

A solution of **8a** (6.15 g, 13 mmol) or **8b** (5.49 g, 13 mmol) and Bu'OK (2.89 g, 26 mmol) in THF (40 cm³) was stirred at rt for 1.5 h. To this solution was added 2-chlorocyclohepta-2,4,6trienone **10** (1.81 g, 13 mmol) in THF (10 cm³), and the mixture was stirred for 17 h at rt. After the THF was removed *in vacuo*, the resulting residue was dissolved in AcOEt and filtered through Celite to remove insoluble materials. The filtrate was concentrated and the residue was purified through column chromatography on Al₂O₃ by using hexane–AcOEt (3 : 1) as the eluent to give the products **7a** (838 mg, 13%) and **7b** (800 mg, 14%), respectively.

For **7a**: yellow prisms; mp 152–153 °C (decomp. from hexane–AcOEt); $\delta_{\rm H}$ (CDCl₃) 0.74 (3H, t, *J* 7.1, –CH₃), 3.52 (2H, q, *J* 7.1, –CH₂–), 6.48 (1H, d, *J* 11.0, H7), 6.51 (1H, dd, *J* 9.4, 9.5, H5), 6.90 (1H, dd, *J* 9.4, 11.0, H6), 7.09 (1H, dd, *J* 9.5, 11.0, H4), 8.37 (1H, d, *J* 11.0, H3), 7.30–7.40 (9H, m, *m*, *p*-Ph), 7.50–7.60 (6H, m, *o*-ph); $\delta_{\rm C}$ (CDCl₃) 13.8 (–CH₃), 59.2 (–CH₂–), 75.8 (C=As), 123.8 (C5), 124.9 (C7), 125.2 (C3), 128.7 (Ph), 129.9 (Ph), 131.1 (Ph), 135.7 (Ph), 136.4 (C6), 137.0 (quart), 155.5 (quart), 168.4 (quart), 178.3 (quart); $\nu_{\rm max}/\rm{cm}^{-1}$ (CHCl₃) 1664, 1586, 1496, 1437, 1422, 1413, 1367; $\lambda_{\rm max}$ (MeCN)/nm (log ε/\rm{dm}^3 mol⁻¹ cm⁻¹) 267 (4.31), 404 (4.08), 475 (4.09); *m*/*z* 496 (M⁺, 10%), 77 (100) (Found M⁺ + 1, 496.1052; C; 69.9, H; 5.0. C₂₉H₂₆O₃As requires *M* + 1, 496.1020; C; 70.16, H; 5.08%).

For **7b**: yellow prisms; mp 197–199 °C (decomp. from PhH– hexane); $\delta_{\rm H}$ (CDCl₃) 6.57 (1H, d, *J* 11.1, H7), 6.61 (1H, dd, *J* 9.4, 9.6, H5), 7.05 (1H, dd, *J* 9.4, 11.1, H6), 7.14 (1H, dd, *J* 9.6, 10.8, H4), 7.65 (1H, d, *J* 10.8, H3), 7.32–7.75 (15H, br, –Ph); $\delta_{\rm C}$ (CDCl₃) 54.8 (C=As), 123.0 (–CN), 123.4 (C7), 123.8 (C3), 123.9 (C5), 129.0 (Ph), 131.0 (Ph), 131.3 (Ph), 135.0 (Ph), 136.3 (C4), 137.4 (C6), 157.7 (C2), 176.6 (C1); $\nu_{\rm max}/\rm{cm^{-1}}$ (CHCl₃) 2169, 1588, 1501, 1442, 1436, 1413, 1354; $\lambda_{\rm max}$ (MeCN)/ nm (log $\varepsilon/\rm{dm^3}$ mol⁻¹ cm⁻¹) 401 (4.08), 479 (4.13), 567 (2.71), 586 (2.70), 664 (1.94); *m*/*z* 496 (M⁺, 10%), 77 (100) (Found M⁺ + 1, 450.0861; C; 71.8, H; 4.4, N; 3.1. C₂₇H₂₁NOAs requires *M* + 1, 450.0839; C; 72.15, H; 4.49, N; 3.12%).

Attempted hydrolysis of 7a,b in acidic conditions

A solution of **7a** (60 mg, 0.12 mmol) or **7b** (54 mg, 0.12 mmol) in 1.0 M H₂SO₄ (H₂O–EtOH 1 : 1; 5 cm³) was stirred at rt for 12 h. This mixture was neutralized with aqueous NaHCO₃, extracted with CH₂Cl₂, and the extract was dried over Na₂SO₄. After evaporation of the solvent, the residue was purified through column chromatography on Al₂O₃ by using AcOEt as the eluent to recover **7a** (60 mg, 100%), **7b** (49 mg, 91%).

Reaction of 7a,b with carbon disulfide 14

A solution of **7a** (100 mg, 0.2 mmol) or **7b** (90 mg, 0.2 mmol) in a mixture of carbon disulfide **14** (3 cm³) and PhH (3 cm³) was heated under reflux for the period indicated in Table 1. After evaporation of the solvent, the residue was purified through column chromatography on SiO₂ by using hexane– AcOEt (1 : 1) as the eluent to give **20a** (23 mg, 49%) or the starting material **7b** (81 mg, 90%) (Table 1, Entries 1 and 2).

Attempted reaction of 2a,b with carbon disulfide 14

A solution of **2a** (90 mg, 0.2 mmol) or **2b** (81 mg, 0.2 mmol) in a mixture of carbon disulfide **14** (3 cm³) and PhH (3 cm³) was heated under reflux for the period indicated in Table 1. After evaporation of the solvent, the residue was purified through column chromatography on Al_2O_3 by using hexane–AcOEt (2 : 1) as the eluent to give **2a** (78 mg, 87%), **2b** (54 mg, 67%) (Table 1, Entries 3 and 4).

Reaction of 7a,b with phenyl isothiocyanate 15

A solution of **7a** (50 mg, 0.1 mmol) or **7b** (45 mg, 0.1 mmol) and phenyl isothiocyanate **15** in xylene (3 cm³) was heated under reflux for the period indicated in Table 1. After evaporation of the solvent, the residue was purified through column chromatography on SiO₂ by using hexane–AcOEt (3 : 1) as the eluent to give the product (*Z*)-**21a** (8 mg, 27%) or (*Z*)-**21b** (12 mg, 49%) (Table 1, Entries 5 and 6).

Reaction of 7a,b with diphenylcarbodiimide 16

A solution of phenyl isocyanate 17 (238 mg, 2.0 mmol) and Ph₃AsO (6 mg, 0.02 mmol) in xylene (5 cm³) was heated at 90 °C for 1.5 h to give 16 *in situ*. The generation of 16 was confirmed by IR spectrum. To this solution was added 7a (198 mg, 0.4 mmol) or 7b (90 mg, 0.2 mmol), and the mixture was heated under reflux for the period indicated in Table 1. After evaporation of the solvent, the residue was purified through column chromatography on SiO₂ by using hexane–AcOEt (3 : 1) as the eluent to give the product 24a (7 mg, 5%) or 24b (44 mg, 61%) (Table 1, Entry 9 and 10).

For 24a: dark red prisms; mp 160-161 °C (from hexane-AcOEt); δ_H (CDCl₃) 0.89 (3H, t, J7.2, -CH₃), 3.54 (2H, q, J7.2, -CH₂-), 6.41 (1H, d, J 9.5, H8), 6.64 (1H, dd, J 9.7, 10.5, H6), 6.85-6.89 (3H, m, Ph), 6.86 (1H, dd, J 9.5, 9.7, H7), 7.02 (1H, dd, J 10.5, 11.3, H5), 7.17 (2H, dd, J 7.7, 7.9, Ph), 7.37 (2H, d, J 7.5, Ph), 7.45 (1H, t, J 7.7, Ph), 7.54 (2H, dd, J 7.5, 7.7, Ph), 7.83 (1H, d, J 11.3, H4); δ_c (CDCl₃) 13.8 (-CH₃), 60.6 (-CH₂-), 104.4 (quart), 112.1 (C8), 120.3 (Ph), 121.8 (Ph), 125.8 (C4), 127.8 (C6), 128.8 (Ph), 128.8 (Ph), 129.1 (Ph), 129.6 (Ph), 134.2 (C7), 135.0 (C5), 135.5 (quart), 146.7 (quart), 150.0 (quart), 151.5 (quart), 154.4 (quart), 164.8 (quart); v_{max}/cm^{-1} (CHCl₃) 1692, 1619, 1583, 1532, 1475, 1433; λ_{max} (MeCN)/nm $(\log \varepsilon/dm^3 mol^{-1} cm^{-1})$ 279 (4.48), 456 (4.25); m/z 368 (M⁺, 12%), 77 (100) (Found M⁺ + 1, 369.1595; C; 77.9, H; 5.3, N; 7.5. $C_{24}H_{21}N_2O_2$ requires M + 1, 369.1603; C; 78.23, H; 5.48, N; 7.61%).

For **24b**: yellow prisms; mp 211–214 °C (from hexane–AcOEt); $\delta_{\rm H}$ (CDCl₃) 6.53 (1H, d, J 9.7, H8), 6.78 (1H, dd, J 9.7, 10.0, H6), 6.94 (2H, d, J 7.6, Ph), 7.03 (1H, dd, J 9.7, 10.0, H7), 7.08 (1H, t, J 7.5, Ph), 7.17 (1H, dd, J 9.7, 11.0, H5), 7.28 (2H, dd, J 7.5, 7.6, Ph), 7.40 (2H, d, J 7.6, Ph), 7.46 (1H, d, J 11.0, H4), 7.50 (1H, t, J 7.6, Ph), 7.58 (2H, dd, J 7.6, 7.6, Ph); $\delta_{\rm C}$ (CDCl₃) 113.4 (C8), 113.8 (quart), 122.1 (Ph), 123.7 (Ph), 125.9 (Ph), 128.7 (Ph), 128.7 (Ph), 128.8 (C6), 129.3 (Ph), 130.0 (Ph), 134.5 (quart), 135.9 (C7), 136.5 (C5), 149.4 (quart), 150.2 (quart), 151.5 (quart), 154.4 (quart) and one carbon overlapped; $v_{\rm max}$ /cm⁻¹ (CHCl₃) 2209, 1631, 1583, 1534, 1470, 1438, 1272; $\lambda_{\rm max}$ (MeCN)/nm (log ε /dm³ mol⁻¹ cm⁻¹) 212 (4.24), 283 (4.71), 452 (4.56); *m*/*z* 321 (M⁺, 15%), 69 (100) (Found M⁺ + 1, 322.1364; C; 82.0, H; 5.0, N; 13.0. C₂₂H₁₇N₃ requires *M* + 1, 322.1344; C; 81.95, H; 5.01, N; 13.04%).

Reaction of 7a,b with phenyl isocyanate 17

A solution of **7a** (198 mg, 0.4 mmol) or **7b** (90 mg, 0.2 mmol) and **17** (238 mg, 2.0 mmol) in xylene (4 cm³) was heated under reflux for the period indicated in Table 1. After evaporation of the solvent, the residue was purified through column chromatography on SiO₂ by using hexane–AcOEt (1 : 1) as the eluent to give the product **24a** (8 mg, 5%), **26a** (37 mg, 42%), **24b** (37 mg, 57%), and **26b** (8 mg, 23%), respectively (Table 1, Entry 13 and 14).

Reaction of 7a,b with DMAD 27

A solution of **7a** (198 mg, 0.4 mmol) or **7b** (180 mg, 0.4 mmol) and DMAD **27** (284 mg, 2.0 mmol) in xylene (5 cm³) was heated under reflux for the period indicated in Table 1. After evaporation of the solvent, the residue was purified through column chromatography on SiO₂ by using hexane–AcOEt (3 : 1) as the eluent to give the product **31a** (28 mg, 22%) and **31b** (76 mg, 71%) (Table 1, Entry 17–19).

For **31a**: reddish violet needles; mp 168–169 °C (from hexane–AcOEt) (lit.²⁵ 175–176 °C); $\delta_{\rm H}$ (400 MHz) 1.42 (3H, t, J 7.2, –CH₃), 3.95 (3H, s), 4.03 (3H, s), 4.41 (2H, q, J 7.1, –CH₂–), 7.83 (2H, dd, J 9.9, 9.9, H5, H6), 8.05 (1H, dd, J 9.8, 9.9, H7), 9.84 (1H, d, J 9.8, H8), 9.89 (1H, d, J 9.9, H4); $\nu_{\rm max}/{\rm cm}^{-1}$ (CHCl₃) 1738, 1695, 1457, 1443, 1426, 1263, 1238; *m*/*z* 316 (M⁺, 100%), 69 (100) (Found M⁺, 316.0916. C₁₇H₁₆O₆ requires 316.0885).

For **31b**: reddish purple needles; mp 92.5–93.5 °C (from hexane–AcOEt) (lit.²⁵ 93–94 °C); $\delta_{\rm H}$ (400 MHz) 3.97 (3H, s), 4.07 (3H, s), 7.85 (1H, dd, *J* 10.0, 10.0, H6), 7.88 (1H, dd, *J* 10.0, 10.0, H7), 8.14 (1H, dd, *J* 9.8, 10.0, H5), 8.86 (1H, d, *J* 9.8, H4), 9.64 (1H, d, *J* 10.0, H8); $v_{\rm max}/{\rm cm}^{-1}$ (CHCl₃) 2200, 1734, 1704, 1457, 1444, 1443, 1425, 1264; *m*/*z* 269 (M⁺, 31%), 237 (100) (Found M⁺ + 1, 270.0783. C₁₅H₁₂NO₄ requires 270.0767).

Attempted reaction of 2a,b with DMAD 27

A solution of **2a** (181 mg, 0.4 mmol) or **2b** (162 mg, 0.4 mmol) and DMAD **25** (284 mg, 2.0 mmol) in xylene (5 cm³) was heated under reflux for the period indicated in Table 1. After evaporation of the solvent, the residue was purified through column chromatography on SiO₂ by using hexane–AcOEt (2 : 1) as the eluent to give **2b** (37 mg, 23%) (Table 1, Entry 21 and 22).

X-Ray structure determination of 7a,b †

Single crystal of 7a was recrystallized from AcOEt.

Crystal data for 7a. All the measurements were performed using a Rigaku AFC5S radiation diffractometer with graphite monochromated Cu-K α radiation: yellow prisms, C₂₉H₂₅O₃As, M = 496.00, orthorhombic, a = 15.780(6), b = 17.021(6), c =8.864(5) Å, V = 2380.760010(2) Å³, T = 298 K, space group $Pna2_1$ (no. 33), Z = 4, μ (Cu-K α) = 2.1415 mm⁻¹, 9937 reflections measured, 2341 unique ($R_{int} = 0.036$). The final $R(F^2)$ and $wR(F^2)$ were 0.048 and 0.152 for observed reflections [$F^2 > 2\sigma(F^2)$] used in all calculations.²⁷

Single crystal of 7b was recrystallized from PhH-hexane.

Crystal data for 7b. All the measurements were performed using a Rigaku RAXIS-RAPID radiation diffractometer with graphite monochromated Mo-K α radiation: yellow prisms, $C_{27}H_{20}ONAs$, M = 449.38, monoclinic, a = 8.0799(5), b =13.4532(7), c = 9.6920(4) Å, V = 1043.41(8) Å³, T = 298 K, space group $P2_1$ (no. 4), Z = 2, μ (Mo-K α) = 16.48 cm⁻¹, 2490 reflections measured, 2517 unique ($R_{int} = 0.068$). The final $R(F^2)$ and $wR(F^2)$ were 0.036 and 0.103 for observed reflections [$F^2 > 3.0\sigma(F^2)$] used in all calculations.²⁸

Acknowledgements

Financial support from Waseda University Grants for Special Research Projects is gratefully acknowledged. The authors also thank the Materials Characterization Central Laboratory, Waseda University, for technical assistance with spectral data, elemental analyses, and X-ray structure analyses.

References

 (a) D. Lloyd, I. Gosney and R. A. Ormiston, *Chem. Soc. Rev.*, 1987, 16, 45; (b) H. Yaozeng and S. Yanchang, *Adv. Organomet. Chem.*, 1982, 20, 115; (c) D. Lloyd and I. Gosney, *The Chemistry of Organic Arsenic, Antimony, and Bismuth Compounds*, ed. S. Patai, John Wiley

[†] CCDC reference number(s) 187524–187525. See http://www.rsc.org/ suppdata/p1/b2/b205612g/ for crystallographic files in .cif or other electronic format.

& Sons, Chichester, New York, Brisbane, Toronto, Singapore, 1994, pp. 657–693.

- 2 M. Shao, X. Jin, Y. Tang, Q. Huang and Y. Huang, *Tetrahedron Lett.*, 1982, 5343.
- 3 X. Zong-Xiang and Z. Zhi-ming, *Acta Chim. Sin. (Engl. Ed.)*, 1983, **41**, 148.
- 4 F. Zhao-Chang and S. Yang-Chang, Acta Chim. Sin., 1984, 42, 759.
- 5 I. Kawamoto, T. Hata, Y. Kishida and C. Tamura, *Tetrahedron Lett.*, 1972, 1611.
- 6 I. Kawamoto, T. Hata, Y. Kishida and C. Tamura, *Tetrahedron Lett.*, 1971, 2417.
- 7 I. Kawamoto, Y. Sugiyama and Y. Kishida, *Tetrahedron Lett.*, 1973, 877.
- 8 I. Kawamoto, Y. Sugiyama, N. Soma and Y. Kishida, Chem. Lett., 1972, 931.
- 9 M. Nitta and S. Naya, J. Chem. Res. (S), 1998, 522 ((M), 1998, 2363).
- 10 H. Yamamoto, M. Ohnuma and M. Nitta, J. Chem. Res. (S), 1999, 173 ((M), 1999, 901).
- 11 M. Nitta, Y. Mitsumoto and H. Yamamoto, J. Chem. Soc., Perkin Trans. 1, 2001, 1901.
- 12 T. Takayasu and M. Nitta, J. Chem. Soc., Perkin Trans. 1, 1997, 681.
- 13 H. Depoorter, J. Lieer and G. Van. Mierlo, Bull. Soc. Chim. Belg., 1968, 77, 521.
- 14 (a) C. M. Moorhoff, Synth. Commun., 1998, 28, 2925; (b) A. J. Bellamy, G. Howat and I. S. MacKirdy, J. Chem. Soc., Perkin Trans. 2, 1978, 786.
- 15 (a) T. Nozoe, K. Takase and T. Sato, *Proc. Jpn. Acad.*, 1954, **30**, 473;
 (b) G. Biggi, F. Del Cima and F. Pietra, *J. Am. Chem. Soc.*, 1973, **95**, 710; (c) F. Pietra, *Acc. Chem. Res.*, 1979, **12**, 132.
- 16 (a) R. O. Day, J. M. Holmes, A. C. Sau, J. R. Deviller, R. R. Holmes and J. A. Delters, *J. Am. Chem. Soc.*, 1982, **104**, 2127; (b) R. H. Fish and R. S. Tannous, *Organometallics*, 1982, **1**, 1238; (c) C. A.

Poutasse, R. O. Day, J. M. Holmes and R. R. Holmes, *Organometallics*, 1985, **4**, 708; (*d*) R. R. Holmes, R. O. Day and A. C. Sau, *Organometallics*, 1985, **4**, 714.

- 17 A. Bondi, J. Phys. Chem., 1964, 68, 441.
- 18 (a) G. Ferguson, D. F. Rendle, D. Lloyd and M. I. C. Singer, J. Chem. Soc., Chem. Commun., 1971, 1647; (b) G. Ferguson and D. F. Rendle, J. Chem. Soc., Dalton Trans., 1975, 1284; (c) G. Ferguson and D. F. Rendle, J. Chem. Soc., Dalton Trans., 1976, 171.
- 19 L. Pauling, *The Naure of the Chemical Bond*, Cornell University Press, Ithaca, New York, 1948.
- A. Krebs and B. Schrader, Justus Liebigs Ann. Chem., 1967, 709, 46.
 M. J. Barrow, O. S. Mills and G. Filippini, J. Chem. Soc., Chem. Commun., 1973, 66.
- 22 (a) M. Regitz, Methoden der organische chemie: Organische phosphorverbindungen Bande 1, Georg Thieme Verlag, Stuttgart, New York, 1982; H. Staudinger and J. Meyer, Helv. Chim. Acta, 1919, 2, 619; (c) P. Frøyen, Acta Chem. Scand. Ser. B, 1974, 28, 586; (d) T. Saito, M. Nakane, T. Miyazaki and S. Motoki, J. Chem. Soc., Perkin Trans. 1, 1989, 2140; (e) S. Trippet and D. M. Walker, J. Chem. Soc., 1959, 3874; (f) Y. Ohshiro, Y. Mori, T. Minami and T. Agawa, J. Org. Chem., 1970, 35, 2076; (g) H. Bestmann and S. Pfohl, Angew. Chem., Int. Ed. Engl., 1969, 10, 762.
- 23 (a) J. J. Monagle, J. Org. Chem., 1962, 27, 3851; (b) P. Frøyen, Acta Chem. Scand., 1969, 23, 2935.
- 24 I. Ueda, M. Matsuo, K. Taniguchi, S. Shigenaga and T. Ogawa, Jpn Kokai Tokkyo Koho, 1989, Jp 6403, 172 [8903, 172] (Chem. Abstr., 1989, 111, 57734g).
- 25 M. Oda and Y. Kitahara, Chem. Ind., 1969, 27, 920.
- M. Oda and Y. Kitahara, J. Chem. Soc., Chem. Commun., 1969, 352.
 CRYSTAN, Crystal Structure Analysis Package, MAC Science Co., Ltd., Yokohama, Kanagawa 222–0033, Japan, 1994.
- 28 G. M. Sheldrick, SHELXL-97, Program for the Refinement of Crystal Structures, University of Göttingen, Germany, 1997.