

PMe₃ at 20 °C to soluble derivatives which then remain in solution when pure **5** (60% yield) is precipitated with hexanes.

Significantly if Mo(N₂)₂(PMePh₂)₄ is used in place of **1** in the reaction then a mixture of **3** (35% based on the starting Mo complex **1**), **5** (20%), and WCl₂(PMePh₂)₃ (30%) is formed. Thus the regioselectivity of the reaction could be attributed to the ability of **1** to coordinate via the η⁶-ligand to the tungsten and hence favor heterobimetallic bonding.¹² The complex W₂Cl₄(PMePh₂)₄ (**6**)¹⁰ is not produced in any of these reactions. Preliminary work shows that MoWCl₄(PMe₂Ph)₄¹⁸ is produced when Mo(η⁶-PhPMe₂)(PMe₂Ph)₃¹⁹ is used in place of **1** in Scheme I.

Complex **5** is most readily identified by the coupling constants of the inequivalent phosphorus atoms on the ¹⁸³W⁴-Mo isotopomer in the ³¹P NMR spectrum. The *J* values are similar to those observed for the ¹⁸³W⁴-W isotopomer of complex **6** (Table I). The ¹H NMR spectrum of **5** in C₆D₆ shows two methyl resonances as virtual triplets at 1.90 and 2.07 ppm and two sets of ortho phenyl proton multiplets at 7.56 and 7.73 ppm consistent with two types of phosphines, whereas **3** gives only one methyl peak at 1.98 ppm and one ortho proton multiplet at 7.65 ppm. The λ_{max}(visible) of **5** in benzene at 650 nm falls in the range of δ → δ* transitions of homonuclear complexes,^{5,8,14,20} including Mo₂Cl₄(PMe₃)₄ at 582 nm⁵ and W₂Cl₄(PMe₃)₄ at 657 nm.⁵ It is interesting that the Mo⁴-W complex with bridging carboxylates² has a yellow color; perhaps a δ → π* (ligand) transition, recently observed in the 500–600 nm region for W₂(O₂CR)₄ complexes,²¹ complicates the visible spectrum when these bridging ligands are present. No frequency (IR or Raman) assignable to ν(Mo⁴-W) is observed in the range 250–400 cm⁻¹. Solutions of **5** are sensitive to water and oxygen. Complex **5** decomposes when refluxed in degassed toluene; it does not disproportionate to **3** and **6**.

The substitution reactions of **5** are proving to be fascinating. For example, **5** reacts with excess PMe₃ at 20 °C for 3 h to give (Me₃P)₂Cl₂Mo⁴WCl₂(PMePh₂)₂ (**7**)²² where only the phosphines on the molybdenum have been substituted. Further reaction (45 °C, 19 h) gives MoWCl₄(PMe₃)₄.²³ The course of reaction differs from that reported for Mo₂Me₄(PEt₃)₄.²⁴

The structure of **5** (Scheme I) is proposed on the basis of the D_{2d} geometry of the homonuclear complexes.⁵ Work is under way to obtain crystals of **5** or its derivatives suitable for X-ray diffraction in order to verify the structure and obtain further data on this unique heteronuclear bond.

Note Added in Proof. Preliminary X-ray crystallographic analysis of complex **7** confirms the geometry shown in Scheme I. The structure has refined to *R* = 3.8% in the space group I₂/a. The molybdenum-tungsten distance is 2.209 (1) Å. (Personal communication from Jeffrey F. Sawyer.)

Acknowledgment. This work is supported by the Natural Sciences and Engineering Research Council of Canada. We thank Douglas Blue and Dr. William M. Reichert for recording the Raman spectra and Dr. Jeffrey Zubkowski for the preparation of Mo(η⁶-PhPMe₂)(PMe₂Ph)₃.

Supplementary Material Available: Preparation of **3** and **5** and ¹H and ³¹P NMR spectra of **5** (4 pages). Ordering information is given on any current masthead page.

(17) Identified by ³¹P NMR (C₆H₆) δ -16.7 (1 P, *J*_{PW} = 416, *J*_{PP} < 5 Hz), 0.0 (2 P, *J*_{PW} = 339, *J*_{PP} < 5 Hz).¹⁶

(18) ³¹P NMR (C₆H₆) δ -20.1 (2 PMo, *J*_{PP}³ = 24.0, *J*_{PW}² = 50 ± 5 Hz), 17.4 (2PW, *J*_{PP}³ = 24.0, *J*_{PW}¹ = 280 Hz).

(19) Anker, M. W.; Chatt, J.; Leigh, G. J.; Wedd, A. G. *J. Chem. Soc., Dalton Trans.* 1975, 2639–2645.

(20) McGinnis, R. N.; Ryan, T. R.; McCarley, R. E. *J. Am. Chem. Soc.* 1978, 100, 7900–7902.

(21) Cotton, F. A.; Wang, W. *Inorg. Chem.* 1984, 23, 1604–1610.

(22) ³¹P NMR (C₆H₆) δ -28.9 (2 PMe₃Mo, *J*_{PP}³ = 24.4 Hz), +21.7 (2 PMePh₂W, *J*_{PP}³ = 24.4, *J*_{PW}¹ = 273 Hz).

(23) ³¹P NMR (C₆H₆) δ -28.9 (2 PMe₃Mo, *J*_{PP}³ = 25.6 Hz), +9.5 (2 PMe₃W, *J*_{PP}³ = 25.6 Hz, PMe₃¹⁸³W, *J*_{PW}¹ = 271, *J*_{PP}³ = 24.4 Hz).

(24) Girolami, G. S.; Mainz, V. V.; Andersen, R. A.; Vollmer, S. H.; Day, V. W. *J. Am. Chem. Soc.* 1981, 103, 3953–3955.


Synthesis of Anatoxin-a: Very Fast Death Factor

Joseph J. Tufariello,* Harold Meckler, and
K. Pushpananda A. Senaratne

Department of Chemistry
State University of New York at Buffalo
Buffalo, New York 14214

Received July 12, 1983

A class of microalgae, the cyanophytes, contains several toxic strains, including *Anabaena flos-aquae*.¹ Graphic descriptions of the death of animals induced by such blue-green algae have been recorded.^{2,3} An alkaloidal toxin identified from these sources was shown⁴ to be 2-acetyl-9-azabicyclo[4.2.1]non-2-ene (anatoxin-a, **1**), also designated "Very Fast Death Factor" (VFDF).³



This structural assignment was confirmed by X-ray crystallography.⁵ Anatoxin-a has been shown to be a potent muscarinic¹ and nicotinic agonist⁶ and has engendered synthetic interest.^{7–9}

We report herein an efficient, nitrene-based entry to this interesting natural product. It was anticipated^{10,11} that the addition of 1-pyrroline 1-oxide (**2**) to *trans*-3,5-hexadien-2-ol (**3**)^{12–14} would exhibit the desired site selectivity and regioselectivity (cf., Scheme I) to afford the isoxazolidine **4a**.¹⁵ The latter, formed in 70% yield, upon oxidation with manganese dioxide (Celite, methylene chloride), produces the ketone **6a**, which exhibits a clean quartet at δ 4.71 (*J* = 6.71 Hz), assignable to the C-5 proton (isoxazolidine numbering).

Oxidative cleavage^{16,17} of the isoxazolidine ring with *m*-chloroperbenzoic acid gives the nitrene **5** as the sole identifiable product in 79% yield. The product is a single regioisomer exhibiting a broad singlet in its ¹H NMR spectrum (CDCl₃) 7.0 ppm (1 H) characteristic of the proton at C-2 of the nitrene function. Warming of a solution containing the nitrene to 45 °C leads to the formation of a single cycloadduct, **6a** in 71% overall

(1) Gorham, P. R.; Carmichael, W. W. *Pure Appl. Chem.* 1980, 52, 165–174.

(2) Rose, E. T. *Proc. Iowa Acad. Sci.* 1953, 60, 738–745.

(3) Gorham, P. R.; McLachlan, J.; Manner, U. T.; Kim, W. K. *Verh.—Int. Ver. Theor. Angew. Limnol.* 1964, 15, 796–804.

(4) Devlin, J. P.; Edwards, O. E.; Gorham, P. R.; Hunter, N. R.; Pike, P. K.; Stavric, B. *Can. J. Chem.* 1977, 55, 1357–1371.

(5) Huber, L. S. *Acta Crystallogr., Sect. B* 1972, B28, 2577–2582.

(6) Spivak, C. E.; Witkop, B.; Albuquerque, E. X. *Mol. Pharmacol.* 1980, 18, 384–394.

(7) Campbell, H. F.; Edwards, O. E.; Kolt, R. *Can. J. Chem.* 1977, 55, 1372–1379.

(8) Campbell, H. F.; Edwards, O. E.; Elder, J. W.; Kolt, R. *J. Pol. J. Chem.* 1979, 53, 27–37.

(9) Rapoport, H.; Bates, H. A. *J. Am. Chem. Soc.* 1979, 101, 1259–1267.

(10) Tufariello, J. J. In "1,3-Dipolar Cycloaddition Chemistry"; Padwa, A., Ed.; Wiley-Interscience: New York, 1984; Chapter 9.

(11) Houk, K. N.; Sims, J.; Watts, C. R.; Laskus, L. J. *J. Am. Chem. Soc.* 1973, 95, 7301–7315.

(12) Crombie, L.; Harper, S. H.; Thompson, D. *J. Chem. Soc.* 1951, 2906–2915.

(13) Woods, G. F.; Sanders, H. *J. Am. Chem. Soc.* 1946, 68, 2483–2485.


(14) (a) Woods, G. F.; Sanders, H. *J. Am. Chem. Soc.* 1947, 69, 2926–2928. (b) Woods, G. F.; Schwartzman, L. H. *J. Am. Chem. Soc.* 1948, 70, 3394–3396.

(15) The stereochemical relationship of the hydrogens at C-3 and C-5 of the isoxazolidine ring are assigned by analogy with work done in related systems. Tufariello, J. J.; Puglis, J. M., unpublished observations.

(16) LeBel, N. A.; Post, M. E.; Huang, D. J. *J. Org. Chem.* 1979, 44, 1819–1823.


(17) Tufariello, J. J.; Mullen, G. B.; Tegeler, J. J.; Trybulski, E. J.; Wong, S. E.; Ali, S. K.; A. *J. Am. Chem. Soc.* 1979, 101, 2435.

Scheme I



(a) heat, benzene (70%); (b) MnO_2 , celite, CH_2Cl_2 (96%); (c) MCPBA, CH_2Cl_2 (71%); (d) $(\text{CH}_3\text{OH})_2$, *p*-TsOH (96%); (e) MsCl , Et_3N (94%); (f) LiAlH_4 , NiCl_2 , THF (-40°); (g) *p*-TsOH, acetone; (h) NaHCO_3 , H_2O ; (i) $(t\text{-BuCO})_2\text{O}$, CHCl_3 ; (j) 3NHCl , EtOAc .

yield from **4b**. It has already been reported¹⁸ that such *N*-alkenyl nitrones can cyclize to produce either the product of 6-ring closure (e.g., **10**) or that of 7-ring closure (e.g., **6a**). A preference was observed in most cases for the transition state resulting in 6-ring closure. This was attributed to diminished strain when compared to its counterpart leading to 7-ring closure.¹⁸ Clearly, nitronium **5** can undergo either 6- or 7-ring closure (cf. eq 1 and 2).



We believed that **5** might overcome the normal predilection for 6-ring closure because of the natural tendency of nitrones, in intermolecular cycloaddition reactions, to afford the adduct with the nitronium oxygen bound to the β -carbon of the dipolarophilic α,β -unsaturated carbonyl system.^{17,19-21}

We could only identify a single cycloadduct from **5** (i.e., **6a**).²² This adduct appears to be of kinetic origin since it is formed conveniently at 45°C . This fortunate circumstance results in the construction of the ring system of anatoxin-*a*. Subsequent ketalization and mesylation results in the formation of ketal mesylate **6b**. Finally, treatment of **6b** with a 1:1 (molar) mixture of

(18) Oppolzer, W.; Siles, S.; Snowden, R. L.; Baker, B. H.; Petrziika, M. *Tetrahedron Lett.* **1979**, 4391-4394.

(19) (a) Huisgen, R. *Angew. Chem., Int. Ed. Engl.* **1963**, 2, 565-568. (b) Black, D. St. C.; Crozier, R.; Davis, Y. *Synthesis* **1975**, 205-221.

(20) Tufariello, J. J.; Tette, J. P. *J. Org. Chem.* **1975**, 40, 3866-3869.

(21) Tufariello, J. J. *Acc. Chem. Res.* **1979**, 12, 396-403.

(22) The ^1H NMR evidence strongly supports this assignment. This data will be considered in detail in the full report.

$\text{LiAlH}_4/\text{NiCl}_2$ in THF (-40°C) leads to **7**. When direct conversion into anatoxin-*a* by acid hydrolysis proved to be problematic, the hydroxy ketal **7** was treated with a stoichiometric amount of *p*-toluenesulfonic acid in acetone to induce both trans ketalization and dehydration, thereby affording the *p*-toluenesulfonic acid salt (i.e., **18**) of the natural product. For purposes of purification, the crude salt **8** was treated with 2 equiv of sodium bicarbonate and di-*tert*-butyl dicarbonate.²³ The resultant *tert*-butyl carbamate **9** (43% overall yield from mesylate **6b**) was subjected to acid hydrolysis²⁴ to give anatoxin-*a* hydrochloride. The ^1H NMR, IR, and mass spectral comparisons of the synthetic material with the natural product confirmed the successful outcome of the synthetic effort.

Acknowledgment. We thank Nicholas Saccomano, Joseph Koslowski, David Scherer, Bridget McCourtney, and John Brinkman for technical assistance. We appreciate the efforts of Dr. O. E. Edwards of the National Research Council of Canada in supplying the ^1H NMR and IR spectra of authentic anatoxin-*a*. Finally, we thank the National Institutes of Health (GM 25303) for financial support.

Registry No. (\pm)-**1-HCl**, 70470-07-4; **2**, 24423-88-9; (\pm)-**3**, 3280-51-1; **4a**, 92844-74-1; (\pm)-**4b**, 92844-75-2; **5**, 92844-79-6; (\pm)-**6a**, 92844-76-3; (\pm)-**6b**, 92844-77-4; (\pm)-**7**, 92844-78-5; (\pm)-**8**, 92844-80-9; (\pm)-**9**, 92998-50-0.

Supplementary Material Available: Experimental section for **4a, b**, **6a, b**, **7**, **9**, and **1-HCl** and tables of mass spectral data for **4a, 6a, b**, **7**, **9**, and **1-HCl** (9 pages). Ordering information is given on any current masthead page.

(23) Tarbell, D. S.; Yamato, Y.; Pope, B. M. *Proc. Natl. Acad. Sci. U.S.A.* **1972**, 69, 730-732.


(24) Stahl, G. L.; Walter, R.; Smith, C. W. *J. Org. Chem.* **1978**, 43, 2285-2286.

1,8-Biphenylenediol Forms Two Strong Hydrogen Bonds to the Same Oxygen Atom

Jack Hine,* Kyunghye Ahn, Judith C. Gallucci, and Shwn-Meei Linden

Department of Chemistry, The Ohio State University
Columbus, Ohio 43210
Received August 6, 1984

The molecular geometry of 1,8-biphenylenediol is such that the two hydroxyl groups should be capable of forming hydrogen bonds simultaneously to the same basic atom. This expectation is now supported by the isolation and X-ray structure determination of crystalline adducts of the diol¹ with *N,N,N',N',N'',N''*-hexamethylphosphoramide (**1**)² and with 1,2,6-trimethyl-4-pyridone^{3,4} and 2,6-dimethyl- γ -pyrone.⁵



(1) Blatchly, J. M.; Garner, D. V.; McOmie, J. F. W.; Watts, M. L. *J. Chem. Soc. C* **1968**, 1545-1549.

(2) Prepared by dissolving the diol (mp 222°C) in the amide (HMPA) and removing excess amide by heating in a high vacuum. Recrystallized from cyclohexane containing a little chloroform: mp $132-134^\circ\text{C}$.

(3) Van Allan, J. A.; Reynolds, G. A.; Alessi, J. T.; Chang, S. C.; Joines, R. C. *J. Heterocycl. Chem.* **1971**, 8, 919-922.

(4) Adduct prepared by dissolving equimolar amounts of the pyridone (mp $245-245.5^\circ\text{C}$) and diol in acetonitrile and allowing the solvent to evaporate slowly. Decomposes without melting at $268-271^\circ\text{C}$.

(5) Adduct prepared by dissolving equimolar amounts of pyrone (mp $133-137^\circ\text{C}$) and diol in chloroform and crystallized by slow evaporation, first from chloroform-cyclohexane and then from ethyl acetate: mp $182-183^\circ\text{C}$.