


**Table I.** Comparison of  $^{13}\text{C}$  NMR Chemical Shifts<sup>a</sup> of Trihalomethyl Cations<sup>b</sup> and the Corresponding Trihalomethanes<sup>c</sup>

trihalomethyl cation	$\delta^{13}\text{C}$	trihalomethane	$\delta^{13}\text{C}$	$\Delta\delta^{13}\text{C}$
$^+\text{CCl}_3$	236.3	$\text{HCCl}_3$	77.7	158.6
$^+\text{CBr}_3$	207	$\text{HCCl}_3$	+12.3	194.7
$^+\text{CI}_3$	95	$\text{HCl}_3$	-139.7	234.7

<sup>a</sup>Chemical shifts in ppm from external tetramethylsilane signal. <sup>b</sup>In  $\text{SbF}_5/\text{SO}_2\text{ClF}$  solution at  $-80^\circ\text{C}$ . <sup>c</sup>Data taken from ref 9.

**Figure 1.** A plot of  $^{13}\text{C}$  NMR chemical shifts of halomethyl cations vs  $^{11}\text{B}$  NMR chemical shifts of isoelectronic isomethanes.

then reacts with benzene to give benzotrifluoride. Benzotrifluoride on subsequent reaction with benzene and aluminum trichloride gives triphenylmethyl chloride.

In the gas phase the  $\text{CF}_3^+$  ion is stable and is an abundant species, specially in the mass spectra<sup>11</sup> of organofluorine compounds. However, in solution the great strength of the C-F bond in  $\text{CF}_4$  (ca. 140 kcal/mol) leads to rapid quenching of  $\text{CF}_3^+$  to  $\text{CF}_4$  even in low nucleophilicity fluorinated superacid media.

We have compared the  $^{13}\text{C}$  NMR chemical shifts of the trihalomethyl cations to those of the respective trihalomethanes and found a consistent trend (see Table I). The  $\Delta\delta^{13}\text{C}$  chemical shift differences<sup>12</sup> are 158.6, 194.7, and 234.7 for the trichloro-, tribromo-, and triiodomethyl derivatives, respectively. This trend is in agreement with the positive charge stabilization (increased back-bonding) order of  $\text{Cl} > \text{Br} > \text{I}$ . The  $^{13}\text{C}$  shifts of trihalomethyl cations correlate well with the electronegativities of the halogen atoms.<sup>14</sup>

The  $^{13}\text{C}$  NMR chemical shifts of the trihalomethyl cations and related reported methyl substituted mono- and dihalocarbenium ions<sup>3</sup> also correlate well with the  $^{11}\text{B}$  NMR chemical shifts of the corresponding isomethanes, isoelectronic boron halides<sup>13</sup> (see Figure 1). An excellent linear relationship is obtained with a correlation coefficient of unity.<sup>15</sup> From this plot, with the known  $^{11}\text{B}$  NMR chemical shift of  $\text{BF}_3$  (10.4 ppm) we can estimate the corresponding  $^{13}\text{C}$  NMR shift of the yet unknown trifluoromethyl cation as to be 140 ppm which is only 31.2 ppm more deshielded

than trifluoromethane at  $\delta^{13}\text{C}$  118.8. This indicates substantial stabilization of the trifluoromethyl cation by fluorine back-bonding. Experimental verification, however, must await preparation of the still elusive  $\text{CF}_3^+$  ion under long-lived conditions.

**Acknowledgment.** Support of our work by the National Institutes of Health is gratefully acknowledged.

### Anatoxin-a(s), a Potent Anticholinesterase from *Anabaena flos-aquae*

Shigeki Matsunaga, Richard E. Moore,\* and Walter P. Niemczura


Department of Chemistry, University of Hawaii  
Honolulu, Hawaii 96822

Wayne W. Carmichael

Department of Biological Sciences  
Wright State University, Dayton, Ohio 45435

Received June 1, 1989

Anatoxin-a(s) is a neurotoxic alkaloid associated with the blue-green alga *Anabaena flos-aquae*.<sup>1</sup> Its potent toxicity ( $\text{LD}_{50}$  20–40  $\mu\text{g}/\text{kg}$  mice) is attributed to exceptional anticholinesterase activity.<sup>2</sup> We report here the isolation of anatoxin-a(s) from a cultured strain NRC 525-17 and a field-collected bloom implicated in animal poisonings<sup>3</sup> and the determination of its structure as 1.



Freeze-dried alga was extracted with 0.05 N  $\text{AcOH}/\text{EtOH}$ . The filtered extract was partitioned between water and  $\text{CH}_2\text{Cl}_2$ , the aqueous layer was washed with  $n\text{-BuOH}$  and evaporated in vacuo, and the residue was extracted successively with small portions of 0.05 N  $\text{AcOH}/\text{MeOH}$  and 0.05 N  $\text{AcOH}/\text{EtOH}$  to give a toxic concentrate. Gel filtration on Toyopearl HW40F (Supelco)<sup>4</sup> followed by HPLC on CN and ODS columns gave pure anatoxin-a(s) as a colorless solid in 0.05% yield. Toxin isolation was followed by assaying fractions for anticholinesterase activity.<sup>5</sup> Anatoxin-a(s) decomposed rapidly in basic solution but was

(11) Murdoch, H. D.; Weiss, E. *Helv. Chim. Acta* **1962**, *45*, 1927.  
(12) Schleyer, P. v. R.; Lenoir, D.; Mison, P.; Liang, G.; Prakash, G. K. S.; Olah, G. A. *J. Am. Chem. Soc.* **1980**, *102*, 683.

(13) Nuclear Magnetic Resonance Spectroscopy of Boron Compounds. Nöth, H.; Wrackmeyer, B. A volume of *NMR Basic Principles and Progress*; Diehl, P.; Fluck, E.; Kosfeld, R., Eds.; Springer-Verlag: New York, 1978.

(14) Correlation with Pauling's electronegativities of the halogens were excellent ( $r^2 = 1$ ). From this correlation the  $^{13}\text{C}$  NMR shift of  $^+\text{CF}_3$  is estimated to be  $\approx 150$  ppm.


(15) The equation from the plot is  $\delta^{13}\text{C} = 112.4272 + 2.6504 \delta^{11}\text{B}$ ,  $r^2 = 1$ .

(1) Mahmood, N. A.; Carmichael, W. W. *Toxicon* **1986**, *24*, 425.  
(2) (a) Mahmood, N. A.; Carmichael, W. W. *Toxicon* **1987**, *25*, 1221. (b) Cook, W. O.; Beasley, V. R.; Dahlem, A. M.; Dellinger, J. A.; Harlin, K. S.; Carmichael, W. W. *Toxicon* **1988**, *26*, 750.

(3) Mahmood, N. A.; Carmichael, W. W.; Pfahler, D. *Am. J. Vet. Res.* **1988**, *49*, 500.

(4) Harada, K.; Kimura, Y.; Suzuki, M.; Dahlam, A. M.; Beasley, V. R.; Carmichael, W. W. Abstracts, Annual Meeting of the Pharmaceutical Society of Japan, Hiroshima, 1988; p 288.

(5) Ellman, G. L.; Courtney, K. D.; Andres, V., Jr.; Featherstone, R. M. *Biochem. Pharm.* **1961**, *7*, 88. A modification of the assay described in this paper was used to observe the enzyme inhibitory activity. Solutions of samples to be tested were first spotted on filter paper or a TLC plate and sprayed with a mixture of acetylthiocholine (5 mg/mL) and 5,5'-dithiobis(2-nitrobenzoic acid) (5 mg/mL) in ethanol. After drying in a stream of air, a solution of electric el acetylcholinesterase (EC 3.1.1.7, 0.5 units/mL) was next applied to the paper or plate. Active samples showed a white zone on a dense yellow background.

Scheme 1<sup>a</sup>

<sup>a</sup> Reagents and conditions: (a) *N*-hydroxysuccinimide (1.2 equiv), DCC (1.2 equiv), dioxane, 0 °C 10 min → room temperature 15 h; (b) Me<sub>2</sub>NH (2 equiv) in ether, room temperature 10 min; (c) CF<sub>3</sub>CO<sub>2</sub>H, room temperature 1 h; (d) 10% Pd-C, H<sub>2</sub>, MeOH; (e) excess BH<sub>3</sub>/Me<sub>2</sub>S, THF, reflux 15 h; (f) *S,S*-dimethyl-*N*-tosyliminodithiocarbonimidate (1 equiv), EtOH, reflux 15 h; (g) 48% HBr, reflux 4 h.

relatively stable in neutral or acidic (pH 3–5) media.<sup>6</sup> Anatoxin-a(s) from cultured and field-collected *A. flos-aquae* exhibited identical chemical and spectral properties, including optical (CD in H<sub>2</sub>O: [θ]<sub>207</sub> −3300, [θ]<sub>232</sub> +3900).

Mass spectral analysis of anatoxin-a(s) [positive FABMS (*m/z* 253.1067, MH<sup>+</sup>), negative FABMS (*m/z* 251, M-H<sup>−</sup>), FDMS (*m/z* 253, MH<sup>+</sup>)] indicated the molecular formula C<sub>7</sub>H<sub>17</sub>N<sub>4</sub>O<sub>4</sub>P. The <sup>1</sup>H and <sup>13</sup>C NMR spectra<sup>7,8</sup> revealed the presence of dimethylamino and P-OMe (*J*<sub>H-P</sub> 11.0 Hz; *J*<sub>C-P</sub> 6.7 Hz) groups, a 1,2,3-trisubstituted propane unit, and an sp<sup>2</sup> carbon that was fully substituted by heteroatoms (δ<sub>C</sub> 163.7). The methoxyl protons and carbon were the only ones showing distinct coupling to phosphorus. Only one signal was seen in the <sup>31</sup>P NMR spectrum and its chemical shift (δ 6.16) agreed well for either a phosphate ester or phosphoramidate.<sup>9</sup> The *J*<sub>gem</sub> (−10.1 Hz) and <sup>1</sup>*J*<sub>H,C</sub> values for the protons in one of the methylenes of the propane unit suggested that this CH<sub>2</sub> was in a five-membered ring, along with the adjacent CH.

More information was obtained from NMR analysis of anatoxin-a(s) that had been uniformly enriched to 50% <sup>13</sup>C and 90+% <sup>15</sup>N<sup>10,11</sup> (See also Supplementary Material). The following conclusions could be made: (1) The sp<sup>2</sup> carbon at 163.7 ppm was connected to three nitrogens of a guanidine group and that two of these nitrogens were attached to the CH and CH<sub>2</sub> in the five-membered ring. (2) The NMe<sub>2</sub> group was connected to the side-chain CH<sub>2</sub> on the resulting imidazoline.<sup>12</sup> (3) No nitrogens

were connected to the phosphorus; a methyl phosphate group was therefore present in the toxin. (4) The methyl phosphate group was attached to one of the nitrogens (<sup>2</sup>*J*<sub>P-N</sub> 4 Hz); the toxin was therefore zwitterionic.

Anatoxin-a(s) slowly decomposed during storage at −20 °C into a mixture of 2, 3 (sometimes), and monomethyl phosphate, separable by Toyopearl HW40F chromatography. Compound 2 (FABMS, MH<sup>+</sup> *m/z* 159.1245; CD in H<sub>2</sub>O, [θ]<sub>235</sub> +2400), which differed from compound 3 by an oxygen, could be converted into 3 (FABMS, MH<sup>+</sup> *m/z* 143.1298; CD in H<sub>2</sub>O, [θ]<sub>198</sub> +11 000) by catalytic hydrogenation (Pd-C/MeOH). Hydrolytic removal of the monomethyl phosphate group caused a diamagnetic shift of the H-5 signal from 4.71 ppm in 1 to 4.48 ppm in 2; the methylene <sup>1</sup>H chemical shifts, however, were essentially identical for the two compounds. Although the <sup>1</sup>H chemical shifts for 3 were similar to those for 2, except for one of the H-6 signals which was shifted upfield appreciably (−0.37 ppm), the <sup>13</sup>C chemical shifts were significantly different, i.e., upfield for C-5 (−7.8 ppm) and C-2 (−1.9 ppm) and downfield for C-4 (+2.7 ppm) and C-6 (+2.4 ppm).<sup>13,14</sup> These chemical shift differences were consistent with placements of the hydroxyl group on N-1 in 2<sup>15</sup> and the methyl phosphate group on N-1 in 1. Anatoxin-a(s) therefore had to have structure 1.

To elucidate the absolute configuration at C-5, *R*- and *S*-3 were prepared from *D*- and *L*-asparagine, respectively (Scheme I). *N*-(2-Benzyloxycarbonyl)-*N*-(3-*tert*-butoxycarbonyl)-*L*-2,3-diaminopropionic acid (4),<sup>16</sup> for example, was converted to dimethylamide 5 via the *N*-hydroxysuccinimide ester.<sup>17</sup> After removal of the amino-protecting groups (trifluoroacetic acid; H<sub>2</sub>/Pd-C), the resulting diamine was reduced with BH<sub>3</sub>-Me<sub>2</sub>S complex<sup>18</sup> to give the triamine 6, which was then treated with *S,S*'-dimethyl-*N*-tosyliminodithiocarbonimidate<sup>19</sup> to furnish the tosylguanidine 7. Removal of the *N*-tosyl group was accomplished by refluxing 7 in 48% HBr.<sup>20</sup> Synthetic 3 showed identical chromatographic properties and <sup>1</sup>H and <sup>13</sup>C NMR spectra with the degradation product. The CD spectrum of 3 derived from anatoxin-a(s) was identical with that of synthetic 3 from *L*-Asn ([θ]<sub>198</sub> +13 000), which meant that C-5 was *S*.

Anatoxin-a(s) is a unique phosphate ester of a cyclic *N*-hydroxyguanidine. The structure and reactivity is reminiscent of an ester of *N*-hydroxysuccinimide or 1-hydroxybenzotriazole. Cholinesterase inactivation may proceed by nucleophilic attack of Ser at the esteratic site of the enzyme on the phosphate group of 1 with concomitant elimination of 2.<sup>21</sup>

**Acknowledgment.** This research was supported by NSF Grant CHE-8800527 (R.E.M.) and in part by U.S. Army Medical Research Acquisition Activity, Contract No. 17-87-C-7019, Department of Army Medical Defense (W.W.C.). S.M. thanks the Naito Foundation for supplemental fellowship support. FABMS and MS-MS studies were carried out at the Midwest Center for Mass Spectrometry (M. L. Gross, director) by R. L. Cerny. Preliminary FABMS studies were conducted at the Institute of Applied Microbiology, University of Tokyo by H.

(6) Air evaporation of a methanolic solution of the toxin results in significant hydrolysis to 2.

(7) All NMR spectra have been determined in D<sub>2</sub>O with 5 μL of acetic acid-d<sub>4</sub> added per 0.5 mL of D<sub>2</sub>O, unless otherwise noted.

(8) 1: <sup>1</sup>H NMR δ 3.00 (s, NMe<sub>2</sub>), 3.47 (dd, *J* = 2.9 and −13.9 Hz, H-6), 3.51 (dd, *J* = 9.7 and −10.1 Hz, H-4), 3.75 (dd, *J* = 9.3 and −13.9 Hz, H-6), 3.79 (d, <sup>3</sup>*J*<sub>H-P</sub> = 11.0 Hz, P-OMe), 4.01 (dd, *J* = 9.4 and −10.1 Hz, H-4), 4.71 (m, H-5); exchangeable proton signals seen in acidic 80% H<sub>2</sub>O/20% D<sub>2</sub>O, δ 8.21 and 7.72 (two v br s, NH<sub>2</sub> on C-2), 7.44 (br s, H-3); <sup>13</sup>C NMR δ 43.9 (v br, NMe<sub>2</sub>), 45.3 (dd, *J*<sub>C-P</sub> = 6.7 Hz, P-OMe), 58.7 (C-6), 60.3 (C-5), 163.7 (C-2).

(9) (a) Tebb, J. C.; In *Phosphorus-31 NMR Spectroscopy in Stereochemical Analysis*; Verkade, J. G.; Quin, L. D., Eds.; VCH Publishers: Florida, 1987; p 1. (b) Gorenstein, D. G. *Prog. NMR Spectrosc.* **1983**, *16*, 1.

(10) The labeled toxin was isolated from *A. flos-aquae* NRC 525-17 that had been grown in culture on NaH<sup>13</sup>CO<sub>3</sub> (99 atom %) and Na<sup>15</sup>NO<sub>3</sub> (99 atom %) by using the procedure described in the following: Moore, R. E.; Bornemann, V.; Niemczura, W. P.; Gregson, J. M.; Chen, J.-L.; Norton, T. R.; Patterson, G. M. L.; Helms, G. L. *J. Am. Chem. Soc.* **1989**, *111*, 6128.

(11) 1 uniformly enriched with <sup>13</sup>C to 50+% and <sup>15</sup>N to 90+%: <sup>1</sup>H NMR signals for NMe<sub>2</sub> and H-6 (<sup>1</sup>*J*<sub>H-C</sub> = 144 Hz), OMe (<sup>1</sup>*J*<sub>C-H</sub> = 148 Hz), H-4 and H-5 (<sup>1</sup>*J*<sub>H-C</sub> = 151 Hz); <sup>13</sup>C NMR δ 43.9 (v br → doublet at 43.8 ppm in MeOH-d<sub>3</sub> or two doublets at 42.0 and 45.7 ppm in D<sub>2</sub>O/CF<sub>3</sub>CO<sub>2</sub>H, *J*<sub>C-N</sub> = 4.1 Hz, NMe<sub>2</sub>), 45.3 (dd, *J*<sub>C-P</sub> = 33.5 Hz, *J*<sub>C-N</sub> = 8.7 Hz, *J*<sub>C-N</sub> = 0, C-4), 56.1 (d, *J*<sub>C-P</sub> = 6.7 Hz, P-OMe), 58.7 (br d, *J*<sub>C-P</sub> = 40.5 Hz, *J*<sub>C-N</sub> = 2–3 Hz, C-6), 60.3 (br t, *J*<sub>C-N</sub> = 0 Hz, C-5), 163.7 (td, *J*<sub>C-N</sub> = 23.5, 23.5, and 11.5 Hz, C-2); <sup>15</sup>N NMR (acidic 4:1 H<sub>2</sub>O/D<sub>2</sub>O) δ 350.6 (NMe<sub>2</sub>), 306.9 and 306.7 (br, NH<sub>2</sub> on C-2 and N-4), 300.2 (N-1); <sup>31</sup>P NMR (acidic 4:1 H<sub>2</sub>O/D<sub>2</sub>O) δ 6.15 (d, <sup>2</sup>*J*<sub>P-N</sub> = 4 Hz).

(12) Also supported by FAB MS-MS and high-resolution data. Fragmentation of MH<sup>+</sup>, for example, leads to *m/z* 58 (Me<sub>2</sub>N<sup>+</sup>=CH<sub>2</sub>) as the base peak.

(13) 2: <sup>1</sup>H NMR δ 3.00 (s, NMe<sub>2</sub>), 3.46 (dd, *J* = 8.0 and −10.0 Hz, H-4), 3.47 (dd, *J* = 4.5 and −13.9 Hz, H-6), 3.75 (dd, *J* = 6.8 and −13.9 Hz, H-6), 3.93 (dd, *J* = 8.5 and −10.0 Hz, H-4), 4.48 (dddd, H-5); <sup>13</sup>C NMR δ 44.7 (q, NMe<sub>2</sub>), 44.9 (t, C-4), 58.7 (t, C-6), 58.8 (d, C-5), 162.5 (s, C-2).

(14) 3: <sup>1</sup>H NMR δ 2.96 (s, NMe<sub>2</sub>), 3.38 (dd, *J* = 4.8 and −13.4 Hz, H-6), 3.48 (dd, *J* = 8.1 and −13.4 Hz, H-6), 3.50 (dd, *J* = 5.4 and −10.1 Hz, H-4), 3.98 (dd, *J* = 9.7 and −10.1 Hz, H-4), 4.55 (dddd, H-5); <sup>13</sup>C NMR δ 44.3 (q, NMe<sub>2</sub>), 47.6 (t, C-4), 51.0 (d, C-5), 61.1 (t, C-6), 160.6 (s, C-2).

(15) When the hydroxyl group is removed from a *N*-hydroxyguanidine, the signals for the carbons α to the relevant nitrogen shift upfield, whereas the ones for the β-carbons shift downfield, e.g., neosaxitoxin and saxitoxin (Shimizu, Y.; Hsu, C.; Fallon, W. E.; Ohima, Y. *J. Am. Chem. Soc.* **1978**, *100*, 6791) and δ-*N*-hydroxyarginine and arginine (Seto, H.; Koyama, M.; Ogino, H.; Tsuruoka, T. *Tetrahedron Lett.* **1983**, *24*, 1805).

(16) Waki, M.; Kitajima, Y.; Izumiya, N. *Synthesis* **1981**, 266.

(17) Anderson, G. W.; Zimmerman, J. E.; Callahan, F. M. *J. Am. Chem. Soc.* **1964**, *86*, 1839.

(18) Brown, H. C.; Narasimhan, S.; Choi, Y. M. *Synthesis* **1981**, 441.

(19) (a) Tsuji, S.; Kusumoto, S.; Shiba, T. *Chem. Lett.* **1975**, 1281. (b) Todricks, J. V.; Rapoport, H. *J. Org. Chem.* **1971**, *36*, 46.

(20) Snyder, H. R.; Heckert, R. J. *J. Am. Chem. Soc.* **1952**, *74*, 2006.

(21) Hucho, F. *Neurochemistry*; VCH Publishers: Weinheim, FRG, 1986.

Morisaki and S. Iwasaki. We thank P. Thorn (WSU) for assistance in toxin isolation, V. Bornemann and B. Moore (UH) for culturing *A. flos-aquae* in  $^{13}\text{C}$  and  $^{15}\text{N}$  enriched media, J. Stewart (UH) and N. A. Mahmood (WSU) for carrying out initial purification and characterization studies on **1** and **2**, and M. A. Tius (UH) for helpful discussions on the synthesis of **3**.

**Supplementary Material Available:**  $^1\text{H}$ ,  $^{13}\text{C}$ , and  $^{31}\text{P}$  NMR spectra of **1**, **2**, and **3**;  $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^{15}\text{N}$ , and  $^{31}\text{P}$  NMR spectra of 50%  $^{13}\text{C}$  and 90+%  $^{15}\text{N}$  enriched **1**, and experimental details for the synthesis of *R*- and *S*-**3** from *D*- and *L*-Asn (16 pages). Ordering information is given on any current masthead page.

## A New Macrobicyclic (Cryptand) Siderophore Containing Three Endocyclic Hydroxamate Donor Groups

Yizhen Sun and Arthur E. Martell\*


Department of Chemistry, Texas A&M University  
College Station, Texas 77843-3255

Received June 14, 1989


The natural hydroxamate siderophores contain bidentate hydroxamate donor groups in acyclic, exocyclic, and endocyclic arrangements.<sup>1</sup> Of these, the sexadentate endocyclic ligands, such as desferrioxamine E, have the highest affinities for iron(III), because of the involvement of the macrocyclic rings in coordination of the metal ion. The only synthetic endocyclic hydroxamate ligand previously reported is a diamino bishydroxamate macrocycle containing pendant carboxylate donors.<sup>2</sup> Up to the present time, no synthetic or natural endocyclic trishydroxamate cryptand has been reported, although the potential of such ligands for effective binding of trivalent metal ions has been pointed out.<sup>3</sup>

The orientation of oxygen donor groups in the hydroxamates places stringent demands on the polyatomic chains which link them together in a manner that places three pairs of oxygen donors symmetrically around a six-coordinate (octahedral) metal ion. Molecular models indicate that eight or more connecting atoms are needed to accomplish this effectively. It is the purpose of this paper to report the synthesis and properties of the first trishydroxamate cryptand. In this case the ligand has nine and 11 atoms between the bidentate hydroxamate units.

Cyanoethylation at 25–35 °C in *p*-dioxane of the 1,1,1-tris(hydroxymethyl)ethane gives the tricyano ether **1** in 83% yield. Hydrolysis of **1** with hydrogen chloride gas in methanol leads to the tricarboxylic acid trimethyl ester **2** (yield = 64%) (Calcd for  $\text{C}_{17}\text{H}_{30}\text{O}_9$ : C, 53.97; H, 7.94. Found: C, 53.63; H, 7.93. FAB MS ( $M + H$ )<sup>+</sup> = 379). Treatment of **2** with lithium aluminum hydride affords the triol **5** in 83% yield (Calcd for  $\text{C}_{14}\text{H}_{30}\text{O}_6$ : C, 57.12; H, 10.27. Found: C, 57.24; H, 10.20. FAB MS ( $M + H$ )<sup>+</sup> = 295). The tosylate **6**, prepared from tosyl chloride and **5**, was treated with *O*-benzylhydroxylamine in 1,2-dimethoxyethane to give the tris(*O*-benzylhydroxylamine) **7** (Calcd for  $\text{C}_{35}\text{H}_{51}\text{N}_3\text{O}_6$ : C, 68.96; H, 8.37; N, 6.90. Found: C, 68.68; H, 8.45; N, 6.69. FAB MS ( $M + H$ )<sup>+</sup> = 610, yield = 37%). The triacid chloride **4** was obtained by allowing **3**, which was prepared from its methyl ester, to react with oxalyl chloride in benzene, yield >95%. High dilution acylation of **7** with **4** in benzene gives the protected macrobicyclic trishydroxamate **8** (Calcd for  $\text{C}_{49}\text{H}_{69}\text{N}_3\text{O}_{12}\cdot 4\text{H}_2\text{O}$ : C, 61.06; N, 4.36; H, 8.00. Found: C, 60.95; N, 4.18; H, 7.10. FAB MS ( $M + H$ )<sup>+</sup> = 893) in 30% yield. The



- |                          |                          |
|--------------------------|--------------------------|
| 1 X = CN                 | 5 X = CH <sub>2</sub> OH |
| 2 X = COOCH <sub>3</sub> | 6 X = OTs                |
| 3 X = COOH               | 7 X = NHOBz              |
| 4 X = COCl               |                          |



- |          |
|----------|
| 8 Y = Bz |
| 9 Y = H  |

macrobicyclic trishydroxamate cryptand **9** was obtained from **8** by catalytic hydrogenation. This cryptand (1,13-dimethyl-3,11,15,23,26,34-hexaoxa-6,20,29-trioxo-7,19,30-tris(hydroxy-aza)bicyclo[11.11.11]pentatricontane,  $\text{H}_3\text{THX}$ ) was characterized by  $^1\text{H}$  NMR,  $^{13}\text{C}$  NMR, elemental analysis, and FAB MS (Calcd for  $\text{C}_{28}\text{H}_{51}\text{N}_3\text{O}_{12}\cdot 1/2\text{H}_2\text{O}$ : C, 53.27; H, 8.40; N, 6.59. Found: C, 53.16; H, 8.19; N, 6.38. FAB MS ( $M + H$ )<sup>+</sup> = 622). This cryptand is very soluble in methanol and sparingly soluble in water. Its sodium salt is very water soluble.

The 1:1 Fe(III) complex of the cryptand was prepared by combining Fe(III) chloride with a slight excess of the neutral (acid) form of the ligand to form a  $2.0 \times 10^{-4}$  molar solution in 2:8 v/v methanol–water and gradually increasing the pH to the desired neutral value. The absorbance spectra of the cryptate (Figure 1) compare well with those of the Fe(III) chelate of desferrioxamine B, DFB, reported by Anderegg et al.<sup>4</sup> At pH 4.0 the Fe(III) cryptate has a molar absorbance of 2700 at  $\lambda_{\text{max}}$  423 nm (see Figure 1), while that of Fe<sup>III</sup>-DFB is 2640 ( $\lambda_{\text{max}}$  440 nm) at pH 4.<sup>4</sup> The absorbance shifts with pH are also similar for the cryptate and DFB complexes, with an isosbestic point at 480 nm for the cryptate and 481 for the DFB complex. The close similarity in the magnitudes of the molar absorbances of the iron(III) complexes of DFB and the cryptand provide assurance that all three bidentate donor groups of the latter are coordinated to the Fe(III) center. The observed isosbestic point indicates the conversion of one pure Fe(III) complex to another as the pH increases. It is suggested that the reaction corresponds to a monoprotated complex  $\text{FeHTHX}^+$ , having two coordinated hydroxamate groups, and one protonated, non-coordinated hydroxamate group, which is converted at higher pH to the octahedral Fe(III) cryptate  $\text{FeTHX}$ , with three coordinated hydroxamate groups arranged in an octahedral fashion around the metal ion. At pH 4.4 and above there is little further increase in absorbance, and one therefore concludes the cryptate to be fully formed, with a molar absorbance of 2750 at  $\lambda_{\text{max}}$  = 430 nm.

Because the Ga(III) ionic radius is only slightly smaller than that of Fe(III), the new cryptand would also be expected to complex Ga(III) strongly in an octahedral fashion. The 1:1 Ga(III) complex was prepared by the reaction of molar equivalents of  $\text{Ga}(\text{OH})_4^-$  and the ligand in aqueous solution at pH 8.9. This is above the pH at which Ga(III) precipitates as  $\text{Ga}(\text{OH})_3$ . The white solid which separated was characterized by elemental analysis and mass spectra (Calcd for  $\text{C}_{28}\text{H}_{48}\text{N}_3\text{O}_{12}\cdot \text{Ga}\cdot 2\text{H}_2\text{O}$ : C,

(1) Nielands, J. B. In *Development of Iron Chelators for Clinical Use*; Martell, A. E., Anderson, W. F., Badman, D. G., Eds.; Elsevier North Holland: New York, 1981; pp 13–31.

(2) Sun, Y.; Martell, A. E.; Motekaitis, R. J. *Inorg. Chem.* **1985**, *24*, 4343.

(3) Martell, A. E. In *Development of Iron Chelators for Clinical Use*; Martell, A. E., Anderson, W. F., Badman, D. G., Eds.; Elsevier North Holland: New York, 1981; pp 102–108.

(4) Anderegg, G.; L'Eplattenier, F.; Schwarzenbach, G. *Helv. Chim. Acta* **1963**, *46*, 1409.