Accepted Manuscript

Synthesis of New Indolylpyrrole Derivatives Via a Four-Component Domino Reaction Between Arylglyoxals, Acetylacetone, Indole and Aliphatic Amines in Aqueous Media

Fereshteh Mousavizadeh, Mahdiyeh Talebizadeh, Mohammad Anary-Abbasinejad





Please cite this article as: Mousavizadeh, F., Talebizadeh, M., Anary-Abbasinejad, M., Synthesis of New Indolylpyrrole Derivatives Via a Four-Component Domino Reaction Between Arylglyoxals, Acetylacetone, Indole and Aliphatic Amines in Aqueous Media, *Tetrahedron Letters* (2018), doi: https://doi.org/10.1016/j.tetlet. 2018.06.043

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Synthesis of New Indolylpyrrole Derivatives Via a Four-Component Domino Reaction Between Arylglyoxals, Acetylacetone, Indole and Aliphatic Amines in Aqueous Media

Fereshteh Mousavizadeh, Mahdiyeh Talebizadeh, Mohammad Anary-Abbasinejad*

Department of chemistry, Vali-e-Asr University, Rafsanjan, 77176, Iran

Corresponding Author E-mail: m.anary@vru.ac.ir



ABSTRACT: A novel synthesis of indolylpyrrole derivatives is described by a four-component domino reaction between arylglyoxals, acetylacetone, indole and aliphatic amines in water as solvent at 60°C without using any catalyst or promoter. The FT-IR, ¹H NMR, ¹³C NMR spectral and elemental analysis confirm the structures of the products.

Keywords. Indolylpyrrole, Four-component reaction, Arylglyoxal, Aqueus media, Synthesis

Pyrrole ring is one of the most important chemical subunits which is widely found in the structure of many therapeutically active compounds.¹ Pyrrole ring also exists in the structure of a wide variety of natural products such as vitamin B12, pigments like bilirubine and biliverdine and the porphyrine of heme.² Access to pyrrole derivatives using multi-component reactions is particularly attractive from the synthetic efficiency and environmentally point of views.³

Indole derivatives having a heterocyclic ring at C3 have recently attracted much attention from synthetic point of view for their presence at natural products such as martefagin A6⁴ and alkaloids, which are potent inhibitors of HIV-1 integrase.⁵⁻⁹ Some indole derivatives containing a pyrrole substituent at their C3 position have been recently synthesized and proven to possess interesting biological activities.¹⁰⁻¹² Multi-component domino reactions (MDRs), particularly those performed in water have attracted much attraction from chemists for their synthetic power point of view and green characteristics. These reactions have been successfully utilized for synthesis of a wide variety of chemically and biologically important compounds.¹³

Arylglyoxals have gained much attention for synthesis of heterocycles using multi-component reactions. The application of arylglyoxals for preparation of a wide range of three- to six-membered heterocycles has been recently reviewed.¹⁴ In continuation of our previous studies on the application of arylglyoxals in synthesis of heterocyclic compounds,¹⁵⁻¹⁷ we report herein a green MDR approach for synthesis of some indole derivatives having a pyrrole functionality at their 3-position. Thus the four-component domino

2

ACCEPTED MANUSCRIPT

Tetrahedron

reaction between arylglyoxals, acetylacetone, indole and aliphatic amines in water as solvent in the absence of any catalyst or promoter afforded indole derivatives in excellent yields (Scheme 1).





Table 1.





At the beginning of our investigation, we carried out the reaction between 4-chlorophenylglyoxal monohydrate, acetylacetone, indole and benzylamine in water as solvent. After stirring for 1h at 60°C, the analysis of the reaction mixture by TLC showed the presence of only one product. Filtering the reaction mixture and simple washing of the product by diethyl ether afforded the indolylpyrrole derivative **5a** in nearly quantitative yield. The NMR, IR spectral, and combustion analytical data deduced the structure of product. The ¹H NMR spectrum of compound **5a** showed three sharp single signals at 1.7, 2.4 and 5.1 ppm which are related to two methyl and one methylene groups, respectively. The aromatic protons resonated between 6.8 and 7.3 ppm. The NH proton was observed at 10.9 ppm. The ¹³C NMR spectrum of **5a**

4

CCEPTED V IUSCRIPT

Tetrahedron

exhibited twenty-four distinct signals in agreement with the proposed structure, the carbonyl carbon resonated at 196.8 ppm. The proposed structure was also confirmed by the IR spectrum of **5a**, which showed the absorption bands at 1635 cm⁻¹ for carbonyl group and at 3359cm⁻¹ for NH bond. To define the scope and generality of the reaction, a series of substituted arylglyoxals and aliphatic amines were examined. To our delight, in each case the desired indolylpyrrole derivative was obtained in excellent yield. This method has some advantages, as the starting materials are simply available, the reaction conditions are simply accessible and the products were isolated in high yields by simple filtration and washing with diethyl ether.

When the aromatic amines such as aniline or 4-methoxyaniline were used as the amine component, complex mixtures were obtained and no pure product could be isolated. Similarly, no product could be isolated when cyclic diketones like dimedone or ketoesters such as ethyl acetoacetate were used instead of acetylacetone.

A plausible mechanism for the formation of compounds 5 is shown in Scheme 2. The Knovenagel condensation of arylglyoxal and acetylacetone afforded enone 6 which under the Michael addition of indole converted to trione derivative 7. The reaction of benzyl amine with intermediate 7, known as Paal-Knorr pyrrole synthesis, afforded the indolylpyrrole derivative 5.



Continuing the process of our study, we decided to investigate the possibility of using the reaction conditions for preparing and isolating intermediate 7. Thus, the reaction of arylglyoxals, acetylacetone and indole was carried out in water at 60 °C in the absence of any catalyst or promoter. As shown in Scheme 3 and Table 2 the indolyltrione derivatives 7a-g were obtained in excellent yields. As shown in Table 1 the reaction is compatible for different arylglyoxals with electron deficient or electron poor aromatic rings. Compounds 7a-g were also isolated by simple filtering the product and washing with diethyl ether. The structures of compounds 7a-g were proved by NMR and IR spectral and analytical data. For example, the ¹H NMR spectrum of **7a** showed two single signals at 1.1 and 2.2 ppm for methyl groups. Two methine protons were observed as two double signals (${}^{3}J_{HH} = 11 \text{ H}_{Z}$) at 5.0 and 5.6 ppm. The aromatic protons resonated between 6.9-7.9 ppm and the NH proton was observed at 11.1 ppm. ¹³C NMR spectrum of **7a** exhibited nineteen distinct signals in agreement with the proposed structure, the carbonyl carbons resonated

at 196.58, 202.91 and 204.08 ppm. The proposed structure was also confirmed by the IR spectrum of **7a**, which showed the absorption bands at 1725, 1700, 1678 cm⁻¹ for carbonyl bonds and at 3396 for NH bond. To prove that indolyltrione derivative **7** is an intermediate for formation of indolylpyrrole derivatives **5**, the reaction of **7a** with benzyl amine was carried out at in water at 60° C and the indolylpyrrole derivative **5a** was obtained in nearly quantitative yield.



r

Entery	Ar	Indole	Intermediate product	Yield%
1	4-BrC ₆ H ₄		$Br \xrightarrow{O} \xrightarrow{O} \xrightarrow{O} \xrightarrow{CH_3} \xrightarrow{O} \xrightarrow{O} \xrightarrow{CH_3}$	92
2	4-ClC ₆ H ₄	ZE	$CI \xrightarrow{O} \xrightarrow{O} \xrightarrow{O} \xrightarrow{O} \xrightarrow{CH_3} \xrightarrow{O} \xrightarrow{CH_3}$	93
3	4-MeC ₆ H ₄	N N N N N N N N N N N N N N N N N N N	$H_{3}C \xrightarrow{O} O \xrightarrow{O} CH_{3}$ $\downarrow O \xrightarrow{O} CH_{3}$	95
4	4-NO ₂ C ₆ H ₄	N H	$\begin{array}{c} O_2 N & \overbrace{}^{O} & O \\ & \overbrace{}^{O} & -CH_3 \\ & \overbrace{}^{O} & -CH_3 \\ & O \\ & H \\ \end{array}$	95

6

Tetrahedron								
5	4-BrC ₆ H ₄	Br N H	$ \begin{array}{c} $	93				
6	C ₆ H ₅	N H	$\begin{array}{c} & & O & O \\ & & & & \\ & & & & \\ & & & &$	94				
7	2-naphthyl	N H		93				

ASSOCIATED CONTENT

Supporting Information

Experimental procedures, ¹H, ¹³C, NMR characterization data of all novel products (PDF)

AUTHOR INFORMATION

Corresponding Author

Corresponding Author E-mail: m.anary@vru.ac.ir

Present Addresses

Department of Chemistry, Faculty of Science, Vali-e-Asr University of Rafsanjan, Rafsanjan 77176, Iran

ACKNOWLEDGMENT

We gratefully acknowledge, the Vail-e-Asr University of Rafsanjan Faculty Research Grant for financial support

Conclusion

In conclusion, we have achieved the synthesis of indolylpyrrole derivatives by a four-component domino reaction between readily accessible starting materials in excellent yield. This synthesis has the advantages that the reactions were carried out in water as solvent without using any catalyst or promoter and products were purified by simple filtration and washing with diethyl ether.

References

1. Bhardwaj, V.; Gumber, D.; Abbot, V.; Dhimana, S.; Sharmaa, P. RSC Adv. 2015, 5, 15233-15266.

- 2. physical chemistry and chemical physics, Issue 10, 2000, 2, 2145-2151
- 3. Chemical society reviews, Issue 13, 2014, 43, 4633-4657, V. Estevez, M. Villacampa, J. C. Menendez.
- 4. Takahashi, S.; Matsunaga, T.; Hasegawa, C.; Saito, H.; Fujita, D.; Kiuchi, F.; Tsuda, Y. *Chem Pharm Bull.* **1998**, 46, 1527-1529.
- 5. Joshi, B. S.; Taylor, w. I.; Bhate, D. S.; Karmarkar, S. S. Tetrahedron. 1963, 19, 1437-1439.
- 6. Noltenmeyer, M.; Sheldrick, G. M.; Hoppe, H.U.; Zeeck, A. J. Antibiot. 1982, 35, 549-555.
- 7. Umehara, K.; Yoshida, K.; Okamoto, M.; Iwami, M.; Tanaka, H.; Kohsaka, M.; Imanaka, H. *J. Antibiot*.**1984**, 37, 1153-1160.
- 8. Kelly, T. R.; Fu, Y.; Xie, R. L. Tetrahedron Lett. 1999, 40, 1857-1860.
- 9. Davies, J. R.; Kane, P. D.; Moody, C. J.; Slawin, A. M. Z. J. Org Chem. 2005, 70, 5840-5851.
- 10. Sani, R.; Dharavath, N.; Eligeti, R.; Malladi, S. R. J. Heterocyclic Chem. 2017. DOI 10.1002/jhet.2888.
- 11. Vimala, G.; kamal Raja, J.; Perumalb, P. T.; SubbiahPandi, A. IUCrData. 2016, 1, x160724
- 12. Lopchuk, J. M.; Gribble, G. W. Heterocycles. 2011,82,1617-1631.
- 13. Na, N.; Jiang, B.; Zhang, G.; Tu, S.J.; Wever, V.; Li, G.GreenChem.2010, 12, 1357-1361.
- 14. Eftekhari-Sis, B.; Zirak, M.; akbari, A.; Chem. Rev. 2013, 113, 2958-3043.
- 15. Masoudi, M.; Anary-Abbasinejad, M.; Mohammadi, M. J. Iran. Chem. Soc. 2016, 13, 315-321.
- 16. Masoudi, M.; Anary-Abbasinejad, M. Tetrahedron Lett. 2016, 57, 103-104.
- 17. Amiranipoor, M.; Anary-Abbasinejad, M.; Darehkordi, A. Arkivoc, 2017, part (v), 141-147.
- 18. Riley, H. A.; Gray, A. R. Organic Syntheses Collect. New York. 1943, 2, 509.

Colif

Tetrahedron

Scheme 1. Synthesis of indolylpyrrole derivatives by reaction between arylglyoxals, acetylacetone, indole and aliphatic amines

Scheme 2. The possible reaction mechanism for synthesis of indolylpyrroles 5a-i.

Scheme 3. Synthesis of 3-acetyl-1-aryl-2-(1H-indol-3-yl)pentane-1,4-dione 7

Table 1. Synthesis of indolylpyrrole derivatives (**5a- i**) by reaction of arylglyoxals, acetylacetone, indole and aliphatic amines

 Table 2. Synthesis of 3-acetyl-1-aryl-2-(1H-indol-3-yl)pentane-1,4-diones 7a-g by reaction between arylglyoxals, acetylacetone , indole and aliphatic amine.

Pyrrole ring is one of the most important chemical subunits Indoles with a pyrrole ring at their C3 position possess biological activities Multi-component reactions in water have attracted much attraction Developing new efficient methods for synthesis of heterocycles are of interest

Accepting

NUSCRIPT ACCEPTED

Tetrahedron

Accembra