

An Easy Construction of 8,12-Dioxa-13-azatricyclo[8.3.1.0^{2,7}]tetradeca-2(7),3,5,13-tetraen-14-ones


Surya K. De and Asok K. Mallik*

Department of Chemistry, Jadavpur University, Calcutta-700 032, INDIA


Received 19 August 1997; revised 16 January 1998; accepted 23 January 1998

Abstract: 8,12-Dioxa-13-azatricyclo[8.3.1.0^{2,7}]tetradeca-2(7),3,5,13-tetraen-14-ones are obtained in moderate yield by treatment of oximes of *o*-allyloxyacetophenones with [hydroxy(tosyloxy)iodo]benzene in acetonitrile. © 1998 Elsevier Science Ltd. All rights reserved.

Deoxygenation of ketoximes utilising (diacetoxyiodo)benzene (DAIB)^{1a,b} or (dichloroiodo)benzene (DCIB)^{1c} is one of the many important synthetic applications of hypervalent iodine reagents reported in recent years.² The suggested mechanism for the reaction with DAIB^{1a,b} is as follows :




We envisaged that if the reaction is done by using a hypervalent iodine reagent unable to supply nucleophile of sufficient nucleophilicity, then the 1,4-elimination reaction shown below would be a competitive process. The product of such reaction, a nitrosoalkene, can undergo [4+2]-cycloaddition³ with a double bond (intramolecular or intermolecular), constituting a simple method for synthesis of interesting 5,6-



dihydro-4H-1,2-oxazine derivatives. We report here the preliminary interesting results obtained from the reaction on oximes (1) of *o*-allyloxyacetophenones utilising [hydroxy(tosyloxy)iodo]benzene (HTIB) as the reagent.

On addition of HTIB to a solution of 1 in dry acetonitrile, the colour changed to bluish green, indicating the formation of nitroso compound. Therefore, solid HTIB (1 equiv.) was added in portions during 1 hr. to a solution of 1 in the said solvent and the mixture was stirred for further 6 hr. Usual work up of the reaction mixture⁴ followed by chromatographic resolution of the product mixture gave 8,12-dioxa-13-azatricyclo[8.3.1.0^{2,7}]tetradeca-2(7),3,5,13-tetraen-14-ones (2)⁵ as a new type of product (yield : 30-34%)

along with the deoximation product (22-26%).



The compounds **2a-c** are obviously the product of the desired cycloaddition followed by an oxidation. The oxidation may be effected either by hypervalent iodine⁶ or by aerial oxygen⁷. The interesting structural features of **2a-c** are that they contain a bridgehead double bond and a bridging $>\text{C}=\text{O}$. Another interesting aspect of such molecules is the possibility of their conversion to biologically important vicinal amino alcohols⁸ through reduction.^{3c}

Thus, we describe a new method for generation of nitrosoalkenes and their cycloaddition reaction in intramolecular fashion, which, to our knowledge, was not known previously. The most widely used method for generation of nitrosoalkenes involves a base-catalysed 1,4-elimination of hydrogen halide from the oximes of α -haloketones.³ The present method appears to be a good substitute to that because it does not require the handling of lachrymatory and toxic α -haloketones. Furthermore, the observed regioselectivity in the cycloaddition process and the facile oxidation of the cycloaddition product are noteworthy.

Acknowledgements : The authors are thankful to the UGC, New Delhi and the authorities of Jadavpur University for financial assistance.

References and Notes:

- (a) Moriarty, R. M. ; Prakash, O. ; Vavilikolanu, P. R. *Synth. Commun.* **1986**, *16*, 1247. (b) Prakash, O. ; Pahuja, S. ; Sawhney, S. N. *Indian J. Chem.* **1989**, *28B*, 73. (c) Radhakrishna, A. S. ; Augustine, B. ; Sivaprakash, K. ; Singh, B. B. *Synth. Commun.* **1991**, *21*, 1473.
- (a) Varvoglis, A. *Chem. Soc. Rev.* **1981**, *10*, 377. (b) Varvoglis, A. *Synthesis* **1984**, 709. (c) Moriarty, R. M. ; Prakash, O. *Acc. Chem. Res.* **1986**, *19*, 244. (d) Merkushev, E. B. *Russ. Chem. Rev.* (Engl. Transl.) **1987**, *56*, 826. (e) Moriarty, R. M. ; Vaid, R. K. *Synthesis* **1990**, 431. (f) Prakash, O. ; Saini, N. ; Sharma, P. K. *Synlett* **1994**, 221. (g) Prakash, O. ; Saini, N. ; Tanwar, M. P. ; Moriarty, R. M. *Contemp. Org. Synth.* **1995**, *2*, 121. (h) Stang, P. J. ; Zhdankin, V. V. *Chem. Rev.* **1996**, *96*, 1123. (i) Varvoglis, A. *Tetrahedron* **1997**, *53*, 1179.
- (a) Gilchrist, T. L. *Chem. Soc. Rev.* **1983**, *12*, 53. (b) Venugopalan, B. ; Sathe, K. M. ; Pinto de Souza, E. *Indian J. Chem.* **1996**, *35B*, 475. (c) Faragher, R. ; Gilchrist, T. L. *J. Chem. Soc. Perkin Trans. I* **1979**, 249.
- De, S. K. ; Mallik, A. K. *Indian J. Chem.* **1997**, *36A&B*, 536.
- Selected data of **2** : **2a**, mp 93-94° (light yellow needles). Anal: Found C, 64.91; H, 4.30; N, 6.62, Calcd for $\text{C}_{11}\text{H}_9\text{NO}_3$ C, 65.02; H, 4.43; N, 6.89%. IR (v, KBr, cm^{-1}): 1650 (C=O), 1608 (C=N). ^1H NMR (300 MHz, CDCl_3) : δ 3.87-3.98 (1H, m, H-10), 4.24 (1H, dd, J= 11.7 and 7.4 Hz), 4.32 (1H, t, J=9.1 Hz), 4.39 (1H, dd, J= 11.7 and 5.5 Hz), 4.76 (1H, dd, J=11.1 and 8.9 Hz), 7.08 (1H, dd, J=8.2 and 0.9 Hz, H-6), 7.18 (1H, ddd, J=8.7, 7.9 and 0.9 Hz, H-4), 7.52 (1H, ddd, J=8.7, 8.2 and 1.7 Hz, H-5), 7.97 (1H, dd, J=7.9 and 1.7 Hz, H-3). ^{13}C NMR (75 MHz, CDCl_3) : δ 47.30 (C-10), 71.86 (C-11), 74.80 (C-9), 122.57 (C-6), 124.05 (C-4), 127.90 (C-2), 130.77 (C-3), 135.91 (C-5), 158.18 (C-1), 159.52 (C-7), 182.41 (C-14). MS : M^+ 203. **2b**, mp 110-111°. **2c** mp 157-158°.
- Moriarty, R. M. ; Vaid, R. K. ; Duncan, M. P. ; Ochiai, M. ; Inenaga, M. ; Nagao, Y. *Tetrahedron Lett.* **1988**, *29*, 6913.
- Biswas, K. M. ; Jackson, A. H. *J. Chem. Soc. Perkin Trans. I* **1989**, 1981.
- (a) Ding, C. Z. *Tetrahedron Lett.* **1996**, *37*, 945. (b) Lindstrom, U. M. ; Franckowiak, R. ; Pinault, N. ; Somfai, P. *ibid* **1997**, *38*, 2027.