Complexes of the Bidentate Ligands $PPh_2CH_2C(Bu^t)=NNR_2$ (R = H or Me) and $PPh_2CH_2C(Bu^t)=N-N=CHPh$ with Palladium and Platinum. X-Ray Crystal Structure of cis-[Pt{PPh_2CH_2C(Bu^t)=NNH}_2][†]

King Kuok Hii, Sarath D. Perera, Bernard L. Shaw^{*} and Mark Thornton-Pett School of Chemistry, The University of Leeds, Leeds LS2 9JT, UK

tert-Butyl diphenylphosphinomethyl ketone dimethylhydrazone L¹ reacts with $[PtCl_2(cod)]$ (cod = cycloocta-1,5-diene) to give the chelate [PtCl₃{PPh₂CH₂C(Bu')=NNMe₂}] with a six-membered chelate ring; the corresponding palladium complex was also prepared. The phosphino dimethylhydrazone L¹ or the phosphino hydrazone L² also react with [PtMe₂(cod)] to give [PtMe₂{PPh₂CH₂C(Bu')=NNR₂}], (R = Me or H); these platinum(II) complexes react with methyl iodide to give the corresponding trimethylplatinum(IV) complexes [PtMe₃I{PPh₂CH₂C(Bu')=NNR₂}]. Treatment of [PtCl₂(cod)] with 2 equivalents of L² gave the dication cis-[Pt{PPh₂CH₂C(Bu')=NNH₂}²⁺ isolated as the chloride or chloride hexafluorophosphate salts. The dicationic dichloride reacted with 2 moles of sodium ethoxide to give the neutral *cis*-[Pt{PP₂CH₂C(Bu')=NNH}₂]. Treatment of [PdCl₂(NCPh)₂] with L² gave $[\dot{P}d\dot{P}h_2CH_2C(Bu')=N\dot{N}H_2]_2]Cl_2$, which with sodium ethoxide gave the neutral *trans*- $[Pd{Ph_2CH_2C(Bu')=NNH_2]}$. The mixed azine monophosphine PPh_2CH_2C(Bu')=N-N=CHPh L³ with [PdCl₂(NCPh)₂] gave the chelate complex [PdCl₂{PPh₂CH₂C(Bu')=N-N=CHPh}] and with [PtMe₂(cod)] it gave [PtMe₂{PPh₂CH₂C(Bu')=N-N=CHPh}]. Treatment of [PtCl₂(NCMe)₂] with 2 equivalents of L³ gave trans-[PtCl₂{PPh₂CH₂C(Bu¹)=N-N=CHPh₂] which on heating gave the dicationic dichloride salt cis-[Pt{Ph2CH2C(Bu')=N-N=CHPh22Cl2. Successive treatments of this salt with 1 mole of sodium methoxide gave cis-[{Ph2CH2C(Bu')=N-R=CHPh}Pt{Ph2CH2C(Bu')N-N=CHPh}]⁺ and cis-[Pt{PP₂CH=C(Bu')N-N=CHPh}]. Proton, ¹³C-{¹H}, ³¹P-{¹H} NMR and infrared data have been obtained. Crystals of cis-[$Pt{Ph_2CH_2C(Bu')=NNH_2}$] are monoclinic, space group $P2_1/c$, with a = 1261.5(2), b = 1794.6(3), c = 1792.0(4) pm and Z = 4; final R = 0.0451 for 4088 observed reflections.

In a previous paper we have described the synthesis of tertbutyl diphenylphosphinomethyl ketone dimethylhydrazone L^1 by deprotonating *tert*-butyl methyl ketone dimethylhydrazone with *n*-butyllithium and treating the resultant carbanion with chloro(diphenyl)phosphine.1 We established that the configuration of this phosphine around the C=N bond as Z since, when treated with $[Mo(CO)_4(nbd)]$ (nbd = norbornadiene) under mild conditions (20 °C), it gave $[\dot{M}o(CO)_4{\dot{P}Ph_2CH_2C(Bu')=N\dot{N}Me_2}]$ the crystal structure of which was determined;¹ this showed the presence of a sixmembered chelate ring, i.e. the configuration around the C=N is Z. When heated strongly (180 °C) this six-membered chelate ring complex isomerised to the five-membered chelate ring complex with an E-configuration around C=N. We also showed that the phosphino dimethylhydrazone (=NNMe₂) L¹ underwent an exchange reaction with hydrazine in the presence of acetic acid as catalyst to give a phosphino hydrazone (=NNH₂) L^2 which probably also had the Z-configuration; we also studied this phosphine as a ligand for chromium, molybdenum and tungsten carbonyls in the zero-valent state. It was of interest to study the behaviour of these hydrazone phosphines, L¹ and L², as ligands towards some more electropositive and labile metals viz. platinum(II) and palladium(II) and in the present paper we describe the results of this study. For convenience the work is summarised in Scheme 1.

Results and Discussion

We first attempted to prepare a chelate of L^1 with platinum dichloride by treating the labile olefin complex [PtCl₂(cod)] (cod = cycloocta-1,5-diene) with 1 equivalent of the phosphino dimethylhydrazone L^1 at ca. 20 °C. This gave the hoped for chelate complex 1a in ca. 80% yield; preparative details and elemental analytical data for this and other new complexes are in the Experimental section, infrared and ³¹P NMR data are in Table 1 and ¹H data are in Table 2. The complex showed a single resonance in its ³¹P-{¹H} NMR spectrum, with satellites due to platinum-195, and the large value of ${}^{1}J(PtP)$ of 4126 Hz is consistent with a phosphorus trans to a chloride ligand.² The infrared spectrum showed two strong bands, at 340 and 285 cm⁻¹ for the cis-PtCl₂ moiety³ due to v(Pt-Cl). The proton NMR data for the NMe₂ methyls are consistent with the proposed structure for 1a, having a sixmembered chelate ring; the value of $J(PtNCH_3)$ of 26.9 Hz is consistent with a three-bond coupling^{3,4} and too large for a four-bond coupling, which it would have to be if the chelate ring were five-membered. The carbon-13 data (see Experimental section) are also consistent with the assigned structure. In particular, the low δ_{C} value of δ 17.9 for the methylene carbon is in agreement with the observed low δ_{C} value of δ 24.0 for the sixmembered $[\dot{M}o(CO)_4 \{\dot{P}Ph_2CH_2C(Bu')=N\dot{N}Me_2\}]$ complex whereas the five-membered $[Mo(CO)_4]{PPh_2CH_2C(Bu')=N-}$ NMe₂] complex has a higher δ_{c} value of δ 42.1. In extensive studies of azine diphosphines acting as terdentates with metals such as Cr, Mo, W, Pd, Pt and Ir, we have found that the methylene carbons in the six-membered rings give lower $\delta_{\rm C}$

[†] Supplementary data available: see Instructions for Authors, J. Chem. Soc., Dalton Trans., 1994, Issue 1, pp. xxiii-xxviii.



Scheme 1 (*i*) $[PtCl_2(cod)]$, $[PdCl_2(NCPh)_2]$ or $[PtMe_2(cod)]$; (*ii*) 1c or 1d with MeI; (*iii*) 0.5 equivalent $[PtCl_2(cod)]$; (*iv*) $[PdCl_2(NCPh)_2]$; (*v*) NH₄PF₆; (*vi*) NaOMe

Table 1 Infr	ared and	${}^{31}P-{}^{1}H$ NM	Rª data	
Compound	$v(C=N)^b$	v(M-Cl) ^c	v(N–H) ^c	δ _P
L ¹	1610m		_	-9.7
L ²	1640w		3330m, 3350w -	-22.6
L ³	1585w	_	g	59.1
1a	1640m	285m, 340m		16.5 (4126)
1b	1620m	275w, 325w		43.2
1c	1610m	_		43.2 ^{<i>d</i>} (2186)
1d	1630m		3100w, 3300m	41.5 ^d (2199)
2a	1640w	_	—	7.2 ^d (1217)
2b	1630m		3100m, 3300m	7.1 ^{<i>d</i>} (1215)
3a	1630m		3100s, 3350w	36.4 ^e (3600)
3b	1630m	_	3100s, 3350w	36.9 ^{<i>f</i>} (3607)
4	1595s		g	38.4 ^{<i>h</i>} (3313)
5a	1630m		3130s, 3330w	63.3 ^e
5b	1640m		3130s, 3330w	64.1 ^h
6	1615m	_		- 10.1
7a	1600s	280w, 340w		49.3
7Ъ	1600s	_		31.3 (2136)
8	1630m	340w	_	11.2 (2575)
9a	—	_		21.5 ^e (3412)
9b	1620m		AL	21.7 ^f (3361)
11	1595m		—	21.9 ^d (3146)
12a ⁱ				20.2 ^{<i>d</i>} (1703)
12b ⁱ			—	9.06 ^d (1540)
12c ^{<i>i</i>}				18.15 (1657)

^{*a*} Recorded at 36.2 MHz, chemical shifts δ (±0.1 ppm) relative to 85% H₃PO₄, solvent CDCl₃ unless otherwise stated, ¹*J*(PtP)/Hz in parentheses. ^{*b*} As compressed KBr disc. ^cAs Nujol mull between polythene plates. ^{*d*} In C₆D₆. ^{*e*} In ethanol with C₆D₆ as external reference. ^{*f*} In (CD₃)₂CO. ^{*g*} Not observed. ^{*h*} In CD₂Cl₂. ^{*i*} v(C=O) 2040 cm¹ in C₆H₆.

values (δ 20–25) than the ones in five-membered chelate rings (δ 40–45).^{5,6} Similarly, treatment of [PdCl₂(NCPh)₂] with L¹ gave the corresponding palladium dichloride chelate **1b**; characterising data are given in the Experimental section and in Tables 1 and 2. We prepared the chelate complex of PtMe₂ with L¹ *i.e.* [PtMe₂{PPh₂CH₂C(Bu¹)=NNMe₂}] **1c** by displacing cod from [PtMe₂(cod)] under mild conditions (*ca.* 20 °C). As would be expected,² the value of ¹J(PtP), *viz.* 2186 Hz was much less than with the corresponding dichloride **1a**. The proton NMR data (Table 2) are consistent with the assigned structure **1c**, *e.g.* the NMe₂ methyls are equivalent, J(PtH) = 15.9 Hz, consistent with a three-bond coupling.^{3,4} We also treated [PtMe₂(cod)] with the phosphino hydrazone L² and obtained the corresponding dimethylplatinum complex **1d**; the characterising data are consistent with a six-membered chelate ring, in particular, the values of ²J(PtNH₂) (29.3 Hz) and ³J(PNH₂) (6.1 Hz), excluded a five-membered ring chelate.

A characteristic reaction of complexes of the type [PtMe₂-(phosphine)₂] is the oxidative addition of methyl iodide to give platinum(IV) adducts of the type [PtMe₃I(phosphine)₂]⁴ and it was of interest to see whether the chelates 1c or 1d would react similarly. Treatment of 1c with an excess of iodomethane gave an adduct 2a which was fully characterised. The small value of ¹J(PtP) in the ³¹P-{¹H} NMR spectrum (Table 1) is typical for platinum(IV)^{2,7} and the occurrence of three chemically inequivalent platinum methyl resonances in the proton NMR spectrum indicated a *fac*-PtMe₃ arrangement. It is known that in trimethylplatinum(IV) complexes, methyl *trans* to iodine or nitrogen have ²J(PtCH₃) = ca. 70 Hz⁸ and for methyl *trans* to phosphorus ²J(PtCH₃) = ca. 57 Hz.⁴ The two *N*-methyls are not equivalent, chemically, excluding a five-membered chelate ring and the values of J(PtNCH₃) of 11.7 and 8.5 Hz are

Table 2 Proton NMR data⁴

105

Compound	δ(Bu')	δ(CH ₂)	Others
L^1	1.12 (9 H. s)	3.05 [2 H. d. ² J(PH) 2.9]	$2.10 (6 H, s, NMe_2)$
\overline{L}^2	0.98 (9 H. s)	3.10 [2 H, d, ² J(PH) 2.2]	4.75 (2 H, br s, NH ₂)
L ³	1.22 (9 H. s)	3.52 $[2 H, d, ^2 J(PH) 3.2]$	8.04 (1 H, s, CHPh)
1a	0.74 (9 H. s)	3.20 [2 H, d, ² J(PH) 13.2, ³ J(PtH) 45.4]	$3.55 [6 \text{ H}, \text{ s}, {}^{3}J(\text{PtH}) 26.9, \text{NMe}_{2}]$
16	0.73 (9 H, s)	3.01 [2 H, d, ² J(PH) 13.9]	$3.41 (6 H, s, NMe_2)$
le	0.63 (9 H. s)	2.80 [2 H, d, ² J(PH) 10.3, ³ J(PtH) 16.4]	$1.10[3 \text{ H}, d, {}^{3}J(\text{PH}) 8.0, {}^{2}J(\text{PtH}) 89.9, \text{PtMe}]$
		- , <u>-</u>	1.28 [3 H, d, ³ J(PH) 7.8, ² J(PtH) 64.9, PtMe]
			2.94 [6 H, s, ³ J(PtH) 15.9, NMe ₂]
1d ^b	0.64 (9 H, s)	2.65 [2 H, d, ² J(PH) 10.0, ³ J(PtH) 17.6]	$1.20 [3 H, d, {}^{3}J(PH) 7.8, {}^{2}J(PtH) 67.4, PtMe]$
			$1.36 [3 H, d, {}^{3}J(PH) 7.1, {}^{2}J(PtH) 90.8, PtMe]$
			5.34 [2 H, br d, ${}^{3}J(PH)$ 6.1, ${}^{2}J(PtH)$ 29.3, NH ₂]
2a ^b	0.70 (9 H, s)	2.5 (1 H) ^c	$1.09 [3 H, d, {}^{3}J(PH) 7.1, {}^{2}J(PtH) 75.7, PtMe]$
		3.80 [1 H, dd, ² J(PH) 11.2, ² J(HH) 13.6]	$1.85 [3 H, d, {}^{3}J(PH) 7.3, {}^{2}J(PtH) 78.1, PtMe]$
			$1.95 [3 H, d, {}^{3}J(PH) 7.6, {}^{2}J(PtH) 56.6, PtMe]$
			2.51 [3 H, s, ${}^{3}J(PtH)$ 11.7, NMe ₂]
			$3.17 [3 H, s, {}^{3}J(PtH) 8.5, NMe_{2}]$
2b ^b	0.85 (9 H, s)	3.19 [1 H, dd, ² J(PH) 9.7, ² J(HH) 13.9]	$1.42[3 \text{ H}, d, {}^{3}J(\text{PH}) 6.3, {}^{2}J(\text{PtH}) 70.3, \text{PtMe}]$
		3.39 [1 H, dd, ² J(PH) 12.0, ² J(HH) 14.4]	$1.95[3 \text{ H}, d, {}^{3}J(\text{PH}) 7.3, {}^{2}J(\text{PtH}) 73.7, \text{PtMe}]$
			$2.03[3 \text{ H}, \text{d}, {}^{3}J(\text{PH}) 8.5, {}^{2}J(\text{PtH}) 57.8, \text{PtMe}]$
			5.72 [2 H, br d, ${}^{3}J(PH)$ 5.1, ${}^{2}J(PtH)$ 20.0, NH ₂]
3b ^{<i>d</i>,<i>e</i>}	0.67 (18 H, s)	4.14 [4 H, br d, ${}^{2}J(PH)$ 13.0, ${}^{3}J(PtH)$ 45.0]	—
4 ^{e,j}	0.56 (18 H, s)	2.96 [4 H, d, ² J(PH) 12.2, ³ J(PtH) 44.0]	
5b ^{b,e}	0.65 (18 H, s)	3.92 (4 H, filled-in d, N = 13.2)	
6 ^e	0.61 (18 H, s)	2.79 (4 H, vt, N = 11.0)	
7 a	0.87 (9 H, s)	2.66 [1 H, dd, ² J(PH) 11.5, ² J(HH) 13.0]	8.25[1 H, d, J(PH) 1.7, CHPh]
		$3.19 [1 H, dd, {}^{2}J(PH) 15.0, {}^{2}J(HH) 13.0]$	1 22 52 11 1 3 (D11) 7 9 2 (D11) 70 4 D1 4-1
7b °	0.79 (9 H, s)	$2.28 [1 H, dd, ^2J(PH) 10.7, ^2J(HH) 12.6,$	1.23 [3 H, d, J(PH) 7.8, J(PH) 70.4, PIMe]
		$^{\circ}J(PtH) 10.3$	1.53 [3 H, 0, J(PH) 1.0, J(P(H) 09.4, P(M))]
		3.11 [1 H, dd, 2(PH) 10.7, 2(HH) 12.0,	8.40[1 H, a, J(PH) 1.9, J(PH) 57.0, CHPh]
0.4	1 05 (10 II -)	$^{\circ}J(P(H) 18.8]$	
8" 01. d. h	1.25(18 H, s)	4.43 [4 fr, vi, N = 9.2, J(rin) 24.9]	$0.10(2 H \otimes CHPh)$
90	1.10 (10 п, 8)	4.71[2.11, dd, J(F11) 13.4, J(H11) 19.5] 5 15 [2 H dd ² I(PH) 7.8 ² I(HH) 18 5]	<i>9.10 (2 11, 3, C11 11)</i>
11.6	161(19 U a)	5.15 [2 11, dd, 5(111) 7.8, 5(111) 18.5]	4 42 [2 H d ² /(PH) 8 8 ³ /(PtH) 22 2 CHP]
11	1.01 (10 11, 5)	—	$\frac{4.42}{2.2.2}$, $\frac{11}{2.2.2}$, $\frac{11}{2.2.$
12a ^b	112 (0 H a)	$3.01 (7 H) d^{-2} (/PH) 10.7 (^{3} //PtH) 11.87$	$0.54[3 \text{ H} \text{ d}^{-3} l(\text{PH}) 10.3 ^{2} l(\text{PtH}) 69.6 \text{ PtMe}]$
128	1.12 (9 11, 8)	5.01 [2 11, d, 5(1 11) 10.7, 5(1 11) 11.8]	1.27 [3 H d ³ /(PH) 7 3 ² /(PtH) 74 4 PtMe]
			1.27 [511, 0, 5(111) 7.5, 5(111) 7.5, 1000]
12h ^b	0.88 (9 H s)	3 45 [2 H d ² ((PH) 11 2 ³ ((PtH) 21 5]	$0.93 \Gamma_3 H d^{-3} I(PH) 10.3, {}^2 I(PtH) 70.1, PtMe]$
120	0.00 (7 11, 3)	5.45 [2 II, d, V(III) II.2, V(I II) 21.5]	$1.56 [3 H. d. {}^{3}J(PH) 7.3, {}^{2}J(PtH) 74.5, PtMe]$
			5.54 (2 H, br s, NH ₂)
120	1 18 (9 H s)	4.07 [2 H. d. ² J(PH) 11.5, ³ J(PtH) 17.3]	$0.25 [3 H, d, {}^{3}J(PH) 10.25, {}^{2}J(PtH) 69.6, PtMe]$
			0.81 [3 H, d, ³ J(PH) 7.3, ² J(PtH) 74.7, PtMe]
			7.99 (1 H, s, CHPh)

^{*a*} Recorded at 100 MHz, chemical shifts $\delta(\pm 0.01 \text{ ppm})$ relative to SiMe₄, solvent CDCl₃ unless otherwise stated, coupling constants J in Hz; s = singlet, d = doublet, vt = virtual triplet and br = broad. ^{*b*} In C₆D₆. ^{*c*} Obscured by the NMe₂ signal. ^{*d*} In (CD₃)₂CO. ^{*e*} NH resonance not observed. ^{*f*} In CD₂Cl₂. ^{*g*} CHPh obscured by the aromatic protons. ^{*b*} Recorded at -50 °C. $N = {}^{4}J(PH) + {}^{2}J(PH)$.

consistent with a three-bond coupling in a platinum(IV) complex.⁴

When we treated [PtCl₂(cod)] with 1 equivalent of the phosphino hydrazone L^2 in dichloromethane solution in an attempt to make the chelate [$PtCl_2$ { $PPh_2CH_2C(Bu')=NNH_2$ }] we obtained a mixture of two platinum-phosphine complexes, the more intense resonances occurring at δ_P 37.9 with ¹J(PtP) = 3796 Hz and the less intense at δ_P 33.6 with ¹J(PtP) = 4143 Hz; the proportions of the two complexes being ca. 3:1. Subsequently, when we treated [PtCl₂(cod)] with 2 equivalents of L^2 we produced the complex with δ_P 37.9, ¹J(PtP) 3796 Hz, exclusively and identified it as the cis-bis(chelate) dication chloride salt 3a (see below). The minor product with δ_P 33.6, ${}^{1}J(PtP) = 4143$ Hz was probably the chelate dichloride $[PtCl_2{PPh_2CH_2C(Bu')=NNH_2}]$ but we were unable to isolate this pure or characterise it. When we treated [PtCl₂-(cod) with 2 equivalents of L² in dichloromethane we prepared the bis(chelate) dication, isolated as its dichloride salt $[\dot{P}t{\dot{P}Ph_2CH_2C(Bu^t)=N\dot{N}H_2}_2]Cl_2$, 3a, characterised by elemental analysis (Experimental section) and ³¹P NMR data (Table 1). The dication was also isolated as the mixed chloride/

hexafluorophosphate salt $[Pt{PP_2CH_2C(Bu')=NNH_2}_2]$ -Cl[PF₆] **3b**, which was similarly characterised; the large value of ¹J(PtP) of 3607 Hz excludes the *trans* stereochemistry, but is consistent with phosphorus being *trans* to a more electronegative ligand such as nitrogen.^{9,10}

We anticipated that the bis(chelate) dication should have NH₂ groups which could be deprotonated with base. We therefore treated the dication dichloride salt **3a** with 2 equivalents of sodium ethoxide in ethanol. This gave the hoped for neutral complex *cis*- $[Pt{PPh_2CH_2C(Bu')NNH}_2]$ **4**, the crystal structure of which was determined and is discussed below. This neutral complex was also characterised by elemental analysis and by its infrared spectrum for which the value of v(C=N) of 1595 cm⁻¹ was significantly less than the value for the free ligand or for the values for the complexes 1–3 (Table 1). The value of ${}^{1}J(PtP)$ of 3313 Hz (Table 1) is in agreement with the assigned structure for **4** although we could not observe the NH resonance which was presumably too broad



Scheme 2 R = -N=CHPh. (i) $[PdCl_2(NCPh)_2]$ or $[PtMe_2(cod)]$; (ii) $[PtCl_2(NCMe)_2]$; (iii) reflux in ethanol for 4 h; (iv) NH_4PF_6 ; (v) 1 mole equivalent of NaOMe

because of quadrupole relaxation or chemical exchange effects, or it might be obscured by the aromatic proton signals.¹⁰

Treatment of $[PdCl_2(NCPh)_2]$ with 2 equivalents of L² gave bis(chelate) dication cis-[Pd{PPh_2CH_2C(Bu')=NNthe $H_2_{2}^{2^+}$, isolated as the dichloride 5a, and the bis(hexafluorophosphate) 5b, salts. The ¹H NMR pattern of the methylene hydrogens in both cases consisted of a second order or 'filled-in doublet with $N = {}^{4}J(PH) + {}^{2}J(PH) = 13.2$ Hz. We tentatively assign these to have *cis* geometry.^{11,12} We also treated the dicationic dichloride salt 5a with 2 moles of sodium ethoxide in ethanol in order to effect deprotonation. An orange product was obtained, which appeared to be essentially pure from the phosphorus-31 and proton NMR spectra although it was too unstable to prepare a sample pure enough for elemental analysis. However, the phosphorus-31 NMR spectrum showed a singlet resonance at δ_p 59.1 and the proton NMR spectrum showed a well-defined 'virtual triplet' pattern for the CH₂ protons with $N = {}^{2}J(PH) + {}^{4}J(PH) = 11.0$ Hz, suggesting that the phosphorus atoms were mutually trans. The fast atom bombardment (FAB) mass spectrum showed a parent molecular ion at m/z = 701, in agreement with M^+

We have prepared and described the crystalline mixed azine monophosphine $L^{3,1}$ formed by condensing benzaldehyde with the phosphino hydrazone, L^2 . This azine monophosphine probably has the Z,E-configuration; for comparison purposes the infrared and NMR data are given in Tables 1 and 2. The coordination chemistry of this ligand has been summarised in Scheme 2. Treatment of *trans*-[PdCl₂(NCPh)₂] with L^3 in dichloromethane at 20 °C gave the required chelated palladium dichloride complex 7a in over 90% yield. We formulate this as having a six-membered chelate ring, *i.e.* with the Zconfiguration around CH₂C(Bu¹)=N-N since it was formed under mild conditions and in the proton NMR spectrum coupling between phosphorus and =CHPh was observed $[{}^{4}J(PH) = 1.7 \text{ Hz}]$. The six-membered chelate ring structure is also supported by the carbon-13 NMR data (see Experimental section). We also prepared the dimethylplatinum complex with this ligand and formulate this with a six-membered chelate ring, as in 7b; in this case the values of ${}^{4}J(PCHPh) = 1.9 \text{ Hz}$ and ${}^{3}J(PtCHPh) = 37.0 \text{ Hz}$ exclude the five-membered chelate ring.

Treatment of *trans*-[PtCl₂(NCMe)₂] with 2 equivalents of the mixed azine phosphine L³ at 20 °C gave a bis(phosphine) complex which we formulate as the dichloride **8** with *trans* monodentate phosphines. Electric conductivity measurements in acetone solution showed this complex to be a non-electrolyte (specific molar conductivity, $\Lambda_m = 1.95 \times 10^{-2} \Omega^{-1} \text{ mol}^{-1}$ cm²).¹³ The coupling constant of phosphorus to platinum [¹J(PtP) = 2575 Hz] is typical of mutually *trans* phosphines co-ordinated to platinum(II).² Additionally, in the proton NMR spectrum, the CH₂ protons gave a virtual triplet with $N = {}^2J(PH) + {}^4J(PH) = 9.2$ Hz, with satellites due to platinum-195 coupling, ${}^3J(PtCH_2) = 24.9$ Hz, typical of mutually *trans* phosphines.¹²

When *trans*-[PtCl₂(NCMe)₂] was heated with 2 equivalents of L³ in ethanol-dichloromethane for 4 h, the hoped for dichloro salt **9a** was formed (Table 1), and was isolated as the bis(hexafluorophosphate) salt **9b**. Electric conductivity measurements in acetone solution showed this complex to be a 2:1 electrolyte (specific molar conductivity, $\Lambda_m = 0.2 \Omega^{-1} \text{ mol}^{-1}$ cm²).¹³ The coupling constant of phosphorus to platinum-195 of 3361 Hz is consistent with the phosphorus atoms being coordinated mutually *cis* and is much too high for mutually *trans* co-ordination.^{2,9} We found that at *ca*. 20 °C the resonance of the methylene protons (CH₂P) of **9b** was broad, and at -50 °C, an ABX pattern was observed but coupling to platinum was not ic, id or 7b

resolved. The broadness was possibly due to exchange or a fluxional process. When we treated a methanol solution of **9a** with 1 mole equivalent of sodium methoxide a product was formed which showed an AB phosphorus pattern with satellites $[\delta(P_A) \ 17.2, \ ^1J(PtP_A) = 3363$ and $\delta(P_B) \ 31.5, \ ^1J(PtP_B) = 3363, \ ^2J(P_AP_B) = 11 \ Hz]$, which we suggest was due to the monodeprotonated complex **10**. When we treated **10** with 1 mole equivalent of sodium methoxide we obtained the orange neutral complex *cis*-[$Pt{PPh_2CH=C(Bu^t)N-N=CHPh}_2$] **11** with δ_P 21.9 and $\ ^1J(PtP) = 3146$ Hz was obtained. In the proton NMR spectrum of **11**, the methine proton showed a doublet at δ_H 4.42, $\ ^2J(PH) = 8.8$ Hz, with satellites, $\ ^3J(PtH) = 22.2 \ Hz$ (Table 2). As expected complex **11** was found to be a non-electrolyte in acetone (specific molar conductivity, $\Lambda_m = 6.1 \times 10^{-3} \ \Omega^{-1} \ mol^{-1} \ cm^{2}$).¹³

We have also studied the action of carbon monoxide on the three dimethylplatinum complexes 1c, 1d and 7b (Scheme 3). When carbon monoxide was bubbled through a solution of 1c in C_6D_6 at 20 °C the ³¹P-{¹H} NMR spectrum due to 1c disappeared and was replaced by a singlet with satellites, δ_P 20.2, ¹J(PtP) = 1703 Hz. Moreover, the solution showed a strong infrared absorption at 2040 cm⁻¹ typical of carbon monoxide co-ordinated to platinum(II).¹⁴ The product proved to be too soluble to isolate but was further characterised by its proton NMR spectrum (Table 2). We formulate the complex as 12a with the ligand L¹ being monodentate through phosphorus. The other two chelates 1d and 7b reacted similarly and were similarly characterised as 12b and 12c, respectively, data in Tables 1 and 2.

C_cD_c

L 12a L¹ 12b L² 12c L³ 107

Crystal Structure of cis- $[Pt{PP_2CH_2C(Bu')=NNH}_2]$ 4.— The crystal structure of complex 4 is shown in Fig. 1, with selected bond lengths and angles in Table 3 and atom coordinates in Table 4. As can be seen from the figure and Table 3, the arrangement around the platinum is strictly planar, the phosphorus atoms are *cis* to each other as are the nitrogens and the six-membered chelate rings are puckered. The bond lengths and angles are not unusual.

Experimental

All the reactions were carried out in an inert atmosphere of dry nitrogen or dry argon. Infrared spectra were recorded using a Perkin-Elmer model 457 grating spectrometer. NMR spectra were recorded using a JEOL FX-90Q spectrometer (operating frequencies for ¹H and ³¹P of 89.5 and 36.2 MHz, respectively), a JEOL FX-100 spectrometer (operating frequencies for ¹H and ³¹P of 99.5 and 40.25 MHz, respectively) or a Bruker AM400 spectrometer (operating frequencies for ¹H, ³¹P and ¹³C of 400.13, 161.9 and 100.6 MHz, respectively; ¹H and ¹³C chemical shifts are relative to tetramethylsilane and ³¹P shifts are

 Table 3
 Selected bond lengths (pm) and angles (°) for complex 4 with estimated standard deviations (e.s.d.s) in parentheses

P(1)Pt	224.2(4)	P(6)-Pt	223.4(5)
N(5)-Pt	207.8(10)	N(10)-Pt	206.3(11)
C(2) - P(1)	183.1(11)	C(7) - P(6)	184.7(11)
C(3)-C(2)	150.1(16)	C(8) - C(7)	150.1(16)
C(31)-C(3)	154.7(15)	C(81)-C(8)	151.3(16)
N(4)-C(3)	129.7(13)	N(9)-C(8)	128.0(14)
N(5)-N(4)	139.0(12)	N(10)-N(9)	140.7(12)
P(6)Pt-P(1)	100.2(2)	N(10)-Pt-N(5)	81.8(4)
N(5)-Pt-P(1)	89.0(4)	N(10) - Pt - P(6)	89.2(3)
N(10)-Pt-P(1)	170.2(3)	P(6) - Pt - N(5)	170.1(3)
C(2)-P(1)-Pt	106.3(4)	C(7) - P(6) - Pt	107.8(4)
C(3)-C(2)-P(1)	113.0(7)	C(8)-C(7)-P(6)	113.3(8)
N(4)-C(3)-C(2)	121.3(10)	N(9)-C(8)-C(7)	123.4(10)
N(5)-N(4)-C(3)	119.7(10)	N(10)-N(9)-C(8)	119.7(10)
N(4)N(5)Pt	125.0(8)	N(9)-N(10)-Pt	125.7(7)



Fig. 1 Crystal structure of $cis-[Pt{PPh_2CH_2C(Bu')=NNH}_2]$ 4

Table 4Fractional non-hydrogen atomic coordinates ($\times 10^4$) for compound 4 with e.s.d.s in parentheses

Atom	x	у	Z	Atom	x	у	Z
Pt	663.3(3)	444.0(2)	2018.0(2)	C(611)	2695(4)	1750(3)	1966(3)
P(1)	233(2)	1196(1)	2850(2)	C(612)	3069(4)	2333(3)	1622(3)
C(111)	1410(5)	1794(3)	3539(4)	C(613)	4178(4)	2644(3)	2058(3)
C(112)	1475(5)	2534(3)	3322(4)	C(614)	4913(4)	2373(3)	2840(3)
C(113)	2452(5)	2966(3)	3798(4)	C(615)	4538(4)	1791(3)	3184(3)
C(114)	3363(5)	2658(3)	4490(4)	C(616)	3429(4)	1479(3)	2747(3)
C(115)	3297(5)	1918(3)	4707(4)	C(621)	263(5)	2021(3)	836(3)
C(116)	2321(5)	1487(3)	4232(4)	C(622)	361(5)	2733(3)	1173(3)
C(121)	-1037(4)	1803(3)	2397(3)	C(623)	-527(5)	3256(3)	793(3)
C(122)	-1302(4)	2349(3)	2845(3)	C(624)	-1512(5)	3067(3)	76(3)
C(123)	-2357(4)	2739(3)	2481(3)	C(625)	-1610(5)	2356(3)	-262(3)
C(124)	- 3147(4)	2584(3)	1670(3)	C(626)	-723(5)	1833(3)	119(3)
C(125)	-2882(4)	2037(3)	1222(3)	C(7)	1542(9)	794(6)	560(6)
C(126)	-1827(4)	1647(3)	1586(3)	C(8)	2289(9)	109(6)	856(6)
C(2)	-106(9)	585(6)	3538(6)	C(81)	3331(10)	23(6)	654(7)
C(3)	-1074(9)	44(5)	3081(6)	C(82)	4273(9)	645(6)	1124(7)
C(31)	-2163(8)	9(5)	3267(5)	C(83)	3932(11)	- 748(7)	934(8)
C(32)	-2717(10)	757(6)	3194(8)	C(84)	2926(11)	125(7)	-281(7)
C(33)	-1759(10)	-224(7)	4166(6)	N(9)	2055(7)	- 427(5)	1233(5)
C(34)	- 3018(10)	- 575(6)	2697(8)	N(10)	1049(7)	- 391(5)	1383(5)
N(4)	-985(7)	-437(5)	2569(5)	C(1S)	6874(13)	4494(8)	1364(9)
N(5)	22(8)	-451(5)	2435(6)	C(1S)	5663(4)	4476(2)	1619(3)
P(6)	1309(2)	1282(1)	1385(2)	C(2S)	6399(5)	4439(3)	296(3)

relative to 85% phosphoric acid. Mass spectra were recorded on an VG Autospec mass spectrometer with 8 kV acceleration.

The ligands PPh₂CH₂C(Bu^t)=NNR₂ (R = Me L¹ or H L²) and PPh₂CH₂C(Bu^t)=N-N=CHPh L³ were prepared according to our published procedure.¹ ¹³C-{¹H} NMR (100.6 MHz, CDCl₃): L¹ δ_c 177.1 (1 C, s, C=N), 141.6 [2 C, d, ¹J(PC) 16.9, C_{*ipso*}], 132.4 [4 C, d, ²J(PC) 20.1, C_{ortho}], 128.1 [4 C, d, ³J(PC) 7.0, C_{*meta*}], 128.1 (2 C, s, C_{*para*}), 45.8 (2 C, s, NMe₂), 38.4 (1 C, s, CMe₃), 29.0 [1 C, d, ¹J(PC) 21.6 Hz, CH₂] and 28.3 (3 C, s, CMe₃), 29.0 [1 C, d, ²J(PC) 19.7, C_{ortho}], 129.0 (2 C, s, C_{*para*}), 128.5 [4 C, d, ³J(PC) 7.0, C_{*meta*}], 38.1 [1 C, d, ³J(PC) 1.4, CMe₃], 28.6 (3 C, s, CMe₃) and 26.2 [1 C, d, ¹J(PC) 22.7 Hz, CH₂]; L³ δ_c 175.6 [1 C, d, ²J(PC) 4.4, Bu^tC=N], 157.3 [1 C, d, ⁵J(PC) 1.6, HC=N], 139.1 [2 C, d, ¹J(PC) 17.3, C_{*ipso*}], 134.6 [1 C, s, C¹ of CHPh], 133.0 [4 C, d, ²J(PC) 20.4, C_{ortho}], 131.4 (1 C, s, CHPh), 131.3 (1 C, s, CHPh), 130.8 (1 C, s, CHPh), 130.7 (1 C, s, CHPh), 130.1 (1 C, s, C⁴ of CHPh), 128.6 (2 C, s, C_{*para*}), 128.3 [4 C, d, ³J(PC) 6.7, C_{*meta*}], 38.7 (1 C, s, CMe₃), 28.9 [1 C, d, ¹J(PC) 22.5, CH₂] and 28.1 [3 C, d, ⁴J(PC) 1.9 Hz, CMe₃].

Preparations of Complexes.—[PtCl₂{PPh₂CH₂C(Bu¹)=NN-Me₂}] **1a**. A solution of [PtCl₂(cod)] (60 mg, 0.16 mmol) and the phosphino dimethylhydrazone L¹ (53 mg, 0.16 mmol) in dichloromethane (1.5 cm³) was put aside for 1 h at *ca*. 20 °C. The solvent was then removed under reduced pressure and methanol added to the residue gave the required product **1a** as a pale yellow microcrystalline solid. Yield 74 mg, 78% (Found: C, 39.55; H, 4.5; N, 4.5. C₂₀H₂₇Cl₂N₂PPt-0.25CH₂Cl₂ requires C, 39.65; H, 4.5; N, 4.55%). ¹³C-{¹H} NMR (25.2 MHz, CDCl₃): $\delta_{\rm C}$ 170.1 (1 C, s, C=N), 134.2 [4 C, d, ²J(PC) 10.8, ³J(PtC) 22, C_{ortha}], 132.5 [2 C, d, ⁴J(PC) 2.7, C_{para}], 128.8 [4 C, d, ³J(PC) 11.8, C_{meta}], 126.8 [2 C, d, ¹J(PC) 60.5, C_{ipso}], 55.6 [2 C, s, ²J(PtC) 14.7, NMe₂], 40.9 [1 C, d, ³J(PC) 1.7, CMe₃], 26.9 (3 C, s, CMe₃) and 17.9 [1 C, d, ¹J(PC) 24.0, ²J(PtC) 6.6 Hz, CH₂].

The dichloropalladium analogue **1b** was prepared similarly from [PdCl₂(NCPh)₂] and L¹ in 89% yield (Found: C, 46.65; H, 5.3; Cl, 15.55; N, 5.55. $C_{20}H_{27}Cl_2N_2PPd\cdot0.15CH_2Cl_2$ requires C, 46.85; H, 5.3; Cl, 15.8; N, 5.4%).

[$PtMe_2$ { $PPh_2CH_2C(Bu')=NNMe_2$ }] 1c. A solution of [Pt-Me_2(cod)] (50 mg, 0.15 mmol) and the phosphino dimethylhydrazone L¹ (50 mg, 0.15 mmol) in benzene (3 cm³) was put aside for 1 h. The dimethylplatinum complex 1c crystallized out as a white solid after 1 h and was collected. Yield: 45 mg, 55%. (Found: C, 47.75; H, 6.2; N, 5.0. $C_{22}H_{33}N_2PPt$ requires C, 47.90; H, 6.0; N, 5.1%).

The dimethylplatinum complex 1d was prepared in a similar manner from equimolar amounts of the phosphino hydrazone L^2 and [PtMe₂(cod)]. Yield 40% (Found: C, 46.05; H, 5.55; N, 5.35. $C_{20}H_{29}N_2PPt$ requires C, 45.9; H, 5.60; N, 5.35%). ¹³C-{¹H} NMR (100.6 MHz, C₆D₆): δ_C 169.5 (1 C, s, C=N), 134.1 [4 C, d, ²J(PC) 12.0, C_{ortho}], 133.1 [2 C, d, ¹J(PC) 39.0, C_{ipso}], 130.1 [2 C, d, ⁴J(PC) 1.8, C_{para}], 128.3 [4 C, d, ³J(PC) 7.2, C_{meta}], 53.0 (2 C, s, NMe₂), 40.7 [1 C, d, ²J(PC) 2.3, CMe₃], 27.8 (3 C, s, CMe₃), 21.3 [1 C, d, ¹J(PC) 16.7, CH₂], 16.0 [1 C, d, ¹J(PC) 117.7, PtMe trans to P] and -21.9 [1 C, d, ¹J(PC) 3.9 Hz, PtMe, cis to P].

[$\dot{P}tMe_3I{\dot{P}Ph_2CH_2C(Bu^{1})=N\dot{N}Me_2}$] **2a**. An excess of iodomethane (0.1 cm³) was added to a solution of the dimethylplatinum complex **1c** (38 mg, 0.07 mmol) in benzene (1.0 cm³). The solvent was then removed, trituration of the resultant yellow oil with methanol gave the required complex **2a** as a white microcrystalline solid. Yield: 37 mg, 77% (Found: C, 40.4; H, 5.25; N, 4.05. C₂₃H₃₆IN₂PPt•0.1C₆H₆ requires C, 40.4; H, 5.25; N, 4.0%).

[PtMe₃I{PPh₂CH₂C(Bu⁴)=NNH₂}] **2b**. Iodomethane (0.5 cm³) was added to a solution of the dimethylplatinum complex **1d** (80 mg, 0.15 mmol) in benzene (2 cm³). The reaction mixture was put aside for 1.5 h, filtered, and the solvent was removed under reduced pressure. Addition of methanol to the residue gave the required complex **2b** as colourless prisms (71 mg, 70%) (Found: C, 38.3; H, 4.85; N, 4.35. C₂₁H₃₂IN₂PPt requires C, 37.9; H, 4.85; N, 4.20%).

cis-[Pt{PPh₂CH₂C(Bu^t)=NNH₂}₂]Cl₂ **3a**. The complex [PtCl₂(cod)] (125 mg, 0.33 mmol) was added to a stirred solution of the phosphino hydrazone L² (200 mg, 0.67 mmol) in dichloromethane. After 30 min, the required product **3a** was formed in the reaction mixture as a white solid. This was filtered off, and washed with dichloromethane. Yield: 200 mg, 70% (Found: C, 48.9; H, 5.6; N, 6.15. C₃₆H₄₆Cl₂N₄P₂Pt•0.35CH₂Cl₂ requires C, 48.9; H, 5.3; N, 6.25%).

cis-[$\dot{P}t{\dot{P}Ph_2CH_2C(Bu')=N\dot{N}H_2}_2$]Cl[PF₆] **3b**. This complex was prepared by addition of a solution of ammonium hexafluorophosphate(v) in methanol to **3a** in methanol in 64% yield (Found: C, 43.85; H, 4.9; Cl, 4.15; N, 5.55. C₃₆H₄₆-

 $ClF_6N_4P_3Pt.0.15CH_2Cl_2$ requires C, 44.05; H, 4.8; Cl, 4.65; N, 5.7%).

cis-[Pt{PPh₂CH₂C(Bu¹)=NNH}₂] 4. The complex [PtCl₂-(cod)] (120 mg, 0.32 mmol) was added to the phosphino hydrazone L² (200 mg, 0.67 mmol) in dichloromethane (5 cm³). A white solid was obtained which was filtered off, dried and then dissolved in hot ethanol (4 cm³). Sodium ethoxide solution (1.2 cm³, 0.48 mol dm⁻³ in ethanol, 0.58 mmol) was added. After 1.5 h, the solvent was reduced to a low volume under reduced pressure. The required product 4 was obtained as a pale yellow solid. Yield 205 mg, 81% (Found: C, 54.55; H, 5.65; N, 7.05. C₃₆H₄₄N₄P₂Pt requires C, 54.75; H, 5.65; N, 7.10%).

 $[Pd{PPh_2CH_2C(Bu')=NNH_2}_2]Cl_2$ 5a. A mixture of the phosphino hydrazone L² (200 mg, 0.5 mmol) and $[PdCl_2(NCPh)_2]$ (95 mg, 0.25 mmol) in dichloromethane (3 cm³) was put aside at room temperature for 1 h. Complex 5a was obtained as a white solid, which was filtered off and washed with a little dichloromethane. Yield 183 mg, 95% (Found: C, 52.65; H, 6.0; N, 7.05. C₃₆H₄₆Cl₂N₄P₂Pd·0.75CH₂Cl₂ requires C, 52.70; H, 5.7; N, 7.0%).

 $[Pd{PPh_2CH_2C(Bu')=NNH_2}_2][PF]_2$ **5b**. Complex **5b** was prepared by the addition of a methanolic solution of ammonium hexafluorophosphate(v) to a solution of **5a** in hot methanol. The required salt was obtained as a pale yellow solid in 60% yield (Found: C, 44.0; H, 4.8; N, 6.05. C₃₆H₄₆F₁₂N₄P₄Pd requires C, 43.6; H, 4.7; N, 5.65%).

trans-[Pd{PPh₂CH₂C(Bu')=NNH₂] **6**. A solution of the palladium salt **5a** (80 mg, 0.20 mmol) in ethanol (1.5 cm³) was treated with a solution of sodium ethoxide (0.48 mol dm⁻³ in ethanol, 0.45 cm³, 0.21 mmol) to give the required compound **6** as an orange solid. Yield 80 mg, 57%; m/z 701 (M^+). The complex decomposed on attempted recrystallization.

[$^{h}dCl_{2}{^{h}Ph_{2}CH_{2}C(Bu^{i})=N-\dot{N}=CHPh}$] 7a. A solution of the azine monophosphine L³ (60 mg, 0.16 mmol) and [$^{P}dCl_{2}(NCPh)_{2}$] (59 mg, 0.15 mmol) in dichloromethane was put aside for 30 min. The solvent was then removed under reduced pressure and methanol added to the residue; this gave the required palladium complex 7a as a yellow microcrystalline solid. Yield 79 mg, 91% (Found: C, 50.7; H, 4.75; N, 5.05. C_{2s}H_{27}Cl_{2}N_{2}N_{2}PPd\cdot1.5CH_{2}Cl_{2} requires C, 50.5; H, 4.7; N, 4.6%). ¹³C-{¹H} NMR (100.6 MHz, CDCl_{3}): δ_{C} 175.8 (1 C, s, Bu'C=N), 162.8 [1 C, $^{3}J(PC)$ 3.3, HC=N], 134.8 [2 C, d, $^{2}J(PC)$ 12.1, C_{ortho}], 133.8 (1 C, s, C_{para}), 133.4 [2 C, d, $^{1}J(PC)$ 57.8, C_{ipso}], 132.7 (1 C, s, CHPh), 132.3 [2 C, d, $^{2}J(PC)$ 9.9, C_{ortho}], 131.3 (1 C, s, C_{para}), 129.9 (1 C, s, C⁴ of CHPh), 129.4 [2 C, d, $^{3}J(PC)$ 11.2, C_{meta}], 128.9 [2 C, d, $^{3}J(PC)$ 10.2, C_{meta}], 128.8 (1 C, s, CHPh), 33.2 [1 C, d, $^{3}J(PC)$ 10.9, CMe₃], 27.0 (3 C, s, CMe₃) and 25.2 [1 C, d, $^{1}J(PC)$ 25.3 Hz, CH₂] (C¹ of CHPh was obscured).

[$\dot{P}tMe_2(\dot{P}Ph_2CH_2C(Bu')=N-\dot{N}=CHPh$]] 7b.—A solution of the azine monophosphine L³ (120 mg, 0.31 mmol) and [$PtMe_2(cod)$] (100 mg, 0.30 mmol) in degassed benzene (2 cm³) was put aside for 2 h. The solvent was then removed under reduced pressure and methanol added to the residue to give the required compound 7b as a yellow solid. Yield 130 mg, 71% (Found: C, 51.85; H, 5.7; N, 4.25. C₂₇H₃₃N₂PPt-1.5CH₃OH requires C, 51.9; H, 5.95; N, 4.25%).

trans-[PtCl₂{PPh₂CH₂C(Bu¹)=N-N=CHPh}₂] 8. A solution of the azine monophosphine L³ (90 mg, 0.23 mmol) and [PtCl₂(NCMe)₂] (40 mg, 0.11 mmol) in dichloromethane (2 cm³) was stirred for 1 h, until a clear solution was obtained. The solvent was then removed under reduced pressure and the residue triturated with methanol to give the required product 8 as a white solid. Yield 52 mg, 43% (Found: C, 54.25; H, 4.85; N, 4.95. C₅₀H₅₄Cl₂N₄P₂Pt·1.1CH₂Cl₂ requires C, 54.2; H, 5.00; N, 4.95%).

cis-[Pt{PPh₂CH₂C(Bu^t)=N-N=CHPh}₂][PF₆]₂ 9b. A solution of the azine monophosphine L³ (90 mg, 0.23 mmol) in

ethanol (2 cm³) was added to a suspension of $[PtCl_2(NCMe)_2]$ (40 mg, 0.11 mmol) in dichloromethane (1 cm³). The reaction mixture was then refluxed under a nitrogen atmosphere for 4 h after which, a saturated solution of ammonium hexafluorophosphate(v) in methanol was then added to precipitate the required product **9b** as a white solid (76 mg, 53%) (Found: C, 44.85; H, 3.9; N, 4.4. C₅₀H₅₄F₁₂N₄P₂Pt·1.4CH₂Cl₂ requires C, 44.85; H, 4.1; N, 4.1%).

cis-[Pt{PPh₂CH=C(Bu')-N-N=CHPh}₂] 11. The complex [PtCl₂(CNMe)₂] (40 mg, 0.11 mmol) and the azine monophosphine L³ (90 mg, 0.23 mmol) were dissolved in dichloromethane (1.5 cm³). Ethanol (1 cm³) was then added and the reaction mixture was refluxed for 3.5 h to give a pale yellow solution. Sodium methoxide (0.673 mol dm⁻³ in methanol, 0.34 cm³, 0.23 mmol) was then added, the resultant dark red solution filtered, and the solvents removed under reduced pressure. The required product 11 was obtained as an orange solid. Yield 43 mg, 40% (Found: C, 59.85; H, 5.25; N, 5.4. $C_{50}H_{52}N_4P_2Pt$ -0.6CH₂Cl₂ requires C, 59.80; H, 5.25; N, 5.5%).

Single-crystal X-Ray Diffraction Analysis of Complex 4.—All crystallographic measurements were carried out on a Stoe STADI4 diffractometer operating in the ω - θ scan mode using graphite-monochromated Mo-K α X-radiation ($\lambda = 71.069$ pm). The data set was corrected for absorption using azimuthal ψ -scans.

The structure was determined by standard heavy-atom methods and was refined by full-matrix least squares using SHELX 76.¹⁵ All non-hydrogen atoms were refined with anisotropic thermal parameters and the phenyl groups were treated as rigid bodies with idealised hexagonal symmetry (C–C 139.5 pm). All hydrogen atoms were included in calculated positions (C–H 96 pm) and were refined with an overall isotropic thermal parameter. The weighting scheme $w = [\sigma^2(F_o) + 0.0004(F_o)^2]^{-1}$ was used. Final non-hydrogen atomic coordinates are given in Table 4. An ORTEP¹⁶ diagram of the molecular structure of 4 is given in Fig. 1.

Crystal data. $C_{36}H_{44}N_4P_2Pt$ ·CH₂Cl₂, 0.65 × 0.5 × 0.45 mm, M = 874.73 (includes solvent molecule), monoclinic, space group $P2_1/c$, a = 1261.5(2), b = 1794.6(3), c = 1792.0(4) pm, U = 3.6675(12) nm³, Z = 4, $D_c = 1.58$ Mg m⁻³, $\mu = 39.08$ cm⁻¹, F(000) = 1752.

Data collection. $4.0 < 2\theta < 50.0^\circ$, 6843 data collected, 4088 with $I > 2.0\sigma(I)$ considered observed, T = 200 K.

Structure refinement. Number of parameters = 381, R = 0.0451, R' = 0.0443.

Additional material available from the Cambridge Crystallographic Data Centre comprises H-atom coordinates, thermal parameters and remaining bond lengths and angles.

Acknowledgements

We thank the SERC for a post-doctoral research fellowship (to S. D. P.) and for other support. We also thank the Committee of Vice-Chancellors and Principals for an ORS award (to K. K. H.), the trustees of the Lord Edward Boyle Foundation for a research scholarship (to K. K. H.) and the University of Leeds for the Tetley and Lupton Scholarship (to K. K. H.).

References

- I K. K. Hii, S. D. Perera, B. L. Shaw and M. Thornton-Pett, J. Chem. Soc., Dalton Trans., 1992, 2361.
- 2 P. S. Pregosin and R. W. Kunz, ³¹P and ¹³C NMR of Transition Metal Phosphine Complexes, Springer, Berlin, Heidelberg, New York, 1979.
- 3 S. D. Perera, B. L. Shaw and M. Thornton-Pett, J. Chem. Soc., Dalton Trans., 1992, 999 and refs therein.
- 4 J. D. Ruddick and B. L. Shaw, J. Chem. Soc. A, 1969, 2801.
- 5 S. D. Perera, B. L. Shaw and M. Thornton-Pett, J. Chem. Soc., Dalton Trans., 1992, 1469.

109

J. CHEM. SOC. DALTON TRANS. 1994

- 6 S. D. Perera, B. L. Shaw and M. Thornton-Pett, J. Chem. Soc.,
- Dalton Trans., 1993, 3653.
 J. D. Kennedy, W. McFarlane, R. J. Puddaphatt and P. J. Thompson, J. Chem. Soc., Dalton Trans., 1976, 874.
- 8 D. E. Clegg, J. R. Hall and G. A. Swile, J. Organomet. Chem., 1972, 38, 403.
- 9 A. Albanati, P. S. Pregosin and F. Wombacher, Inorg. Chem., 1990, **29**, 1812.
- 10 J.-T. Chen, T.-M. Huang, M.-C. Cheng, Y.-C. Lin and Y. Wang, Organometallics, 1992, 11, 1761.
- 11 D. A. Redfield, J. H. Nelson and L. W. Cary, Inorg. Nucl. Chem. Lett., 1974, 10, 727.

- J. M. Jenkins and B. L. Shaw, J. Chem. Soc. A, 1966, 770.
 W. J. Geary, Coord. Chem. Rev., 1971, 7, 81.
 G. K. Anderson and R. J. Cross, J. Chem. Soc., Dalton Trans., 1979, 1246.
- 15 G. M. Sheldrick, SHELX 76, program system for X-ray structure determination, University of Cambridge, 1976. 16 C. K. Johnson, ORTEP II, Report ORNL-5138, Oak Ridge National
- Laboratory, TN, 1976.

Received 23rd July 1993; Paper 3/04366E

[©] Copyright 1994 by the Royal Society of Chemistry