


Dyotropic Rearrangements of Fused Tricyclic β -Lactones: Application to the Synthesis of (–)-Curcumanolide A and (–)-CurcumalactoneCarolyn A. Leverett,^{†,¶} Vikram C. Purohit,^{†,§} Alex G. Johnson,^{†,⊥} Rebecca L. Davis,[‡] Dean J. Tantillo,[‡] and Daniel Romo^{*,†}[†]Department of Chemistry, Texas A&M University, P.O. Box 30012, College Station, Texas 77842-3012, United States[‡]Department of Chemistry, University of California-Davis, One Shields Avenue, Davis, California 95616, United States

S Supporting Information

ABSTRACT: Dyotropic rearrangements of fused, tricyclic β -lactones are described that proceed via unprecedented stereospecific, 1,2-acyl migrations delivering bridged, spiro- γ -butyrolactones. A unique example of this dyotropic process involves a fused bis-lactone possessing both β - and δ -lactone moieties which enabled rapid access to the core structures of curcumanolide A and curcumalactone. Our current mechanistic understanding of the latter dyotropic process, based on computational studies, is also described. Other key transformations in the described divergent syntheses of (–)-curcumanolide A and (–)-curcumalactone from a common intermediate (11 and 12 steps from 2-methyl-1,3-cyclopentanedione, respectively), include a catalytic, asymmetric nucleophile (Lewis base)-catalyzed aldol-lactonization (NCAL) leading to a tricyclic β -lactone, a Baeyer–Villiger oxidation in the presence of a β -lactone, and highly facial-selective and stereocomplementary reductions of an intermediate spirocyclic enoate. The described dyotropic rearrangements significantly alter the topology of the starting tricyclic β -lactone, providing access to complex spirocyclic cyclopentyl- γ -lactones and bis- γ -lactones in a single synthetic operation.



INTRODUCTION

Curcumanolides A (1) and B (2) are sesquiterpenoid natural products previously isolated from *Curcuma zedoaria* and related species (Figure 1).¹ Both of these spiro-lactones are present in the crude drug Zedoary, the ground rhizome of *C. zedoaria* Roscoe, which has been used medicinally in China for a variety of indications.² Curcumalactone (3), a dihydro derivative of curcumanolide A, was isolated from *C. aromatica* Silib³ and *C. wenyujin*⁴ which have long been used as remedies for cervical cancer in China⁵ and were reported to exhibit anti-inflammatory activity.³ While no information regarding the biological activity of curcumanolides A and B is available, the structural similarity with curcumalactone elevates their interest as targets for further biological studies. In particular, the enoate and spirocyclic γ -lactone moieties attracted our interest since these constitute potential electrophilic sites, which could render these natural products as covalent modifiers of their putative cellular targets.

In addition to their potential bioactivity, the unique, functionalized cyclopentane bearing a spiro-lactone found in these natural products has attracted significant interest from the synthetic community, leading to diverse strategies to access this common core. The first total synthesis of curcumanolide A and the only reported synthesis of curcumalactone to date, both in racemic form, were accomplished by Kato employing a bromonium-induced cyclization/ring contraction route from geraniol.⁶ Subsequently, Honda reported the first enantio-




Figure 1. Spirolactone natural products isolated from extracts of *Curcuma* species.

selective synthesis of curcumanolide A from (–)-carvone establishing the absolute configuration,⁷ and Fujita utilized an enzymatic transesterification process to access (–)-curcumanolide A.⁸ However, the reported syntheses have led to little additional biological information, and there is no information regarding possible cellular targets.

Dyotropic rearrangements were originally defined as reactions in which “two σ -bonds simultaneously migrate intramolecularly” (Figure 2a),⁹ but a variety of examples have since been reported in which the simultaneity of the two σ -bond migrations varies from synchronous (and concerted) to asynchronous (and concerted) to stepwise (separated by a discrete intermediate).¹⁰ Several types of dyotropic rearrangements have been employed in various synthetic contexts,

Received: April 17, 2012





Figure 2. (a) Prototypical dyotropic rearrangement wherein two σ -bonds simultaneously migrate intramolecularly (b) Dyotropic rearrangements of β -lactones involving hydrogen, electron-rich groups, and amino migrations. (c) β -Lactone dyotropic rearrangements involving acyl and carboxylate migrations.

ranging from the early report of a dyotropic interchange of vicinal Br atoms on a steroid skeleton,¹¹ to more recent applications in the synthesis of *N*-cis-propenyl amides,¹² trisubstituted olefins,¹³ and tetraazaphthalenes.¹⁴ A widely studied dyotropic process, of particular interest to us, is the Lewis acid-mediated rearrangement of β -lactones leading to various γ -butyrolactones (Figure 2b). This area of dyotropic rearrangement chemistry commenced with seminal reports by Mulzer on the ring expansion of β -lactones through a dyotropic process in which the β -lactone oxygen and electron-rich migrating groups exchanged positions.¹⁵ Related reports include Reetz's demonstration that acyclic amino migrating groups allow for the generation of 3-amino- γ -lactones,¹⁶ Cossio's study of the dyotropic rearrangement of propiolactones to γ -butyrolactones under similar conditions,¹⁷ and Black's examination of the dyotropic rearrangement of spirobicyclic β -lactones to a variety of fused γ -butyrolactone systems.¹⁸

While β -lactone dyotropic reactions with electron-rich migrating groups have been studied extensively, we recently reported the first example of a 1,2-acyl migration in a β -lactone dyotropic process (Figure 2c).¹⁹ This Lewis-acid mediated, stereospecific, dyotropic rearrangement of fused tricyclic β -lactones enables the synthesis of spirocyclic, bridged γ -butyrolactones. Subsequent studies revealed plausible mechanistic pathways for this rearrangement, formulated on the basis of both experimental and theoretical results.²⁰ We projected that a related dyotropic process involving a bis-lactone bearing both a δ - and a β -lactone moiety would enable efficient access to the spirocyclic core structure of members of the *Curcuma* family of natural products. Herein, we describe concise synthetic routes to (–)-curcumanolide A (1) and (–)-curcumlactone (3) that demonstrate the utility of these dyotropic processes for complex natural product synthesis with the latter being the first reported asymmetric synthesis.

Retrosynthetic Analysis of Curcumanolides and Curcumlactone. Access to the spirocyclic cores (e.g., 4, Scheme 1) of the curcumanolides (1, 2) and curcumlactone

Scheme 1. Retrosynthetic Analysis of Curcumlactone and Curcumanolide A




(3) was envisioned employing our recently described nucleophile-catalyzed aldol-lactonization (NCAL) process involving desymmetrization of dione 7 with homobenzo-tetramisole (HBTM) as the Lewis base promoter.²¹ This process would deliver tricyclic β -lactone 6 and subsequent Baeyer–Villiger oxidation and dyotropic rearrangement would provide bis-lactone 4 bearing the common spirocyclic core of the targeted natural products. Completion of the synthesis of (–)-curcumlactone and (–)-curcumanolide A would require further transformations of the tricycle 4 including two stereochemistry setting steps that would be dictated by the topologically rich spirocyclic core. In particular, a facially selective hydrogenation of an intermediate cyclopentene would set the C4-methyl-bearing stereocenter of the two targets and a diastereoselective reduction of an enoate would introduce the C7-isopropyl-bearing stereocenter of curcumlactone.

RESULTS AND DISCUSSION

The synthesis of the core structure of both curcumanolide A (1) and (–)-curcumlactone (3) began with the multigram-scale synthesis of dione acid 7 from commercially available 2-methyl-1,3-cyclopentanedione 8 (Scheme 2). We previously

Scheme 2. Multigram Synthesis of Dione Ketoacid 7 from 1,3-Cyclopentanedione^a




^aTBAI, tetrabutylammonium iodide.


utilized a four-step sequence to access this substrate;¹⁹ however, scale-up of this sequence proved problematic and inefficient. After extensive optimization, a streamlined synthesis of this key substrate was developed involving a one-pot alkylation/hydrolysis sequence to directly deliver dione acid 9 employing commercially available methyl 4-bromocrotonate in 1 M NaOH as solvent under phase transfer conditions. This method enabled production of >20 g of dione acid 9 in a single run and is useful for the large-scale synthesis of a number of related dione acids following hydrogenation of the alkene.²²

With a scaleable route to dione 7 in hand, extensive optimization was required to develop a practical, catalytic, asymmetric NCAL process to access tricyclic β -lactone (+)-6. Initial studies of this transformation employed 1.5 equiv of (–)-tetramisole-HCl (10)²³ as a promoter in this desymmet-

rization process and the modified Mukaiyama reagent **14**, which led to the desired product with high optical purity but only in moderate yields (Table 1, entry 1). Several other

Table 1. Optimization of the NCAL Process with Dione Acid **7 Leading to Tricyclic β -Lactone (+)-**6****



entry	activating agent	Lewis base	% yield ^{a,b}	% ee ^c
1		(-)-Tet·HCl (10 , 150 mol%)	35	98 ^d
2	<i>p</i> -NO ₂ C ₆ H ₄ SO ₂ Cl	(-)-Tet·HCl (10 , 25 mol%)	11	67 ^d
3	CDI	"	34	1.8 ^d
4	3,5-(NO ₂) ₂ BzCl	"	25	ND
5	<i>p</i> -TsCl	"	23	53 ^d
6	<i>p</i> -TsCl	(<i>S</i>)-BTM (11 , 20 mol%)	33	20 ^d
7	<i>p</i> -TsCl	(<i>S</i>)-HBTM (12 , 20 mol%)	58	96
8	14	(<i>S</i> , <i>R</i>)-Naphth/Me HBTM (13 , 20 mol%)	40	91

^aSingle diastereomer observed by ¹H NMR. ^bRefers to isolated, purified products. ^cDetermined by chiral GC analysis. ^d(-)-Tetramisole·HCl (**10**) and (*S*)-BTM (**11**) provide (-)-**6**. ^eWith 1.0 equiv LiCl added. CDI, carbonyldiimidazole; Ts, tosyl; Bz, benzoyl; Naphth, naphthyl.


carboxylic acid activating agents were explored but led to inferior yields and generally significantly decreased enantioselectivities, especially when catalytic amounts of (-)-tetramisole·HCl were employed (Table 1, entries 2–5). We next

studied the use of alternative chiral isothiourea catalysts (**11**–**13**, Table 1) developed by Birman and related to tetramisole (**10**), as Lewis base promoters for the NCAL process.²⁴ The use of 20 mol % (*S*)-HBTM (**11**)^{24a} provided good yields and excellent enantioselectivities (58%, 96% ee) of tricyclic β -lactone (+)-**6** in conjunction with *p*-toluenesulfonyl chloride (*p*-TsCl) as activating agent. A further significant increase in yield was realized with use of the mild Lewis acid, LiCl (1.0 equiv), with no sacrifice in enantioselectivity (77%, 96% ee), and this process was utilized to access a number of bicyclic and tricyclic β -lactones with good to excellent yields and enantioselectivities.²¹


Toward the synthesis of curcumlactone and congeners, conducting the NCAL process on gram scale with (*R*)-HBTM provided a 65% yield of the desired tricyclic β -lactone (-)-**6** in high optical purity (98% ee, Scheme 3). The absolute stereochemistry of (-)-**6** was confirmed indirectly by conversion of the enantiomeric tricyclic β -lactone (+)-**6** derived from (*S*)-HBTM to the corresponding oxazolidinone via a Curtius rearrangement (Scheme 3, inset). Following conversion to the *N*-*p*-bromobenzoate (+)-**15**, assignment of absolute configuration of tricyclic β -lactone (+)-**6** was made possible by X-ray analysis using heavy atom anomalous dispersion.²¹ Thus, the relative and absolute configuration of (-)-**6** was secured by comparison of NMR and optical rotations with the enantiomeric series. A subsequent Baeyer–Villiger oxidation of tricyclic β -lactone (-)-**6** under buffered conditions led to slow conversion to the ring-expanded δ -lactone (-)-**5** in moderate yield, setting the stage for the key dyotropic rearrangement. It is worth noting that this oxidative rearrangement is possible, notwithstanding the presence of the potentially reactive β -lactone.

Dyotropic Rearrangements of Tricyclic Acyl- β -lactones and Bis-lactones. We began our studies of the dyotropic rearrangement of acyl- β -lactones with tricyclic β -lactone **16a** (Table 2), available by the NCAL reaction of a cyclohexanedione precursor. We surmised that Lewis acid activation of the β -lactone would promote a dyotropic rearrangement via 1,2-acyl migration leading to simultaneous and stereospecific ring expansion of the β -lactone to a γ -lactone

Scheme 3. Gram-Scale Synthesis of Optically Active Bis-lactone (-)-6** and Confirmation of Absolute Stereochemistry^{a,25}**



^aHBTM, homobenzotetramisole; TsCl, *p*-toluenesulfonyl chloride; *m*-CPBA, *m*-chloroperoxybenzoic acid; NCAL, nucleophile-catalyzed aldol lactonization; DPPA, diphenylphosphoryl azide. Inset: ORTEP representation of the X-ray crystal structure of tricyclic carbamate (+)-**15**; hydrogens removed for clarity; thermal ellipsoids shown at 50% probability.

Table 2. Dyotropic 1,2-Acyl Migration of Tricyclic β -Lactones


entry	tricyclic- β -lactone	spirocycle	% yield 17^a
1	(±)- 16a	(±)- 17a	94
2	(±)- 16b	(±)- 17b + (±)- 18 (3%)	90
3	(±)- 16c	(±)- 17c	85
4	(±)- 16d	(±)- 17d	63
5	(±)- 16e	(±)- 17e	55
6	(±)- 16a	(±)- 17f	73 ^b


^aRefers to isolated yields. ^bReaction performed using 0.2 equiv TMSOTf at 23 °C for 12 h instead of Zn(OTf)₂. Isolated yield of diacid **17f** accompanied by only trace amounts of dyotropic product.

with concomitant ring contraction of the cyclohexanone. It should be noted that in these rigid tricyclic systems, stereospecificity is enforced due not solely to synchronicity of migration but also to ring constraints. Importantly, β -lactone substrate **16** possesses the required antiperiplanar relationship between the migrating C₄–O₁ bond of the β -lactone and the C₅–C=O bond of the cyclohexanone. After screening a variety of Lewis acids, including AlBr₃, Et₂AlCl, LaCl₃, PrCl₃, YCl₃, MgBr₂, Yb(OTf)₃, In(OTf)₃, and Mg(OTf)₂, the best yields for this transformation were realized using Zn(II) salts, providing up to 94% yield of **17a** when employing 1.1 equiv of Zn(OTf)₂ (Table 2, entry 1). Substoichiometric amounts of Zn(OTf)₂ led to significant amounts of recovered starting material. Other tricyclic β -lactones **16b–d** were also studied under these conditions and gave excellent yields of bridged γ -lactones **17b–d** (Table 2, entries 2–4) with high stereospecificity (dr >19:1). It is important to note that in the case of tricyclic β -lactone **16e** (entry 5), which lacks the antiperiplanar relationship between the β -lactone C₄–O bond and the C₅–acyl carbon, only olefin acid **17e** was isolated (55%), likely generated from β -lactone opening and subsequent elimination. By a presumed similar pathway, the tricyclic β -lactone **16b** produced trace amounts (3%) of β,γ -unsaturated acid **18** in addition to the expected dyotropic product **17b** in high yield. These two examples provided our initial evidence for carbocation formation in the Zn(II)-promoted dyotropic rearrangement, which was later

supported computationally, as detailed below. Interestingly, while Zn(II) salts led to good yields of dyotropic products in most cases, subjecting β -lactone **16a** to catalytic TMSOTf provided only ring-opened diacid **17f** (73%, entry 6), suggestive of an intervening Grob-type fragmentation, which was subsequently confirmed to occur due to adventitious TfOH and not TMSOTf under these conditions.²⁰ The first dyotropic rearrangement of an optically active tricyclic β -lactone was also performed with lactone (–)-**16a** (97% ee) using Zn(OTf)₂ and this provided γ -lactone (–)-**17a** in excellent yield (91%) with high stereochemical fidelity (99% ee, chiral GC).

Toward application of the dyotropic process to the targeted natural products, we studied the rearrangement of tricyclic bis-lactone (–)-**5** bearing both a δ - and β -lactone (Scheme 4).

Scheme 4. Enantioselective Synthesis of Spirolactone (+)-**4** via a Bis-Lactone Dyotropic Rearrangement^a



^aTMSOTf, trimethylsilyl trifluoromethanesulfonate.

While use of Zn(OTf)₂ gave only recovered bis-lactone (–)-**5**, a brief survey of Lewis acids led to use of substoichiometric TMSOTf (15 mol %), which slowly produced the desired spiro- γ -lactone (+)-**4** in good yield (73%) and with complete stereochemical fidelity (98% ee).

Mechanistic Studies of the Dyotropic Rearrangement of Tricyclic β -Lactones Involving 1,2-Acyl Migrations.

The described dyotropic rearrangements of β -lactones involving 1,2-acyl migrations were unprecedented in type I dyotropic processes.^{10a} Our initial hypotheses for this rearrangement invoked a 2-electron-3-centered acylium intermediate or a carboxylate-stabilized, four-membered transition state. Both pathways would benefit from a homoconjugated, carbonyl-assisted migration. However, subsequent theoretical (B3LYP/6-31+G(d,p)) and experimental studies of this dyotropic process (e.g., **16** → **17**, Table 2) revealed that the concertedness of this rearrangement was dependent on the nature of the Lewis acid employed.²⁰ Theoretical calculations on both the Zn(OTf)₂- and ZnCl₂-promoted rearrangements of **16a** to **17a** revealed that these reactions occur in a stepwise manner, involving an intermediate with a Zn(II)-complexed carboxylate and a tertiary carbocation center (cf. **19**, Figure 3). However, attempts to trap this carbocation intermediate by both inter- and intramolecular nucleophiles were unsuccessful. These observations are consistent with the computational results, which suggested that the 1,2-acyl shift/carbocation capture process is extremely facile for [6-5] bicyclic




Figure 3. (a) Intermediate for Zn(II)-based Lewis acid-mediated and (b) transition state structure for TMSOTf-mediated, dyotropic rearrangement of tricyclic β -lactone **16a**.




Figure 4. Computed geometries (B3LYP/6-31+G(d,p))²⁷ of species involved in the rearrangement of TMS-ent-5 to TMS-ent-4. Energies shown are in kcal/mol and are relative to that of the reactant complex (ZPE-corrected energies in normal text; free energies at 25 °C in italics). Selected bond lengths are given in Å.

carbocations, making this intermediate difficult to intercept. Overall, the ΔG^\ddagger for the reaction with $\text{Zn}(\text{OTf})_2$ was calculated to be 16 kcal/mol in the gas phase (<10 kcal/mol in CH_2Cl_2), and the product was calculated to be favored by 12 kcal/mol relative to the β -lactone– $\text{Zn}(\text{OTf})_2$ reactant.

In contrast, examination of the TMSOTf-mediated rearrangement of the tricyclic β -lactone **16a** indicated that, unlike the $\text{Zn}(\text{II})$ -bound species, this species can undergo a low-barrier *concerted* dyotropic rearrangement (cf. **20**, Figure 3).²⁰ This reaction is predicted to have a barrier of only 6–7 kcal/mol in the gas phase (5 kcal/mol in CH_2Cl_2). These results indicate that the reaction can proceed readily at room temperature to produce the desired γ -lactone, despite the fact that only trace amounts of the γ -lactone product were observed experimentally using TMSOTf.

Dyotropic Rearrangements of Bis-lactones. We also examined the mechanism of the bis-lactone dyotropic rearrangement, which delivers the curcumlactone core (cf. Scheme 4). Related dyotropic rearrangements have been used in total synthesis efforts, with the most closely related being that by Fuchs in his strategy to cephalostatin analogues.²⁶ The TMSOTf-promoted rearrangement of (–)-**5** to (+)-**4** is predicted to be a concerted dyotropic rearrangement (Figure 4).²⁷ The barrier for this rearrangement is approximately 7 kcal/mol higher than that predicted previously for the analogous rearrangement of **16a**,²⁰ reflecting the different propensities of ester and acyl groups for migration. In addition, the transition state structure shown in Figure 4 also has shorter partial bonds to the migrating groups and a longer central C–C bond (1.48 vs 1.39 Å) than found for **16a**.²⁰ Thus, in contrast to the rearrangement of **16a**, in which substantial charge separation occurs (the transition state structure for this process resembles an acyl cation and a TMS-complexed carboxylate migrating across a partial double bond, cf. Figure 3, **20**), the rearrangement shown in Figure 4 more closely resembles a “double $\text{S}_\text{N}2$ ” process, in which a lone pair on each migrating oxygen attacks the backside of the other C–O bond. This picture is borne out by computed electrostatic potential maps for the structures in Figure 4. As shown in Figure 5, charge separation actually appears to decrease slightly as the transition state structure is reached. Silylation of the δ -lactone carbonyl is also possible and is actually favored by approximately 8 kcal/mol. Silylation is associated with an increase in double bond





Figure 5. Electrostatic potential surfaces for the stationary points shown in Figure 4 (blue is most positive; charge range: 6.0×10^{-2} to 14.5×10^{-2} ; isovalue: 0.0200).

character for the O–C bond within the δ -lactone ring and is accompanied by greater angle strain for the smaller β -lactone ring. However, the transition state structure for rearrangement of the resulting species is predicted to be approximately 5 kcal/mol higher in energy than that shown in Figure 4, and there appears to be greater strain associated with the former transition state structure, in contrast to the situation for the corresponding reactants.²⁸ If the β -lactone-complexed and δ -lactone-complexed reactants can interconvert under the reaction conditions, which seems plausible, then the Curtin–Hammett principle²⁹ is expected to apply, and the transition state structure in Figure 3 is expected to correspond to the predominant route to product.


Final Stages of the Synthesis of (–)-Curcumanolide A (1) and (–)-Curcumlactone (3). We recognized that the reactivity of the two γ -lactones present in the bis-lactone (+)-**4** would require further differentiation to enable selective ring cleavage of the bridged γ -lactone. We thus chose to introduce the required β,β' -dimethyl enoate at an early stage to further differentiate the reactivity of these two γ -lactones. Thus, aldol condensation with acetone in a two stage, one-pot reaction provided the desired enoate (+)-**25** in 65% yield (Scheme 5). In efforts to directly form the isopropylidene precursor **27** by nucleophilic addition to the more strained, bridged, γ -lactone over the less electrophilic α,β -unsaturated lactone, we first tested a variety of conditions to generate tertiary alcohol **27** via methylolithium or methyl Grignard addition. In early studies with racemic bis-lactone (\pm)-**25**, treatment with 1.1 equiv MeLi

Scheme 5. Aldol Addition/Elimination and Selective Ring-Opening of Bis-lactone (+)-25^a



^aLHMDS, lithium bis(trimethylsilyl)amide; MsCl, methanesulfonyl chloride.

Scheme 6. Conversion of Methyl Ester (-)-28 to the Common Core (-)-31 and Synthesis of (-)-Curcumanolide A (1)^a




^aInset: ORTEP representation of the X-ray crystal structure of spirolactone (-)-30; hydrogens removed for clarity; thermal ellipsoids shown at 50% probability.³²

did not deliver tertiary alcohol 27 but rather yielded dieny tetrahydrofuran 26, presumably formed by 1,2-addition and subsequent alcohol elimination during the mild acid work-up. Reaction with Grignard reagents (MeMgI or MeMgBr) gave a mixture of recovered starting material and both mono- and bis-addition products as judged by analysis of crude ¹H NMR. As an alternative, we next considered ring cleavage to a methyl ester for subsequent conversion to the tertiary alcohol. Employment of standard methanolysis conditions (NaOMe, MeOH) provided some desired ester, but also the diester from cleavage of both γ -lactones. However, use of Et₃N in MeOH under reflux conditions gave the desired hydroxy ester (-)-28 in modest yield with only trace amounts of the diester, and finally a decrease in reaction temperature to 50 °C provided exclusively the desired product in serviceable yield (51%) and good recovery of starting material (34%).³⁰ Longer reaction times and increased amounts of Et₃N led to similar results, suggestive of reversible ring-opening of the two γ -lactones under these conditions.

To install the C4-methyl-bearing stereocenter, we envisioned dehydration of the secondary alcohol of (-)-28 at C4 followed by a facially selective hydrogenation of the more sterically accessible face syn to the C–O bond of the spirolactone (Scheme 6). A number of standard alcohol elimination products were screened at various temperatures including SOCl₂/Et₃N, SOCl₂/pyridine, MsCl/Et₃N, MsCl/Et₃N/DBU, or MsCl/DMAPI/pyridine and performed well on small scale;


Table 3. Conditions Tested for the Diastereoselective, Substrate-Directed Reduction of Enolate (-)-31 to (-)-33



entry	conditions	% yield (dr, 33:33')
1	5 mol % PtO ₂ , EtOH, 35 bar H ₂ , 48 h ^a	55 (<1:19) ^b
2	CuBr, Red-Al, THF, 0 °C, 2 h	ND (3:1) ^c
3	L-Selectride, THF, -78 °C → 0 °C, 1.5 h	83 (>19:1) ^b

^aUse of EtOAc as solvent gave similar results. ^bIsolated yield after column chromatography. ^cRatio determined by ¹H NMR analysis of crude reaction mixture. Yield not determined (ND).

Scheme 7. Dehydration of Tertiary Alcohol 33 Leading to (-)-Curcumlactone (3)




however, these reagents were inferior, leading to low conversion on scale-up. Ultimately, Burgess' reagent (29)³¹ in refluxing benzene was found to give high yields (84%) of an inseparable *exo/endo* mixture (2.4:1) of olefinic products. Several conditions were screened for the required olefin reduction, including diimide reduction, and hydrogenation with Pd/C and Wilkinson's catalyst (Rh(PPh₃)₃Cl); however, these conditions gave low conversion. Ultimately, hydrogenation with 5 mol % PtO₂ led to complete reduction at 1 atm H₂ after 12 h with good facial selectivity (83% yield, ~9:1 dr). The stereochemistry of the major diastereomer (-)-30 was confirmed by X-ray analysis supporting the anticipated facial selectivity providing the desired C4 stereochemistry. The minor diastereomer obtained during this hydrogenation possesses the relative stereochemistry found in curcumanolide B.

We next turned our attention to conversion of the methyl ester to an isopropylidene group (Scheme 6). Selective addition of Grignard reagents to the methyl ester over the spirolactone was initially studied with MeMgI and MeMgBr under a variety of conditions but all led to complex mixtures of products that included monoaddition to the ester, and both 1,4- and 1,2-addition to the spirolactone. The use of CeCl₃ as an additive³³ and subsequently CeCl₃/LiCl³⁴ had profoundly beneficial effects, and ultimately the use of MeLi rather than MeMgI or MeMgBr³⁵ led to efficient conversion to the desired tertiary alcohol (-)-31 in 79% yield. The final dehydration toward (-)-curcumanolide A was accomplished by employing Burgess' reagent (29) once again under conditions similar to those previously reported by Kato,^{6b} providing a 93% yield of a ~3:1 mixture of curcumanolide A (1, major product) and the corresponding exocyclic olefin regioisomer 32 (not shown). Partial separation of this mixture could be achieved by column chromatography providing pure (-)-curcumanolide A, and characterization data including optical rotation correlated well with previously reported data.

In strategies toward curcumlactone, a key challenge posed in setting the stereochemistry at C7 (cf. 33 in Table 3). Previous synthetic studies typically led to the incorrect diastereomer at

Scheme 8. ^1H NMR Data (300 MHz) Supporting Intramolecular Protonation during Reduction with L-Selectride: (a) ^1H NMR Expansion (δ 1.7–3.15) of Saturated Lactone **33** for Comparison and Same Expansion Following Initial Deuterium Exchange of Tertiary Alcohol with CD_3OD for (b) 17 h and (c) 4 h, Leading to Deuterated **33** (D-**33**) in Different Ratios^a



^aRatio of D incorporation based on integration: (b) ~50% and (c) ~30% D. Diastereotopic C6-protons going from two doublet of doublets to two doublets due to deuteration at C7 are marked by asterisks.

C7 through reduction of related enoate intermediates or direct alkylation leading primarily to C7-*epi*-curcumalactone. Thus, a subsequent epimerization was required to achieve, at best, a mixture of diastereomers.^{6b} Given the presence of the tertiary alcohol in intermediate **31**, we considered two approaches to set the C7-stereochemistry. The first involved a possible hydroxyl-directed hydrogenation³⁶ of the tetra-substituted olefin, a process not without precedent.³⁷ However, the tetrasubstituted and electron-poor olefin in **31** was recalcitrant to hydrogenation with multiple catalysts under various conditions (Pd/C, Pd(OH)₂/C, Pd/BaSO₄,^{37a} and Crabtree's catalyst) including high pressure (up to 35 bar H₂). Only when employing PtO₂ at 35 bar H₂ was reduction possible, providing 55% yield with 41% recovered starting material (Table 3, entry 1). This led to high diastereoselectivity for **33'** based on comparison with previously described C7-*epi*-curcumalactone,^{6b} obtained following alcohol elimination of **33'**, indicating that the hydroxyl group was not directing the hydrogenation. A second strategy would make use of the tertiary alcohol in a different capacity. We considered the possibility of a 1,4-conjugate reduction followed by intramolecular protonation of the intermediate enolate by the tertiary alcohol in a facially selective fashion to deliver curcumalactone (cf. **34** in Table 3). Initially, reaction with the copper-hydride species generated in situ from CuBr and Red-Al³⁸ led to an ~3:1 mixture of **33/33'**, indicating that some intramolecular protonation had likely occurred, but also some possible coordination of the tertiary

alcohol with the cuprate species as described previously in the literature.³⁹ Ultimately, we discovered that 1,4-reduction with L-selectride in THF at –78 °C followed by warming to 0 °C and quenching at that temperature indeed led to the desired diastereomer **33** in high diastereoselectivity and yield (dr >19:1, 83%).

The synthesis of (–)-curcumalactone was completed by dehydration employing SOCl₂ following the method of Kato^{6b} to deliver an ~6.2:1 mixture of *endo/exo* olefin regioisomers in 57% yield (Scheme 7). This elimination could also be performed with Burgess' reagent in higher yields with slightly decreased ratios of curcumalactone and the corresponding *endo* regioisomer (92%, ~5.5:1 **3/35**).^{6b} Purification by column chromatography delivered pure (–)-curcumalactone and both spectral and optical rotation data correlated well with reported values.⁶

The high diastereoselectivity observed from conjugate reduction of enoate **31** with L-Selectride is noteworthy and points to the likelihood of an *intramolecular* protonation of an intermediate boron enolate since the diastereomeric product **33'** would be expected from an *intermolecular* protonation of the enolate **34** (Table 3). We designed an experiment to probe the hypothesis that intramolecular protonation by the adjacent tertiary alcohol (i.e. via enolate **34**, Table 3) was indeed occurring following conjugate reduction. Intramolecular protonation of enolates by alcohols has been described in the literature,⁴⁰ and the preferred conformation of the spirocyclic

system (i.e., 31) appeared conducive for such an intramolecular protonation (see 34, Table 3). We reasoned that alcohol OH \rightarrow OD exchange with excess CD₃OD, followed by azeotropic drying with xylenes and reduction with L-selectride, would lead to deuterated species D-33 (Scheme 8). This process gave varying amounts of D-33 depending on the time allowed for H \rightarrow D exchange (4 vs 17 h, Scheme 8b,c); however, high diastereoselectivity was consistently observed, suggestive of an intramolecular protonation/deuteration. Incorporation of deuterium at C7 is evidenced by simplification of the diastereotopic C6-protons (indicated by *) from two sets of doublet of doublets (dd, $J = 10.5, 13.2$) to a doublet (d, $J = 13.2$) centered at δ 2.47, due to exchange of a vicinal proton at C7 for deuterium. Similar simplification is also observed with the diastereotopic C6' proton centered at δ 1.77 but is obscured by other protons. Further confirmation of deuterium incorporation came from HRMS analysis of the mixture of 33/D-33.²⁸ The inability to achieve complete deuterium incorporation is not easily explained, given the high facial selectivity that excludes intermolecular processes, but the simplest explanation is incomplete deuterium exchange of the tertiary alcohol. Importantly, this sequence provides a highly stereocontrolled introduction of the C7 stereocenter, a challenge encountered in the previous synthesis of curcumlactone.^{6b}

CONCLUSIONS

In summary, we provided a full account of our recently disclosed stereospecific dyotropic process to spirocyclic, bridged γ -butyrolactones via Lewis-acid-mediated 1,2-acyl and δ -lactone migrations. By employing these methods, concise asymmetric routes to the spirocyclic sesquiterpene natural products (–)-curcumanolide A and (–)-curcumlactone have been accomplished through a key dyotropic process of a tricyclic β,δ -bis-lactone, which establishes the spirocyclic stereocenter. DFT calculations on this dyotropic rearrangement suggest a double ‘S_N2-like’ process of the two migrating carboxylates with a diminution in charge separation as the transition state structure is reached. The synthesis demonstrates the utility of our catalytic, asymmetric NCAL process and the ability to perform a Baeyer–Villiger oxidation in the presence of a β -lactone. A highly diastereoselective 1,4-reduction involving intramolecular protonation by a resident tertiary alcohol established the final stereocenter found in curcumlactone. The described enantioselective synthesis of curcumanolide A and curcumlactone was accomplished in 11 and 12 steps, respectively, from 2-methyl-1,3-cyclopentanedione and did not require the use of protecting groups. The synthetic strategy described herein provides efficient routes to both natural products, enabling further biological studies of these potentially bioactive spirocyclic lactones.

ASSOCIATED CONTENT

Supporting Information

Complete ref 27d. Complete experimental and characterization details for all new compounds reported, along with additional details on calculations. Crystallographic files. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

romo@tamu.edu

Present Addresses

[§]Teva North America, 1090 Horsham Rd., North Wales, PA 19454

[†]Department of Molecular Microbiology and Immunity, Oregon Health and Science University, 3181 S.W. Sam Jackson Park Rd., Portland, OR 97239

[¶]Pfizer, Inc., 558 Eastern Point Rd., Groton, CT 06340

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We are grateful for generous support from the National Science Foundation (CHE-0809747 to D.R., CHE-0755207 REU supporting A.G.J., CHE-0449845 and CHE-030089 (Pittsburgh Supercomputer Center) to D.J.T., and the Welch Foundation (A-1280 to D.R.). We thank Dr. Joe Reibenspies for the X-ray analyses and Dr. Arjun Raghuraman for technical assistance.

REFERENCES

- (1) Shiobara, Y.; Asakawa, Y.; Kodama, M.; Yasuda, K.; Takemoto, T. *Phytochemistry* **1985**, *24*, 2629.
- (2) Matthes, H. W. D.; Luu, B.; Ourisson, G. *Phytochemistry* **1980**, *19*, 2643.
- (3) Jiehan, H.; Xiuwen, H.; Tao, J.; Zhenyun, Y.; Xiuying, W.; Jihong, X.; Yongtian, G. *Kexue Tongbao* **1987**, *32*, 816.
- (4) (a) Inayama, S.; Gao, J.-F.; Hariyama, K.; Hikichi, M.; Iitaka, Y.; Guo, Y.-T.; Kawamata, T. *Chem. Pharm. Bull.* **1985**, *33*, 2179. (b) Gao, J.-F.; Ohkura, T.; Harimaya, K.; Hikichi, M.; Kawamata, T.; Ying, W.-X.; Iitaka, Y.; Inayama, S. *Chem. Pharm. Bull.* **1986**, *34*, 5122.
- (5) See ref 3 and references cited therein.
- (6) (a) Hirukawa, T.; Oguchi, M.; Yoshikawa, N.; Kato, T. *Chem. Lett.* **1992**, 2343. (b) Kato, T.; Mutoh, M.; Oguchi, M.; Yasuoka, H. *Sci. Iran.* **1997**, *4*, 94.
- (7) Honda, T.; Ishige, H. *J. Chem. Soc., Perkin Trans. 1* **1994**, 3567.
- (8) Fujita, T.; Tanaka, M.; Norimine, Y.; Suemune, H. *J. Org. Chem.* **1997**, *62*, 3824.
- (9) (a) Reetz, M. T. *Angew. Chem., Int. Ed. Engl.* **1972**, *11*, 129. (b) Reetz, M. T. *Angew. Chem., Int. Ed. Engl.* **1972**, *11*, 130. (c) Reetz, M. T. *Tetrahedron* **1973**, *29*, 2189. (d) Reetz, M. T. *Adv. Organomet. Chem.* **1977**, *16*, 33. (e) Hoffmann, R.; Williams, J. E., Jr. *Helv. Chim. Acta* **1972**, *55*, 67.
- (10) For a review and leading references, see: (a) Fernández, I.; Cossio, F. P.; Sierra, M. A. *Chem. Rev.* **2009**, *109*, 6687–6711. (b) Davis, R. L.; Tantillo, D. J. *J. Org. Chem.* **2010**, *75*, 1693–1700.
- (11) (a) Grob, C. A.; Winstein, S. *Helv. Chim. Acta* **1952**, *35*, 782–802. (b) Barton, D. H. R.; Head, A. J. *J. Chem. Soc.* **1956**, 932.
- (12) (a) Lin, S.; Danishefsky, S. J. *Angew. Chem., Int. Ed.* **2002**, *41*, 512. (b) Zhang, X.; Houk, K. N.; Lin, S.; Danishefsky, S. J. *J. Am. Chem. Soc.* **2003**, *125*, 5111.
- (13) Kocienski, P.; Barber, C. *Pure Appl. Chem.* **1990**, *62*, 1933.
- (14) Stöckner, F.; Käßlinger, C.; Beckert, R.; Görls, H. *Synlett* **2005**, 643.
- (15) Mulzer, J.; Brüntrup, G. *Angew. Chem., Int. Ed. Engl.* **1979**, *18*, 793.
- (16) Reetz, M. T.; Schmitz, A.; Holdgrün, X. *Tetrahedron Lett.* **1989**, *30*, 5421.
- (17) Arrastia, I.; Lecea, B.; Cossio, F. P. *Tetrahedron Lett.* **1996**, *37*, 245–248.
- (18) (a) Black, T. H.; DuBay, W. J., III. *Tetrahedron Lett.* **1988**, *29*, 1747. (b) Black, T. H.; McDermott, T. S. *J. Chem. Soc., Chem Commun.* **1991**, 184.
- (19) Purohit, V. C.; Matla, A. S.; Romo, D. *J. Am. Chem. Soc.* **2008**, *130*, 10478.
- (20) Davis, R. L.; Leverett, C. A.; Romo, D.; Tantillo, D. J. *J. Org. Chem.* **2011**, *76*, 7167.
- (21) Leverett, C. A.; Purohit, V. C.; Romo, D. *Angew. Chem., Int. Ed.* **2010**, *49*, 9479.

- (22) The scope of this concise approach to dione acids will be disclosed in a separate report.
- (23) Birman, V. B.; Li, X. *Org. Lett.* **2006**, *8*, 1351.
- (24) (a) Birman, V. B.; Li, X. *Org. Lett.* **2008**, *6*, 1115. (b) Padakanti, P. K. Ph.D. Thesis, Washington University: St. Louis, MO, USA, 2010.
- (25) CCDC 875167 (15) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- (26) (a) Li, W.; LaCour, T. G.; Fuchs, P. L. *J. Am. Chem. Soc.* **2002**, *124*, 4548. For selected dyotropic rearrangements in total synthesis efforts, see: (b) Zhang, X.; Houk, K. N.; Lin, S.; Danishefsky, S. J. *J. Am. Chem. Soc.* **2003**, *125*, 5111. (c) Denmark, S. E.; Montgomery, J. I. *Angew. Chem., Int. Ed.* **2005**, *44*, 3732. For additional examples, see ref 10a.
- (27) Calculations were performed at the B3LYP/6-31+G(d,p) level of theory: (a) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648. (b) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 1372. (c) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785. (d) Stephens, P.; Devlin, F. J.; Chabalowski, C. F.; Frisch, M. J. *J. Phys. Chem.* **1994**, *98*, 11623 using GAUSSIAN03 (complete reference in Supporting Information). (e) All structures were characterized by frequency analysis, and reported energies from B3LYP/6-31G+(d,p) calculations include zero-point energy (ZPE) corrections. Free energies at 25 °C are also reported. Intrinsic reaction coordinate (IRC) calculations were also used to characterize the transition state structure shown (see Supporting Information for plot). (f) Gonzalez, C.; Schlegel, H. B. *J. Phys. Chem.* **1990**, *94*, 5523. (g) Fukui, K. *Acc. Chem. Res.* **1981**, *14*, 363.
- (28) See Supporting Information for further details.
- (29) For leading references, see: Seeman, J. I. *J. Chem. Educ.* **1986**, *63*, 42.
- (30) Henry-Riyad, H.; Lee, C.; Purohit, V. C.; Romo, D. *Org. Lett.* **2006**, *8*, 4363.
- (31) Burgess, E. M.; Penton, H. R., Jr.; Taylor, E. A. *J. Org. Chem.* **1973**, *38*, 26.
- (32) CCDC 842815 (30) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- (33) (a) Imamoto, T.; Takiyama, N.; Nakamura, K.; Hatajima, T.; Kamiya, Y. *J. Am. Chem. Soc.* **1989**, *111*, 4392. (b) Conlon, D. A.; Kumke, D.; Moeder, C.; Hardiman, M.; Hutson, G.; Sailer. *Adv. Synth. Catal.* **2004**, *346*, 1307.
- (34) Krasovsky, A.; Kopp, F.; Knochel, P. *Angew Chem. Int. Ed.* **2006**, *45*, 497.
- (35) (a) Dunn, T. B.; Ellis, J. M.; Kofink, C. C.; Manning, J. R.; Overman, L. E. *Org. Lett.* **2009**, *11*, 5658. (b) Martin, C. L.; Overman, L. E.; Rohde, J. M. *J. Am. Chem. Soc.* **2010**, *132*, 4894.
- (36) Hoveyda, A. H.; Evans, D. A.; Fu, G. C. *Chem. Rev.* **1993**, *93*, 1307.
- (37) (a) Nishi, T.; Kataoka, M.; Morisawa, Y. *Chem. Lett.* **1989**, 1993. (b) Hanessian, S.; Abad-Grillo, T.; McNaughton-Smith, G. *Tetrahedron* **1997**, *53*, 6281. (c) Lee, N.; Kim, Y.-W.; Chang, K.; Kim, K. H.; Jew, S.-S.; Kim, D.-K. *Tetrahedron Lett.* **1996**, *37*, 2429.
- (38) Semmelhack, M. F.; Stauffer, R. D.; Yamashita, A. *J. Org. Chem.* **1977**, *42*, 3180.
- (39) Yang, J.; Long, Y. O.; Paquette, L. A. *J. Am. Chem. Soc.* **2003**, *125*, 1567.
- (40) For representative examples, see: (a) Diba, A. K.; Noll, C.; Richter, M.; Gieseler, M. T.; Kalesse, M. *Angew Chem. Int. Ed.* **2010**, *49*, 8367. (b) Jousseau, T.; Wurz, N. E.; Glorius, F. *Angew Chem. Int. Ed.* **2011**, *50*, 1410. (c) Kanoh, N.; Sakanishi, K.; Limori, E.; Nishimura, K.; Iwabuchi, Y. *Org. Lett.* **2011**, *13*, 2864.