pubs.acs.org/joc

Metal- and Photocatalyst-Free 23 Examples

# Metal-Free Trifluoromethylthiolation of Arylazo Sulfones

Ankun Li,<sup> $\perp$ </sup> Yuxuan Li,<sup> $\perp$ </sup> Junjie Liu, Jingqi Chen, Kui Lu, Di Qiu, Maurizio Fagnoni, Stefano Protti,\* and Xia Zhao\*



Downloaded via AUCKLAND UNIV OF TECHNOLOGY on December 22, 2020 at 17:32:38 (UTC) See https://pubs.acs.org/sharingguidelines for options on how to legitimately share published articles.

he formation of an Ar-SCF<sub>3</sub> bond is important in life sciences for the synthesis of bioactive molecules. In fact the SCF<sub>3</sub> group, when present in an aromatic compound, strongly affected its physicochemical properties, mostly its lipophilicity. The latter is a key parameter in drug design since the bioavailability of the drug is enhanced when lipophilicity increases.<sup>1</sup> In this case, a combined effect of the presence of fluorine atoms<sup>2</sup> with heteroatoms imparts a good lipophilicity as witnessed by the high values of the Hansch parameters ( $\pi_{\rm R}$ = 1.44).<sup>3</sup> Some drugs, such as the antiprotozoal agent Toltrazuril and the stimulant amphetamine Tiflorex, contain the Ar-SCF<sub>3</sub> moiety. Accordingly, a more reliable synthetic procedure in the forging of the Ar-S bond could increase a wider application of trifluoromethylthiolated aromatics in medicinal chemistry. In the last years, several synthetic protocols for the introduction of a SCF<sub>3</sub> group in a (hetero)aromatic core have been developed.<sup>4</sup> An interesting approach is the direct trifluoromethylthiolation of (hetero)arenes, but only certain electron-rich derivatives led to the desired product in a clean fashion, avoiding the concomitant formation of undesired regioisomers (Scheme 1a).<sup>5</sup> Thus, several approaches have been developed by replacing an aromatic substituent with a SCF3 group making use of a metal catalyst (e.g., Cu,<sup>6</sup> Ni,<sup>7</sup> Pd,<sup>8</sup> and Ag<sup>9</sup>). The reaction is, in most cases, an ipso-substitution that starts from aryl diazonium salts,  $^{6a,d,h}$  aryl halides,  $^{6c,7a,c,d,9}$  aryl boronic acids,  $^{6b-g}$  aryltrifluoroborates,<sup>6e</sup> di(hetero)aryl- $\lambda^3$ -iodanes,<sup>6f</sup> aryl sulfonates,<sup>7b</sup> and arylmercaptodifluoroacetic acids (Scheme 1b).<sup>8a</sup> Metalfree alternatives, however, are limited to the alkylation of aryl sulfides by using CF<sub>3</sub>I under basic conditions<sup>10</sup> (Scheme 1c) except the trifluoromethylation of diaryl sulfides with trifluoromethyltriisopropylsilane  $(TIPSCF_3)^{11}$  and the radical

# Scheme 1

a) Ar-H + "SCF<sub>3</sub> source"  $\xrightarrow{\Delta}$  Ar-SCF<sub>3</sub>

b) Ar-Y + "SCF<sub>3</sub> source" 
$$\xrightarrow{Cu, Ni, Pd} \Delta$$
 Ar-SCF<sub>3</sub>  
Y = N<sub>2</sub><sup>\*</sup>, I, Br, CI, B(OH)<sub>2</sub>, BF<sub>3</sub>K,  
IAr(OTf), OSO<sub>2</sub>R, SCF<sub>2</sub>COOH

c) Metal-free trifluoromethylation of sulfides

$$Ar-SH + CF_3I \xrightarrow{Base}{\Delta} Ar-SCF_3$$

d) Trifluoromethylthiolation of arene diazonium salts

Ar-N<sub>2</sub><sup>+</sup> 
$$\xrightarrow{PC}$$
 Ar<sup>-</sup>  $\xrightarrow{Ar'SO_2SCF_3 \text{ or}}$  Ar-SCF<sub>3</sub>  
CF<sub>3</sub>S-SCF<sub>3</sub> Ar-SCF<sub>3</sub>

e) This work

Ar-N<sub>2</sub>SO<sub>2</sub>Me 
$$\xrightarrow{hv}$$
 Ar  $\xrightarrow{Ar'SO_2SCF_3}$  Ar-SCF<sub>3</sub>

trifluoromethylthiolation of arenediazonium salts with  $Me_4NSCF_3$ .<sup>12</sup> In recent years, a handful of examples were reported where photochemistry was applied for the synthesis of ArSCF<sub>3</sub> thanks to the easy photogeneration of aryl radicals

Received: November 9, 2020

from aryl diazonium salts and ensuing reaction with a *S*-trifluoromethyl arylsulfonothioate<sup>13 a,b</sup> or 1,2-bis-(trifluoromethyl)disulfane  $(CF_3S)_2$  (Scheme 1d).<sup>13c</sup> Unfortunately, the latter approach requires the presence of a photocatalyst (PC) and a sensitive aryl radical precursor complicated by the special caution required on the use and storage of arenediazonium salts.<sup>14</sup>

We then envisioned that a straightforward route to  $Ar-SCF_3$  could involve again aryl radicals but was generated by visiblelight irradiation of arylazo sulfones ( $ArN_2SO_2Me$ , Scheme 1e). Such derivatives exhibit a wavelength selective behavior<sup>15</sup> and release an aryl radical upon visible-light irradiation<sup>16</sup> by cleavage of the N-S bond followed by nitrogen loss of the resulting aryl diazenyl radical.<sup>17</sup> The thus generated aryl radicals were recently applied for the forging of various Ar-C and Ar-heteroatom bonds.<sup>18</sup> In view of these premises, we planned to use such sulfones for the easy arylation of a  $SCF_3$ -containing derivative to form the desired  $Ar-SCF_3$ . As the trifluoromethylthiolating agent, we exclude the use of ( $CF_3S$ )<sub>2</sub> due to its difficult handling,<sup>13c</sup> focusing our attention on *S*-trifluoromethyl arylsulfonothioates.

Preliminary experiments were carried out on compound 1a, and the obtained results are summarized in Table 1. Irradiation





of 1a (0.125 M, in 1,2-dichloroethane, DCE) in the presence of two equivalents of S-(trifluoromethyl) 4-difluorobenzenesulfonothioate (2a) by means of a 21 W green LED afforded the arylated compound 3a in low yields (21%, entry 1). Moving to blue light (entries 2 and 3) resulted in an improvement of the efficiency of the process, and a 53% yield was reached with a 21W blue LED lamp. The use of a different arylazo sulfone (1a', entry 4) as well as the adoption of either electron-rich (2b) or difluorinated (2c) benzenesulfonothioates (entries 5 and 6) did not afford better results, while, among the different media tested (entries 7-9), DCE furnished the most satisfactory performance. Hydrodeaminated biphenyl can be competitively formed as the main product by changing the reaction medium (e.g., MeCN). Gratifyingly, when doubling the concentration of the reactants, **3a** was isolated in 63% yield (entry 10), and this yield slightly decreased by using a 0.5 M amount of **1a** (entry 11). Notably, increasing the loading of the trifluoromethylthiolating agent **2a** from 2.0 equiv to 4.0 equiv improved the yield up to 75% (entry 12). The formation of **3a** was completely inhibited in the absence of light (entry 13). Dedicated on–off experiments confirmed that the reaction did not proceed in the dark (see Figure S1).

The conditions described in entry 12 have been thus adopted to investigate the scope of the synthetic protocol.

Compound 3a was also synthesized on a gram scale and isolated in 45% yield (see Table 2). We then investigated the

#### Table 2. Trifluoromethylthiolation of Arylazo Sulfones<sup>a</sup>



<sup>*a*</sup>A solution of 1a-x in DCE in the presence of 2a (4 equiv) irradiated for 36 h by means of a 21 W Blue Light LED. <sup>*b*</sup>Reaction carried out on 10 mmol 1a, 48 h irradiation (gram scale = 1.15 g). <sup>*c*</sup>2 equiv of 2aemployed. <sup>*d*</sup>Quinoline (30% yield) was observed as the byproduct via GC-MS analyses. <sup>*c*</sup>Benzothiazole (24% yield) was observed as the byproduct via GC-MS analyses.

scope of the protocol, as shown in Table 2. The desired trifluoromethylthiolated products have been isolated in discrete to satisfactory yields, and the process showed a good tolerance to both electron-donating and electron-withdrawing substituents present on the aromatic ring, including (thio)-alkoxy groups (see products 3c-e, 3p), halogens (3i,n), and carbonyls (3k-m). In some cases, good results were obtained in the presence of only 2 equiv of 2a (as in the synthesis of 3f and 3l). The process was found to be also suitable to prepare naphthyl derivative 3q, whereas heteroaryl trifluoromethyl thioethers (3r-x) were mainly obtained in a lower yield. In the

latter cases, the hydrodeamination product (see the case of 3u and 3x) was detected as the main byproduct (up to 30% yield).

A tentative mechanism is shown in Scheme 2. Sulfones 1a-x possess a weak  $n\pi$  band between 400 and 500 nm ( $\epsilon$  ca. 500

Scheme 2. Proposed Mechanism for the Thiotrifluoromethylation of Arylazo Sulfones



 $M^{-1}$ ) sufficient to allow the absorption in the visible region (465–470 nm blue LEDs have been used as inexpensive light source).<sup>16</sup> A direct photolysis of **2a** is safely excluded since its absorption at the wavelength used is negligible (see Figure S2). The labile S–N bond underwent a smooth photocleavage (path a) liberating the desired aryl radical Ar<sup>•</sup> that, as previously described, <sup>13a,b</sup> gave a substitution reaction with **2a** (path b) to form derivatives **3a–x** along with the stable 4-fluorophenylsulfonyl radical. Hydrogen atom abstraction of Ar<sup>•</sup> from the solvent to form the hydrodeaminated product Ar–H (path c) is the main competitive path.<sup>18c</sup> The intermediacy of an aryl radical has been further confirmed by the formation of adduct **4k** that was isolated in 30% yield when the reaction of **1k** with **2a** was carried out in the presence of TEMPO (4 equiv, path d).

In summary, we proposed a radical trifluoromethylthiolation reaction for the formation of  $aryl-SCF_3$  bonds via simple visible-light activation of bench-stable arylazo sulfones under both metal-free and (photo)catalyst-free conditions. The process employed *S*-(trifluoromethyl) 4-difluorobenzenesulfonothioate as the trifluoromethylthiolating agent and was found suitable for the preparation of both electron-rich and electronpoor SCF<sub>3</sub>-substituted aromatics in discrete to satisfactory yields. In analogy with other metal-free<sup>12</sup> or photocatalyzed<sup>13a-c</sup> trifluoromethylthiolation protocols, a low efficiency in the synthesis of heteroaryl trifluoromethyl thioethers was observed, with the only exception of benzofuran **3r**.

#### EXPERIMENTAL SECTION

**General Remarks.** All solvents were distilled prior to use. For chromatography, 200–300 mesh silica gel (Qingdao, China) was employed. <sup>1</sup>H, <sup>13</sup>C, and <sup>19</sup>F NMR spectra were recorded at 400, 100 and 375 MHz with a Bruker ARX 400 spectrometer. Chemical shifts are reported in ppm using tetramethylsilane as an internal standard. HRMS was performed on an Thermo Scientific LTQ Orbitrap XL

(ion trap) or Bruker Solarix XR FTMS (Q-TOF) mass instrument. All arylazo sulfones were prepared from the corresponding arenediazonium tetrafluoroborates according to literature procedures.<sup>18e-g</sup> The light-promoted reactions were carried out by using a standard blue LED lamp with 20 one-light-emitting diodes (12–28 V, 21 W, 465–470 nm). The distance from the light source to the irradiation vessel was 3 cm.

Typical Procedure for the Preparation of Arenediazonium Tetrafluoroborates. *Method A*. In a 100 mL round-bottom flask, the aromatic amine (20 mmol) was dissolved in a mixture of  $H_2O$  (10 mL) and HBF<sub>4</sub> (48% aq, 6 mL). After the mixture was stirred for 15 min, a solution of NaNO<sub>2</sub> (1.5 g, 1.1 equiv, in 4 mL of  $H_2O$ ) was added dropwise at 0 °C (in an ice bath). The mixture was stirred for another 30 min at 0 °C. Then, the arenediazonium tetrafluoroborate was removed by filtration and washed with diethyl ether (2 × 10 mL). The crude product was dried in vacuo for 20 min and was then directly used without further purification.

Method B. In a 100 mL round-bottom flask, the aromatic amine (20 mmol) was dissolved in a mixture of ethanol (8 mL) and HBF<sub>4</sub> (48% aq, 6 mL). Subsequently, *tert*-butyl nitrite (4.7 mL, 2.0 equiv) was added dropwise to the solution at 0 °C. The reaction mixture was stirred for 30 min at 0 °C, and anhydrous diethyl ether (20 mL) was added to precipitate the corresponding arenediazonium tetrafluoroborate. The solid was filtered off and washed with diethyl ether (2 × 10 mL). The product was then dried in vacuo for 20 min and used without further purification.

Typical Procedure for the Preparation for Arylazo Sulfones. In a 100 mL round-bottom flask, the suspension of the freshly prepared arenediazonium tetrafluoroborate (20 mmol) in  $CH_2Cl_2$  (20 mL) was stirred at 0 °C; then sodium methanesulfinate (2.24 g, 1.1 equiv) was added to the reaction system in one portion. The temperature was risen to rt; the mixture was stirred overnight and then filtered, and the obtained solution evaporated. The crude product was thus purified by dissolution in  $CH_2Cl_2$  and precipitation by adding cold petroleum ether. The obtained arylazo sulfones were filtered and dried in vacuo for 30 min.

Characterization Data for the Arylazo Sulfones 1a–t. 1-([1,1'-Biphenyl]-4-yl)-2-(methylsulfonyl)diazene (1a).<sup>18d</sup> Yellow solid (3.6 g, 69%). Mp (dec): 107–109 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.03 (d, J = 8.6 Hz, 2H), 7.80 (d, J = 8.6 Hz, 2H), 7.68– 7.66 (m, 2H), 7.52–7.48 (m, 2H), 7.46–7.42 (m, 1H), 3.24 (s, 3H).

1-([1,1'-Biphenyl]-4-yl)-2-((4-nitrophenyl)sulfonyl)diazene (1*a*'). Yellow solid (4.0 g, 54%). Mp (dec): 100–101 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.45 (d, *J* = 8.88 Hz, 2H), 8.19 (d, *J* = 8.92 Hz, 2H), 7.89 (d, *J* = 8.64 Hz, 2H), 7.75 (d, *J* = 8.68 Hz, 2H), 7.64–7.62 (m, 2H), 7.51–7.41 (m, 3H). <sup>13</sup>C {<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 151.3, 148.7, 147.9, 139.4, 138.9, 131.8, 129.1, 129.0, 128.2, 127.3, 125.5, 124.1. IR (neat, *v* cm<sup>-1</sup>): 3413, 1616, 1526, 1352, 1149, 1074, 952, 858, 620, 517. The compound was found to decompose upon HRMS analysis.

4-(4-((Methylsulfonyl)diazenyl)phenyl)morpholine (**1b**). Orange solid (2.1 g, 39%). Mp (dec): 139–140 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.86 (d, *J* = 9.2 Hz, 2H), 6.90 (d, *J* = 9.2 Hz, 2H), 3.86 (t, *J* = 4.8 Hz, 4H), 3.46 (t, *J* = 5.1 Hz, 4H), 3.14 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 155.7, 140.9, 127.8, 113.1, 66.3, 46.8, 35.0. IR (neat, *v* cm<sup>-1</sup>): 3413, 1607, 1379, 1318, 1234, 1141, 1067, 619, 533. HRMS (ESI) *m/z*:  $[M + H]^+$  calcd for C<sub>11</sub>H<sub>16</sub>N<sub>3</sub>O<sub>3</sub>S, 270.0907; found, 270.0904.

1-(Methylsulfonyl)-2-(4-phenoxyphenyl)diazene (1c). Brown solid (4.4 g, 80%). Mp (dec): 110–111 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.94 (d, J = 9.0 Hz, 2H), 7.47–7.43 (m, 2H), 7.27 (t, J = 7.4 Hz, 1H), 7.13–7.01 (m, 4H), 3.20 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 164.3, 154.5, 144.0, 130.2, 127.1, 125.4, 120.6, 117.7, 34.8. IR (neat, v cm<sup>-1</sup>): 3413, 1615, 1486, 1338, 1243, 1137, 949, 853, 773, 617, 502. HRMS (ESI) m/z: [M + H]<sup>+</sup> calcd for C<sub>13</sub>H<sub>13</sub>N<sub>2</sub>O<sub>3</sub>S, 277.0641; found, 277.0641.

(Benzyloxy)phenyl)-2-(methylsulfonyl)diazene (1d). Pale yellow solid (3.0 g, 52%). Mp (dec): 150-151 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.95 (d, J = 9.8 Hz, 2H), 7.45–7.36 (m, 5H), 7.11 (d, J = 9.1 Hz, 2H), 5.19 (s, 2H), 3.19 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz,

CDCl<sub>3</sub>):  $\delta$  164.8, 143.3, 135.5, 128.8, 128.5, 127.5, 127.3, 115.7, 70.7, 34.9. IR (neat,  $v \text{ cm}^{-1}$ ): 3413, 1614, 1485, 1338, 1137, 949, 852, 617, 502. HRMS (ESI) m/z: [M + H]<sup>+</sup>calcd for C<sub>14</sub>H<sub>15</sub>N<sub>2</sub>O<sub>3</sub>S, 291.0795; found, 291.0798.

1-(Methylsulfonyl)-2-(4-(methylthio)phenyl)diazene (1e). Yellow solid (2.9 g, 63%). Mp (dec): 130–132 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.87 (d, J = 8.8 Hz, 2H), 7.35 (d, J = 8.0 Hz, 2H) 3.20 (s, 3H), 2.58 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 150.0, 145.8, 125.4, 124.9, 34.8, 14.7. IR (neat, v cm<sup>-1</sup>): 3413, 1614, 1486, 1337, 1137, 949, 617, 502. HRMS (ESI) m/z: [M + H]<sup>+</sup> calcd for C<sub>8</sub>H<sub>11</sub>N<sub>2</sub>O<sub>2</sub>S<sub>2</sub>, 231.0257; found, 231.0256.

1-(4-(tert-Butyl)phenyl)-2-(methylsulfonyl)diazene (1f).<sup>18d</sup> Yellow solid (2.6 g, 55%). Mp (dec): 78–80 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.88 (d, J = 8.8 Hz, 2H), 7.59 (d, J = 8.8 Hz, 2H), 3.20 (s, 3H), 1.36 (s, 9H).

*Ethyl* 3-(4-((*E*)-(*Methylsulfonyl*)*diazenyl*)*phenyl*)*acrylate* (**1g**).<sup>18e</sup> Yellow solid (5.5 g, 98%). Mp (dec): 98–100 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.95(d, *J* = 8.5 Hz, 2H), 7.68–7.72 (m, 3H), 6.57 (d, *J* = 16.1 Hz, 1H), 4.28 (q, *J* = 7.1 Hz, 2H), 3.22 (s, 3H), 1.34 (t, *J* = 7.1 Hz, 3H).

1-(Methylsulfonyl)-2-(4-((triisopropylsilyl)ethynyl)phenyl)diazene (1h).<sup>18e</sup> Yellow solid (2.7 g, 37%). Mp (dec): 63–64 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.89 (d, J = 8.5 Hz, 2H), 7.65 (d, J = 8.5 Hz, 2H), 3.22 (s, 3H), 1.16–1.12 (m, 21H).

1-(4-Bromophenyl)-2-(methylsulfonyl)diazene (1i).<sup>18f</sup> Yellow solid (3.2 g, 60%). Mp (dec): 131–133 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.83 (d, J = 8.8 Hz, 2H), 7.74 (d, J = 8.8 Hz, 2H), 3.22 (s, 3H).

4-((Methylsulfonyl)diazenyl)benzonitrile (**1j**).<sup>18h</sup> Yellow solid (3.3 g, 79%). Mp (dec): 114–115 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.04 (d, J = 8.4 Hz, 2H), 7.89 (d, J = 8.4 Hz, 2H), 3.26 (s, 3H).

Ethyl 4-((Methylsulfonyl)diazenyl)benzoate (1k). <sup>18d</sup> Yellow solid (3.7 g, 73%). Mp (dec): 83–84 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.25(d, J = 8.6 Hz, 2H), 7.99 (d, J = 8.6 Hz, 2H), 4.44 (q, J = 7.1 Hz, 2H), 3.25 (s, 3H), 1.43 (t, J = 7.1 Hz, 3H).

1-(4-((Methylsulfonyl)diazenyl)phenyl)ethanone (11).<sup>18d</sup> Yellow solid (3.6 g, 80%). Mp (dec): 119–120 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.13 (d, J = 8.5 Hz, 2H), 8.00 (d, J = 8.5 Hz, 2H), 3.24 (s, 3H), 2.67 (s, 3H).

(4-((Methylsulfonyl)diazenyl)phenyl)(phenyl)methanone (1m). Yellow solid (4.3 g, 75%). Mp (dec): 130 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.01 (dd, *J* = 8.6, 31.4 Hz, 4H), 7.81 (d, *J* = 7.04 Hz, 2H), 7.67–7.62 (m, 1H), 7.52 (t, *J* = 7.88 Hz, 2H), 3.27 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  195.1, 150.6, 142.8, 136.4, 133.2, 131.0, 130.0, 128.5, 124.2, 34.8. IR (neat, *v* cm<sup>-1</sup>): 3413, 1659, 1617, 1340, 1276, 1159, 1069, 954, 861, 700. HRMS (ESI) *m*/*z*: [M + H]<sup>+</sup> calcd for C<sub>14</sub>H<sub>13</sub>N<sub>2</sub>O<sub>3</sub>S, 289.0638; found, 289.0641.

1-(3-Bromophenyl)-2-(methylsulfonyl)diazene (1n).<sup>18e</sup> Yellow solid (2.5 g, 47%). Mp (dec): 95–96 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.06 (s, 1H), 7.90 (d, J = 8.0 Hz, 1H), 7.77 (d, J = 8.0 Hz, 1H), 7.47 (t, J = 8.0 Hz, 1H), 3.22 (s, 3H).

1-(*Methylsulfonyl*)-2-(3,4,5-trimethoxyphenyl)diazene (**10**). Pale yellow solid (2.7 g, 49%). Mp (dec): 122–124 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.26 (s, 2H), 4.00 (s, 3H), 3.94 (s, 6H), 3.22 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 153.5, 144.4, 102.5, 61.1 (d, J = 6.4 Hz), 56.3 (d, J = 5.4 Hz), 34.9. IR (neat, v cm<sup>-1</sup>): 3413, 1614, 1486, 1338, 1137, 949, 617, 502. HRMS (ESI) m/z: [M + H]<sup>+</sup>calcd for C<sub>10</sub>H<sub>14</sub>N<sub>2</sub>O<sub>5</sub>S, 275.0696; found, 275.0696.

1-(Benzo[d][1,3]dioxol-5-yl)-2-(methylsulfonyl)diazene (1p). Dark green solid (0.91 g, 20%). Mp (dec): 113–114 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.69 (dd, J = 2.0, 8.2 Hz, 1H), 7.34 (d, J = 1.9, Hz, 1H), 6.99 (d, J = 8.2 Hz, 1H), 6.14 (s, 2H), 3.18 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  154.4, 149.5, 144.8, 129.3, 108.4, 102.8, 98.8, 34.9. IR (neat,  $v \text{ cm}^{-1}$ ): 3414, 1617, 1477, 1414, 1152, 618. HRMS (EI) m/z: [M + H]<sup>+</sup> calcd for C<sub>8</sub>H<sub>9</sub>N<sub>2</sub>O<sub>4</sub>S, 229.0278; found, 229.0281.

1-(Methylsulfonyl)-2-(naphthalen-2-yl)diazene (**1q**).<sup>18e</sup> Yellow solid (2.2 g, 47%). Mp (dec): 108–109 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.56 (s, 1H), 8.03–8.01 (m, 1H), 7.92–7.89 (m, 3H), 7.67 (t, *J* = 7.5 Hz, 1H), 7.61 (t, *J* = 7.5 Hz, 1H), 3.27 (s, 3H).

Note

Ethyl 5-((Methylsulfonyl)diazenyl)benzofuran-2-carboxylate (1r).<sup>18e</sup> Yellow solid (4.6 g, 78%). Mp (dec): 117–118 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.35 (d, J = 2.0 Hz, 1H), 8.05 (dd, J = 2.0, 9.0 Hz, 1H), 7.73 (d, J = 9.0 Hz, 1H), 7.64 (d, J = 0.8 Hz, 1H), 4.47 (q, J = 7.1 Hz, 2H), 3.24 (s, 3H), 1.44 (t, J = 7.1 Hz, 3H).

2-Methyl-5-((methylsulfonyl)diazenyl)benzo[d]thiazole (1s).<sup>18e</sup> Brown solid (3.9 g, 77%). Mp (dec): 123–124 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.50 (d, J = 1.9 Hz, 1H), 7.95 (d, J = 8.6 Hz, 1H), 7.92 (dd, J = 1.9, 8.6 Hz, 1H), 3.25 (s, 3H), 2.89 (s, 3H).

Methyl 5-((Methylsulfonyl)diazenyl)nicotinate (1t).<sup>18e</sup> Yellow solid (4.1 g, 85%). Mp (dec): 103-105 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.46 (d, J = 1.9 Hz, 1H), 9.38 (d, J = 2.3 Hz, 1H), 8.73 (t, J = 2.1 Hz, 1H), 4.02 (s, 3H), 3.29 (s, 3H).

7-((Methylsulfonyl)diazenyl)quinoline (1u). Red solid (3.1 g, 66%). Mp (dec): 140–142 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.06 (dd, *J* = 1.5, 4.2 Hz, 1H), 8.79 (s, 1H), 8.25 (d, *J* = 8.2 Hz, 1H), 7.99–7.92 (m, 2H), 7.59–7.56 (m, 1H), 3.31 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 152.1, 149.0, 148.1, 136.0, 133.7, 132.0, 129.8, 123.8, 116.2, 34.8. IR (neat, *v* cm<sup>-1</sup>): 3440, 3045, 3020, 1500, 1463, 1328, 1151, 946, 867, 772, 746, 616, 566, 443. HRMS (ESI) *m/z*: [M + Na]<sup>+</sup> calcd for C<sub>10</sub>H<sub>9</sub>N<sub>3</sub>O<sub>2</sub>SNa, 258.0308; found, 258.0308.

4-Methyl-7-((methylsulfonyl)diazenyl)-2H-chromen-2-one (1v). Orange solid (2.3 g, 43%). Mp (dec): 102–104 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.88–7.85 (m, 2H), 7.80 (d, *J* = 9.0 Hz, 1H), 6.46 (d, *J* = 1.3 Hz, 1H), 3.27 (s, 3H), 2.51 (d, *J* = 1.3 Hz, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  159.04, 153.9, 151.1, 150.2, 126.0, 125.0, 118.9, 118.0, 113.6, 35.0, 18.7. IR (neat, *v* cm<sup>-1</sup>): 3440, 3045, 3006, 2920, 1715, 1624, 14110, 1340, 1259, 1158, 979, 903, 558. HRMS (ESI) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>11</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub>S, 267.0434; found, 267.0434.

9-Ethyl-3-((methylsulfonyl)diazenyl)-9H-carbazole (1w). Brown solid (4.9 g, 81%). Mp (dec): 145–147 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.70 (d, J = 1.9 Hz, 1H), 8.12 (d, J = 7.8 Hz, 1H), 8.08 (dd, J = 1.9, 8.9 Hz, 1H), 7.58–7.54 (m, 1H), 7.48–7.45 (m, 2H), 7.37–7.33 (m, 1H), 4.41 (q, J = 7.2 Hz, 2H), 3.24 (s, 3H), 1.49 (t, J = 7.3 Hz, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 144.3, 142.3, 140.9, 127.3, 123.7, 123.3, 122.2, 121.0, 120.5, 111.4, 109.6, 109.2, 38.1, 35.0, 13.8. IR (neat, v cm<sup>-1</sup>): 3442, 3032, 2975, 1593, 1498, 1418, 1119, 958, 835, 489. HRMS (ESI) m/z: [M + H]<sup>+</sup> calcd for C<sub>15</sub>H<sub>16</sub>N<sub>3</sub>O<sub>2</sub>S, 302.0958; found, 302.0958.

5-((Methylsulfonyl)diazenyl)benzo[d]thiazole (1**x**). Yellow solid (1.5 g, 31%). Mp (dec): 152–154 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.16 (s, 1H), 8.74 (d, *J* = 1.8 Hz, 1H), 8.12 (d, *J* = 8.1 Hz, 1H), 8.02 (dd, *J* = 1.9, 8.7 Hz, 1H), 3.28 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 156.6, 153.8, 147.7, 140.7, 123.4, 122.9, 118.2, 34.9. IR (neat,  $\nu$  cm<sup>-1</sup>): 3438, 3049, 1477, 1326, 1151, 865, 823, 531. HRMS (ESI) *m*/*z*: [M + H]<sup>+</sup> calcd for C<sub>8</sub>H<sub>8</sub>N<sub>3</sub>O<sub>2</sub>S<sub>2</sub>, 242.0052; found, 242.0052.

Synthesis of Compound 2c. Compound 2c was obtained by following a procedure previously reported for the synthesis of compounds 2a,b.<sup>19</sup> A mixture of 3,5-difluorobenzenesulfinate sodium salt (20 mmol, 1 equiv), N-[(trifuoromethyl)thio] aniline (1 equiv),<sup>20</sup> and paratoluenesulfonic acid (2.5 equiv) in DCE (130 mL) was stirred at room temperature. After the completion of the reaction, indicated by TLC, the mixture was filtered with a sand core funnel with silica gel, washed with CH2Cl2, and dried over Na2SO4. After concentration, the residue was purified by flash column chromatography to obtain the final product 2c (light yellow oil, 3.8 g, 68%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.55 (dd, J = 1.92, 5.68 Hz, 2H), 7.19 (tt, J = 2.28, 8.20 Hz, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$ 164.1 (d, J = 11.52 Hz, 1C), 161.5 (d, J = 11.36 Hz, 1C), 147.0 (t, J = 9.46 Hz, 1C), 120.7 (q, J = 311.66 Hz, 1C), 111.3 (d, J = 29.06 Hz, 1C), 111.3 (d, J = 11.31 Hz, 1C), 110.9 (t, J = 24.89 Hz, 1C). <sup>19</sup>F NMR (375 MHz,  $CDCl_3$ ):  $\delta$  -38.1 (s, 3F), -103.4 (s, 2F). The present compound was found to decompose upon HRMS analysis.

Synthesis of Thiotrifluoromethyl Arenes (3a-x). A solution of the chosen arylazo sulfone (1a-x, 0.25 M) in DCE was placed in a 10 mL microwave tube, and then S-(trifluoromethyl) 4-fluorobenzenesulfonothioate (2a, 1 mmol, 2.0-4.0 equiv, see Table 1) was added. The resulting solution was irradiated under stirring at room temperature

for 36 h by means of a 21 W blue LED, and then the reaction mixture was concentrated under reduced pressure to evaporate the solvent. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate mixture as the eluent).

Gram-Scale Preparation of **3a**. 1-([1,1'-Biphenyl]-4-yl)-2-(methylsulfonyl) diazene (1a, 10 mmol, 2.6 g) was placed in a round-bottom flask and dissolved in DCE (100 mL) under aerated conditions. Compound**2a**(20 mmol, 2.0 equiv) was added, and the resulting solution was irradiated under stirring for 48 h by means of a 21 W blue LED. The reaction mixture was concentrated under reduced pressure to evaporate the solvent, and the residue was purified by silica gel column chromatography (petroleum ether as the eluent) to obtain [1,1'-biphenyl]-4-yl (trifluoromethyl) sulfane**3a**(1.15 g, 45%) as a white solid.

**Characterization Data for the Isolated Products (3a–x).** [1,1'-Biphenyl]-4-yl(trifluoromethyl)sulfane (**3a**).<sup>13</sup> After purification by silica gel column chromatography (PE), compound **3a** was isolated as a white solid (50 mg, 75%).  $R_f$  (PE) = 0.8. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.74 (d, J = 8.3 Hz, 2H), 7.66–7.60 (m, 4H), 7.49 (t, J = 7.4 Hz, 2H), 7.42 (d, J = 7.4 Hz, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  143.8, 139.7, 136.7, 129.6 (q, <sup>1</sup> $_J$ (C,F) = 306.4 Hz), 128.9, 128.1, 127.2, 123.1 (q, <sup>2</sup> $_J$ (C,F) = 1.9 Hz). <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  –42.7 (s, 3F).

4-(4-((Trifluoromethyl)thio)phenyl)morpholine (**3b**).<sup>9</sup> After purification by silica gel column chromatography (PE/EA = 30:1), compound **3b** was isolated as a white solid (39 mg, 60%).  $R_f$  (PE/EA = 20:1) = 0.3. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.52 (d, J = 8.3 Hz, 2H), 6.88 (d, J = 8.3 Hz, 2H), 3.85 (t, J = 5.1 Hz, 4H), 3.23 (t, J = 5.0 Hz, 4H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  152.9, 137.9, 129.7 (q, <sup>1</sup>J(C,F) = 306.3 Hz), 115.2, 112.5, 66.6, 48.0. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  -44.1 (s, 3F).

(4-Phenoxyphenyl)(trifluoromethyl)sulfane (3c).<sup>13</sup> After purification by silica gel column chromatography (PE), compound 3c was isolated as a white solid (39 mg, 58%).  $R_f$  (PE) = 0.55. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.60 (d, J = 8.7 Hz, 2 H), 7.40 (t, J = 7.6 Hz, 2H), 7.22 (t, J = 7.4 Hz, 1H), 7.07 (d, J = 7.7 Hz, 2H), 7.00 (d, J = 8.8 Hz, 2H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  160.4, 155.6, 138.3, 129.6 (q, <sup>1</sup>*J*(C,F) = 306.3 Hz), 130.0, 124.5, 120.1, 118.6, 117.3 (q, <sup>2</sup>*J*(C,F) = 2.0 Hz). <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  -43.5 (s, 3F).

(4-(Benzyloxy)phenyl)(trifluoromethyl)sulfane (**3d**).<sup>21</sup> After purification by silica gel column chromatography (PE), compound **3d** was isolated as a white solid (37 mg, 52%).  $R_f$  (PE) = 0.6. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.59 (d, J = 8.8 Hz, 2H), 7.45–7.34 (m, 5H), 7.01 (d, J = 8.9 Hz, 2H), 5.09 (s, 2H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  161.0, 138.3, 136.2, 129.6 (q, <sup>1</sup>J(C,F) = 306.2 Hz), 128.7, 128.2, 127.5, 115.8, 115.2 (q, <sup>2</sup>J(C,F) = 2.0 Hz), 70.2. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  –43.8 (s, 3F).

*Methyl*(4-((*trifluoromethyl*)*thio*)*phenyl*)*sulfane* (**3e**).<sup>13</sup> After purification by silica gel column chromatography (PE), compound **3e** was isolated as a yellow oil (26 mg, 46%).  $R_f$  (PE) = 0.6. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.55 (d, *J* = 8.4 Hz, 2H), 7.25 (d, *J* = 8.8 Hz, 2H), 2.50 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  143.3, 136.6, 129.5 (q, <sup>1</sup>*J*(C,F) = 306.4 Hz), 126.4, 119.7, 15.0. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  -43.3 (s, 3F).

(4-(tert-Butyl)phenyl)(trifluoromethyl)sulfane (**3f**).<sup>13</sup> After purification by silica gel column chromatography (PE/EA = 100:1), compound **3f** was isolated as a colorless oil (35 mg, 60%).  $R_f$  (PE/EA = 50:1) = 0.6. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.58 (d, J = 8.0 Hz, 2H), 7.43 (d, J = 8.4 Hz, 2H), 1.33 (s, 9H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  154.4, 136.1, 129.7 (q, <sup>1</sup>J(C,F) = 306.0 Hz), 126.6, 120.9 (q, <sup>2</sup>J(C,F) = 1.8 Hz), 34.9, 31.1. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  -43.0 (s, 3F).

*Ethyl (E)-3-(4-((Trifluoromethyl)thio)phenyl)acrylate (3g).* After purification by silica gel column chromatography (PE/EA = 50:1), compound **3g** was isolated as a white solid (22 mg, 32%).  $R_f$  (PE/EA = 40:1) = 0.4. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.68–7.65 (m, 3H), 7.55 (d, *J* = 8.3 Hz, 2H), 6.49 (d, *J* = 16.0 Hz, 1H), 4.28 (q, *J* = 7.2 Hz, 2H), 1.34 (t, *J* = 7.1 Hz, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  166.4, 142.7, 136.9, 136.5, 129.4 (q, <sup>1</sup>*J*(C,F) = 306.6 Hz), 128.7, 126.2 (q, <sup>2</sup>*J*(C,F) = 2.0 Hz), 120.6, 60.8, 14.3. <sup>19</sup>F NMR (375

MHz, CDCl<sub>3</sub>):  $\delta$  –42.3 (s, 3F). HRMS (EM) m/z: [M – e]<sup>+</sup>calcd for C<sub>12</sub>H<sub>11</sub>F<sub>3</sub>O<sub>2</sub>S, 276.0432; found, 276.0417.

*Triisopropyl((4-((trifluoromethyl)thio)phenyl)ethynyl)silane (3h).* After purification by silica gel column chromatography (PE), compound **3h** was isolated as a colorless oil (40 mg, 45%).  $R_f$  (PE) = 0.8. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.59 (d, J = 8.2 Hz, 2H), 7.51 (d, J = 8.2 Hz, 2H), 1.13 (s, 21H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  136.0, 134.0, 132.9, 124.4 (q, <sup>1</sup>J(C,F) = 306.4 Hz), 126.4, 124.2 (q, <sup>2</sup>J(C,F) = 2.1 Hz), 105.5, 94.1, 18.6, 11.3. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  -42.6 (s, 3F). HRMS (EM) m/z: [M – e]<sup>+</sup>calcd for C<sub>18</sub>H<sub>25</sub>F<sub>3</sub>SSi, 358.1398; found, 358.1385.

(4-Bromophenyl)(trifluoromethyl)sulfane (3i).<sup>13</sup> After purification by silica gel column chromatography (PE), compound 3i was isolated as a colorless oil (36 mg, 56%).  $R_f$  (PE) = 0.85. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.57 (d, J = 8.8 Hz, 2H), 7.52 (d, J = 8.5 Hz, 2H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  137.7 132.8 129.2 (q, <sup>1</sup>J(C,F) = 306.6 Hz), 126.0, 123.4 (q, <sup>2</sup>J(C,F) = 2.1 Hz). <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  -42.7 (s, 3F).

4-((*Trifluoromethyl*)*thio*)*benzonitrile* (**3***j*).<sup>13</sup> After purification by silica gel column chromatography (PE/EA = 100:1), compound **3***j* was isolated as a white solid (27 mg, 53%).  $R_f$  (PE/EA = 50:1) = 0.35. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.77 (d, J = 8.4 Hz, 2H), 7.72 (d, J = 8.4 Hz, 2H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 136.0 132.9, 130.6 (q, <sup>2</sup>*J*(C,F) = 2.1 Hz), 129.0 (q, <sup>1</sup>*J*(C,F) = 307.0 Hz<sub>1</sub>), 117.6, 114.7. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>): δ -41.5 (s, 3F).

*Methyl* 4-((*Trifluoromethyl*)*thio*)*benzoate* (**3***k*).<sup>13</sup> After purification by silica gel column chromatography (PE/EA = 50:1), compound **3***k* was isolated as a yellow oil (52 mg, 83%).  $R_f$  (PE/EA = 50:1) = 0.5. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.08 (d, J = 8.5 Hz, 2H), 7.71 (d, J = 8.0 Hz, 2H), 4.40 (q, J = 7.1 Hz, 2H), 1.40 (t, J = 7.1 Hz, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  165.5, 135.5, 132.6, 130.8, 129.7 (q, <sup>2</sup>*J*(C,F) = 2.0 Hz), 129.3 (q, <sup>1</sup>*J*(C,F) = 306.5 Hz), 61.5, 14.2. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  -41.9 (s, 3F). 1-(4-((*Trifluoromethyl*)*thio*)*phenyl*)*ethanone* (**3***J*).<sup>13</sup> After purifi-

1-(4-((Trifluoromethyl)thio)phenyl)ethanone (**3**).<sup>13</sup> After purification by silica gel column chromatography (PE/EA = 50:1), compound **3**I was isolated as a yellow soil (36 mg, 66%).  $R_f$  (PE/EA = 50:1) = 0.25. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.98 (d, *J* = 8.4 Hz, 2H), 7.74 (d, *J* = 8.3 Hz, 2H), 2.63 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 197.0, 138.5, 135.7, 130.0 (q, <sup>2</sup>*J*(C,F) = 2.0 Hz), 129.3 (q, <sup>1</sup>*J*(C,F) = 306.6 Hz), 129.1, 26.7. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>): δ -41.8 (s, 3F).

*Phenyl*(4-((trifluoromethyl)thio)phenyl)methanone (**3m**).<sup>13</sup> After purification by silica gel column chromatography (PE/EA = 100:1), compound **3m** was isolated as a white solid (48 mg, 68%).  $R_f$  (PE/EA = 50:1) = 0.4. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.84–7.76 (m, 6H), 7.63 (t, J = 7.4 Hz, 1H), 7.51 (t, J = 8.4 Hz, 2H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 195.5, 139.5, 136.8, 135.5, 133.0, 130.6, 130.7, 129.3 (q, <sup>1</sup>J(C,F) = 306.7 Hz), 129.1 (q, <sup>2</sup>J(C,F) = 2.0 Hz), 128.5. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>): δ –41.8 (s, 3F).

(3-Bromophenyl)(trifluoromethyl)sulfane (3n).<sup>22</sup> After purification by silica gel column chromatography (PE), compound 3n was isolated as a white solid (31 mg, 48%):  $R_f$  (PE) = 0.8. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.82 (s, 1H), 7.64–7.59 (m, 2H), 7.31 (t, *J* = 8.0 Hz, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  138.7, 134.7, 134.0, 130.7, 129.3 (q, <sup>1</sup>*J*(C,F) = 306.5 Hz), 126.3 (q, <sup>2</sup>*J*(C,F) = 2.1 Hz), 122.9. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  –42.4 (s, 3F).

(*Trifluoromethyl*)(*3*, *4*, *5*-trimethoxyphenyl)sulfane (**30**).<sup>23</sup> After purification by silica gel column chromatography (PE/EA = 40:1), compound **30** was isolated as a yellow solid (30 mg, 44%).  $R_f$  (PE/EA = 40:1) = 0.35. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  6.86 (s, 2H), 3.88(s, 9H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  153.4, 140.6, 129.6 (q, <sup>1</sup>J(C,F) = 306.4 Hz), 118.5 (q, <sup>2</sup>J(C,F) = 2.1 Hz), 113.7, 60.9, 56.3. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  -43.0 (s, 3F).

5-((*Trifluoromethyl*)*thio*)*benzo*[*d*][1,3]*dioxole* (**3***p*).<sup>21</sup> After purification by silica gel column chromatography (PE), compound **3***p* was isolated as a white solid (16 mg, 29%).  $R_f$  (PE) = 0.6. Mp = 113–114 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.17 (dd, J = 1.72, 8.08 Hz, 1H), 7.09 (d, J = 1.56 Hz, 1H), 6.84 (d, J = 8.04 Hz, 1H), 6.04 (s, 2H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  150.3, 148.3, 131.6, 129.5 (q,

<sup>1</sup>J(C,F) = 306.3 Hz), 116.2, 116.0 (q, <sup>2</sup>J(C,F) = 2.1 Hz), 109.0. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  –43.9 (s, 3F).

Naphthalen-2-yl(trifluoromethyl)sulfane (3q).<sup>10c</sup> After purification by silica gel column chromatography (PE), compound 3q was isolated as a white solid (27 mg, 48%).  $R_f$  (PE) = 0.9. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.21 (s, 1H), 7.89–7.87 (m, 3H), 7.67 (d, J = 8.5 Hz, 1H), 7.61–7.55 (m, 2H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$ 137.0, 133.9, 133.4, 131.8, 129.7 (q, <sup>1</sup>J(C,F) = 306.6 Hz), 129.2, 128.2, 127.9, 127.8, 127.0, 121.5 (q, <sup>2</sup>J(C,F) = 2.3 Hz). <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  –42.5 (s, 3F).

*Ethyl 5-((Trifluoromethyl)thio)-2,3-dihydrobenzofuran-2-carboxylate* (**3***r*). After purification by silica gel column chromatography (PE/EA = 50:1), compound **3***r* was isolated as a white solid (49 mg, 67%).  $R_f$  (PE/EA = 50:1) = 0.4. Mp = 102–103 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.03 (s, 1H), 7.73–7.70 (m, 1H), 7.64–7.62 (m, 1H), 7.53 (s, 1H), 4.46 (q, *J* = 7.16 Hz, 2H), 1.43 (t, *J* = 7.1 Hz, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  159.0, 156.6, 147.2, 135.3, 131.8, 129.5 (q, <sup>1</sup>*J*(C,F) = 306.5 Hz), 128.2, 119.5 (q, <sup>2</sup>*J*(C,F) = 2.1 Hz), 113.5, 113.2, 61.8, 14.3. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  –43.3 (s, 3F). HRMS (EM) *m/z*: [M –e]<sup>+</sup> calcd for C<sub>12</sub>H<sub>9</sub>F<sub>3</sub>O<sub>3</sub>S, 290.0224; found, 290.0213.

2-Methyl-5-((trifluoromethyl)thio)benzo[d]thiazole (**3s**). After purification by silica gel column chromatography (PE/EA = 20:1), compound **3s** was isolated as a white solid (19 mg, 30%).  $R_f$  (PE/EA = 20:1) = 0.33. Mp = 87–88 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$ 8.26 (d, J = 1.36 Hz, 1H), 7.87 (d, J = 8.28 Hz, 1H), 7.61 (dd, J =1.48, 8.32 Hz, 1H), 2.86 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  168.9, 153.9, 138.7, 131.8, 130.5, 129.6 (q, <sup>1</sup>*J*(C,F) = 306.4 Hz), 122.1, 131.7, 20.2 (q, <sup>1</sup>*J*(C,F) = 4.4 Hz). <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  – 42.9 (s, 3F). HRMS (EI) m/z: [M + H]<sup>+</sup> calcd for C<sub>9</sub>H<sub>7</sub>F<sub>3</sub>NS<sub>2</sub>, 249.9967; found, 249.9970.

*Methyl* 5-((*Trifluoromethyl*)*thio*)*nicotinate* (**3t**).<sup>24</sup> After purification by silica gel column chromatography (PE/EA = 10:1), compound **3t** was isolated as a white solid (25 mg, 43%).  $R_f$  (PE/EA = 5:1) = 0.5. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.33 (s,1H), 9.01 (s, 1H), 8.59 (s, 1H), 4.00 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  164.4, 159.0, 152.5, 144.3, 128.9 (q, <sup>1</sup>*J*(C,F) = 307.1 Hz), 126.9, 122.4, 52.9. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  -42.0 (s, 3F).

*7-((Trifluoromethyl)thio)quinoline* (**3***u*). After purification by silica gel column chromatography (PE/EA = 30:1), compound **3***u* was isolated as a yellow oil (17 mg, 30%).  $R_f$  (PE/EA = 10:1) = 0.3. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.00 (d, J = 2.8 Hz, 1H), 8.48 (s, 1H), 8.21 (d, J = 8.2 Hz, 1H), 7.87 (d, J = 8.5 Hz, 1H), 7.75 (d, J = 8.5 Hz, 1H), 7.52–7.49 (s, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  151.5, 147.8, 137.8, 136.0, 132.3, 129.5 (q, <sup>1</sup>*J*(C,F) = 306.8 Hz), 129.0, 128.9, 125.9 (q, <sup>2</sup>*J*(C,F) = 1.8 H), 122.7. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  –41.9 (s, 3F). HRMS (ESI) m/z: [M + H]<sup>+</sup> calcd for C<sub>10</sub>H<sub>7</sub>F<sub>3</sub>NS, 230.0246; found, 230.0246.

4-Methyl-7-((trifluoromethyl)thio)-2H-chromen-2-one (**3v**).<sup>25</sup> After purification by silica gel column chromatography (PE/EA = 10:1), compound **3v** was isolated as a white solid (19 mg, 29%).  $R_f$  (PE/EA = 10:1) = 0.2. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.66–7.64 (m,2H), 7.55 (dd, J = 1.6, 8.2 Hz, 1H), 6.38 (d, J = 1.2 Hz,1H), 2.46 (d, J = 1.2 Hz, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  159.6, 153.3, 151.3, 130.8, 129.2 (q, <sup>1</sup>J(C,F) = 306.8 Hz), 128.3 (q, <sup>2</sup>J(C,F) = 2.1 Hz), 125.4, 123.9, 121.7, 116.7, 18.5. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  –41.8 (s, 3F).

9-Ethyl-3-((trifluoromethyl)thio)-9H-carbazole (**3**w).<sup>6d</sup> After purification by silica gel column chromatography (PE), compound **3**w was isolated as a colorless solid (19 mg, 25%).  $R_f$  (PE) = 0.4. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.39 (d, *J* = 1.6 Hz, 1H), 8.12 (d, *J* = 7.8, Hz, 1H), 7.72 (dd, *J* = 1.6, 8.5 Hz, 1H), 7.55–7.51 (m, 1H), 7.45–7.72 (m, 2H), 7.31–7.27 (m, 1H), 5.14 (q, *J* = 7.2 Hz, 2H), 1.46 (t, *J* = 7.2 Hz, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  141.0, 140.3, 133.7, 129.9 (q, <sup>1</sup>*J*(C,F) = 306.4 Hz), 129.6, 126.6, 123.9, 122.2, 120.7, 119.8, 112.5 (q, <sup>2</sup>*J*(C,F) = 2.0 Hz), 109.2, 108.8, 37.7, 13.7. <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  –44.1 (s, 3F).

5-((*Trifluoromethyl*)*thio*)*benzo*[*d*]*thiazole* (**3***x*). After purification by silica gel column chromatography (PE/EA = 50:1), compound **3***x* was isolated as a colorless oil (12 mg, 20%).  $R_f$  (PE/EA = 50:1) = 0.5.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.09 (s,1H), 8.47 (d, *J* = 1.6 Hz, 1H), 8.02 (d, *J* = 8.4 Hz, 1H), 7.71 (dd, *J* = 1.6, 8.4 Hz, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  155.7, 153.7, 136.7, 132.6, 131.9, 129.5 (q, <sup>1</sup>*J*(C,F) = 306.4 Hz), 122.7, 122.2 (q, <sup>2</sup>*J*(C,F)= 2.1 Hz). <sup>19</sup>F NMR (375 MHz, CDCl<sub>3</sub>):  $\delta$  -42.7 (s, 3F). HRMS (ESI) *m*/*z*: [M + H]<sup>+</sup> calcd for C<sub>8</sub>H<sub>3</sub>F<sub>3</sub>NS<sub>2</sub>, 235.9810; found, 235.9811.

Irradiation of 1k in the Presence of TEMPO. A solution of arylazo sulfone 1k (0.25 M) in DCE (1 mL) was placed in a 10 mL microwave tube, and then S-(trifluoromethyl) 4-fluorobenzenesulfonothioate (2a, 1 mmol, 4.0 equiv) and TEMPO (4 equiv) were added. The resulting solution was irradiated under stirring at room temperature for 36 h by means of a 21 W blue LED, and then the reaction mixture was concentrated under reduced pressure to evaporate the solvent. Purification by silica gel column chromatography (PE/EA = 30:1), afforded ethyl 4-((2,2,6,6-tetramethylpiperidin-1-yl)oxy)benzoate 4k as a white solid (23 mg, 30% yield).  $R_f$  (PE/EA = 30:1) = 0.4. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.93 (d, J = 9.1 Hz, 2H), 7.21 (s, 2H), 4.33 (q, J = 7.2 Hz, 2H), 1.64–1.56 (m, SH), 1.44–1.41 (m, 1H), 1.36 (t, J = 7.2 Hz, 3H), 1.23 (s, 6H), 0.99 (s, 6H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  167.4, 166.5, 131.0, 122.4, 113.7, 60.6, 60.4, 39.7, 32.4, 20.4, 17.0, 14.4.

## ASSOCIATED CONTENT

#### **Supporting Information**

The Supporting Information is available free of charge at https://pubs.acs.org/doi/10.1021/acs.joc.0c02669.

Mechanistic studies and spectroscopic data for compounds 1, 2c, and 3 (PDF)

#### AUTHOR INFORMATION

#### **Corresponding Authors**

- Stefano Protti PhotoGreen Lab, Department of Chemistry, University of Pavia, Pavia 27100, Italy; o orcid.org/0000-0002-5313-5692; Email: stefano.protti@unipv.it
- Xia Zhao Tianjin Key Laboratory of Structure and Performance for Functional Molecules, College of Chemistry, Tianjin Normal University, Tianjin 300387, People's Republic of China; orcid.org/0000-0001-9477-5912; Email: hxxyzhx@mail.tjnu.edu.cn

#### Authors

- Ankun Li Tianjin Key Laboratory of Structure and Performance for Functional Molecules, College of Chemistry, Tianjin Normal University, Tianjin 300387, People's Republic of China
- Yuxuan Li Tianjin Key Laboratory of Structure and Performance for Functional Molecules, College of Chemistry, Tianjin Normal University, Tianjin 300387, People's Republic of China; College of Biotechnology, Tianjin University of Science & Technology, Tianjin 300457, China
- Junjie Liu Tianjin Key Laboratory of Structure and Performance for Functional Molecules, College of Chemistry, Tianjin Normal University, Tianjin 300387, People's Republic of China; College of Biotechnology, Tianjin University of Science & Technology, Tianjin 300457, China
- Jingqi Chen Tianjin Key Laboratory of Structure and Performance for Functional Molecules, College of Chemistry, Tianjin Normal University, Tianjin 300387, People's Republic of China
- Kui Lu College of Biotechnology, Tianjin University of Science & Technology, Tianjin 300457, China; October 0000-0002-9685-2691
- **Di Qiu** Tianjin Key Laboratory of Structure and Performance for Functional Molecules, College of Chemistry,

Tianjin Normal University, Tianjin 300387, People's Republic of China; orcid.org/0000-0001-9706-6302 Maurizio Fagnoni – PhotoGreen Lab, Department of

Chemistry, University of Pavia, Pavia 27100, Italy; orcid.org/0000-0003-0247-7585

Complete contact information is available at: https://pubs.acs.org/10.1021/acs.joc.0c02669

## **Author Contributions**

<sup>⊥</sup>A.L. and Y.L. contributed equally to this work.

### Notes

The authors declare no competing financial interest.

# ACKNOWLEDGMENTS

The authors sincerely thank the National Natural Science Foundation of China (21572158) and the Tianjin Natural Science Foundation (18JCQNJC76600) for financial support.

# REFERENCES

(1) (a) Lipinski, C. A.; Lombardo, F.; Dominy, B. W.; Feeney, P. J. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv. Drug Delivery Rev.* **1997**, *23*, 3–25. (b) Lipinski, C. A. Drug-like properties and the causes of poor solubility and poor permeability. *J. Pharmacol. Toxicol. Methods* **2000**, *44*, 235–249. (c) Avdeef, A. *Absorption and Drug Development: Solubility, Permeability, and Charge State*; Wiley: Hoboken, NJ, 2012. (d) Johnson, T. W.; Gallego, R. A.; Edwards, M. P. Lipophilic Efficiency as an Important Metric in Drug Design. *J. Med. Chem.* **2018**, *61*, 6401–6420.

(2) (a) Smart, B. E. Fluorine substituent effects (on bioactivity). J. Fluorine Chem. 2001, 109, 3–11. (b) Muller, K.; Faeh, C.; Diederich, F. Fluorine in Pharmaceuticals: Looking Beyond Intuition. Science 2007, 317, 1881–1886. (c) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. Fluorine in medicinal chemistry. Chem. Soc. Rev. 2008, 37, 320–330. (c) Ojima, I. Fluorine in Medicinal Chemistry and Chemical Biology; Wiley: Chichester, United Kingdom, 2009. (d) Hagmann, W. K. The Many Roles for Fluorine in Medicinal Chemistry. J. Med. Chem. 2008, 51, 4359–4369. (e) Landelle, G.; Panossian, A. R.; Leroux, F. Trifluoromethyl Ethers and – Thioethers as Tools for Medicinal Chemistry and Drug Discovery. Curr. Top. Med. Chem. 2014, 14, 941.

(3) Hansch, C.; Leo, A.; Taft, R. W. A survey of Hammett substituent constants and resonance and field parameters. *Chem. Rev.* **1991**, *91*, 165–195.

(4) (a) Boiko, V. N. Aromatic and heterocyclic perfluoroalkyl sulfides. Methods of preparation. Beilstein J. Org. Chem. 2010, 6, 880-921. (b) Toulgoat, F.; Alazet, S.; Billard, T. Direct Trifluoromethylthiolation Reactions: The "Renaissance" of an Old Concept. Eur. J. Org. Chem. 2014, 2014, 2415-2428. (c) Chachignon, H.; Cahard, D. State-of-the-Art in Electrophilic Trifluoromethylthiolation Reagents. Chin. J. Chem. 2016, 34, 445-454. (d) Barata-Vallejo, S.; Bonesi, S.; Postigo, A. Late stage trifluoromethylthiolation strategies for organic compounds. Org. Biomol. Chem. 2016, 14, 7150-7182. (e) Toulgoat, F.; Billard, T. Towards CF<sub>3</sub>S Group: From Trifluoromethylation of Sulfides to Direct Trifluoromethylthiolation. In Modern Synthesis Processes and Reactivity of Fluorinated Compounds: Progress in Fluorine Science; Groult, H., Leroux, F., Tressaud, A., Eds.; Elsevier Science: London, 2017; p 141. (f) Srivastava, V.; Singh, P. K.; Srivastava, A.; Singh, P. P. Recent application of visible-light induced radicals in C-S bond formation. RSC Adv. 2020, 10, 20046.

(5) (a) Liu, J.; Zhao, X.; Jiang, L.; Yi, W.  $Tf_2O$ -Promoted Trifluoromethylthiolation of Various Arenes Using  $NaSO_2CF_3$ . Adv. Synth. Catal. **2018**, 360, 4012–4016. (b) Zhao, X.; Wei, A.; Yang, B.; Li, T.; Li, Q.; Qiu, D.; Lu, K. Transition-Metal-Free Direct Trifluoromethylthiolation and Trifluoromethylsulfoxidation of Electron-rich Aromatics with  $CF_3SO_2Na$  in the presence of PCl<sub>3</sub>. J. Org.

Chem. 2017, 82, 9175–9181. (c) Lu, K.; Deng, Z.; Li, M.; Li, T.; Zhao, X. Transition Metal-Free Direct Trifluoromethylthiolation of Indoles using Trifluoromethanesulfonyl Chloride in the Presence of Triphenylphosphine. Org. Biomol. Chem. 2017, 15, 1254–1260. (d) Glenadel, Q.; Alazet, S.; Billard, T. First and second generation of trifluoromethanesulfenamide reagent: A trifluoromethylthiolating comparison. J. Fluorine Chem. 2015, 179, 89–95. (e) Shao, X.; Xu, C.; Lu, L.; Shen, Q. Structure-Reactivity Relationship of Trifluoromethanesulfenates: Discovery of an Electrophilic Trifluoromethylthiolating Reagent. J. Org. Chem. 2015, 80, 3012–3021.

(6) For representative examples of the trifluoromethylthiolation reaction, see: (a) Zheng, C.; Liu, Y.; Hong, J.; Huang, S.; Zhang, W.; Yang, Y.; Fang, G. Copper(I)-promoted trifluoromethylthiolation of arenediazonium salts with AgSCF<sub>3</sub>. Tetrahedron Lett. 2019, 60, 1404-1407. (b) Zhao, M.; Zhao, X.; Zheng, P.; Tian, Y. Cu-mediated oxidative trifluoromethylthiolation of arylboronic acids with (bpy)-CuSCF3. J. Fluorine Chem. 2017, 194, 73-79. (c) Yang, Y.; Xu, L.; Yu, S.; Liu, X.; Zhang, Y.; Vicic, D. A. Triphenylphosphine-Mediated Deoxygenative Reduction of CF<sub>3</sub>SO<sub>2</sub>Na and Its Application for Trifluoromethylthiolation of Aryl Iodides. Chem. - Eur. J. 2016, 22, 858-863. (d) Matheis, C.; Wagner, V.; Goossen, L. J. Sandmeyer-Type Trifluoromethylthiolation and Trifluoromethylselenolation of (Hetero)Aromatic Amines Catalyzed by Copper. Chem. - Eur. J. 2016, 22, 79-82. (e) Dubbaka, S.; Atthunuri, A.; Prakash, K.; Rangabashyam, P.; Gadde, S.; Kothandaraman, R. Copper-Mediated Oxidative Trifluoromethylthiolation of Potassium Aryltrifluoroborates with Elemental Sulfur and Ruppert-Prakash Reagent. Synthesis 2016, 48, 1246-1252. (f) Saravanan, P.; Anbarasan, P. Copper-Catalyzed Trifluoromethylthiolation of Di(hetero)aryl- $\lambda_3$ -iodanes: Mechanistic Insight and Application to Synthesis of (Hetero)Aryl Trifluoromethyl Sulfides. Adv. Synth. Catal. 2015, 357, 3521-3528. (g) Glenadel, Q.; Alazet, S.; Tlili, A.; Billard, T. Mild and Soft Catalyzed Trifluoromethylthiolation of Boronic Acids: The Crucial Role of Water. Chem. - Eur. J. 2015, 21, 14694-14698. (h) Danoun, G.; Bayarmagnai, B.; Gruenberg, M. F.; Goossen, L. J. Sandmeyer trifluoromethylthiolation of arenediazonium salts with sodium thiocyanate and Ruppert-Prakash reagent. Chem. Sci. 2014, 5, 1312-1316.

(7) (a) Kalvet, I.; Guo, Q.; Tizzard, G. J.; Schoenebeck, F. When Weaker Can Be Tougher: The Role of Oxidation State (I) in P- vs N-Ligand-Derived Ni-Catalyzed Trifluoromethylthiolation of Aryl Halides. ACS Catal. 2017, 7, 2126–2132. (b) Duerr, A. B.; Yin, G.; Kalvet, I.; Napoly, F.; Schoenebeck, F. Nickel-catalyzed trifluoromethylthiolation of Csp<sup>2</sup>-O bonds. Chem. Sci. 2016, 7, 1076–1081. (c) Yin, G.; Kalvet, I.; Englert, U.; Schoenebeck, F. Fundamental Studies and Development of Nickel-Catalyzed Trifluoromethylthiolation of Aryl Chlorides: Active Catalytic Species and Key Roles of Ligand and Traceless MeCN Additive Revealed. J. Am. Chem. Soc. 2015, 137, 4164–4172. (d) Zhang, C.-P.; Vicic, D. A. Nickel-Catalyzed Synthesis of Aryl Trifluoromethyl Sulfides at Room Temperature. J. Am. Chem. Soc. 2012, 134, 183–185.

(8) (a) Krishanmoorthy, S.; Schnell, S. D.; Dang, H.; Fu, F.; Prakash, G.K. S. Fluorodecarboxylation: Synthesis of aryl trifluoromethyl ethers (ArOCF<sub>3</sub>) and thioethers (ArSCF<sub>3</sub>). *J. Fluorine Chem.* **2017**, 203, 130–135. (b) Ma, J.-j.; Liu, Q.-r.; Lu, G.-p.; Yi, W.-b. Silver-catalyzed fluoroalkylation of thiols using fluoroalkanesulfinates. *J. Fluorine Chem.* **2017**, *193*, 113–117.

(9) Yin, G.; Kalvet, I.; Schoenebeck, F. Trifluoromethylthiolation of Aryl Iodides and Bromides Enabled by a Bench-Stable and Easy-To-Recover Dinuclear Palladium(I) Catalyst. *Angew. Chem., Int. Ed.* **2015**, *54*, 6809–6813.

(10) (a) Sumii, Y.; Taniguchi, M.; Xu, X.-H.; Tokunaga, E.; Shibata, N. Ortho-lithiation reaction of aryl triflones. *Tetrahedron* **2018**, *74*, 5635–5641. (b) Ma, J.-J.; Yi, W.-B.; Lu, G.-P.; Cai, C. Trifluoromethylation of thiophenols and thiols with sodium trifluoromethanesulfinate and iodine pentoxide. *Catal. Sci. Technol.* **2016**, *6*, 417–421. (c) Yamaguchi, K.; Sakagami, K.; Miyamoto, Y.; Jin, X.; Mizuno, N. Oxidative nucleophilic strategy for synthesis of thiocyanates and trifluoromethyl sulfides from thiols. *Org. Biomol.* 

*Chem.* **2014**, *12*, 9200–9206. (d) Harsányi, A.; Dorkó, E.; Csapó, A.; Bakó, T.; Peltz, C.; Rábai, J. Convenient synthesis and isolation of trifluoromethylthio-substituted building blocks. *J. Fluorine Chem.* **2011**, *132*, 1241–1246.

(11) Surya Prakash, G. K.; Wang, F.; Zhang, Z.; Haiges, R.; Rahm, M.; Christe, K. O.; Mathew, T.; Olah, G. A. Long-Lived Trifluoromethanide Anion: A Key Intermediate in Nucleophilic Trifluoromethylations. *Angew. Chem., Int. Ed.* **2014**, *53*, 11575–11578.

(12) Bertoli, G.; Exner, B.; Evers, M. V.; Tschulik, K.; Gooßen, L. J. Metal-free trifluoromethylthiolation of arenediazonium salts with  $Me_4NSCF_3$ . J. Fluorine Chem. **2018**, 210, 132–136.

(13) (a) Zhao, X.; Zheng, X.; Tian, M.; Tong, Y.; Yang, B.; Wei, X.; Qiu, D.; Lu, K. Visible-light photocatalytic trifluoromethylthiolation of aryldiazonium salts: conversion of amino group into trifluoromethylthiol group. Org. Chem. Front. 2018, 5, 2636-2640. (b) Ghiazza, C.; Monnereau, C.; Khrouz, L.; Billard, T.; Tlili, A. Visible-Light-Mediated Synthesis of Trifluoromethylthiolated Arenes. Synthesis 2019, 51, 2865-2870. (c) Koziakov, D.; Majek, M.; von Wangelin, A. J. Radical Aromatic Trifluoromethylthiolation: Photoredox Catalysis vs. Base Mediation. Eur. J. Org. Chem. 2017, 2017, 6722-6725. The Eosin Y photocatalyzed trifluoromethylation of alkyl sulfides was also reported: (d) Srivastava, V.; Singh, P. K.; Singh, P. P. Trifluoromethylation of Disubstituted Morpholines by Metal-Free Visible Light Photoredox Catalysis. Asian J. Chem. 2016, 28, 2159. See for review: (e) Ghiazza, C.; Billard, T.; Tlili, A. Merging Visible-Light Catalysis for the Direct Late-Stage Group-16-Trifluoromethyl Bond Formation. Chem. - Eur. J. 2019, 25, 6482-6495.

(14) Firth, J. D.; Fairlamb, I. J. S. A Need for Caution in the Preparation and Application of Synthetically Versatile Aryl Diazonium Tetrafluoroborate Salts. *Org. Lett.* **2020**, *22*, 7057–7059.

(15) Protti, S.; Ravelli, D.; Fagnoni, M. Wavelength-dependence and wavelength-selectivity in photochemical reactions. *Photochem. Photobiol. Sci.* **2019**, *18*, 2094–2101.

(16) (a) Crespi, S.; Protti, S.; Fagnoni, M. Wavelength Selective Generation of Aryl Radicals and Aryl Cations for Metal-free Photoarylations. *J. Org. Chem.* **2016**, *81*, 9612–9619.

(17) Abdulla, H. O.; Scaringi, S.; Amin, A. A.; Mella, M.; Protti, S.; Fagnoni, M. Aryldiazenyl Radicals from Arylazo Sulfones: Visible Light-Driven Diazenylation of Enol Silyl Ethers. *Adv. Synth. Catal.* **2020**, 362, 2150.

(18) For recent applications of aryl radicals photogenerated from arylazo sulfone, see: (a) da Silva Júnior, P. E.; Amin, H. I. M.; Nauth, A. M.; da Silva Emery, F.; Protti, S.; Opatz, T. Flow Photochemistry of Azosulfones: Application of "Sunflow" Reactors. ChemPhotoChem. 2018, 2, 878-883. (b) Jung, H. I.; Lee, J. H.; Kim, D. Y. Photocatalyst-free Photoredox Arylation of Quinoxalin-2(1H)-Ones with Aryldiazo Sulfones. Bull. Korean Chem. Soc. 2018, 39, 1003-1006. (c) Amin, H. I. M.; Raviola, C.; Amin, A. A.; Mannucci, B.; Protti, S.; Fagnoni, M. Hydro/deutero Deamination of Arylazo Sulfones Under Metal- and (Photo)Catalyst-Free Conditions. Molecules 2019, 24, 2164. (d) Xu, Y.; Yang, X.; Fang, H. Additiveand Photocatalyst-Free Borylation of Arylazo Sulfones under Visible Light. J. Org. Chem. 2018, 83, 12831. (e) Lian, C.; Yue, G.; Mao, J.; Liu, D.; Ding, Y.; Liu, Z.; Qiu, D.; Zhao, X.; Lu, K.; Fagnoni, M.; Protti, S. A Visible Light Driven Synthesis of Arylstannanes from Arylazo Sulfones. Org. Lett. 2019, 21, 5187-5191. (f) Malacarne, M.; Protti, S.; Fagnoni, M. A visible light driven, metal-free route to aromatic amides via radical arylation of isonitriles. Adv. Synth. Catal. 2017, 359, 3826. (g) Abdulla, H. O.; Amin, A. A.; Raviola, C.; Opatz, T.; Protti, S.; Fagnoni, M. Smooth Metal-Free Photoinduced Preparation of Valuable 8-Arylxanthines. Eur. J. Org. Chem. 2020, 2020, 1448. (h) Qiu, D.; Lian, C.; Mao, J.; Fagnoni, M.; Protti, S. Dyedauxiliary Groups, an Emerging Approach in Organic Chemistry. The Case of Arylazo Sulfones. J. Org. Chem. 2020, 85, 12813.

(19) Li, Y.; Qiu, G.; Wang, H.; Sheng, J. Generation of trifluoromethyl thiolsulphonate through one-pot reaction of sulforyl

chloride and trifluoromethanesulfanylamides. *Tetrahedron Lett.* 2017, *58*, 690.

(20) Liu, J.; Yao, H.; Li, X.; Wu, H.; Lin, A.; Yao, H.; Xu, J.; Xu, J. Organocatalytic 1,5-trifluoromethylthiosulfonylation of vinylcyclopropane mediated by visible light in the water phase. *Org. Chem. Front.* **2020**, *7*, 1314.

(21) Jouvin, K.; Matheis, C.; Goossen, L. Synthesis of Aryl Tri- and Difluoromethyl Thioethers via a C-H-Thiocyanation/Fluoroalkylation Cascade. *Chem. - Eur. J.* **2015**, *21*, 14324.

(22) Kieltsch, I.; Eisenberger, P.; Togni, A. Mild Electrophilic Trifluoromethylation of Carbon- And Sulfur-Centered Nucleophiles by a Hypervalent iodine(III)-CF<sub>3</sub> Reagent. *Angew. Chem., Int. Ed.* **2007**, *46*, 754–757.

(23) Chen, C.; Xie, Y.; Chu, L.; Wang, R.; Zhang, X.; Qing, F. Copper-Catalyzed Oxidative Trifluoromethylthiolation of Aryl Boronic Acids with TMSCF<sub>3</sub> and Elemental Sulfur. *Angew. Chem., Int. Ed.* **2012**, *51*, 2492.

(24) Zhang, M.; Weng, Z. Copper-Mediated Trifluoromethylthiolation of Heteroaryl Bromides. *Adv. Synth. Catal.* **2016**, *358*, 386.

(25) Khotavivattana, T.; Verhoog, S.; Tredwell, M.; Pfeifer, L.; Calderwood, S.; Wheelhouse, K.; Lee Collier, T.; Gouverneur, V. <sup>18</sup>F-Labeling of Aryl-SCF<sub>3</sub>, -OCF<sub>3</sub> and -OCHF<sub>2</sub> with [<sup>18</sup>F]Fluoride. *Angew. Chem., Int. Ed.* **2015**, *54*, 9991.