

Note

Palladium-Catalyzed Borylation of Primary Alkyl Bromides

Amruta Joshi-Pangu, Xinghua Ma, Mohamed Diane, Sidra Iqbal,
Robert J Kribs, Richard Huang, Chao-Yuan Wang, and Mark R. Biscoe

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/jo301156e • Publication Date (Web): 10 Jul 2012

Downloaded from <http://pubs.acs.org> on July 13, 2012

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications
High quality. High impact.


The Journal of Organic Chemistry is published by the American Chemical Society.
1155 Sixteenth Street N.W., Washington, DC 20036
Published by American Chemical Society. Copyright © American Chemical Society.
However, no copyright claim is made to original U.S. Government works, or works
produced by employees of any Commonwealth realm Crown government in the course
of their duties.

Palladium-Catalyzed Borylation of Primary Alkyl Bromides

Amruta Joshi-Pangu, Xinghua Ma, Mohamed Diane, Sidra Iqbal, Robert J. Kribs, Richard Huang, Chao-Yuan Wang, and Mark R. Biscoe*

Department of Chemistry, The City College of New York, 160 Convent Ave., New York, NY 10031

Email of corresponding author: mbiscoe@ccny.cuny.edu



Abstract: A mild Pd-catalyzed process for the borylation of alkyl bromides has been developed using bis(pinacolato)diboron as a boron source. This process accommodates the use of a wide range of functional groups on the alkyl bromide substrate. Primary bromides react with complete selectivity in the presence of a secondary bromide. The generality of this approach is demonstrated by its extension to the use of alkyl iodides and alkyl tosylates, as well as borylation reactions employing bis(neopentyl glycolato)diboron as the boron source.


Over the past few decades, the Suzuki-Miyaura cross-coupling reaction has become a workhorse for the construction of carbon-carbon bonds.¹⁻³ The majority of such reactions involves the use of aryl or vinylboron nucleophiles. In comparison, methods that allow the general use of alkylboron nucleophiles in Suzuki-Miyaura cross-coupling reactions have remained far less developed.^{4,5} A major focus of the research in our laboratory is the development of new methods for carbon-carbon bond formation via metal-catalyzed cross-coupling reactions using alkyl nucleophiles.⁶⁻⁸ For such transformations, alkylboron reagents constitute ideal nucleophiles. The high covalency of the carbon-boron bond typically makes alkylboron reagents air and moisture stable, configurationally stable, and compatible with a diverse range of functional groups.^{9,10} Thus, the development of new cross-coupling methods that enable the general use of alkylboron reagents is an attractive synthetic goal.

Classically, organoboron reagents have been generated via borylation of the corresponding lithium or magnesium reagents, or via the hydroboration of a terminal olefin.¹¹ More recently, significant progress has been made towards the development of new, milder methods to generate organoboron reagents for use in synthesis. Particular attention has been focused on the preparation of alkyl and aryl boronic ester derivatives via transition metal-catalyzed processes. Notable examples include the metal-catalyzed hydroboration of alkenes,¹²⁻¹⁶ metal-catalyzed C-H activation/borylation sequences,¹⁷⁻²⁵ and metal-catalyzed β -borylation of α,β -unsaturated compounds.²⁶⁻³⁰ With the aim of developing a mild and general method to generate primary alkylboronate esters for use in cross-coupling reactions, we explored the possibility of using bis(pinacolato)diboron as a boron source in the Pd-catalyzed borylation of alkyl halides. An analogous strategy has been successfully applied to the formation of arylboronic esters from aryl halides.³¹⁻³³ However, a method to effect the direct Pd-catalyzed borylation of alkyl halides has not been reported.³⁴⁻³⁶ Herein, we report a Pd-catalyzed method for the selective borylation of primary alkyl bromides. This process displays excellent functional group tolerance and can be extended to the use of alkyl iodides and alkyl tosylates. Additionally, we have demonstrated that bis(neopentyl glycolato)diboron can be successfully employed as the boron source with nominal variation of the standard reaction conditions.

Initial ligand screening revealed that trialkylphosphines were most effective in the Pd-catalyzed borylation of (3-bromopropyl)benzene. This is consistent with the reported ability of trialkylphosphine-ligated palladium(0) complexes to undergo oxidative addition into primary alkyl halides.³⁷⁻³⁹ Products resulting from alkyl halide reduction and β -hydride elimination constituted the major side-products observed during reaction optimization. There was no evidence for protodeboration of the alkylboronic ester products over the course of the reactions. Our final, optimized conditions are displayed in Table 1. The use of $t\text{-Bu}_2\text{MeP}\cdot\text{HBF}_4$ as a supporting ligand resulted in the highest yields. $t\text{-BuOH}$ and DMA were the most effective solvents for this transformation. K_3PO_4 was the most effective base, though K_2CO_3 could also be used successfully. While the addition of exogenous water was found

to be essential for efficient conversion, only a nominal variation in GC yield was observed when 3 equiv H₂O was used in place of 15 equiv H₂O.

Table 1. Optimization of Standard Reaction Conditions




Entry	Variation from standard conditions above	Yield (%) ^a
1	None	97
2	3 equiv H ₂ O	93
3	NMP used as solvent, 3 equiv H ₂ O	29
4	Toluene used as solvent, 3 equiv H ₂ O	<10
5	DMA used as solvent, 3 equiv H ₂ O	70
6	DMF used as solvent, 3 equiv H ₂ O	44
7	DMF used as solvent, no water	<10
8	PCy ₃ ·HBF ₄ used as ligand, 3 equiv H ₂ O	34
9	P <i>n</i> -Bu ₃ ·HBF ₄ used as ligand, 3 equiv H ₂ O	<10
10	P <i>t</i> -Bu ₃ ·HBF ₄ used as ligand, 3 equiv H ₂ O	<10

^a Yields and selectivities determined by GC with dodecane as an internal standard.

Using the optimized conditions displayed in Table 1, alkylboronate esters were generated cleanly in high yields from a wide range of functionalized primary alkyl bromides (Table 2). Alkyl bromide substrates bearing functional groups including nitriles, alcohols, ethers, esters, amides, and imides were successfully borylated.⁴⁰ The presence of heterocyclic substituents on the alkyl bromide substrate was also well tolerated. 1-Bromo-5-chloropentane underwent selective borylation at the bromide. Although primary alkyl chlorides did not undergo efficient borylation under these conditions, alkyl iodides and alkyl tosylates were viable electrophiles. As these reactions proceeded, we observed no evidence of transmetallation of the pinacol alkylboronate ester produced in the reaction.


Table 2. Pd-Catalyzed Borylation of Primary Alkyl Bromides^a



^a Alkyl bromide (1 mmol), bis(pinacolato)diboron (1.2 mmol); average yield of 2 runs. ^b Using alkyl iodide. ^c Using alkyl tosylate. ^d 0.75 mmol scale. ^e 3 mol % PCy₃•HBF₄. ^f 1.5 mol % Pd₂(dba)₃, 9 mol % PCy₃•HBF₄.

In contrast to nickel- and copper-catalyzed processes which can undergo oxidative addition via radical processes,^{35,36,41,42} Pd(0) complexes generally undergo oxidative addition with alkyl halides via direct nucleophilic displacement of the halide leaving group.³⁷⁻³⁹ As a result, the use of Pd catalysis is generally limited to primary electrophiles. Fortunately, this mechanism for oxidative addition can be exploited to achieve the chemoselective borylation of molecules bearing multiple halide groups. Using 1,4-dibromopentane (**4**) as an electrophile (Scheme 1), we observed that a primary bromide could indeed be selectively borylated in the presence of a remote secondary bromide. This reaction occurred efficiently, with no detectable trace of borylation at the secondary bromide.


Scheme 1. Selective Borylation of Primary Bromide^a



^a Alkyl bromide (1 mmol), bis(pinacolato)diboron (1.2 mmol); average yield of 2 runs.


To broaden the utility of our borylation method, we have developed methods to convert the crude pinacol boronate products to other boronate derivatives (Table 3). After borylation of (3-bromopropyl)benzene via the standard conditions of Table 1, transformation of the resulting pinacol boronate ester to a trifluoroboronate (**6a**), boronic acid (**6b**), and propylene glycol boronate (**6c**) was readily achieved without isolation of the pinacol boronate intermediate.

Table 3. Direct Preparation of Boronate Ester Derivatives^a



^a Alkyl bromide (0.75 mmol); average yield of 2 runs. ^b KHF₂, MeOH, rt, 2 h. ^c NaIO₄, HCl(aq), THF/H₂O, rt, 3 h. ^d footnote c, then 1,3-propanediol, benzene, reflux, 12 h.

It has been demonstrated that transmetallation of an aryl group from a pinacol boronate ester to palladium occurs significantly more slowly than transmetallation of an aryl group from a neopentyl glycol boronate ester.⁴³ Consistent with this trend, Pd-catalyzed borylation reactions that form pinacol alkylboronate esters showed no sign of post-borylation transmetallation of the alkyl group, while neopentyl glycol alkylboronate esters generated by an analogous process displayed evidence of subsequent transmetallation of the alkyl group. However, with minor modifications, we successfully developed a process to achieve the Pd-catalyzed cross-coupling of alkyl bromides and bis(neopentyl glycolato)diboron (**7**) with minimal transmetallation of the generated alkylboronate ester (**8**) (Table 4). Although yields for the reactions were lower than those for the formation of the corresponding pinacol alkylboronate esters, the general success of this reaction underscores the broad utility of our process, and suggests that alkyl boronate compounds may be successfully formed from the Pd-catalyzed cross-coupling reaction of alkyl electrophiles with other diboron reagents.

Table 4. Pd-Catalyzed Borylation of Alkyl Bromides Using Bis(neopentyl glycolato)diboron^a

^a Alkyl bromide (0.2 mmol), bis(pinacolato)diboron (0.22 mmol); average yield of 2 runs.

In conclusion, we have developed a mild process for the Pd-catalyzed borylation of primary alkyl electrophiles using bis(pinacolato)diboron as a boron source. The reaction is highly tolerant of reactive functional groups, requires low catalyst loadings, and accommodates the use of alkyl bromide, iodide, and tosylate electrophiles. Primary bromides react with complete selectivity over secondary bromides. The ability to easily convert the pinacol alkylboronate esters directly into boronic acids, trifluoroboronates, and other boronate esters broadens the utility of this method. Finally, we have demonstrated that our borylation process is not exclusive to the use of bis(pinacolato)diboron as a boron source, and can be extended to the use of other diboron reagents.

Experimental Section.

Toluene and THF (unstabilized) were transferred to separate 20 L solvent-delivery kegs and vigorously purged with argon for 2 h. The solvents were further purified by passing them under argon pressure through two packed columns of neutral alumina (for THF) or through neutral alumina and copper (II) oxide (for toluene). All other reagents and solvents were used as received unless otherwise noted. Flash chromatography was performed using silica gel (ultra pure grade). Nuclear magnetic resonance spectra were recorded on a 500 MHz instrument. All ^1H NMR experiments are reported in δ units, parts per million (ppm), and were measured relative to the signals for residual chloroform (7.26 ppm), dimethyl sulfoxide (2.50 ppm) or acetonitrile (1.94 ppm). All ^{13}C NMR spectra are reported in ppm relative to deuteriochloroform (77.2 ppm), dimethyl sulfoxide (39.5 ppm) or acetonitrile (118.3 ppm) and were obtained with ^1H decoupling. All GC analyses were performed on a gas chromatograph with an FID

1 detector using a 25 m x 0.20 mm capillary column with cross-linked methyl siloxane as the stationary
2 phase.
3
4
5
6

7 **General procedure for borylation of primary alkyl bromides with bis(pinacolato)diboron.**

8
9
10 $\text{Pd}_2(\text{dba})_3$ (4.6 mg, 0.005 mmol), di-*tert*-butyl(methyl)phosphonium tetrafluoroborate (7.4 mg, 0.03
11 mmol), $(\text{BPin})_2$ (305 mg, 1.2 mmol) and $\text{K}_3\text{PO}_4 \cdot \text{H}_2\text{O}$ (460 mg, 2 mmol) were weighed out on the
12 benchtop in an oven-dried 10 mL screw top test tube with stir bar. The test tube was sealed using a
13 screw cap lined with a teflon septum. With stirring begun, the test tube was evacuated (100 mTorr), and
14 backfilled three times with argon using a needle attached to a vacuum manifold. The alkyl bromide (1
15 mmol) was then added to the test tube via a microsyringe, followed by degassed tertiary butyl alcohol (3
16 mL) and degassed water (0.25 mL). If the alkyl bromide was a solid, it was weighed out after
17 $\text{K}_3\text{PO}_4 \cdot \text{H}_2\text{O}$ before evacuating the test tube. The test tube was sealed with electrical tape and the reaction
18 mixture was stirred overnight on the benchtop at 60 °C with no additional argon pressure. *N.B.: Ensure*
19 *that $\text{K}_3\text{PO}_4 \cdot \text{H}_2\text{O}$ does not clump while the reaction is in progress. If clumping becomes problematic,*
20 *K_2CO_3 may be used in place of $\text{K}_3\text{PO}_4 \cdot \text{H}_2\text{O}$.* The reaction mixture was quenched through the addition of
21 saturated aqueous NH_4Cl (5 mL). The resulting mixture was then poured into a separatory funnel and
22 extracted with diethyl ether (3 x 20 mL). The combined organic layers were washed with brine and dried
23 over Na_2SO_4 . The crude product was purified by column chromatography.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 **General procedure for borylation of primary alkyl bromides with bis(neopentylglycolato) diboron.**

46
47 $\text{Pd}_2(\text{dba})_3$ (1.83 mg, 0.002 mmol), di-*tert*-butyl(methyl)phosphonium tetrafluoroborate (3 mg, 0.012
48 mmol), $(\text{BNeop})_2$ (54 mg, 0.24 mmol) and anhydrous K_3PO_4 (85 mg, 0.4 mmol) were weighed out in the
49 glove-box in an oven-dried 10 mL screw top test tube with stir bar. The test tube was sealed using a
50 screw cap lined with a teflon septum. The reaction was stirred on a stir plate and alkyl bromide (0.2
51 mmol) was then added to the vial via a microsyringe. Degassed tertiary butyl alcohol (1mL) and
52 degassed water (20 μL) were then added under argon, outside of the glove box. The test tube was sealed
53
54
55
56
57
58
59
60

with an electrical tape and the reaction mixture was stirred for 5 hours on the benchtop at 80 °C with no additional argon pressure. The reaction mixture was quenched through the addition of saturated aqueous NH₄Cl (ca. 4 mL). The resulting mixture was then poured into a separatory funnel and extracted with diethyl ether (3 x 10 mL). The combined organic layers were washed with brine and dried over Na₂SO₄. The crude product was purified by column chromatography.

4,4,5,5-Tetramethyl-2-(3-phenylpropyl)-1,3,2-dioxaborolane (3a).⁴⁴ The general procedure was employed. A reddish yellow liquid (236 mg, 96%) was isolated by column chromatography (98:2 Hex:Ether). ¹H NMR (500 MHz, CDCl₃) δ: 7.27 (t, *J* = 7.5 Hz, 2H), 7.16-7.20 (m, 3H), 2.63 (t, *J* = 7.5 Hz, 2H), 1.76 (app. quint., *J* = 7.5 Hz, 2H), 1.26 (s, 12H), 0.85 (t, *J* = 7.5 Hz, 2H) ppm. ¹³C NMR (125 MHz, CDCl₃) δ: 142.8, 128.6, 128.3, 125.7, 83.0, 38.69, 26.2, 24.9, 11 (B-CH₂, br) ppm. ¹¹B (MHz, CDCl₃) δ: 33.3 ppm.

2-Isopentyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3b).⁴⁵ The general procedure was employed. A yellow liquid (147 mg, 74%) was isolated by column chromatography (99:1 Hex:Ether). ¹H NMR (500 MHz, CDCl₃) δ: 1.45 (m, *J* = 6.5 Hz, 1H), 1.21-1.3 (m, 2H), 1.23 (s, 12H), 0.84 (d, *J* = 6.5 Hz, 6H), 0.74 (t, *J* = 8.5 Hz, 2H) ppm. ¹³C NMR (125 MHz, CDCl₃) δ: 83.0, 33.1, 30.4, 25.0, 22.4 ppm. The carbon atom directly attached to the boron atom was not detected, likely due to quadrupolar broadening. ¹¹B (MHz, CDCl₃) δ: 33.3 ppm.

6-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)hexan-1-ol (3c).³⁵ The general procedure was employed. A brownish liquid (139 mg, 61%) was isolated by column chromatography (gradient from 80:20 to 75:25 Hex:EtOAc). ¹H NMR (500 MHz, CDCl₃) δ: 3.63 (app. q, *J* = 6.5 Hz, 2H), 1.53-1.59 (m, 2H), 1.38-1.45 (m, 2H), 1.29-1.36 (m, 4H), 1.24 (s, 12H), 0.77 (t, *J* = 7.5 Hz, 2H) ppm. ¹³C NMR (125 MHz, CDCl₃) δ: 83.0, 63.2, 32.8, 32.2, 25.6, 24.9, 24.0 ppm. The carbon atom directly attached to the boron atom was not detected, likely due to quadrupolar broadening. ¹¹B (MHz, CDCl₃) δ: 33.2 ppm.

4,4,5,5-Tetramethyl-2-(2-(thiophen-2-yl)ethyl)-1,3,2-dioxaborolane (3d).³⁵ The general procedure was employed. An orange-yellow liquid (149 mg, 63%) was isolated by column

chromatography (98:2 Hex:Ether). ^1H NMR (500 MHz, CDCl_3) δ : 7.08 (dd, $J = 5$ Hz, 1.5 Hz, 1H), 6.88 (m, 1H), 6.79 (m, 1H), 2.96 (t, $J = 8$ Hz, 2H), 1.24 (m, 14H), ppm. ^{13}C NMR (125 MHz, CDCl_3) δ : 147.9, 126.6, 123.5, 122.7, 83.3, 24.9, 24.5 ppm. The carbon atom directly attached to the boron atom was not detected, likely due to quadrupolar broadening. ^{11}B (MHz, CDCl_3) δ : 32.9 ppm.

6-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)hexanenitrile (3e).³⁵ The general procedure was employed. A golden yellow liquid (214 mg, 96%) was isolated by column chromatography (gradient from 95:5 to 93:7 Hex:Ether). ^1H NMR (500 MHz, CDCl_3) δ : 2.30 (t, $J = 7$ Hz, 2H), 1.63 (quint., $J = 7.0$ Hz, 2H), 1.40-1.43 (m, 4H), 1.22 (s, 12H), 0.76 (t, $J = 7.0$ Hz, 2H) ppm. ^{13}C NMR (125 MHz, CDCl_3) δ : 119.9, 83.1, 31.3, 25.2, 24.9, 23.2, 17.1, 11 (B- CH_2 , br) ppm. ^{11}B (MHz, CDCl_3) δ : 33.1 ppm.

Ethyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)butanoate (3f). The general procedure was employed with following modification: The ligand used in this reaction is tricyclohexylphosphonium tetrafluoroborate (11 mg, 0.03 mmol). A bright yellow liquid (215 mg, 89%) was isolated by column chromatography (95:5 Hex:Ether). ^1H NMR (500 MHz, CDCl_3) δ : 4.10 (q, $J = 7$ Hz, 2H), 2.31 (t, $J = 7.5$ Hz, 2H), 1.74 (app. quint., $J = 7.5$ Hz, 2H), 1.23-1.26 (m, 15H), 0.81 (t, $J = 8$ Hz, 2H) ppm. ^{13}C NMR (125 MHz, CDCl_3) δ : 173.8, 83.1, 60.2, 36.7, 24.8, 19.8, 14.4, 11 (B- CH_2 , br) ppm. ^{11}B (MHz, CDCl_3) δ : 32.8 ppm. HRMS (EI^+): Calcd ($\text{C}_{12}\text{H}_{23}\text{BO}_4^+$) 242.1689; Found 242.1700.

2-(2-Cyclohexylethyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3g).⁴⁶ The general procedure was employed with following modifications: $\text{Pd}_2(\text{dba})_3$ (14 mg, 0.015 mmol), tricyclohexylphosphonium tetrafluoroborate ligand (33 mg, 0.09 mmol). A pale yellow liquid (198 mg, 83%) was isolated by column chromatography (98:2 Hex:Ether). ^1H NMR (500 MHz, CDCl_3) δ : 1.61-1.72 (m, 5H), 1.11-1.31 (m, 18H), 0.83 (app. q, $J = 12$ Hz, 2H), 0.75 (t, $J = 8.5$ Hz, 2H) ppm. ^{13}C NMR (125 MHz, CDCl_3) δ : 83.0, 40.1, 33.1, 31.5, 26.9, 26.6, 24.9, 20 (B- CH_2 , br) ppm. ^{11}B (MHz, CDCl_3) δ : 33.4 ppm.

2-(5-Chloropentyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3h).⁴⁷ The general procedure was employed with following modification: The ligand used in this reaction is

1 tricyclohexylphosphonium tetrafluoroborate (11 mg, 0.03 mmol). A yellowish orange liquid (211 mg,
2 91%) was isolated by column chromatography (98:2 Hex:Ether) ^1H NMR (500 MHz, CDCl_3) δ : 3.50 (t,
3 $J = 7$ Hz, 2H), 1.75 (app. quint., $J = 7$ Hz, 2H), 1.41 (m, 4H), 1.23 (s, 12H), 0.77 (t, $J = 7$ Hz, 2H) ppm.
4 ^{13}C NMR (125 MHz, CDCl_3) δ : 83.1, 45.2, 32.6, 29.6, 24.9, 23.4, 11 (B- CH_2 , br) ppm. ^{11}B (MHz,
5 CDCl_3) δ : 32.9 ppm.
6
7
8
9
10

11 **10-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)decanamide (3i).** The general procedure
12 was employed. A white solid (279 mg, 94%) was isolated by column chromatography (95:5
13 CH_2Cl_2 :MeOH). ^1H NMR (500 MHz, CDCl_3) δ : 5.32 (broad d, 2H), 2.21 (t, $J = 7.5$ Hz, 2H), 1.60-1.66
14 (m, 2H), 1.22-1.26 (m, 22H), 0.76 (t, $J = 7.5$ Hz, 2H) ppm. ^{13}C NMR (125 MHz, CDCl_3) δ : 175.6, 83.0,
15 36.1, 32.6, 29.62, 29.50, 29.46, 29.38, 25.7, 25.0, 24.1 ppm. The carbon atom directly attached to the
16 boron atom was not detected, likely due to quadrupolar broadening. ^{11}B (MHz, CDCl_3) δ : 33.7 ppm.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

4,4,5,5-Tetramethyl-2-(4-phenoxybutyl)-1,3,2-dioxaborolane (3j).⁴⁸ The general procedure
was employed. A bright yellow liquid (259 mg, 94%) was isolated by column chromatography (99:1
Hex:Ether). ^1H NMR (500 MHz, CDCl_3) δ : 7.26 (dt, $J = 7.5$ Hz, 1 Hz, 2H), 6.88-6.93 (m, 3H), 3.95 (t, J
= 6.5 Hz, 2H), 1.80 (quint., $J = 6.5$ Hz, 2H), 1.59 (quint., $J = 8$ Hz, 2H), 1.25 (s, 12H), 0.86 (t, $J = 8$ Hz,
2H) ppm. ^{13}C NMR (125 MHz, CDCl_3) δ : 159.3, 129.5, 120.5, 114.6, 83.1, 67.8, 31.9, 25.0, 20.7 ppm.
The carbon atom directly attached to the boron atom was not detected, likely due to quadrupolar
broadening. ^{11}B (MHz, CDCl_3) δ : 33.1 ppm.

2-(4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)butyl)isoindoline-1,3-dione (3k). The
general procedure was employed with following modification: tricyclohexylphosphonium
tetrafluoroborate (11 mg, 0.03 mmol) was used as ligand. A brown liquid (260 mg, 79%) was isolated
by column chromatography (gradient from 85:15 to 80:20 Hex:Ether). ^1H NMR (500 MHz, CDCl_3) δ :
7.80 (dd, $J = 8.5$ Hz, 3 Hz, 2H), 7.67 (dt, $J = 8.5$ Hz, 2.5 Hz, 2H), 3.64 (t, $J = 7.5$ Hz, 2H), 1.66 (quint., J
= 7.5 Hz, 2H), 1.44 (quint., $J = 8$ Hz, 2H), 1.20 (s, 12H), 0.79 (t, $J = 7.5$ Hz, 2H) ppm. ^{13}C NMR (125

MHz, CDCl₃) δ : 168.5, 133.9, 132.3, 123.2, 83.1, 38.0, 31.2, 24.9, 21.4, 11 (B-CH₂, br) ppm. ¹¹B (MHz, CDCl₃) δ : 33.0 ppm. Anal. Calcd. for C₁₈H₂₄BNO₄: C, 65.67; H, 7.35. Found: C, 65.45; H, 7.28.

2-(4-Bromopentyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (5). The general procedure was employed. A bright yellow liquid (219 mg, 79%) was isolated by column chromatography (98:2 Hex:Ether). ¹H NMR (500 MHz, CDCl₃) δ : 4.13 (app. sextet, *J* = 6.5 Hz, 1H), 1.82-1.90 (m, 1H), 1.73-1.80 (m, 1H), 1.70 (d, *J* = 7 Hz, 3H), 1.48-1.66 (m, 2H), 1.24 (s, 12H), 0.73-0.84 (m, 2H) ppm. ¹³C NMR (125 MHz, CDCl₃) δ : 83.1, 51.6, 43.8, 26.5, 24.9, 22.3, 10 (B-CH₂, br) ppm. ¹¹B (160 MHz, CDCl₃) δ : 33.2 ppm. HRMS (EI⁺): Calcd (C₁₁H₂₂BBrO₂ – CH₃⁺) 261.0661; Found 261.0656.

Potassium 3-phenyl-trifluoroboratopropane (6a).³⁵ The general procedure was employed. A white solid (118mg, 70 %) ¹H NMR (500 MHz, DMSO) δ : 7.20-7.23 (t, *J* = 7.5 Hz, 4H), 7.08- 7.12 (m, 3H), 2.45 (t, *J* = 8.0 Hz, 2H), 1.40 (app. quint., *J* = 7.5 Hz, 2H), 0.02 (m, 2H) ppm. ¹³C NMR (125 MHz, DMSO) δ : 144.1, 128.3, 127.9, 124.9, 28.25, 28.24, 20 (B-CH₂, br) ppm. ¹¹B (160 MHz, DMSO) δ : 9.3 ppm.

3-Phenylpropylboronic acid (6b).⁴⁹ The general procedure was employed. A yellowish white solid (109 mg, 89 %) ¹H NMR (500 MHz, DMSO) δ : 7.40 (s, 2H), 7.25 (t, *J* = 8 Hz, 2H), 7.13-7.16 (m, 3H), 2.50 (m, 2H), 1.60 (app. quint., *J* = 7.5 Hz, 2H), 0.60 (t, *J* = 8.0 Hz, 2H) ppm. ¹³C NMR (125 MHz, CD₃CN) δ : 143.9, 129.4, 129.1, 126.5, 39.2, 27.3 ppm. The carbon atom directly attached to the boron atom was not detected, likely due to quadrupolar broadening. ¹¹B (160 MHz, CD₃CN) δ : 37.0 ppm.

2-(3-Phenylpropyl)-1,3,2-dioxaborinane (6c).⁵⁰ The general procedure was employed. (92 mg, 60%) ¹H NMR (500 MHz, CD₃CN) δ : 7.24-7.27 (m, 2H), 7.13-7.18 (m, 3H), 3.95 (t, *J* = 5.0 Hz, 4H), 2.58 (t, *J* = 7.5 Hz, 2H), 1.90 (quint., *J* = 6.0 Hz, 2H), 1.68 (quint., *J* = 8.0 Hz, 2H), 0.73 (t, *J* = 8.0 Hz, 2H) ppm. ¹³C NMR (125 MHz, CD₃CN) δ : 143.1, 128.7, 128.2, 125.6, 61.7, 38.7, 27.5, 26.1, 11 (B-CH₂, br) ppm. ¹¹B (160 MHz, CDCl₃) δ : 29.9 ppm.

5,5-Dimethyl-2-(3-phenylpropyl)-1,3,2-dioxaborinane (8a). The general procedure was employed. A yellowish liquid (32 mg, 70%) was isolated by column chromatography (90:10 Hex: Ether). ¹H NMR (500 MHz, CD₃CN) δ : 7.24-7.27 (m, 2H), 7.14-7.19 (m, 3H), 3.57 (s, 4H), 2.60 (t, *J* =

8 Hz, 2H), 1.70 (app. quint., $J = 8$ Hz, 2H), 0.95 (s, 6H), 0.76 (t, $J = 8$ Hz, 2H) ppm. ^{13}C NMR (125 MHz, CD_3CN) δ : 143.1, 128.7, 128.3, 125.6, 72.1, 38.9, 31.8, 29.9, 26.3, 22.0, 15 (B- CH_2 , br) ppm. ^{11}B (160 MHz, CDCl_3) δ : 29.7 ppm. Anal. Calcd. for $\text{C}_{14}\text{H}_{21}\text{BO}_2$: C, 72.44; H, 9.12. Found: C, 72.45; H, 9.23.

Ethyl 4-(5,5-dimethyl-1,3,2-dioxaborinan-2-yl)butanoate (8b). The general procedure was employed. A yellow liquid (28 mg, 62%) was isolated by column chromatography (90:10 Hex:Ether). ^1H NMR (500 MHz, CD_3CN) δ : 4.11 (q, $J = 7$ Hz, 2H), 3.58 (s, 4H), 2.29 (t, $J = 7.5$ Hz, 2H), 1.70 (app. quint., $J = 7.5$ Hz, 2H), 1.25 (t, $J = 7$ Hz, 3H), 0.95 (s, 6H), 0.75 (t, $J = 8$ Hz, 2H) ppm. ^{13}C NMR (125 MHz, CD_3CN) δ : 174.1, 72.1, 60.2, 36.9, 31.8, 29.8, 22.0, 19.9, 14.4, 14 (B- CH_2 , br) ppm. ^{11}B (160 MHz, CDCl_3) δ : 29.4 ppm. HRMS (FAB $^+$): Calcd (for $\text{C}_{11}\text{H}_{21}\text{BO}_4 + \text{H}^+$) 229.1611; Found 229.1622.

6-(5,5-Dimethyl-1,3,2-dioxaborinan-2-yl)hexanenitrile (8c). The general procedure was employed. A yellow liquid (18 mg, 42%) was isolated by column chromatography (90:10 Hex:Ether). ^1H NMR (500 MHz, CDCl_3) δ : 3.59 (s, 4H), 2.32 (t, $J = 7.5$ Hz, 2H), 1.65 (app. quint., $J = 7$ Hz, 2H), 1.37-1.47 (m, 4H), 0.95 (s, 6H), 0.72 (t, $J = 8$ Hz, 2H) ppm. ^{13}C NMR (125 MHz, CDCl_3) δ : 120.1, 72.1, 31.8, 31.4, 29.8, 25.4, 23.3, 22.0, 17.2, 14 (B- CH_2 , br) ppm. ^{11}B (160 MHz, CDCl_3) δ : 29.4 ppm. HRMS (FAB $^+$) Calcd (for $\text{C}_{11}\text{H}_{20}\text{BNO}_2 + \text{H}^+$) 210.1665; Found 210.1660.

Acknowledgment. We thank The National Institutes of Health (1SCGM096932-01), The City College of New York (CCNY), The Alfred P. Sloan Foundation, and PSC-CUNY for financial support. We gratefully acknowledge the National Science Foundation for an instrumentation grant (CHE-0840498). Acknowledgement is additionally made to the donors of the American Chemical Society Petroleum Research Fund (50307-DNI1) for partial support of this research. Finally, we thank BASF for their generous gift of (BPin) $_2$.

Supporting Information Available: ^1H , ^{13}C , and ^{11}B NMR spectra for all products. This information is available free of charge via the Internet at <http://pubs.acs.org>.

References and Footnotes.

- ¹ *Metal-Catalyzed Cross-Coupling Reactions*; de Meijere, A., Diederich, F., Eds., Wiley-VCH: New York, 2004.
- ² Molander, G. A.; Ellis, N. *Acc. Chem. Res.* **2007**, *40*, 275.
- ³ Martin, R.; Buchwald, S. L. *Acc. Chem. Res.* **2008**, *41*, 1461.
- ⁴ Chemler, S. R.; Trauner, D.; Danishefsky, S. J. *Angew. Chem. Int. Ed.* **2001**, *40*, 4544.
- ⁵ Jana, R.; Pathak, T. P.; Sigman, M. S. *Chem Rev.* **2011**, *111*, 1417.
- ⁶ Joshi-Pangu, A.; Ganesh, M.; Biscoe, M. R. *Org. Lett.* **2011**, *13*, 1218.
- ⁷ Joshi-Pangu, A.; Wang, C.-Y.; Biscoe, M. R. *J. Am. Chem. Soc.* **2011**, *133*, 8478.
- ⁸ Joshi-Pangu, A.; Biscoe, M. R. *Synlett* **2012**, *23*, 1103.
- ⁹ Boudier, A.; Bromm, L. O.; Lotz, M.; Knochel, P. *Angew. Chem. Int. Ed.* **2000**, *39*, 4414.
- ¹⁰ Crudden, C. M.; Glasspoole, B. W.; Lata, C. J. *Chem. Commun.* **2009**, 6704.
- ¹¹ Hall, D. G. "Structures, Properties, and Preparation of Boronic Acid Derivatives" in *Boronic Acids: Preparation and Applications in Organic Synthesis and Medicine*; Hall, D. G., Ed., Wiley-VCH: Weinheim, 2005, pp. 1-99.
- ¹² Burgess, K.; Ohlmeyer, M. J. *Chem. Rev.* **1991**, *91*, 1179.
- ¹³ Evans, D. A.; Fu, G. C.; Hoveyda, A. H. *J. Am. Chem. Soc.* **1992**, *114*, 6671.
- ¹⁴ Burgess, K.; van der Donk, W. A.; Westcott, S. A.; Marder, T. B.; Baker, R. T.; Calabrese, J. C. *J. Am. Chem. Soc.* **1992**, *114*, 9350.
- ¹⁵ Beletskaya, I.; Pelter, A. *Tetrahedron* **1997**, *53*, 4957.
- ¹⁶ Edwards, D. R.; Crudden, C. M.; Yam, K. *Adv. Synth. Catal.* **2005**, *347*, 50.
- ¹⁷ Chen, H. Y.; Schlecht, S.; Semple, T. C.; Hartwig, J. F. *Science* **2000**, 287, 1995.
- ¹⁸ Shimada, S.; Batsanov, A. S.; Howard, J. A. K.; Marder, T. B. *Angew. Chem.* **2001**, *113*, 2226.
- ¹⁹ Cho, J. -Y.; Tse, M. K.; Holmes, D.; Maleczka, Jr., R. E.; Smith III, M. R. *Science* **2002**, 295, 305.
- ²⁰ Mkhallid, I. A. I.; Barnard, J. H.; Marder, T. B.; Baker, R. T.; Calabrese, J. C. *Chem. Rev.* **2010**, *110*, 890.
- ²¹ Yamamoto, T.; Morita, T.; Takagi, J.; Yamakawa, T. *Org. Lett.* **2011**, *13*, 5766.
- ²² Hartwig, J. F. *Acc. Chem. Res.* **2012**, *6*, 864.
- ²³ Ros, A.; Lopez-Rodriguez, R.; Estepa, B.; Alvarez, E.; Fernandez, R.; Lassaletta, J. M. *J. Am. Chem. Soc.* **2012**, *134*, 4573.
- ²⁴ Yu, J. -Q.; Dai, H. -X. *J. Am. Chem. Soc.* **2012**, *134*, 134.
- ²⁵ Xiao, B.; Li, Y. -M.; Liu, Z. -J.; Yang, H. -Y.; Fu, Y. *Chem. Commun.* **2012**, 48, 4854.
- ²⁶ O'Brien, M.; Lee, K. -S.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2010**, *132*, 10630.
- ²⁷ Schiffner, J. A.; Muthner, K.; Oestreich, M. *Angew. Chem. Int. Ed.* **2010**, *49*, 1194.
- ²⁸ Sasaki, Y.; Zhong, C.; Sawamura, H.; Ito, H. *J. Am. Chem. Soc.* **2010**, *132*, 1226.
- ²⁹ Molander, G. A.; McKee, S. A. *Org. Lett.* **2011**, *13*, 4684.

- ³⁰ Thorpe, S. B.; Calderone, J. A.; Santos, W. *Org. Lett.* **2012**, *14*, 1918.
- ³¹ Ishiyama, T.; Itoh, Y.; Kitano, T.; Miyaura, N. *Tetrahedron Lett.* **1997**, *38*, 3447.
- ³² Furstner, A.; Seidel, G. *Org. Lett.* **2002**, *4*, 541.
- ³³ Billingsley, K. L.; Barder, T. E.; Buchwald, S. L. *Angew. Chem. Int. Ed.* **2007**, *46*, 5359.
- ³⁴ While the work contained in this manuscript was in progress, processes for the Cu-catalyzed borylation of alkyl electrophiles were reported. See references 39-40.
- ³⁵ Yang, C. -T.; Zhang, Z. -Q.; Tajuddin, H.; Wu, C. -C.; Liang, J.; Liu, J. -H.; Fu, Y.; Czyzewska, M.; Steel, P. G.; Marder, T. B.; Liu, L. *Angew. Chem. Int. Ed.* **2012**, *51*, 528.
- ³⁶ Ito, H.; Kubota, K. *Org. Lett.* **2012**, *14*, 890.
- ³⁷ Netherton, M. R.; Dai, C.; Neuschütz, K.; Fu, G. C. *J. Am. Chem. Soc.* **2001**, *123*, 10099.
- ³⁸ Netherton, M. R.; Fu, G. C. *Angew. Chem. Int. Ed.* **2002**, *41*, 3910.
- ³⁹ Hills, I. D.; Netherton, M. R.; Fu, G. C. *Angew. Chem. Int. Ed.* **2003**, *42*, 5749.
- ⁴⁰ When substrates bearing remote alkenes and alkynes were employed, significant background diborylation of the alkene or alkyne was observed. See: Bonet, A.; Pubill-Ulldemolins, C.; Bo, C.; Gulyás, H.; Fernández, E. *Angew. Chem. Int. Ed.* **2011**, *50*, 7158.
- ⁴¹ Gonzalez-Bobes, F.; Fu, G. C. *J. Am. Chem. Soc.* **2006**, *128*, 5360.
- ⁴² Rudolph, A.; Lautens, M. *Angew. Chem. Int. Ed.* **2009**, *48*, 2656.
- ⁴³ Carrow, B. P.; Hartwig, J. F. *J. Am. Chem. Soc.* **2011**, *133*, 2116.
- ⁴⁴ Lata, C. J.; Crudden, C. M. *J. Am. Chem. Soc.* **2010**, *132*, 131.
- ⁴⁵ Chen, H.; Hartwig, J. F. *Angew. Chem. Int. Ed.* **1999**, *38*, 3391.
- ⁴⁶ Guennouni, N.; Lhermitte, F.; Cochard, S.; Carboni, B. *Tetrahedron* **1995**, *51*, 6999.
- ⁴⁷ Pereira, S.; Srebnik, M. *J. Am. Chem. Soc.* **1996**, *118*, 909.
- ⁴⁸ Biedrzycki, M.; Scouten, W. H.; Biedrzycka, Z. *J. Organomet. Chem.* **1992**, *431*, 255.
- ⁴⁹ Falck, J. R.; Bondlela, M.; Venkataraman, S. K.; Srinivas, D. *J. Org. Chem.* **2001**, *66*, 7148-7150.
- ⁵⁰ Pei, W.; Krauss, I. J. *J. Am. Chem. Soc.* **2011**, *133*, 18514.