

# Organic & Biomolecular Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: S. Excopy, Y. Singh and A. V. Demchenko, *Org. Biomol. Chem.*, 2019, DOI: 10.1039/C9OB01610D.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

# Triflic Acid-Mediated Synthesis of Thioglycosides

Samira Escopy, Yashapal Singh, and Alexei V. Demchenko\*

Department of Chemistry and Biochemistry, University of Missouri – St. Louis, One University Boulevard, St. Louis, Missouri 63121, USA; e-mail: [demchenkoa@umsl.edu](mailto:demchenkoa@umsl.edu)

**ABSTRACT:** An efficient synthesis of thioglycosides from per-acetates in the presence of triflic acid is described. The developed protocol features high reaction rates and product yields. Some reactive sugar series give high efficiency in the presence of sub-stoichiometric TfOH in contrast to other known protocols that require multiple equivalents of Lewis acids to reach high conversion rates.


Carbohydrates are the most abundant biomolecules that play crucial roles in many biological processes and their involvement in all diseases has been proven.<sup>1</sup> However, the stereoselective synthesis of complex carbohydrates is still a challenge in glycosciences.<sup>2,3</sup> The key aspect of the oligosaccharide assembly is the attachment of various monosaccharide units via a glycosidic bond.<sup>4-7</sup> The linkage is constructed by a chemical glycosylation reaction that involves a nucleophilic displacement of a leaving group of the glycosyl donor with a hydroxyl moiety of the glycosyl acceptor in the presence of an activator is employed.<sup>8</sup>

Since their invention in 1909 by Fisher,<sup>9</sup> thioglycosides have become key building blocks both for modification of monosaccharides and for construction of glycans.<sup>10-15</sup> Numerous methods for the preparation of thioglycosides have been established.<sup>16-27</sup> The most commonly employed pathway is Lewis acid-mediated thioglycosidation of per-acetylated sugars in the presence of stoichiometric amounts (2-5 equiv) of TMSOTf, BF<sub>3</sub>•Et<sub>2</sub>O, ZrCl<sub>4</sub>, SnCl<sub>4</sub>, etc.<sup>28-30</sup> A variety of other approaches such as one-pot acetylation-thioglycosidation of unprotected sugars,<sup>17,26,27,31,32</sup> including Bronsted-acid-mediated reactions are also known.<sup>33</sup> Although many of these conditions provide good stereoselectivity and yields, excess of promoter requirement indicates the limitation of this methodology. Herein we report that even a sub-stoichiometric amount of triflic acid can promote thioglycosidation of per-acetates of different sugar series.

Our first attempts to standardize the reaction conditions involved common glucose pentaacetate **1** that was set to react with 2.0 equiv of ethanethiol in dichloromethane at 0 °C. The key results of this study are surveyed in Table 1. Thus, when catalytic amount of triflic acid (0.2 equiv) was applied at 0 °C, thioglycoside **2**<sup>34,35</sup> was afforded in 26% yield in 4 h (entry 1). Prolonged experiments showed that the starting material did not consume even after 24 h, and the product yield remained practically the same. Increasing the triflic acid to 50 mol % produced thioglycoside **2** in a respectable yield of 70% in 3 h (entry 2). These experiments were started at 0 °C, and the

temperature was allowed to gradually increase after 1 h. Further increment in the amount of triflic acid to 80 mol % produced thioglycoside **2** in an excellent yield of 94% in 1 h (entry 3). Since many other thioglycosylations demand low temperature to maintain the stereoselectivity we investigated the temperature effect on the outcome of the TfOH-promoted reaction. When essentially the same reaction in the presence of 0.8 equiv of TfOH was set at rt, excellent yield of 97% albeit compromised stereoselectivity ( $\alpha/\beta = 1/5.0$ ) were achieved (entry 4). This result confirms that low reaction temperature is required for maintaining complete  $\beta$ -stereoselectivity of TfOH-catalyzed thioglycosidation of **1**.














**Table 1. Optimization of the reaction conditions for thioglycosidation of penta-acetate **1** with ethanethiol**









					
Entry	T °C	Catalyst (equiv)	Time	Yield of <b>2</b> , Ratio $\alpha/\beta$	
1	0→rt	TfOH (0.2)	4 h	26%, $\beta$ only	
2	0→rt	TfOH (0.5)	3 h	70%, $\beta$ only	
3	0	TfOH (0.8)	1 h	94%, $\beta$ only	
4	rt	TfOH (0.8)	45 min	97%, 1/5.0	
5	0	TfOH (1.0)	35 min	87%, $\beta$ only	
6	0	TfOH (1.2)	20 min	75%, $\beta$ only	
7	0→rt	BF <sub>3</sub> •OEt <sub>2</sub> (0.8)	24 h	55%, $\beta$ only	

Further increase in the amount of triflic acid to 1.0 and 1.2 equiv reduced the reaction time to 35 and 20 min, but the yields declined to 87 and 75%, respectively (entries 5 and 6). We hence concluded that the reaction in the presence of 0.8 equiv TfOH (entry 3) offers the most advantageous combination of the reaction efficiency and rate. For comparison, when the same amount of boron trifluoride diethyl etherate (BF<sub>3</sub>•Et<sub>2</sub>O) was used, thioglycoside **2** was obtained in a modest yield of 55% after 24 h (entry 7). This result is consistent with a common method of thioglycosylation that demands excess BF<sub>3</sub>•Et<sub>2</sub>O to drive this reaction to completion.<sup>21,30</sup>

Having standardized the reaction conditions for the synthesis of ethylthio glucoside **2**, we moved to expand the scope of this reaction to other sugar series and other aglycone types. The key results of this study are summarized in Table 2. Thus, the reaction of galactose pentaacetate **3** with ethanethiol in the presence of 0.8 equiv of TfOH afforded thiogalactoside **4**<sup>36,37</sup> in 90% yield in 30 min (entry 1). Expectedly, the reaction with much less reactive mannose penta-acetate **5** was slow, and ethylthio mannoside **6**<sup>26,37</sup> was obtained in only 26% yield. To achieve the preparative outcome of this reaction, the amount of TfOH was increased to 2.0 equiv. In this case, thiomannoside **6** was isolated in a respectable yield of 85% (entry 2). Even under these fortified conditions, the reaction remained fairly sluggish and required 8 h to complete. Also 2-phthalimido glucose tetra-acetate **7**<sup>38</sup> required similar reaction conditions (2.0 equiv of TfOH) to produce the corresponding ethylthio glycoside **8**<sup>38,39</sup> in 73% yield ( $\beta$ -only) in 4 h. When the amount of TfOH was increased to 2.5 equiv, this reaction produced thiogalactoside **8** in 96% in 45 min (entry 3). Per-acetylated sialic acid **9**<sup>40</sup> produced the corresponding ethylthio sialoside **10**<sup>41,42</sup> in 70% yield ( $\alpha/\beta = 1/1.0$ ) in 45 min (entry 4). Our standard reaction conditions (0.8 equiv of TfOH) were also employed for thioglycosidation of lactose octa-acetate **11**<sup>43-45</sup> to produce thiolactoside **12**<sup>46</sup> in 70% yield in 6 h (entry 5).

**Table 2.** TfOH-promoted thioglycosidation of acetylated hexoses **1**, **3**, **5**, **7**, sialic acid **9**, and lactose **11**

			
Entry	Substrate	Conditions 0 $\rightarrow$ rt	Product (yield, $\alpha/\beta$ )
1		EtSH, TfOH (0.8 equiv), 30 min	 <b>4</b> (90%, $\beta$ only)
2		EtSH, TfOH (2.0 equiv), 8 h	 <b>6</b> (85%, $\alpha$ only)
3		EtSH, TfOH (2.5 equiv), 45 min	 <b>8</b> (96%, $\beta$ only)
4		EtSH, TfOH (2.0 equiv), 45 min	 <b>10</b> (70%, 1/1.0)
5		EtSH, TfOH (0.8 equiv), 6 h	 <b>12</b> (70%, $\beta$ only)
6		PhSH, TfOH (0.8 equiv), 1.5 h	 <b>13</b> (77%, $\beta$ only)

7	3	PhSH, TfOH (0.8 equiv), 30 min	 <b>14</b> (88%, $\beta$ only)
8	5	PhSH, TfOH (2.0 equiv), 9 h	 <b>15</b> (75%, $\alpha$ only)
9	9	PhSH, TfOH (2.0 equiv), 45 min	 <b>16</b> (66%, 1/2.0)
10	1	<i>p</i> - thiocresol, TfOH (1.0 equiv), 30 min	 <b>17</b> (88%, $\beta$ only)
11	3	<i>p</i> - thiocresol, TfOH (1.0 equiv), 30 min	 <b>18</b> (87%, $\beta$ only)
12	5	<i>p</i> - thiocresol, TfOH (1.0 equiv), 2.5 h	 <b>19</b> (88%, $\alpha$ only)
13	9	<i>p</i> - thiocresol, TfOH (2.0 equiv), 30 min	 <b>20</b> (85%, 1/4.0)
14	11	<i>p</i> - thiocresol, TfOH (1.0 equiv), 4 h	 <b>21</b> (80%, 1/10.0)





We then investigated glycosylation of other common thiols, thiophenol and *p*-thiocresol, to generate SPh and STol glycosides, respectively. Glucose per-acetate **1** smoothly reacted with thiophenol (2.0 equiv) under the standard conditions in the presence of 0.8 equiv TfOH at 0 °C. As a result, the desired phenylthio glucoside **13**<sup>34</sup> was obtained in 77% yield in 1.5 h (entry 6). Galactose per-acetate **3** very readily reacted with thiophenol under these conditions affording phenylthio galactoside **14**<sup>18,37</sup> in 88% yield in 30 min (entry 7). Thioglycosidation of mannose per-acetate **5** again required excess TfOH because the reaction under the standard conditions yielded only 31% of thiomannoside **15**.<sup>18,47</sup> In contrast, when this reaction was repeated in the presence of 2.0 equiv TfOH, phenylthio mannoside **15** was obtained in 75% yield (entry 8). As in case of ethanethiol, the reaction was sluggish and required 9 h to complete. The introduction of the SPh anomeric group to sialic acid also required 2.0 equiv of TfOH, but it was rather swift (45 min). As a result, phenylthio sialoside **16**<sup>48,49</sup> was obtained in 66% yield as an anomeric mixture ( $\alpha/\beta = 1/2.0$ , entry 9).

First thioglycosylations with *p*-thiocresol showed that the standard conditions provide somewhat lower efficiency than that seen for reaction with EtSH and PhSH. For instance, the reaction of glucose per-acetate **1** with *p*-thiocresol (2.0 equiv) in the presence of 0.8 equiv TfOH at 0 °C provided STol glucoside **17**<sup>26,34,50</sup> in a modest yield of 70%. The utility of this reaction was enhanced by increasing the amount of TfOH to stoichiometric (1.0

equiv). We have also observed that these reactions can be successfully performed at rt. When glucose penta-acetate **1** was reacted with *p*-thiocresol under these modified conditions, thioglycoside **17** was obtained in 88% yield in 30 min (entry 10). Galactose per-acetate **3** also very readily reacted with *p*-thiocresol affording tolylthio galactoside **18**<sup>50,51</sup> in 87% yield in 30 min (entry 11). Even thioglycosidation of mannose per-acetate **5** was very efficient under these conditions producing tolylthio mannoside **19**<sup>27,33</sup> in 88% yield in 2.5 h (entry 12). The introduction of the STol anomeric group to sialic acid **9** under these reaction conditions produced moderate efficiency for the synthesis of tolylthio sialoside **20**<sup>33</sup> (69%,  $\alpha/\beta = 1/3.0$ ), but it was rather swift (45 min). When this coupling was performed in the presence of excess TfOH (2.0 equiv) at rt, tolylthio sialoside **20** was obtained both in a higher yield and higher stereoselectivity (85%,  $\alpha/\beta = 1/4.0$ , entry 13). Also, lactose octa-acetate **11** reacted smoothly under similar conditions (1.0 equiv of TfOH at rt) producing tolylthio lactoside **21**<sup>33</sup> in 4 h in a good yield of 80% ( $\alpha/\beta = 1/10.0$ , entry 14).

We then briefly investigated a possibility of expanding this methodology to the synthesis glycosyl thioimidates that found some synthetic utility in the recent years. The synthesis *S*-thiazoliny (STaz) imidate, was also deemed possible, but required excess of both HSTaz and TfOH, up to 3.5 equiv each. The key results of this study are summarized in Table 3. Glucose per-acetate **1** produced the desired thioimide **22**<sup>34,52</sup> in 76% yield in 6 h (entry 1). Galactose per-acetate **3** afforded STaz galactoside **23**<sup>52</sup> in 76% yield in 5 h (entry 2). Reaction of mannose per-acetate **5** again required longer reaction time (16 h), but we managed to obtain STaz mannoside **24**<sup>52</sup> in 66% yield (entry 3). It should be noted that all of these reactions were completely stereoselective (Table 3), whereas previous syntheses of STaz imidates from per-acetates in the presence of excess  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  at high temperature often led to anomeric mixtures.<sup>52</sup>

**Table 3.** TfOH-promoted synthesis of STaz imidates 22–24 from per-acetates

			
Entry	Substrate	Time	Product (yield, $\alpha/\beta$ )
1	<b>1</b>	6 h	 <b>22</b> (76%, $\beta$ only)
2	<b>3</b>	5 h	 <b>23</b> (76%, $\beta$ only)
3	<b>5</b>	16 h	 <b>24</b> (66%, $\alpha$ only)

In conclusion, we developed a simple methodology for the preparation of thioglycosides promoted by triflic acid. Many reactions still required stoichiometric TfOH, typical range from 0.8 equiv for SET introduction to 3.5 equiv for STaz imidate synthesis. Our initial attempts to lower the amount of TfOH led to sluggish reactions (16–24 h or longer) and modest yields due to the inability to fully consume the starting material. The scope of this approach was investigated and found to be consistently effective for the synthesis of various thioglycosides in application to different sugar series. Complete stereoselectivity, high yields, and relatively fast reaction rates have been achieved. We have also demonstrated the compatibility of the developed protocol to multi-gram preparation of thioglycosides (see the SI for details). We have also explored a possibility of conducting this reaction in the absence of molecular sieves. While most reactions were successful even without molecular sieves, the reaction yields were generally 10–20% lower due to competing hydrolysis leading to unreactive hemiacetal/hemiketal side products.

## EXPERIMENTAL SECTION

**A general procedure for thioglycosidation of per-acetylated compounds **1**, **3**, **5**, **7**, **9** and **11**** (see the SI for further details). TfOH (0.8–3.5 equiv) was added dropwise to a mixture containing a thiol (2.0 equiv or 3.5 equiv for HSTz), per-acetate (1.0 equiv) in anhydrous  $\text{CH}_2\text{Cl}_2$  (10 mL per gram of per-acetate) and freshly activated molecular sieves (3 Å) and the resulting mixture was stirred under argon for the time and at the temperature specified in tables. After that, the reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$  and subjected to conventional aqueous work-up. The organic layer was separated, dried, and concentrated under reduced pressure. The residue was purified either by crystallization ( $\text{Et}_2\text{O}$ -hexanes, mostly with Glc and Gal derivatives) or by column chromatography for most Man and aminosugar derivatives ( $\text{EtOAc}$ -hexanes gradient elution) to give the target thioglycosides. Anomeric ratios (or anomeric purity) were determined by comparison of the integral intensities of relevant signals in  $^1\text{H}$  NMR spectra.

**Supporting Information.** Additional experimental details and complete NMR spectral data for all synthesized compounds. This material is available free of charge via the Internet at

## ACKNOWLEDGMENT

This work was supported by awards from the NSF (CHE-1800350) and NIGMS (GM120673). SE is indebted to the SACM for providing her with the graduate fellowship.

## REFERENCES

- (1) Varki, A. *Glycobiology* **2017**, 27, 3–49.
- (2) Nigudkar, S. S.; Demchenko, A. V. *Chem. Sci.* **2015**, 6, 2687–2704.
- (3) Crich, D. *Acc. Chem. Res.* **2010**, 43, 1144–1153.
- (4) Adero, P. O.; Amarasekara, H.; Wen, P.; Bohe, L.; Crich, D. *Chem. Rev.* **2018**, 118, 8242–8284.
- (5) Kulkarni, S. S.; Wang, C. C.; Sabbavarapu, N. M.; Podilapu, A. R.; Liao, P. H.; Hung, S. C. *Chem. Rev.* **2018**, 118, 8025–8104.
- (6) Nielsen, M. M.; Pedersen, C. M. *Chem. Rev.* **2018**, 118, 8285–8358.



- (7) Panza, M.; Pistorio, S. G.; Stine, K. J.; Demchenko, A. V. *Chem. Rev.* **2018**, *118*, 8105–8150.
- (8) *Handbook of chemical glycosylation: advances in stereoselectivity and therapeutic relevance*; Demchenko, A. V., Ed.; Wiley-VCH: Weinheim, Germany, 2008.
- (9) Fischer, E.; Delbrück, K. *Ber. Dtsch. Chem. Ges.* **1909**, *42*, 1476–1482.
- (10) Codee, J. D. C.; Litjens, R. E. J. N.; van den Bos, L. J.; Overkleeft, H. S.; van der Marel, G. A. *Chem. Soc. Rev.* **2005**, *34*, 769–782.
- (11) Goswami, M.; Ellern, A.; Pohl, N. L. *Angew. Chem. Int. Ed. Engl.* **2013**, *52*, 8441–8445.
- (12) Goswami, M.; Ashley, D. C.; Baik, M. H.; Pohl, N. L. *J. Org. Chem.* **2016**, *81*, 5949–5962.
- (13) Vibhute, A. M.; Dhaka, A.; Athiyarath, V.; Sureshan, K. M. *Chem. Sci.* **2016**, *7*, 4259–4263.
- (14) Lacey, K. D.; Quarels, R. D.; Du, S.; Fulton, A.; Reid, N. J.; Firesheets, A.; Ragains, J. R. *Organic letters* **2018**, *20*, 5181–5185.
- (15) Saliba, R. C.; Wooke, Z. J.; Nieves, G. A.; Chu, A. A.; Bennett, C. S.; Pohl, N. L. B. *Org. Lett.* **2018**, *20*, 800–803.
- (16) Pakulski, Z.; Pierozynski, D.; Zamojski, A. *Tetrahedron* **1994**, *50*, 2975–2992.
- (17) Agnihotri, G.; Tiwari, P.; Misra, A. K. *Carbohydr. Res.* **2005**, *340*, 1393–1396.
- (18) Weng, S.-S.; Lin, Y.-D.; Chen, C.-T. *Org. Lett.* **2006**, *8*, 5633–5636.
- (19) Dasgupta, F.; Garegg, P. J. *Acta Chem. Scand.* **1989**, *43*, 471–475.
- (20) Fried, J.; Walz, D. E. *J. Am. Chem. Soc.* **1949**, *71*, 140–143.
- (21) Ferrier, R. J.; Furneaux, R. H. *Carbohydr. Res.* **1976**, *52*, 63–68.
- (22) Ferrier, R. J.; Furneaux, R. H.: 1,2-trans-1-Thioglycosides. In *Methods in Carbohydrate Chemistry*; Whistler, R. L., BeMiller, J. N., Eds.; Academic Press: New York - London, 1980; Vol. 8; pp 251–253.
- (23) Pozsgay, V.; Jennings, H. J. *Tetrahedron Lett.* **1987**, *28*, 1375–1376.
- (24) Bennet, S.; von Itzstein, M.; Kiefel, M. J. *Carbohydr. Res.* **1994**, *259*, 293–299.
- (25) Li, P.; Sun, L.; Landry, D. W.; Zhao, K. *Carbohydr. Res.* **1995**, *275*, 179–184 and references therein.
- (26) Mukhopadhyay, B.; Kartha, K. P. R.; Russell, D. A.; Field, R. A. *J. Org. Chem.* **2004**, *69*, 7758–7760.
- (27) Chatterjee, D.; Paul, A.; Rajkamal, R.; Yadav, S. *RSC Adv.* **2015**, *5*, 29669–29674.
- (28) Garegg, P. J. *Adv. Carbohydr. Chem. Biochem.* **1997**, *52*, 179–205.
- (29) Oscarson, S.: Thioglycosides. In *Carbohydrates in Chemistry and Biology*; Ernst, B., Hart, G. W., Sinay, P., Eds.; Wiley-VCH: Weinheim, New York, 2000; Vol. 1; pp 93–116.
- (30) Zhong, W.; Boons, G.-J.: Glycoside synthesis from 1-sulfur/selenium-substituted derivatives: thioglycosides in oligosaccharide synthesis. In *Handbook of Chemical Glycosylation*; Demchenko, A. V., Ed.; Wiley-VCH: Weinheim, Germany, 2008; pp 261–303.
- (31) Tai, C. A.; Kulkarni, S. S.; Hung, S. C. *J. Org. Chem.* **2003**, *68*, 8719–8722.
- (32) Huang, L. C.; Liang, P. H.; Liu, C. Y.; Lin, C. C. *J. Carbohydr. Chem.* **2006**, *25*, 303–313.
- (33) Chao, C.-S.; Chen, M.-C.; Lin, S.-C.; Mong, K.-K. T. *Carbohydr. Res.* **2008**, *343*, 957–964.
- (34) Parameswar, A. R.; Imamura, A.; Demchenko, A. V.: Synthesis of thioglycosides and thioimides from peracetates. In *Carbohydrate Chemistry: Proven Synthetic Methods*; Kovac, P., Ed.; CRC Press, 2012; Vol. 1; pp 187–196.
- (35) Contour, M. O.; Defaye, J.; Little, M.; Wong, E. *Carbohydr. Res.* **1989**, *193*, 283–287.
- (36) Ibatullin, F. M.; Selivanov, S. I.; Shavva, A. G. *Synthesis* **2001**, 419–422.
- (37) Valerio, S.; Iadonisi, A.; Adinolfi, M.; Ravida, A. *J. Org. Chem.* **2007**, *72*, 6097–6106.
- (38) Landström, J.; Bergström, M.; Hamark, C.; Ohlson, S.; Widmalm, G. *Org. Biomol. Chem.* **2012**, *10*, 3019–3032.
- (39) McGeary, R. P.; Wright, K.; Toth, I. *J. Org. Chem.* **2001**, *66*, 5102–5105.
- (40) Marra, A.; Sinay, P. *Carbohydr. Res.* **1989**, *190*, 317–322.
- (41) Tsvetkov, Y. E.; Nifantiev, N. E. *Synlett* **2005**, *9*, 1375–1380.
- (42) Marra, A.; Sinay, P. *Carbohydr. Res.* **1989**, *187*, 35–42.
- (43) Hudson, C. S.; Johnson, J. M. *J. Am. Chem. Soc.* **1915**, *37*, 1270–1275.
- (44) Hronowski, L. J. J.; Szarek, W. A.; Hay, G. W.; Krebs, A.; Depew, W. T. *Carbohydr. Res.* **1989**, *190*, 203–218.
- (45) Hou, S. J.; Saksena, R.; Kováč, P. *Carbohydr. Res.* **2008**, *343*, 196–210.
- (46) Niggemann, J.; Kamerling, J. P.; Vliegenthart, J. F. G. *J. Chem. Soc., Perkin Trans. 1* **1998**, 3011–3020.
- (47) Ahmadipour, S.; Pergolizzi, G.; Rejzek, M.; Field, R. A.; Miller, G. *J. Org. Lett.* **2019**, *21*, 4415–4419.
- (48) Parameswar, A. R.; Mueller, D.; Liu, L.; Meo, C. D.; Demchenko, A. V.: Synthesis of thioglycosides and thioimides from glycosyl halides. In *Carbohydrate Chemistry: Proven Synthetic Methods*; Kovac, P., Ed.; CRC Press, 2012; Vol. 1; pp 181–186.
- (49) Kirchner, E.; Thiem, F.; Dermick, R.; Heukeshoven, J.; Thiem, J. *J. Carbohydr. Chem.* **1988**, *7*, 453–486.
- (50) Kondo, H.; Aoki, S.; Ichikawa, Y.; Halcomb, R. L.; Ritzen, H.; Wong, C. H. *J. Org. Chem.* **1994**, *59*, 864–877.
- (51) Panza, M.; Mannino, M. P.; Baryal, K. N.; Demchenko, A. V.: A facile synthesis and characterization of p-tolyl 2,3,4,6-tetra-O-benzoyl-1-thio-β-D-galactopyranoside. In *Carbohydrate Chemistry: Proven Synthetic Methods*; Wrodnigg, T., Ed.; CRC Press, 2019; Vol. 5; pp in press.
- (52) Pornsuriyasak, P.; Demchenko, A. V. *Chem. Eur. J.* **2006**, *12*, 6630–6646.