ORIGINAL PAPER



Reactivity of substrates with multiple competitive reactive sites toward NBS under neat reaction conditions promoted by visible light

Blaž Grjol¹ · Marjan Jereb²

Received: 23 December 2020 / Accepted: 21 May 2021 $\ensuremath{\mathbb{C}}$ The Author(s) 2021

Abstract

Regioselectivity of visible-light-induced transformations of a range of (hetero)aryl alkyl-substituted ketones bearing several competitive reactive sites (α -carbonyl, benzyl and aromatic ring) with *N*-bromosuccinimide (NBS) was studied under solvent-free reaction conditions (SFRC) and in the absence of inert-gas atmosphere, radical initiators and catalysts. An 8-W energy-saving household lamp was used for irradiation. Heterogeneous reaction conditions were dealt with throughout the study. All substrates were mono- or dibrominated at the α -carbonyl position, and additionally, some benzylic or aromatic bromination was observed in substrates with benzylic carbon atoms or electron-donating methoxy groups, respectively. Surprisingly, *ipso*-substitution of the acyl group with a bromine atom took place with (4-methoxynaphthyl) alkyl ketones. While the addition of the radical scavenger TEMPO (2,2,6,6-tetramethylpiperidin-1-yloxy) decreased the extent of α - and ring bromination, it completely suppressed the benzylic bromination and α,α -dibromination with NBS under SFRC.

Keywords Halogenation · Photochemistry · Solvent-free reaction conditions · NBS · Regioselectivity

Introduction

Solvent-free reaction conditions (SFRC) (Tanaka and Toda 2000; Metzger 1998) are alternative reaction conditions to the transformations in solution (Reichardt 2007) and attracted considerable attention due to important advantages. Shorter reaction times, lower consumption of organic solvents, simple isolation/purification procedures and less reaction waste are some of notable advantages. Nonconventional reaction media, such as water, supercritical CO_2 or ionic liquids, are also a promising replacement for organic solvents (Adams et al. 2004). Another alternative is to perform reactions under highly concentrated reaction conditions (Walsh et al. 2007) or in the presence of a small amount of solvent vapor (Nakamatsu et al. 2005). A great number of different

In memory of Dr. Miha Tišler (1926–2021), Professor Emeritus of University of Ljubljana.

Marjan Jereb marjan.jereb@fkkt.uni-lj.si reactions under (SFRC) have been reviewed in both older and more recent literature (Cave et al. 2001; Martins et al. 2009; Singh and Chowdhury 2012; Gawande et al. 2014; Obst and König 2018; Zagande and Patil 2019).

N-bromosuccinimide (NBS) is one of the most common and versatile reagents in organic chemistry (Djerassi 1948; Paquette et al. 2009). It is easy to handle, inexpensive, readily available, non-toxic, and usually more selective than bromine. Traditional methods of bromination include molecular bromine or bromides under harsh reaction conditions, even though unconventional methods, for instance, in water (Shaw et al. 1997; Tanaka et al. 1999) or under SFRC (Toda and Schmeyers 2003; Pravst et al. 2009; Wang and Gao 2012), have been developed. In a more alternative approach, transformations with NBS are performed under microwave and ultrasound irradiation (Heropoulos et al. 2007), on water (Podgoršek et al. 2006), in ionic liquids (Togo and Hirai 2003; Ganguly et al. 2005), scCO₂ (Tanko and Blackert 1994) and bio-waste extract (Appa et al. 2019). Under SFRC, aromatic (Prakash et al. 2004; Imanzadeh et al. 2006), Wohl-Ziegler (Winkler et al. 2014) and α -bromination of carbonyl compounds (Kumar et al. 2012; Pravst et al. 2008) have been reported, but these methods still require the use of acid catalysts or radical initiators. While more activated substrates react in the absence of catalysts

¹ Present Address: Kilobaser GmbH, Reininghausstrasse 13a, 8020 Graz, Austria

² Faculty of Chemistry and Chemical Technology, University of Ljubljana, Ljubljana, Večna pot 113, 1000, Slovenia

or initiators (Rahman et al. 2005; Pravst et al. 2006a, b, 8; Bose and Mal 2014), other methods make the use of alternative modes of activation, such as microwave irradiation (Goswami et al. 2004) or concentrated solar radiation (Dinda et al. 2013; Deshpande et al. 2015). Photo-induced transformations with NBS have been extensively studied in solvents (Arbuj et al. 2007; Šterk et al. 2013; Cantillo et al. 2014); however, to the best of our knowledge there are rare reports on these reactions under SFRC (Jereb et al. 2009).

Molecular motion in solvent-free reactions is restricted, and the aggregate state of the reaction mixture plays an important role in SFRC transformations. Whereas molecules are more mobile in liquid/solid systems, between two solid reagents, the formation of a melt phase is crucial for increased molecular mobility (Rothenberg et al. 2001). More so than the aggregate state, the structure of the substrates largely affects the reactivity and selectivity of transformations. We have previously shown that a 40-W tungsten lamp promotes benzylic-, ring- or α -carbonyl bromination by NBS under SFRC (Jereb et al. 2009). One might wonder how selective bromination under visible-light conditions would be when a competitive aromatic-, radical- and α -carbonyl substitution was possible in a single substrate. Studies addressing this issue have been reported (Pravst et al. 2006a, b, 47; Podgoršek et al. 2009; Vražič et al. 2013), but in those cases regioselectivity of functionalization was modulated through careful tuning of the reaction conditions. To our knowledge, regioselectivity of visible-light-induced transformations of substrates containing several competitive reactive sites with NBS under SFRC remains unexplored.

In the present study, we investigated photochemical transformations of diverse aryl-substituted ketones with NBS under solvent-free reaction conditions: with an 8-W energy-saving lamp as an activation source, in the absence of any solvents, radical initiators and catalysts and in the presence of air. For testing the substrate scope, we selected a broad range of liquid and solid ketones that contained different functional groups, which could have an influence on the type of transformation.

Experimental

General information

Reagents and solvents were obtained from commercial sources. NBS was recrystallized from a mixture of $CH_2Cl_2/$ petroleum ether, and solvents were used as received. All ketones were prepared by the method of the classical Friedel–Crafts acylation (Vogel et al. 1989). **14** was prepared according to the published procedure (Coan and Becker 1955).

Reactions were performed in closed flasks without inert atmosphere. The experimental procedure was the same for liquid and solid ketones: NBS was added to the substrate and the reaction mixture was stirred with a small stirring bar under solvent-free reaction conditions (SFRC). Reaction mixtures with solid ketones turned into a molten state and the reaction proceeded as efficiently as the reactions with liquid substrates. An 8-W energy-saving lamp was used throughout all the experiments, and the average reaction temperature was 27 °C. The experiments in the dark were carried out at 27 °C. The progress of the reactions was monitored by thin-layer chromatography (TLC). Light source: Philips Genie 8 W energy-saving light bulb. Column chromatography (CC): silica gel 60 (63-200 mm, 70-230 mesh ASTM; Fluka); CH₂Cl₂/petroleum ether (1:3). Preparative thin-layer chromatography (TLC): Merck-60-F₂₅₄ plates; mixtures of light petroleum ether (b.p. 40-60 °C) and CH₂Cl₂. NMR spectra: Bruker DPX 300 (300 MHz) and Bruker AVANCE III (500 MHz) instruments; in $CDCl_2$; δ in ppm relative to TMS as internal standard for ¹H NMR and to the central line of CDCl₃ (δ = 77.00 ppm) for ¹³C NMR, J in Hz. IR spectra: Bruker Alpha FT-IR spectrometer; in cm⁻¹. HRMS: Agilent 6224 Accurate Mass TOF LC/MS system spectrometer; electrospray ionization; in m/z. Elemental analysis: Perkin Elmer CHNS/O Analyzer 2400 Series II elemental analyzer. Melting points: Büchi 535 melting point apparatus; uncorrected.

General procedure for photo-induced reactions with NBS under SFRC

To a ketone (0.3 mmol), NBS (0.33 or 0.66 mmol, 20 or 39 mg) was added and the resulted reaction mixture was stirred under illumination with an 8-W energy-saving light bulb for 19–29 h. At the end of the reaction, conversions and the product distribution were determined by ¹H NMR spectroscopy of the crude reaction mixture. Pure products were obtained after separation by either CC or preparative TLC and were analyzed by ¹H NMR, ¹³C NMR, IR, HRMS and melting point when solid. Spectroscopic data of the known compounds are available in Supporting Information and are in accordance with the literature data.

Characterization of the new products

2-Bromo-1-(4-(isopropyl)phenyl)ethanone 2c.

1c (0.049 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel (R_f =0.55, CH₂Cl₂/petroleum ether 1:3 → 2:1). Colorless oil (0.028 g, 39%). IR (neat): ν 2961, 1676, 1604, 1569, 1461, 1416, 1310, 1278, 1198, 1186, 1056, 1009, 990, 846, 822, 677, 661 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 1.28 (d, J=6.9 Hz, 6H), 2.98 (sept,

 $J=6.9 \text{ Hz}, 1\text{H}), 4.44 (s, 2\text{H}), 7.35 (d, J=8.3 \text{ Hz}, 2\text{H}), 7.93 (d, J=8.3 \text{ Hz}, 2\text{H}). {}^{13}\text{C} \text{ NMR} (125 \text{ MHz}, \text{CDCl}_3): \delta 23.6, 30.9, 34.3, 127.0, 129.2, 131.8, 155.7, 190.9. \text{ HRMS} (ESI-TOF,$ *m/z*): found 241.0223 ([*M*+H]⁺, C₁₁H₁₄BrO⁺; calc. 241.0223).

2-Bromo-1-(3-methyl-4-methoxyphenyl)ethanone 2e.

1e (0.049 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel (R_f =0.47, CH₂Cl₂/petroleum ether 1:3 → 2:1). Colorless solid (0.013 g, 18%). M.p. 45–48 °C. IR (neat): ν 2941, 1683, 1596, 1497, 1438, 1410, 1250, 1212, 1126, 1020, 878, 818, 752, 682, 611 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 2.26 (*s*, 3H), 3.91 (*s*, 3H), 4.40 (*s*, 2H), 6.87 (*d*, *J*=8.6 Hz, 1H), 7.80 (*d*, *J*=1.7 Hz, 1H), 7.86 (dd, *J*=1.7, 8.6 Hz, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 16.3, 30.8, 55.6, 109.4, 126.4, 127.3, 129.3, 131.5, 162.4, 190.2. HRMS (ESI-TOF, *m/z*): found 243.0013 ([*M*+H]⁺, C₁₀H₁₂BrO₂⁺; calc. 243.0015).

2-Bromo-1-(3,5-dimethyl-4-methoxyphenyl)ethanone 2f.

If (0.053 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel (R_f =0.38, CH₂Cl₂/petroleum ether 1:3→2:1). Pale yellowish solid (0.015 g, 19%). M.p. 43–45 °C. IR (neat): ν 2948, 1682, 1595, 1483, 1387, 1313, 1256, 1229, 1140, 1055, 1007, 887, 856, 769, 708, 641 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 2.34 (s, 6H), 3.77 (s, 3H), 4.41 (s, 2H), 7.67 (s, 2H). ¹³C NMR (125 MHz, CDCl₃): δ 16.3, 30.9, 59.7, 129.5, 130.0, 131.6, 162.0, 190.7. HRMS (ESI-TOF, *m/z*): found 257.0172 ([*M*+H]⁺, C₁₁H₁₄BrO₂⁺; calc. 257.0172).

2-Bromo-1-(4-(1-bromoethyl)phenyl)ethanone 3b.

1b (0.044 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Preparative TLC (R_f =0.45, CH₂Cl₂/petroleum ether 1:3). Brownish solid (0.002 g, 4%). M.p. 48–53 °C. IR (neat): ν 2948, 1694, 1603, 1571, 1412, 1388, 1280, 1197, 1162, 1069, 1042, 989, 822, 681 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 2.05 (d, J=6.9 Hz, 3H), 4.42 (s, 2H), 5.20 (q, J=6.9 Hz, 1H), 7.52–7.59 (m, 2H), 7.94–8.00 (m, 2H). ¹³C NMR (75 MHz, CDCl₃): δ 26.4, 30.6, 47.5, 127.4, 129.4, 133.7, 149.1, 190.6. HRMS (ESI-TOF, m/z): found 304.9168 ([M+H]⁺, C₁₀H₁₁Br₂O⁺; calc. 304.9171).

2,2-Dibromo-1-(4-(isopropyl)phenyl)ethanone 4c.

1c (0.049 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel (R_f = 0.68, CH₂Cl₂/ petroleum ether 1:3 → 2:1). Yellowish solid (0.011 g, 21%). M.p. 40–45 °C. IR (neat): ν 2961, 1684, 1599, 1567, 1460, 1414, 1311, 1270, 1223, 1198, 1055, 983, 850, 840,

770, 742, 709, 682 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 1.29 (*d*, *J* = 6.9 Hz, 6H), 2.99 (sept, *J* = 6.9 Hz, 1H), 6.69 (*s*, 1H), 7.36 (*d*, *J* = 8.4 Hz, 2H), 8.02 (*d*, *J* = 8.4 Hz, 2H). ¹³C NMR (75 MHz, CDCl₃): δ 23.5, 34.4, 39.8, 127.1, 128.5, 130.0, 156.3, 185.6. HRMS (ESI-TOF, *m/z*): found 318.9330 ([*M*+H]⁺, C₁₁H₁₄Br₂O⁺; calc. 318.9328).

2,2-Dibromo-1-(4-(cyclohexyl)phenyl)ethanone 4d.

1d (0.061 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel (R_f = 0.75, CH₂Cl₂/ petroleum ether 1:3 → 2:1). Colorless solid (0.003 g, 5%). M.p. 100–104 °C. IR (neat): ν 3027, 2920, 2848, 1683, 1600, 1566, 1445, 1416, 1315, 1273, 1212, 1191, 1145, 988, 841, 774, 707, 630 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 1.21–1.32 (m, 1H), 1.35–1.49 (m, 4H), 1.74–1.81 (m, 1H), 1.82–1.93 (m, 4H), 2.54–2.63 (m, 1H), 6.70 (s, 1H), 7.34 (d, J = 8.3 Hz, 2H), 8.01 (d, J = 8.3 Hz, 2H). ¹³C NMR (125 MHz, CDCl₃): δ 25.9, 26.6, 33.9, 39.9, 44.8, 127.5, 128.4, 129.9, 155.5, 185.6. HRMS (ESI-TOF, m/z): found 358.9634 ($[M + H]^+$, C₁₄H₁₇Br₂O⁺; calc. 358.9641).

2,2-Dibromo-1-(3-methyl-4-methoxyphenyl)ethanone 4e.

1e (0.049 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel (R_f =0.68, CH₂Cl₂/petroleum ether 1:3 → 2:1). Colorless solid (0.015 g, 29%). M.p. 113–115 °C. IR (neat): ν 3033, 2923, 1665, 1591, 1503, 1455, 1438, 1416, 1259, 1217, 1127, 1022, 987, 908, 814, 761, 717, 680, 615 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 2.26 (*s*, 3H), 3.92 (*s*, 3H), 6.70 (*s*, 1H), 6.88 (*d*, *J*=8.7 Hz, 1H), 7.88 (*d*, *J*=2.1 Hz, 1H), 7.97 (dd, *J*=2.1, 8.7 Hz, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 16.3, 40.0, 55.7, 109.5, 122.8, 127.6, 130.1, 132.2, 162.9, 184.9. HRMS (ESITOF, *m/z*): found 320.9117 ([*M*+H]⁺, C₁₀H₁₁Br₂O₂⁺; calc. 320.9120).

2,2-Dibromo-1-(3,5-dimethyl-4-methoxyphenyl)ethanone **4f.**

1f (0.053 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel (R_f =0.48, CH₂Cl₂/ petroleum ether 1:3→2:1). Colorless solid (0.014 g, 25%). M.p. 85–87 °C. IR (neat): ν 2940, 1688, 1593, 1479, 1310, 1221, 1142, 994, 903, 776, 749, 715, 652 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 2.34 (*s*, 6H), 3.79 (*s*, 3H), 6.71 (*s*, 1H), 7.75 (*s*, 2H). ¹³C NMR (125 MHz, CDCl₃): δ 16.3, 39.8, 59.8, 126.2, 130.7, 131.8, 162.5, 185.4. HRMS (ESI-TOF,

m/z): found 334.9276 ($[M + H]^+$, $C_{11}H_{13}Br_2O_2^+$; calc. 334.9277).

2-Bromo-1-(4-(methylthio)phenyl)propan-1-one 6c.

5c (0.054 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel (R_f =0.60, CH₂Cl₂/hexane 1:3 → 2:1). Pale brownish solid (0.053 g, 68%). M.p. 88–90 °C. IR (neat): ν 2988, 1666, 1583, 1405, 1346, 1247, 1192, 1164, 1093, 1058, 992, 944, 830, 768, 745, 689 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 1.89 (d, J=6.6 Hz, 3H), 2.53 (s, 3H), 5.25 (q, J=6.6 Hz, 1H), 7.28 (d, J=8.5 Hz, 2H), 7.93 (d, J=8.5 Hz, 2H). ¹³C NMR (125 MHz, CDCl₃): δ 14.7, 20.1, 41.4, 125.0, 129.3, 130.1, 146.9, 192.3. HRMS (ESI-TOF, m/z): found 258.9785 ([M+H]⁺, C₁₀H₁₂BrOS⁺; calc. 258.9787).

2-Bromo-1-(5-methyl-2-methoxyphenyl)propan-1-one 6d.

5d (0.053 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel (R_f =0.59, CH₂Cl₂/hexane 1:2 → 2:1). Yellowish oil (0.056 g, 73%). IR (neat): ν 2925, 1671, 1608, 1495, 1463, 1441, 1404, 1332, 1283, 1248, 1178, 1160, 1136, 1020, 968, 809, 741, 607 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 1.85 (d, J=6.7 Hz, 3H), 2.31 (s, 3H), 3.89 (s, 3H), 5.53 (q, J=6.7 Hz, 1H), 6.86 (d, J=8.5 Hz, 1H), 7.23–7.31 (m, 1H), 7.49–7.54 (m, 1H). ¹³C NMR (75 MHz, CDCl₃): δ 20.2, 20.3, 48.0, 55.8, 111.5, 125.4, 130.4, 131.7, 134.5, 156.1, 196.6. HRMS (ESI-TOF, m/z): found 257.0164 ([M+H]⁺, C₁₁H₁₄BrO₂⁺; calc. 257.0172).

2-Bromo-1-(3-bromo-4,5-dimethoxyphenyl)propan-1-one **6e.**

5e (0.058 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel (R_f =0.42, CH₂Cl₂/petroleum ether 1:3 → 2:1). Yellowish oil (0.024 g, 42%). IR (neat): ν 2934, 1695, 1592, 1505, 1439, 1372, 1336, 1257, 1204, 1158, 1045, 1024, 972, 897, 855, 790, 760, 630 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 1.91 (d, J=6.7 Hz, 3H), 3.90 (s, 3H), 3.92 (s, 3H), 5.40 (q, J=6.7 Hz, 1H), 7.03 (s, 1H), 7.06 (s, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 19.9, 45.6, 56.2, 56.3, 110.6, 113.2, 115.9, 130.8, 148.4, 151.7, 196.6. HRMS (ESI-TOF, m/z): found 350.9225 ([M+H]⁺, C₁₁H₁₃Br₂O₃⁺; calc. 350.9226).

2-Bromo-1-(4-(bromomethyl)phenyl)propan-1-one 7a.

5a (0.044 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Preparative TLC (R_f =0.51, CH₂Cl₂/petroleum ether 1:3). Colorless solid (0.010 g, 20%). M.p. 80–83 °C. IR (neat): ν 2928, 1675, 1604, 1438, 1417, 1345, 1235, 1205, 1155, 1093, 993, 948, 857, 823, 767, 723, 702, 635 cm⁻¹. ¹H NMR (500 MHz,

CDCl₃): δ 1.90 (*d*, *J* = 6.6 Hz, 3H), 4.50 (*s*, 2H), 5.26 (*q*, *J* = 6.6 Hz, 1H), 7.51 (*d*, *J* = 8.2 Hz, 2H), 8.00 (*d*, *J* = 8.2 Hz, 2H). ¹³C NMR (125 MHz, CDCl₃): δ 20.0, 31.9, 41.4, 129.4, 129.4, 133.8, 143.4, 192.6. HRMS (ESI-TOF, *m/z*): found 304.9171 ([*M*+H]⁺, C₁₀H₁₁Br₂O⁺; calc. 304.9171).

2-Bromo-2-methyl-1-(4-methylphenyl)propan-1-one 9a.

8a (0.049 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Preparative TLC (R_f =0.64, CH₂Cl₂/petroleum ether 1:3). Pale yellowish oil (0.017 g, 23%). IR (neat): ν 2927, 1672, 1606, 1459, 1370, 1267, 1163, 1104, 978, 886, 827, 748, 677 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 2.03 (*s*, 6H), 2.41 (*s*, 3H), 7.24 (*d*, *J*=8.2 Hz, 2H), 8.07 (*d*, *J*=8.2 Hz, 2H). ¹³C NMR (125 MHz, CDCl₃): δ 21.6, 31.7, 60.5, 128.8, 130.4, 131.9, 143.2, 196.3. HRMS (ESI-TOF, *m/z*): found 241.0221 ([*M*+H]⁺, C₁₁H₁₄BrO⁺; calc. 241.0223).

2-Bromo-1-(2,4-dimethylphenyl)-2-methylpropan-1-one **9b.**

8b (0.053 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel (R_f =0.63, CH₂Cl₂/petroleum ether 1:3 → 2:1). Pale yellowish oil (0.041 g, 53%). IR (neat): ν 2972, 2924, 1688, 1612, 1458, 1385, 1369, 1257, 1126, 1102, 973, 875, 817 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 1.97 (*s*, 6H), 2.26 (*s*, 3H), 2.34 (*s*, 3H), 6.98–7.03 (*m*, 1H), 7.06 (*s*, 1H), 7.60 (*d*, *J*=7.9 Hz, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 20.1, 21.3, 30.7, 62.6, 125.5, 126.9, 131.8, 134.9, 135.9, 139.8, 202.2. HRMS (ESI-TOF, *m/z*): found 255.0379 ([*M*+H]⁺, C₁₂H₁₆BrO⁺; calc. 255.0379).

2-Bromo-1-(3-methyl-4-methoxyphenyl)-2-methylpropan-1-one **9d.**

8d (0.058 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Preparative TLC (R_f =0.40, CH₂Cl₂/petroleum ether 1:5). Yellowish oil (0.013 g, 16%). IR (neat): ν 2928, 1664, 1598, 1501, 1458, 1386, 1317, 1298, 1256, 1127, 1104, 1021, 986, 916, 817, 760, 681, 617 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 2.04 (*s*, 6H), 2.25 (*s*, 3H), 3.90 (*s*, 3H), 6.83 (*d*, *J*=8.7 Hz, 1H), 7.99 (*d*, *J*=2.0 Hz, 1H), 8.15 (dd, *J*=2.0, 8.7 Hz, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 16.3, 32.0, 55.5, 60.6, 108.7, 126.4, 126.4, 130.6, 133.1, 161.3, 195.2. HRMS (ESITOF, *m*/*z*): found 271.0329 ([*M*+H]⁺, C₁₂H₁₆BrO₂⁺; calc. 271.0328).

2-Bromo-1-(4-(bromomethyl)phenyl)-2-methylpropan-1-one **10a.**

8a (0.049 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Preparative TLC (R_f =0.48, CH₂Cl₂/petroleum ether 1:3). Yellowish solid (0.012 g, 23%). M.p. 36–38 °C. IR (neat): ν

2979, 1668, 1603, 1451, 1411, 1388, 1264, 1230, 1203, 1163, 1102, 979, 887, 849, 820, 764, 713, 679, 606 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 2.03 (*s*, 6H), 4.50 (*s*, 2H), 7.46 (*d*, *J* = 8.4 Hz, 2H), 8.13 (*d*, *J* = 8.4 Hz, 2H). ¹³C NMR (125 MHz, CDCl₃): δ 31.5, 32.1, 60.2, 128.8, 130.7, 134.6, 142.0, 196.1. HRMS (ESI-TOF, *m/z*): found 318.9329 ([*M*+H]⁺, C₁₁H₁₃Br₂O⁺; calc. 318.9328).

2-Bromo-1-(3-(bromomethyl)-4-methoxyphenyl)-2-methylpropan-1-one **10d.**

8d (0.058 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Preparative TLC (R_f =0.21, CH₂Cl₂/petroleum ether 1:5). Colorless solid (0.011 g, 18%). M.p. 79–81 °C. IR (neat): ν 2929, 1655, 1595, 1503, 1458, 1444, 1265, 1223, 1142, 1105, 1034, 1018, 1001, 937, 916, 842, 819, 763, 617 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 2.04 (*s*, 6H), 3.97 (*s*, 3H), 4.57 (*s*, 2H), 6.91 (*d*, *J*=8.8 Hz, 1H), 8.22 (*d*, *J*=2.3 Hz, 1H), 8.26 (dd, *J*=2.3, 8.8 Hz, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 28.1, 31.8, 55.9, 60.4, 110.0, 125.9, 126.8, 133.4, 133.6, 160.7, 194.5. HRMS (ESI-TOF, *m*/*z*): found 348.9432 ([*M*+H]⁺, C₁₂H₁₅Br₂O₂⁺; calc. 348.9433).

2-Bromo-1-(4-methoxyphenyl)-2-phenylethan-1-one 12b.

11b (0.068 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel (R_f =0.47, CH₂Cl₂/petroleum ether 1:3 → 2:1). Pale yellowish solid (0.055 g, 60%). M.p. 67–70 °C. IR (neat): ν 2928, 1671, 1601, 1571, 1508, 1452, 1422, 1357, 1312, 1264, 1217, 1173, 1027, 846, 824, 774, 734, 688 cm^{-1.} ¹H NMR (500 MHz, CDCl₃): δ 3.85 (*s*, 3H), 6.36 (*s*, 1H), 6.89–6.93 (*m*, 2H), 7.29–7.38 (*m*, 3H), 7.51–7.55 (*m*, 2H), 7.95–8.00 (*m*, 2H). ¹³C NMR (125 MHz, CDCl₃): δ 51.1, 55.5, 114.0, 126.9, 128.9, 129.0, 129.0, 131.5, 136.3, 163.9, 189.6. HRMS (ESI-TOF, *m/z*): found 305.0165 ([*M*+H]⁺, C₁₅H₁₄BrO₂⁺; calc. 305.0172).

2-Bromo-1-(5-methyl-2-methoxyphenyl)-2-phenyle-than-1-one **12c.**

11c (0.120 g, 0.5 mmol), NBS (0.107 g, 0.6 mmol). Column chromatography on silica gel ($R_f = 0.45$, CH₂Cl₂/hexane 1:2 \rightarrow 2:1). Yellowish solid (0.100 g, 63%). M.p. 44–46 °C. IR (neat): ν 2943, 1676, 1607, 1579, 1495, 1453, 1401, 1285, 1246, 1157, 1126, 1020, 810, 695, 626 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 2.27 (*s*, 3H), 3.85 (*s*, 3H), 6.59 (*s*, 1H), 6.81 (*d*, *J* = 8.5 Hz, 1H), 7.21–7.36 (*m*, 4H), 7.44–7.53 (*m*, 3H). ¹³C NMR (75 MHz, CDCl₃): δ 20.2, 55.1, 55.7, 111.5, 125.5, 128.6, 128.6, 129.4, 130.5, 131.9,

134.8, 136.5, 155.9, 193.9. HRMS (ESI-TOF, *m/z*): found 319.0326 ([*M*+H]⁺, C₁₆H₁₆BrO₂⁺; calc. 319.0328).

2-Bromo-2-phenyl-1-(thiophen-2-yl)ethan-1-one 12d.

11d (0.303 g, 1.5 mmol), NBS (0.320 g, 1.8 mmol). Column chromatography on silica gel (R_f =0.53, CH₂Cl₂/hexane 1:2 → 2:1). Yellow solid (0.304 g, 72%). M.p. 36–39 °C. IR (neat): ν 3089, 1660, 1514, 1494, 1454, 1408, 1353, 1285, 1237, 1222, 1178, 1062, 948, 859, 719, 691, 634 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 6.20 (*s*, 1H), 7.11 (dd, *J*=3.9, 4.8 Hz, 1H), 7.30–7.40 (*m*, 3H), 7.53–7.59 (*m*, 2H), 7.68 (dd, *J*=0.9, 4.8 Hz, 1H), 7.76 (dd, *J*=0.9, 3.9 Hz, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 51.4, 128.3, 129.0, 129.0, 129.2, 133.4, 135.2, 135.8, 140.6, 184.1. HRMS (ESI-TOF, *m*/*z*): found 280.9628 ([*M*+H]⁺, C₁₂H₁₀BrOS⁺; calc. 280.9630).

2,2-Dibromo-2-phenyl-1-(thiophen-2-yl)ethan-1-one 13d.

11d (0.303 g, 1.5 mmol), NBS (0.320 g, 1.8 mmol). Column chromatography on silica gel (R_f =0.67, CH₂Cl₂/hexane 1:2 → 2:1). Brownish solid (0.052 g, 16%). M.p. 65–69 °C. IR (neat): ν 3109, 2921, 1667, 1491, 1444, 1406, 1350, 1238, 1222, 1190, 1084, 1064, 961, 845, 801, 749, 722, 704, 689, 648 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 6.92 (dd, J=4.0, 4.9 Hz, 1H), 7.30 (dd, J=1.1, 4.0 Hz, 1H), 7.33–7.42 (m, 3H), 7.56 (dd, J=1.1, 4.9 Hz, 1H), 7.66–7.71 (m, 2H). ¹³C NMR (125 MHz, CDCl₃): δ 68.4, 127.3, 127.8, 128.8, 129.8, 135.1, 136.1, 136.3, 140.7, 180.1. HRMS (ESI-TOF, m/z): found 358.8727 ([M+H]⁺, C₁₂H₉Br₂OS⁺; calc. 358.8735).

2-Bromo-2,4-diphenyl-3-oxobutanenitrile 15.

14 (0.071 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Column chromatography on silica gel ($R_f = 0.70$, CH₂Cl₂/ petroleum ether 1:3 \rightarrow 2:1). Yellowish oil (0.070 g, 75%). IR (neat): ν 2923, 1731, 1495, 1449, 1320, 1031, 1002, 705, 691, 646 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 3.99 (d, J = 16.7 Hz, 1H), 4.03 (d, J = 16.7 Hz, 1H), 7.01–7.06 (m, 2H), 7.22–7.29 (m, 3H), 7.43–7.48 (m, 3H), 7.62–7.68 (m, 2H). ¹³C NMR (125 MHz, CDCl₃): δ 43.1, 54.1, 116.0, 127.5, 127.6, 128.6, 129.4, 129.5, 130.6, 131.3, 132.1, 190.9. HRMS (ESI-TOF, m/z): found 314.0173 ([M +H]⁺, C₁₆H₁₃BrNO⁺; calc. 314.0175).

2-Bromo-2-phenylacetonitrile 18.

16 (0.048 g, 0.3 mmol), NBS (0.059 g, 0.33 mmol). Preparative TLC (R_f =0.35, CH₂Cl₂/petroleum ether 1:3) Yellowish oil (0.031 g, 53%). IR (neat): 2971, 1496, 1455, 1188, 1138, 971, 817, 763, 690, 653, 621 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 5.49 (*s*, 1H), 7.39–7.48 (*m*, 3H), 7.52–7.60 (*m*, 2H). ¹³C NMR (75 MHz, CDCl₃): δ 27.4, 116.2, 127.8, 129.5, 130.3, 133.5.

1-(3-bromo-4-methoxynaphthalen-1-yl)ethanone 21a.

19a (0.200 g, 1.0 mmol), NBS (0.356 g, 2.0 mmol). Column chromatography on silica gel ($R_f = 0.24$, CH₂Cl₂/hexane 1:2 \rightarrow 2:1). Brownish solid (0.008 g, 3%). M.p. 36–39 °C. IR (neat): ν 2940, 2848, 1663, 1568, 1501, 1401, 1352, 1275, 1229, 1134, 1073, 1008, 974, 835, 759, 712, 675, 627 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 2.72 (s, 3H), 4.05 (s, 3H), 7.55–7.68 (m, 2H), 8.07 (s, 1H), 8.16–8.21 (m, 1H), 8.74–8.80 (m, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 29.7, 61.7, 110.8, 122.3, 126.6, 127.3, 128.6, 129.6, 131.1, 132.8, 133.6, 156.9, 199.6. HRMS (ESI-TOF, *m/z*): found 279.0013 ([*M* + H]⁺, C₁₃H₁₂BrO₂⁺; calc. 279.0015).

(3-Bromo-4-methoxynaphthalen-1-yl)(phenyl)methanone 21c.

19c (0.079 g, 0.3 mmol), NBS (0.117 g, 0.66 mmol). Column chromatography on silica gel (R_f =0.67, CH₂Cl₂/petroleum ether 1:3→2:1). Yellowish solid (0.101 g, 99%). M.p. 81–84 °C. IR (neat): ν 2922, 1649, 1567, 1501, 1448, 1372, 1270, 1243, 1152, 1070, 1054, 1022, 963, 863, 832, 763, 710, 693, 670, 656, 613 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 4.08 (*s*, 3H), 7.45–7.51 (*m*, 2H), 7.51–7.56 (*m*, 1H), 7.57–7.65 (*m*, 2H), 7.71 (*s*, 1H), 7.84–7.89 (*m*, 2H), 8.08 (*d*, *J*=8.5 Hz, 1H), 8.22 (*d*, *J*=8.4 Hz, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 61.7, 111.1, 122.4, 126.2, 127.4, 127.8, 128.6, 129.4, 130.4, 131.8, 132.1, 133.5, 133.9, 137.9, 155.6, 196.1. HRMS (ESI-TOF, *m/z*): found 341.0170 ([*M*+H]⁺, C₁₈H₁₄BrO₂⁺; calc. 341.0172).

2-Bromo-1-(3-bromo-4-methoxynaphthalen-1-yl)ethanone 22a.

19a (0.200 g, 1.0 mmol), NBS (0.356 g, 2.0 mmol). Column chromatography on silica gel (R_f =0.33, CH₂Cl₂/hexane 1:2 \rightarrow 2:1). Brownish solid (0.007 g, 2%). M.p. 64–69 °C. IR (neat): ν 2934, 1686, 1566, 1503, 1373, 1265, 1213, 1165, 1124, 1089, 1074, 973, 908, 860, 836, 789, 761, 718, 701, 680, 639 cm⁻¹. ¹H NMR (500 MHz, CDCl₃): δ 4.07 (*s*, 3H), 4.53 (*s*, 2H), 7.60–7.70 (*m*, 2H), 8.07 (*s*, 1H), 8.19–8.23 (*m*, 1H), 8.62–8.67 (*m*, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 33.3, 61.8, 110.6, 122.5, 126.3, 127.7, 129.0, 129.7, 129.8,

131.6, 133.4, 157.7, 192.4. HRMS (ESI-TOF, *m/z*): found 356.9117 ([*M*+H]⁺, C₁₃H₁₁Br₂O₂⁺; calc. 356.9120).

Results and discussion

The ketones under investigation were arranged into the groups of distinct alkyl substituents. First, we investigated the regioselectivity of functionalization at differently phenyl-functionalized model acetophenones 1a-f. Neat ketones were reacted with NBS under stirring and irradiated using an 8-W energy-saving light bulb at 27 °C. The results are summarized in Table 1.

All substrates **1a–f** responded to the above reaction conditions in the same way, mainly affording α -bromoacetophenones **2a–f**, which were accompanied by α, α -dibrominated products **4a–f**. The selectivity was poorer in the cases of 4'-methylacetophenone **1a** and 4'-ethylacetophenone **1b**, where α -bromination was accompanied with benzylic bromination of the corresponding alkyl (Me and Et) group, resulting in a small amount of **3a** and **3b**, respectively (entries 1 and 2).

Surprisingly, selective benzylic bromination of 4'-methylacetophenone 1a into 4'-bromomethylacetophenone took place with NBS in solvent-free reaction under concentrated solar radiation (Deshpande et al. 2015). A comparison of the photochemical functionalization of 4'-methylacetophenone 1a with NBS under different reaction conditions revealed interesting features. On water, 4'-bromomethylacetophenone was the only product formed (Podgoršek et al. 2009), whereas the treatment of 1a with NBS/SiCl₄ in MeCN furnished the compound 3a as the sole product (Salama and Novák 2011). Conversely, compound 1a was selectively α -brominated using NBS/PTSA under ultrasound irradiation in methanol (Adhikari and Samant, 2002). In contrast to the straight-chain functionalized derivatives, at the branched isopropyl 1c and cyclohexyl 1d acetophenones only α -bromination took place with no benzylic bromination observed (entries 4 and 5). Increasing the complexity of the substrate with additional reactive sites, e.g., in 3'-methyl-4'-methoxyacetophenone 1e and 3',5'-dimethyl-4'-methoxyacetophenone 1f, did not diminish the reaction selectivity, furnishing solely α -brominated ketones 2e/4e and 2f/4f, respectively (entries 6 and 8). Conducting the reactions with 1b and 1e in the presence of a radical scavenger TEMPO (2,2,6,6-tetramethylpiperidin-1-yloxy) strongly influenced the selectivity outcome. As indicated in entries 3 and 7, benzylic bromination and introduction of the second bromine atom to the α -position were completely suppressed; hence, the only products formed were 2b and 2e, respectively. Transformation of 1f did not take place in the absence of light (entry 9).

Table 1	Transformation of 1	neat substituted aceto	ophenones 1a-f	with NBS unde	r visible-light irradiation
	Fransision and or i	ieur substituteu ueere	phonones in i	min i up o unde	i vibiolo inglic infadiation

Entry	Substrate	Reaction conditions ^a	Products	Conv. [%] ^b	Selectivity ^b
1	0 1a	<i>hv</i> , 24h	$\begin{array}{c} & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & &$	100	2a , 52 (13) 3a , 10 (4) 4a , 38 (26)
2		<i>hv</i> , 29h	Br Br Br Br	76	2b , 60 (15) 3b , 24 (4) 4b , 16 ^d
3	1b	hv, TEMPO ^c , 24h	2b 3b 4b	20	2b , 100 ^d
4		<i>hv</i> , 21h	$\begin{array}{c} 0 \\ Br \\ Br \\ 2c \\ 4c \end{array}$	78	2c , 66 (39) 4c , 34 (21)
5	Cy 1d	<i>hv</i> , 22h	$\begin{array}{c} 0 \\ Cy \\ Cy \\ 2d \\ \end{array} \begin{array}{c} 0 \\ Br \\ Cy \\ Br \\ B$	70	2d , 89 (36) 4d , 11 (5)
6	MeO	<i>hv</i> , 24h	O Br Br	81	2e , 56 (18) 4e , 44 (29)
7	1e	hv, TEMPO ^c , 24h	MeO Bi MeO Bi	11	2e , 100 ^d
8		<i>hv</i> , 27h	O O Br	71	2f , 55 (19) 4f , 45 (25)
9	MeO´ 丫́ 1f	dark, 21h	MeO Br MeO Br 2f 4f	0	_

^aSubstrate 1 (0.3 mmol), NBS (0.33 mmol), stirring, 8-W energy-saving light bulb, reaction temperature 27 °C

^bConversions and relative distribution of products determined by ¹H NMR spectroscopy of the crude reaction mixtures; values in parentheses refer to isolated yields

^cTEMPO (0.3 equiv.) was added

^dProducts not isolated

Next, the selectivity of photo-transformations of model aryl-substituted propiophenones **5** with NBS under SFRC was examined. As with acetophenones **1**, α -bromination was the main reaction pathway, but in this case no α , α -dibrominated products could be detected. Transformations of most of the propiophenones exhibited a somewhat different selectivity compared to acetophenones. The results are summarized in Table 2.

Functionalization of 4'-methylpropiophenone **5a** yielded α -brominated ketone **6a** as the main product, accompanied with a product of mixed α - and benzylic bromination **7a** (entry 1). Benzylic bromination of **5a** in the presence of

TEMPO was completely suppressed, and only α -bromination could be detected with low conversion into **6a** (entry 2). This might indicate that the main reaction channels are of a radical nature. The reaction of **5a** did not occur in the dark (entry 3). Transformation of 4'-methoxypropiophenone **5b** gave the corresponding α -bromoketone **6b** as the main product. To some extent, this was further brominated at the aromatic ring also yielding a minor product **7b** (entry 4). The sulfur analogue, 4'-methylthiopropiophenone **5c**, exhibited the highest selectivity with only the α -carbon atom being brominated, producing **6c** (entry 5). Unexpectedly, the methylthio group did not affect the type of functionalization as no aromatic

0	5		1 1 1	0	
Entry	Substrate	Reaction conditions ^a	Products	Conv. [%] ^b	Selectivity ^b
1	o L	<i>hv</i> , 27h	o o o o er	87	6a , 61 (30) 7a , 39 (20)
2		<i>hv</i> , TEMPO ^c , 24h		25	6a , 100 ^e
3	5a	dark, 21h	6a ^{Br} 7a	0	_
4	MeO 5b	hv, 23h ^d	MeO 6b Br 7b	92	6b , 90 (68) 7b , 10 ^e
5	MeS 5c	<i>hv</i> , 20h	MeS 6c	83	100 (68)
6	OMe O 5d	<i>hv</i> , 19h	OMe O + 6d Br 7d	98	6d , 93 (73) 7d , 7 ^e
7	° (<i>hv</i> , 24h	Br Br Br	90	6e , 90 (42) 7e , 10 ^e
8	MeO OMe 5e	<i>hv</i> , TEMPO ^c , 24h	MeO OMe 6e 7e	0	-

Table 2 Regioselectivity of functionalization of neat substituted propiophenones with NBS under visible light

^aSubstrate 5 (0.3 mmol), NBS (0.33 mmol), stirring, 8-W energy-saving light bulb, reaction temperature 27 °C

^bConversions and relative distribution of products determined by ¹H NMR spectroscopy of the crude reaction mixtures; values in parentheses refer to isolated yields

^cTEMPO (0.3 equiv.) was added

^dKetone (1.0 mmol), NBS (1.5 mmol)

eProducts not isolated

substitution took place. The sulfur atom, being itself sensitive to oxidation (Ali et al. 2006), remained unchanged. The reaction of 2'-methoxy-5'-methylpropiophenone **5d**, which contained at least three potential reactive sites, gave the related α -bromoketone **6d** as the main product as well as an additional side-chain brominated minor product **7d** (entry 6). Functionalization of 3',4'-dimethoxypropiophenone **5e** yielded α - and ring brominated **6e** as the main product, accompanied by 10% of α -bromoketone **7e** (entry 7). Interestingly, no reaction took place in the presence of TEMPO to **5e** (entry 8). It appears that aromatic substitution of **5e** takes place via a radical pathway. Reactions of α -bromination of acetophenones and propiophenones in the presence of TEMPO suggest involvement of electrophilic and radical reaction pathways without clear boundary.

Subsequently, we selected model isobutyrophenones (aryl isopropyl ketones) (Table 3).

In the group of acetophenones **1**, the isopropyl and cyclohexyl moieties in **1c** and **1d** did not undergo benzylic

bromination (Table 1, entries 4 and 5). In this step, the reactivity of model isopropyl ketones was examined. The general feature of ketones 8 was that, regardless of the nature of the substituents under the studied reaction conditions, α -bromoketones 9 were formed as the main products. Functionalization of 4'-methylbutyrophenone 8a in addition to the α -brominated ketone **9a** also leads to the formation of benzyl bromide 10a (entry 1). The reaction of 8a was completely suppressed in the presence of TEMPO (entry 2) as well as in the dark (entry 3). Reaction of 2',4'-dimethylisobutyrophenone 8b, although being similar to 8a, did not result in benzylic bromination, but the only product formed was α -bromoketone **9b** (entry 4). Transformation of 4'-methoxyisobutyrophenone 8c exclusively produced α -bromoketone **9c** with no ring substitution noted (entry 5). The reaction of 3'-methyl-4'-methoxybutyrophenone 8d furnished a mixture of α -bromoketone 9d and benzyl bromide 10d (entry 6). The addition of TEMPO to the reaction mixture resulted in the recovery of the starting material 8d

Table 3 Regioselectivity of functionalization of heat substituted isobutviobilenones with hybrid under visible

	•		•		
Entry	Substrate	Reaction conditions ^a	Products	Conv. [%] ^b	Selectivity ^b
1	o ↓	<i>hv</i> , 21h		83	9a , 61 (23) 10a , 39 (23)
2		<i>hv</i> , TEMPO ^c , 24h	Br + Br	0	_
3	8a	dark, 21h	9a ^{Br} 10a	0	_
4	, , , , , , , , , , , , , ,	<i>hv</i> , 24h	Br	92	100 (53)
	8b		9b		
5	MeO BC	<i>hv</i> , 20h	MeO Br	91	100 (78)
6	MeO	<i>hv</i> , 20h ^d	9c O MeO MeO Br + MeO Br	78	9d , 69 (16) 10d , 31 (18)
7	. 8d	<i>hv</i> , TEMPO ^c , 24h	9d Br 10d	0	-

^aSubstrate **8** (0.3 mmol), NBS (0.33 mmol), stirring, 8-W energy-saving light bulb, reaction temperature 27 °C ^bConversions and relative distribution of products determined by ¹H NMR spectroscopy of the crude reaction mixtures; values in parentheses refer to isolated yields

^cTEMPO (0.3 equiv.) was added

^dKetone (1.0 mmol), NBS (1.5 mmol)

(entry 7). In contrast to acetophenones and propiophenones, the results of functionalization of isobutyrophenone suggest that α -bromination proceeds exclusively by a radical mechanism. This suggests that functionalization under solvent-free conditions might depend on substrate structure, reminiscent of radical α -bromination of highly substituted ketones with bromine in the presence of 1,2-epoxycyclohexane under photochemical conditions (Caló et al. 1977). Photochemical α -bromination of ketones with NBS under UV–Vis irradiation in diethyl ether is believed to follow a radical pathway (Arbuj et al. 2007).

In the next step, we explored the reactivity of the selected model benzyl ketones **11** (Table 4).

The highest regioselectivity was observed for this group, with all substrates being brominated at the carbonyl carbon atom. This outcome was anticipated, because the site of functionalization was an α -carbonyl and a benzylic position at the same time. The presence of either methyl group **11a**, methoxy group **11b** or both **11c** in the substrate did not influence the selectivity (entries 1, 3 and 4). Also, for the substrate with a sensitive thiophene ring like in **11d**, only α -bromination into **12d** was noticed, even though some α, α -dibrominated product **13d** was formed as well (entry 6).

TEMPO completely stopped any transformation for **11a** and **11d** (entries 2 and 7), and no reaction took place for **11c** in the dark (entry 5). Functionalization of benzyl ketones in the presence of TEMPO indicates that α -bromination might proceed via radical pathway.

To examine the effect of the structure at the carbonyl group on the reactivity of transformations under the applied reaction conditions, two substrates, **14** and **16**, with an additional electron-withdrawing nitrile group bonded to one of the α -carbon atoms, were selected (Table 5).

Clearly, this triple activation in 14 and 16 had a strong influence on regioselectivity and the reaction afforded exclusively products 15 and 17, respectively (entries 1 and 3). Both reactions of 14 and 16 occurred even in the dark (entries 2 and 4). When purifying the product 17 by means of column chromatography, we observed an interesting phenomenon, in which the acetyl group was partly cleaved on the column and a mixture of 17 and 18 was eluted. Attempts to separate these two compounds by a preparative thin-layer chromatography resulted in complete deacetylation of 17 into product 18.

Finally, the reactivity of naphthyl ketones was examined (Table 6). The most interesting and unexpected result was

0	5		•	U	
Entry	Substrate	Reaction conditions ^a	Products	Conv. [%] ^b	Selectivity ^b
1	°,	<i>hv</i> , 24h	O Br	75	100 (35)
2	Ph 11a	<i>hv</i> , TEMPO ^c , 24h	Ph 12a	0	_
3	MeO Ph 11b	<i>hv</i> , 24h	MeO Ph 12b	68	100 (60)
4	OMe O	hv, 22h ^d	OMe O Br	90	100 (63)
5	Ph 11c	dark, 21h	Ph 12c	0	_
6	s	<i>hv</i> , 22h ^e	S Br S Br	100	12d , 80 (72) 13d , 20 (16)
7	Ph 11d	<i>hv</i> , TEMPO ^c , 24h	Ph + Ph Ph 12d 13d	0	_

Table 4 Regioselectivity of functionalization of neat substituted benzyl ketones with NBS under visible light

^aSubstrate 11 (0.3 mmol), NBS (0.33 mmol), stirring, 8-W energy-saving light bulb, reaction temperature 27 °C

^bConversions and relative distribution of products determined by ¹H NMR spectroscopy of the crude reaction mixtures; values in parentheses refer to isolated yields

^cTEMPO (0.3 equiv.) was added

^dKetone (0.5 mmol), NBS (0.6 mmol)

^eKetone (1.5 mmol), NBS (1.8 mmol)

Entry	Substrate	Reaction conditions ^a	Products	Conv. [%] ^b	Selectivity ^b
1	CN	<i>hv</i> , 24h	CN	95	100 (75)
2	14 U	dark, 21h	Br 0 15	95	100 ^d
3	CN	<i>hv</i> , 24h		100	17 , 100 (0) 18 , 0 (53) ^c
4	ڻ 16	dark, 21h	17 18	100	17 , 100 ^d

Table 5 Regioselectivity of functionalization of neat substituted α-cyanoketones with NBS under visible light

^aSubstrate 14/16 (0.3 mmol), NBS (0.33 mmol), stirring, 8-W energy-saving light bulb, reaction temperature 27 °C

^bConversions and relative distribution of products determined by ¹H NMR spectroscopy of the crude reaction mixtures; values in parentheses refer to isolated yields

^cOnly 18 isolated after column and preparative chromatography

^dProducts not isolated

observed by irradiating (4-methoxynaphthyl) methyl ketone **19a** with 2.2 equiv. of NBS under SFRC.

Instead of the anticipated bromination at the alkyl side chain, the main reaction pathway was *ipso*-substitution of the acetyl group by a bromine atom alongside with aromatic and α -carbonyl bromination as minor pathways (entry 1). This observation stimulated us to examine the influence of the acyl moiety structure of (4-methoxynaphthyl) alkyl ketones on potential bromodeacylation. The fact that a similar distribution of the products was found for (4-methoxynaphthyl) ethyl ketone **19b** (entry 4) indicates that the *ipso*substitution under these conditions could be more general. Although the illumination of (4-methoxynaphthyl) benzyl ketone resulted in a complex mixture of products, a significant amount of **20** was found in the crude reaction mixture, indicating that bromodeacylation occurs with NBS under





^aSubstrate 19 (0.3 mmol), NBS (0.66 mmol), stirring, 8-W energy-saving light bulb, reaction temperature 27 °C

^bConversions and relative distribution of products determined by ¹H NMR spectroscopy of the crude reaction mixtures; values in parentheses refer to isolated yields

^cTEMPO (0.3 equiv.) was added

^dSubstrate 23 (0.3 mmol), NBS (0.33 mmol)

^eIncomplete conversion, a complex reaction mixture

^fProducts not isolated

SFRC with this substrate as well. To test the hypothesis that an enolizable carbonyl group must be present for the ipsosubstitution, the non-enolizable (4-methoxynaphthalen-1-yl) (phenyl)methanone 19c was subjected to the same reaction conditions. The reaction exclusively furnished the ring-brominated product 21c with no deacylation observed (entry 5). Interestingly, *ipso*-substitution was also noticed in the case of the considerably electron-rich aryl ketone 2'-methyl-4',5'-dimethoxyacetophenone 23, in which the main bromodeacylated product 24 was accompanied by a small amount of α -bromoketone 25 (entry 8). Bromodeacylation is rather a rare process, known to take place with molecular bromine in a solution (Beirne et al. 1970). The addition of TEMPO to the reaction mixture resulted in different effects for different substrates. While inhibiting α -carbonyl bromination, it had no effect on ipso- and ring substitution at 19a and **19c** (entries 2 and 6). In contrast, TEMPO completely suppressed all transformations of the compound **23** (entry 9). Selectivity of the transformation of **19a** in the dark was decreased considerably resulting in a complex mixture of products (entry 3), while no reaction took place with **19c** in the absence of light (entry 7).

Some further experiments were carried out in order to shed light on mechanism of bromodeacylation. 4-Methoxy-1-acetonaphthone **19a** was reacted as a model substrate under different reaction conditions with NBS and bromine. Transformation of **19a** with NBS/NH₄OAc in diethyl ether (Tanemura et al. 2004) led to 2-bromo-1-(4methoxynaphthalen-1-yl)ethan-1-one with 35% conversion. Reaction of **19a** with NBS/NH₄NO₃ in acetonitrile (Tanemura et al. 2003) furnished a complex mixture of ring-brominated products and a small proportion of bromodeacylated product 20. Functionalization of 19a with bromine in acetic acid furnished 2-bromo-1-(4-methoxynaphthalen-1-yl)ethan-1-one with 65% conversion. This reaction is assumed to take place in an electrophilic fashion via enol form of 19a. The system NBS/pTSA in DCM (Pravst et al. 2008; Vyas et al. 2016) expectedly furnished 2-bromo-1-(4-methoxynaphthalen-1-yl)ethan-1one with 80% conversion. It can be concluded that electrophilic functionalization of α -carbonyl position takes place in the present of an acid (Br₂/AcOH and NBS/pTSA in DCM). The system NBS/NH₄NO₃ in acetonitrile led to the expected electrophilic ring functionalization; in addition, a minor portion of ipso-product 20 was observed. Reactions in the presence of TEMPO suggest that the ring functionalization of 19a might proceed via electrophilic and radical reaction pathways, while formation of 20 is supposed to be attributed to a different reaction channel-potentially a radical ion reaction pathway (Fasani et al. 1994).

Conclusions

In this study, we disclosed regioselectivity of visiblelight-induced transformations of model aryl methyl-, aryl ethyl-, aryl isopropyl-, aryl benzyl-, α -cyano-substitutedand naphthyl alkyl ketones, containing several competitive reactive sites with NBS in the absence of solvents. The entire study was carried out under heterogeneous reaction conditions. In most cases, α -brominated ketones were the main products, accompanied with α, α -dibrominated ketones. Wohl-Ziegler bromination and aromatic ring bromination were noted as minor reaction pathways in the case of several alkyl- and alkoxy-substituted ketones. The α -bromination of ketones under solvent-free reaction conditions appears to depend on the structure of the substrates. While in the case of acetophenones and propiophenones it is proposed to proceed by the electrophilic and radical pathways, in the case of isobutyrophenones and aryl benzyl ketones it is likely to take place by the radical pathway. α -Cyano-substituted ketones were converted into their α -bromo- α -cyano-substituted derivatives even in the dark. An unexpected exception was the reaction of enolizable (4-methoxynaphthyl) alkyl ketones, in which the main product was formed because of an ipso-substitution of the acyl group with a bromine atom. Bromodeacylation is suggested to proceed via a radical cation intermediates. The radical scavenger TEMPO slowed or suppressed α - and ring bromination, whereas benzylic bromination and α . dibromination with NBS were entirely suppressed. It was demonstrated in several cases that reactions might proceed in electrophilic and radical fashion without a distinct boundary. In summary, we contributed to understanding

reaction pathways of these sensitive compounds in reactions with NBS under solvent-free reaction conditions.

Supplementary Information The online version contains supplementary material available at https://doi.org/10.1007/s11696-021-01711-x.

Acknowledgements The authors acknowledge the financial support of the Slovenian Research Agency (Research Core Funding Grant P1-0230) and Projects J1-8147 and J1-9166. The authors thank Dr. Damijana Urankar for HRMS at the Centre for Research Infrastructure at the Faculty of Chemistry and Chemical Technology (IC UL FCCT).

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

- Adams DJ, Dyson PJ, Tavener SJ (2004) Chemistry in alternative reaction media. John Wiley & Sons, Chichester
- Adhikari MV, Samant SD (2002) Sonochemical bromination of acetophenones using p-toluensulfonic acid–N-bromosuccinimide. Ultrason Sonochem 9:107–111. https://doi.org/10.1016/S1350-4177(01)00108-0
- Ali MH, Hartman M, Lamp K et al (2006) Oxidation of sulfides with *N*-bromosuccinimide in the presence of hydrated silica gel. Synth Commun 36:1769–1777. https://doi.org/10.1080/0039791060 0618988
- Appa RM, Naidu BR, Lakshmidevi J et al (2019) Added catalyst-free, versatile and environment beneficial bromination of (hetero)aromatics using NBS in WEPA. SN Appl Sci 1:1281. https://doi.org/ 10.1007/s42452-019-1274-x
- Arbuj SS, Waghmode SB, Ramaswamy AV (2007) Photochemical α -bromination of ketones using *N*-bromosuccinimide: a simple, mild and efficient method. Tetrahedron Lett 48:1411–1415. https://doi.org/10.1016/j.tetlet.2006.12.101
- Beirne JJ, Coyle AM, Donnelly JA (1970) Halodeacylation of aryl ketones. Tetrahedron 26:3809–3814. https://doi.org/10.1016/ S0040-4020(01)92959-X
- Bose A, Mal P (2014) Electrophilic aryl-halogenation using *N*-halosuccinimides under ball-milling. Tetrahedron Lett 55:2154–2156. https://doi.org/10.1016/j.tetlet.2014.02.064
- Caló V, Lopez L, Pece G (1977) Bromination of ketones in the presence of epoxides as a method for regiospecific synthesis of bromoketones. J Chem Soc Perkin Trans 1:501–504. https://doi.org/10. 1039/P19770000501
- Cantillo D, de Frutos O, Rincon JA et al (2014) A scalable procedure for light-induced benzylic brominations in continuous flow. J Org Chem 79:223–229. https://doi.org/10.1021/jo402409k
- Cave GWV, Raston CL, Scott JL (2001) Recent advances in solventless organic reactions: towards benign synthesis with remarkable versatility. Chem Commun. https://doi.org/10.1039/B106677N

- Coan SB, Becker EI (1955) α-(4-chlorophenyl)-γphenylacetoacetonitrile. Org Synth 35:30. https://doi.org/10. 15227/orgsyn.035.0030
- Deshpande S, Gadilohar B, Shinde Y et al (2015) Energy efficient, clean and solvent free photochemical benzylic bromination using NBS in concentrated solar radiation (CSR). Sol Energy 113:332– 339. https://doi.org/10.1016/j.solener.2015.01.008
- Dinda M, Chakraborty S, Samanta S et al (2013) Solar photothermochemical reaction and supercritical CO₂ work up for a fully green process of preparation of pure *p*-nitrobenzyl bromide. Environ Sci Technol 47:10535–10540. https://doi.org/10.1021/es4019282
- Djerassi C (1948) Brominations with N-bromosuccinimide and related compounds. The Wohl-Ziegler Reaction. Chem Rev 43:271–317. https://doi.org/10.1021/cr60135a004
- Fasani E, d'Alessandro N, Albini A, Mariano PS (1994) Set Photochemical Reactions between 1,4-Naphthalenedicarbonitrile and Benzylic Donors. Medium Effects J Org Chem 59:829–835. https://doi.org/10.1021/jo00083a025
- Ganguly NC, De P, Dutta S (2005) Mild regioselective monobromination of activated aromatics and heteroaromatics with *N*-bromosuccinimide in tetrabutylammonium bromide. Synthesis. https:// doi.org/10.1055/s-2005-861866
- Gawande MB, Bonifácio VDB, Luque R et al (2014) Solvent-free and catalyst-free chemistry. Chemsuschem 7:24–44. https://doi.org/ 10.1002/cssc.201300485
- Goswami S, Dey S, Jana S et al (2004) Side chain bromination of mono and dimethyl heteroaromatic and aromatic compounds by solid phase *N*-bromosuccinimide reaction without radical initiator under microwave. Chem Lett 33:916–917. https://doi.org/10. 1246/cl.2004.916
- Heropoulos GA, Cravotto G, Screttas CG et al (2007) Contrasting chemoselectivities in the ultrasound and microwave assisted bromination reactions of substituted alkylaromatics with N-bromosuccinimide. Tetrahedron Lett 48:3247–3250. https://doi.org/10. 1016/j.tetlet.2007.03.023
- Imanzadeh GK, Zamanloo MR, Eskandari H et al (2006) A new ring bromination method for aromatic compounds under solvent-free conditions with NBS/Al₂O₃. J Chem Res. https://doi.org/10.3184/ 030823406776330657
- Jereb M, Zupan M, Stavber S (2009) Visible-light-promoted Wohl— Ziegler functionalization of organic molecules with N-bromosuccinimide under solvent-free reaction conditions. HeIv Chim Acta 92:555–566. https://doi.org/10.1002/hlca.200800308
- Kumar A, Lohan P, Prakash O (2012) Selective bromination of dehydroacetic acid with N-Bromosuccinimide. Synth Commun 42:2739–2747. https://doi.org/10.1080/00397911.2011.566461
- Martins MAP, Frizzo CP, Moreira DN et al (2009) Solvent-free heterocyclic synthesis. Chem Rev 109:4140–4182. https://doi.org/ 10.1021/cr9001098
- Metzger JO (1998) Solvent-free organic syntheses. Angew Chem Int Ed 37:2975–2978. https://doi.org/10.1002/(SICI)1521-3773(19981 116)37:21%3C2975::AID-ANIE2975%3E3.0.CO;2-A
- Nakamatsu S, Toyota S, Jones W et al (2005) The important role of solvent vapor in an organic solid state reaction. Chem Commun. https://doi.org/10.1039/b503922c
- Obst M, König B (2018) Organic synthesis without conventional solvents. Eur J Org Chem. https://doi.org/10.1002/ejoc.201800556
- Paquette LA, Crich D, Fuchs PL (2009) N-Bromosuccinimide. Encyclopedia of reagents for organic synthesis, 2nd edn. John Wiley & Sons, Chichester, pp 1657–1667
- Podgoršek A, Stavber S, Zupan M et al (2006) Visible light induced 'on water' benzylic bromination with *N*-bromosuccinimide. Tetrahedron Lett 7:1097–1099. https://doi.org/10.1016/j.tetlet.2005. 12.040
- Podgoršek A, Stavber S, Zupan M et al (2009) Environmentally benign electrophilic and radical bromination 'on water': H₂O₂–HBr

system versus *N*-bromosuccinimide. Tetrahedron 65:4429–4439. https://doi.org/10.1016/j.tet.2009.03.034

- Prakash GKS, Mathew T, Hoole D et al (2004) N-halosuccinimide/ BF₃·H₂O, efficient electrophilic halogenating systems for aromatics. J Am Chem Soc 126:15770–15776. https://doi.org/10.1021/ ja0465247
- Pravst I, Zupan M, Stavber S (2006a) Directed regioselectivity of bromination of ketones with NBS: solvent-free conditions versus water. Tetrahedron Lett 47:4707–4710. https://doi.org/10.1016/j. tetlet.2006.04.119
- Pravst I, Zupan M, Stavber S (2006b) Solvent-free bromination of 1,3-diketones and β-keto esters with NBS. Green Chem 8:1001– 1005. https://doi.org/10.1039/B608446J
- Pravst I, Zupan M, Stavber S (2008) Halogenation of ketones with N-halosuccinimides under solvent-free reaction conditions. Tetrahedron 64:5191–5199. https://doi.org/10.1016/j.tet.2008.03.048
- Pravst I, Zupan M, Stavber S (2009) Introduction of halogen atoms into organic compounds under solvent-free reaction conditions. Curr Org Chem 13:47–70. https://doi.org/10.2174/1385272097 87193819
- Rahman ANMM, Bishop R, Tan R et al (2005) Solid-state regio- and stereo-selective benzylic bromination of diquinoline compounds using N-bromosuccinimide. Green Chem 7:207–209. https://doi. org/10.1039/B416275G
- Reichardt C (2007) Solvents and solvent effects: an introduction. Org Proc Res Dev 11:105–113. https://doi.org/10.1021/op0680082
- Rothenberg G, Downie AP, Raston CL et al (2001) Understanding solid/solid organic reactions. J Am Chem Soc 123:8701–8708. https://doi.org/10.1021/ja0034388
- Salama TA, Novák Z (2011) N-halosuccinimide/SiCl₄ as general, mild and efficient systems for the α-monohalogenation of carbonyl compounds and for benzylic halogenation. Tetrahedron Lett 52:4026–4029. https://doi.org/10.1016/j.tetlet.2011.05.135
- Shaw H, Perlmutter HD, Gu C et al (1997) Free-radical bromination of selected organic compounds in water. J Org Chem 62:236–237. https://doi.org/10.1021/jo950371b
- Singh MS, Chowdhury S (2012) Recent developments in solvent-free multicomponent reactions: a perfect synergy for eco-compatible organic synthesis. RSC Adv 2:4547–4592. https://doi.org/10. 1039/C2RA01056A
- Šterk D, Jukič M, Časar Z (2013) Application of flow photochemical bromination in the synthesis of a 5-bromomethylpyrimidine precursor of rosuvastatin: improvement of productivity and product purity. Org Process Res Dev 17:145–151. https://doi.org/10.1021/ op300248y
- Tanaka K, Toda F (2000) Solvent-free organic synthesis. Chem Rev 100:1025–1074. https://doi.org/10.1021/cr940089p
- Tanaka K, Shiraishi R, Toda F (1999) A new method for stereoselective bromination of stilbene and chalcone in a water suspension medium. J Chem Soc Perkin Trans 1:3069–3070. https://doi.org/ 10.1039/A906967D
- Tanemura K, Suzuki T, Nishida Y, Satsumabayashi K, Horaguchi T (2003) Halogenation of Aromatic Compounds by N-chloro-, N-bromo- and N-iodosuccinimide. Chem Lett 32:932–933. https:// doi.org/10.1246/cl.2003.932
- Tanemura K, Suzuki T, Nishida Y, Satsumabayashi K, Horaguchi T (2004) A mild and efficient procedure for α-bromination of ketones using *N*-bromosuccinimide. Chem Commun. https://doi.org/10.1039/B315340A
- Tanko JM, Blackert JF (1994) Free-radical side-chain bromination of alkylaromatics in supercritical carbon dioxide. Science 263:203– 205. https://doi.org/10.1126/science.263.5144.203
- Toda F, Schmeyers J (2003) Selective solid-state brominations of anilines and phenols. Green Chem 5:701–703. https://doi.org/10. 1039/B306612F

- Togo H, Hirai T (2003) Environmentally-friendly Wohl-Ziegler bromination: ionic-liquid reaction and solvent-free reaction. Synlett. https://doi.org/10.1055/S-2003-38379
- Vogel AI, Furniss BS, Hannaford AJ et al (1989) Vogel's textbook of practical organic chemistry, 5th edn. Longman Group, Harlow Essex, England
- Vražič D, Jereb M, Laali KK et al (2013) Brønsted acidic ionic liquid accelerated halogenation of organic compounds with N-halosuccinimides (NXS). Molecules 18:74–96. https://doi.org/10.3390/ molecules18010074
- Vyas VK, Bhanage BM (2016) Kinetic resolution driven diastereo-and enantioselective synthesis of cis-β-heteroaryl amino cycloalkanols by ruthenium-catalyzed asymmetric transfer hydrogenation. Org Lett 18:6436–6439. https://doi.org/10.1021/acs.orglett.6b03334
- Walsh PJ, Li H, de Parrodi CA (2007) A green chemistry approach to asymmetric catalysis: solvent-free and highly concentrated reactions. Chem Rev 107:2503–2545. https://doi.org/10.1021/cr050 9556

- Wang GW, Gao J (2012) Solvent-free bromination reactions with sodium bromide and oxone promoted by mechanical milling. Green Chem 14:1125–1131. https://doi.org/10.1039/C2GC1 6606B
- Winkler M, Steinbiß M, Meier MAR (2014) A more sustainable Wohl-Ziegler bromination: versatile derivatization of unsaturated FAMEs and synthesis of renewable polyamides. Eur J Lipid Sci Technol 116:44–51. https://doi.org/10.1002/ejlt.201300126
- Zagande S, Patil P (2019) A review on solvent-free methods in organic synthesis. Curr Org Chem 23:2295–2318. https://doi.org/10.2174/ 1385272823666191016165532

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.