

Journal Pre-proof

Synthesis of DBpin using Earth-abundant metal catalysis

Andrew W.M. Cummins, Shuyang Li, Dominic R. Willcox, Tommi Muilu, Jamie H. Docherty, Stephen P. Thomas



PII: S0040-4020(20)30205-2

DOI: <https://doi.org/10.1016/j.tet.2020.131084>

Reference: TET 131084

To appear in: *Tetrahedron*

Received Date: 24 January 2020

Accepted Date: 25 February 2020

Please cite this article as: Cummins AWM, Li S, Willcox DR, Muilu T, Docherty JH, Thomas SP, Synthesis of DBpin using Earth-abundant metal catalysis, *Tetrahedron* (2020), doi: <https://doi.org/10.1016/j.tet.2020.131084>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd.

Graphical Abstract

To create your abstract, type over the instructions in the template box below.
Fonts or abstract dimensions should not be changed or altered.


Synthesis of DBpin Using Earth-abundant Metal Catalysis

Leave this area blank for abstract info.

Andrew W. M. Cummins^a, Shuyang Li^a,
Dominic R. Willcox^a, Tommi Muilu^b, Jamie H.
Docherty^{a,*}, Stephen P. Thomas^{a,*}

^a EaStCHEM School of Chemistry, University of Edinburgh, Joseph Black Building, David Brewster Road, Edinburgh, EH9 3FJ, UK.

^b EaStCHEM, School of Chemistry, University of St Andrews North Haugh, Fife, KY16 9ST, UK.



• Alkoxide Activation • Safe and Simple Procedure • High Isolated Yields • >99% Deuterium Incorporation



Synthesis of DBpin Using Earth-abundant Metal Catalysis

Andrew W. M. Cummins^a, Shuyang Li^a, Dominic R. Willcox^a, Tommi Muilu^b, Jamie H. Docherty^{a, *}, Stephen P. Thomas^{a, *}

^a EaStCHEM School of Chemistry, University of Edinburgh, Joseph Black Building, David Brewster Road, Edinburgh, EH9 3FJ, UK.

^b EaStCHEM, School of Chemistry, University of St Andrews North Haugh, Fife, KY16 9ST, UK

ARTICLE INFO

Article history:

Received

Received in revised form

Accepted

Available online

Keywords:

Catalysis

Boron

Iron

Cobalt

Deuterium

ABSTRACT

The synthesis of DBpin was achieved using (^{Et}BIP)CoCl₂ or (^tBuPNN)FeCl₂ as pre-catalysts activated with NaO^tBu. (^{Et}BIP)CoCl₂ was used as a pre-catalyst for the hydrogen isotope exchange of HBpin with D₂, and (^tBuPNN)FeCl₂ for deuterogenolysis of B₂pin₂. The one-pot, tandem hydrogenolysis-hydroboration/deuterogenolysis-deuteroboration reaction of terminal alkenes could be catalysed by (^tBuPNN)FeCl₂ to give alkyl boronic esters.

1. Introduction

Deuterated compounds have found use across a wide range of applications from mechanistic analysis to pharmacokinetic studies. The strength of the C-D bond slows metabolism and can therefore facilitate lower pharmaceutical doses and numerous deuterated active pharmaceutical ingredients are currently undergoing clinical trials, with Deutetrabenazine having been fully approved by the FDA [1].

Boronic esters are ubiquitous in modern organic chemistry due to their functional group tolerance and broad applications [2–4]. Boronic esters are commonly prepared by the metal-catalysed hydroboration of alkenes and alkynes, with a wide range of transition metal and main group catalysts having been developed [5,6]. 4,4,5,5-Tetramethyl-1,3,2-dioxaborolane (HBpin) has become a common reagent in organic synthesis, due to its stability and that of alkyl and aryl pinacolboronic esters [7]. The deuterio-isotopologue, DBpin offers a simple means of deuterium-incorporation and has been commonly used for *in situ* reaction monitoring of reactive species, and allows for a B-H kinetic isotope effect to be determined [8–16].

The synthesis of HBpin was first reported by Knochel by reaction of Me₂S·BH₃ with pinacol and distillation to give a high yield of HBpin [17]. A similar synthetic route has been commonly used to prepare DBpin, but requires the prior synthesis of B₂D₆ gas, which is toxic, pyrophoric, and requires a complex reaction setup [18]. DBpin has also been prepared by hydrogen isotope exchange of HBpin, however low deuterium


incorporation is common, and forcing conditions can be needed. The first example of hydrogen isotope exchange of HBpin was reported by Perutz and co-workers using a rhodium catalyst, however the DBpin was not isolated from the reaction [19]. Further examples of hydrogen isotope exchange have been shown with iridium and ruthenium complexes [13,20], but only a single example has been reported using an Earth-abundant metal, a methyl-cobalt complex, which gave good deuterium incorporation at room temperature [8].

The deuterogenolysis of bis-(pincolato)borane (B₂pin₂) provides a synthetically simple route to DBpin, with high deuterium incorporation. The deuterogenolysis of B₂pin₂ was first demonstrated by Hartwig and Hall using an iridium catalyst, however due to low reactivity the reaction required multiple charges of D₂ over the period of 24 h [9]. Marder and Braunschweig demonstrated the hydrogenolysis of B₂pin₂ using a heterogenous catalyst [21]. Huang demonstrated the use of a PNN-ligated cobalt catalyst for the deuterogenolysis of B₂pin₂, with high reactivity and the generated DBpin could be used *in situ* for the deuteroboration of alkenes and alkynes using the same catalyst. NaHBET₃ was required to activate the cobalt pre-catalyst [22].


We have previously demonstrated NaO^tBu as an alternative to organometallic activators in Earth-abundant metal catalysis, thus we sought to extend this to the synthesis of DBpin [23]. Herein we report the use of Earth-abundant metal catalysts for the hydrogen isotope exchange of HBpin and deuterogenolysis of

B_2pin_2 using bench stable pre-catalysts and NaO^tBu as the activator to give DBpin with high deuterium incorporation and simple purification.

Previous Work:



This Work:




Scheme 1. Synthesis of 2-deutero-4,4,5,5-tetramethyl-1,3,2-dioxaboralane (DBpin) catalysed by transition metal complexes.

2. Results and discussion

2.1. Hydrogen Isotope Exchange of HBpin


Chirik reported that the Co(I)-methyl complex, (^{Me}BIP)CoCH₃, could catalyse hydrogen isotope exchange with HBpin, therefore we attempted to access this reactivity by *in situ* alkoxide activation from the parent Co(II) complex (^{Et}BIP)CoCl₂ [8]. We found that hydrogen isotope exchange occurred readily using (^{Et}BIP)CoCl₂ **1** as a catalyst and NaO^tBu as an activator when the reaction was carried out in *neat* HBpin under D₂ (4 bar) giving excellent deuterium incorporation (99%-*d*) and conversion (>99%), but poor isolated yield (25%). Presumably when the reaction was removed from the D₂ atmosphere, the catalyst promoted the degradation of DBpin into redistribution products, such as B₂pin₃.



Scheme 2. Hydrogen isotope exchange of HBpin by (^{Et}BIP)CoCl₂ using NaO^tBu as an activator.

2.2. Hydrogenolysis and deuterogenolysis of B₂pin₂

We next turned our attention to the hydrogenolysis of B₂pin₂. Various iron complexes were assessed, Milstein's PNN and Huang's PONN complexes [(^tBuPNN^tPr)FeCl₂ **7** and (^tBuPNN)FeCl₂ **8** were found to give the only active catalysts using a combination of PhSiH₃ and NaO^tBu for pre-catalyst activation (Scheme 3) [23]. We also found that Ni(COD)₂ in combination with PCy₃ gave excellent conversion [22].



Scheme 3. Hydrogenolysis of B₂pin₂ with various catalysts. Values listed are conversion by ¹¹B NMR. ^a Without PhSiH₃ and NaO^tBu, with THF as the solvent. ^b 5 mol% catalyst loading used.

Using (^tBuPNN)FeCl₂ **8** we tested various organometallic activators as a performance comparison (Table 1) [23]. All organometallic activators were found to give significant amounts of B₂pin₃ along with other degradation products, as observed by ¹¹B NMR spectroscopy, and PhSiH₃ and NaO^tBu remained the activator of choice.

Table 1. Activator screen for the hydrogenolysis of B₂pin₂ by (^tBuPNN)FeCl₂. Conversion calculated from ¹¹B NMR.

Entry	Activator	Conversion (%)
1	NaO ^t Bu + PhSiH ₃	89
2	NaHBET ₃	37
3	EtMgBr	51
4	PhLi	81

We then carried out a series of control reactions to ensure that the iron catalyst and activator were required. Interestingly, it was found the exogenous PhSiH₃ was not required to activate the pre-catalyst, with B₂pin₂ able to perform the same function (Table 2,


entry 2). The same results were obtained regardless of whether the B_2pin_2 was purified by recrystallisation or not. We found that the reaction was complete in 2 hours.

Table 2. Control reactions. Conversion calculated from ^{11}B NMR.

2, (1 mol%)
NaO^tBu (2 mol%)
PhSiH₃ (2 mol%)
Et₂O, r.t., 16 h

Entry	Variation from standard conditions	Conversion (%)
1	None	89
2	No PhSiH ₃	93
3	No NaO ^t Bu	0
4	Only (^t BuPNN)FeCl ₂	0
5	Only NaO ^t Bu	0
6	Only ^t BuPNN Ligand	0
7	Time = 2 h	89

Having optimised hydrogenolysis conditions, we then set out to optimise the purification process. Et₂O was found to be the best reaction solvent, as it could easily be separated from the HBpin by distillation to give good isolated yield (76%) on gram scale. With optimised conditions for hydrogenolysis, we next sought to apply this reaction to the preparation of DBpin. Deuterohydrogenolysis was found to be much slower than hydrogenolysis, presumably due to a kinetic isotope effect and a lower reaction pressure, therefore, the reaction time was extended to 16 hours. This gave DBpin in high isolated yield (86%) on gram-scale with >99% *d* incorporation.




Scheme 4. (a) Optimised hydrogenolysis of B_2pin_2 . (b) Optimised deuterohydrogenolysis of B_2pin_2 .

2.3. One-pot tandem hydrogenolysis-hydroboration and deuterohydrogenolysis-deuteroboration of B_2pin_2

As (^tBuPNN)FeCl₂ is also reported to be a highly efficient hydroboration catalyst [26], we trialed a one-pot hydrogenolysis-hydroboration using the HBpin that was formed *in situ*. This hydroboration showed functional group tolerance to siloxanes (**10b**), esters (**10c**), and internal alkenes (**10d**) (Scheme 5, a). We extended this tandem reaction to the deuterohydrogenolysis-deuteroboration of terminal alkenes (Scheme 5, b). Presumably due to a significant kinetic

isotope effect using DBpin the yields were lower than the analogous protio equivalent for all substrates. Deuterium incorporation was observed at both the terminal and internal position indicating the catalyst is operating by reversible hydrometallation of the alkene. No deuterium incorporation was observed at the internal alkene of a diene (**10d**), suggesting the hydrometallation is chemoselective for terminal alkenes (Scheme 5, b).



Scheme 5. (a) Scope of iron-catalysed one-pot hydrogenolysis-hydroboration of terminal alkenes. (b) Scope of iron-catalysed one-pot deuterohydrogenolysis-deuteroboration of terminal alkenes. Deuterium incorporation determined by integration of the residual signal in the 1H NMR spectrum.

3. Conclusions

We have successfully demonstrated the use of cobalt- and iron-catalysis for the synthesis of DBpin with high deuterium incorporation and isolated yield, using either hydrogen isotope exchange or deuterohydrogenolysis. The one-pot tandem hydrogenolysis-hydroboration and deuterohydrogenolysis-deuteroboration were also developed and applied to terminal alkenes.

4. Experimental section

4.1 General

Reaction setup: All reactions were performed in oven (185 °C) and/or flame dried glassware. All glassware was cleaned using base (KOH, ^tPrOH) and acid (HCl_{aq}) baths. All reported reaction temperatures correspond to ambient room temperature, which was approximately 20 °C.

NMR spectroscopy: ^1H , ^2D , ^{13}C and ^{11}B spectra were recorded on Bruker Avance III 400 and 500 MHz; Bruker PRO 500 MHz; Bruker Avance I 600 MHz spectrometers. Chemical shifts are reported in parts per million (ppm). ^1H and ^{13}C NMR spectra were referenced to the residual solvent peak (CDCl_3 : 7.26 ppm). Multiplicities are indicated by app. (apparent), br. (broad), s (singlet), d (doublet), t (triplet), q (quartet), quin. (quintet), sext. (sextet), sept. (septet), non. (nonet). Coupling constants, J , are reported in Hertz and rounded to the nearest 0.1 Hz. Integration is provided and assignments are indicated.

Chromatography: Column chromatography was carried out on a Teledyne ISCO CombiFlash NextGen 300+ using RediSep Rf Gold normal phase silica flash columns (12, 25, 40, or 80 g; 20–40 microns). Substrates were purified using 40/60 petroleum ether and EtOAc on a gradient of 100:0 to 0:100 with flow rates of 10–110 mL min^{-1} depending on the size of column and ΔR_f .

Mass Spectrometry: Mass spectrometry (MS) was performed by the University of Edinburgh, School of Chemistry, Mass Spectrometry Laboratory. High resolution mass spectra were recorded on a VG autospec, or Thermo/Finnigan MAT 900, mass spectrometer. Electron Impact (EI^+) spectra were performed at 70 eV using methane as the carrier gas, with either a double focusing sector field (DFSF) or time-of-flight (TOF) mass analyzer. Chemical Ionization (CI^+) spectra were performed with methane reagent gas, with either a double focusing sector field (DFSF) or time-of-flight (TOF) mass analyzer. Electrospray Ionization (ESI^+) spectra were performed using a time-of-flight (TOF) mass analyzer. Data are reported in the form of m/z (intensity relative to the base peak = 100).

Chemicals: All reagents and were purchased from Sigma Aldrich, Alfa Aesar, Acros organics, Tokyo Chemical Industries UK, Fluorochem and Apollo Scientific without further purification. Sodium *tert*-butoxide (97%) was purchased from Sigma Aldrich (UK). B_2pin_2 was purchased from Fluorochem and recrystallised before use. All cobalt and iron species were prepared according to reported procedures [23,24,26–28].

4.2. Preparation of DBpin with ($^{\text{Et}}\text{BIP}$) CoCl_2

Under argon, to an autoclave was added ($^{\text{Et}}\text{BIP}$) CoCl_2 (0.14 mmol, 76 mg), and NaO^tBu (0.28 mmol, 30 mg). HBpin (2.0 mL, 14 mmol) was added dropwise *n.b. exothermic*. The vessel was sealed, flushed with D_2 , then pressurised with D_2 (4 bar). The reaction was left to stir at room temperature for 36 h, then the flush/pressurise cycle was repeated, and the reaction was left to stir for another 24 h. The DBpin was distilled *in vacuo* (25 °C, 0.1 mbar), to give DBpin (0.50 g, 25%, 99%-d) as a colourless oil. The deuterium incorporation was determined by integration of the residual HBpin signal. ^1H NMR (500.12 MHz, CDCl_3): 1.27 (12H, s, CH_3). ^{11}B NMR (128.34 MHz, CDCl_3): 28.2 (s br), ^{13}C NMR (125.77 MHz, CDCl_3) 25.0, 83.4. ^2D NMR (76.75 MHz, toluene) 4.15 (br m). Analytical data were in accordance with those previously reported [22].

4.3. Hydrogenolysis of B_2pin_2

B_2pin_2 (10 mmol, 2.54 g), ($^{\text{Bu}}\text{PNN}$) FeCl_2 (0.1 mmol, 44 mg) and NaO^tBu (0.2 mmol, 19 mg) were dissolved in Et_2O (10 mL) in an autoclave. The vessel was purged with H_2 then pressurised with H_2 (10 bar). This was left to stir for 2 hours, then the pressure was released. The reaction mixture was distilled at atmospheric pressure to remove Et_2O (60 °C). The HBpin was distilled *in vacuo* (25 °C, 0.1 mbar) giving HBpin (1.95 g, 76%) as a colourless oil. ^1H NMR (500.12 MHz, CDCl_3): 1.22 (12H, s, CH_3), 3.81 (1H, 1:1:1:1 q, $J = 171.7$ Hz). ^{11}B NMR (128.34 MHz, CDCl_3): 28.2 (d, $J = 174.9$ Hz), ^{13}C NMR (125.77 MHz,

CDCl_3) 25.0, 83.3. Analytical data were in accordance with those previously reported [22].

4.4. General procedure for one-pot hydroboration of terminal alkenes.

B_2pin_2 (0.500 mmol, 127 mg), ($^{\text{Bu}}\text{PNN}$) FeCl_2 (0.005 mmol, 2.2 mg), and NaO^tBu (0.01 mmol, 0.8 mg) were dissolved in Et_2O (0.5 mL) in an autoclave. The vessel was purged with H_2 then pressurised with H_2 (10 bar). The reaction was stirred for 2 hours, then the pressure was released and the vessel transferred into a glovebox. Alkene (1.00 mmol) was added, and the reaction left to stir for 30 mins. The solvent was removed *in vacuo* and the residue purified by flash column chromatography on silica (with a mixture of 40/60 petroleum ether and diethyl ether). Volatiles were removed *in vacuo* to give the alkyl boronic ester.

4.5. Deuterogenolysis of B_2pin_2

B_2pin_2 (10.0 mmol, 2.54 g), ($^{\text{Bu}}\text{PNN}$) FeCl_2 (0.10 mmol, 44 mg) and NaO^tBu (0.20 mmol, 19 mg) were dissolved in Et_2O (10 mL) in an autoclave. The vessel was purged with D_2 then pressurised with D_2 (10 bar). This was left to stir for 16 hours, then the pressure was released. The reaction mixture was distilled at atmospheric pressure to remove Et_2O (60 °C). The DBpin was distilled *in vacuo* (25 °C, 0.1 mbar), to give DBpin (2.22 g, 86%, >99%-d) as a colourless oil. ^1H NMR (500.12 MHz, CDCl_3): 1.27 (12H, s, CH_3). ^{11}B NMR (128.34 MHz, CDCl_3): 28.2 (s br), ^{13}C NMR (125.77 MHz, CDCl_3) 25.0, 83.4. ^2D NMR (76.75 MHz, toluene) 4.15 (br m). Analytical data were in accordance with those previously reported [22].

4.6. General procedure for one-pot deuteroboration of terminal alkenes.

B_2pin_2 (0.500 mmol, 127 mg), ($^{\text{Bu}}\text{PNN}$) FeCl_2 (0.005 mmol, 2.2 mg), and NaO^tBu (0.01 mmol, 0.8 mg) were dissolved in Et_2O (0.5 mL) in an autoclave. The vessel was purged with D_2 then pressurised with D_2 (4 bar). This was left to stir for 16 hours, then the pressure was released and the vessel transferred into a glovebox. Alkene (1.00 mmol) was added, and the reaction left to stir for 4 hours. The solvent was removed *in vacuo* and the residue purified by flash column chromatography on silica (with a mixture of 40/60 petroleum ether and diethyl ether). Volatiles were removed *in vacuo* to give the deuterio alkyl boronic ester.

Acknowledgements

S.P.T. thanks the Royal Society for a University Research Fellowship. D.R.W. thanks the Royal Society for a PhD studentship. J.H.D. and S.P.T. acknowledge GlaxoSmithKline, EPSRC, and the University of Edinburgh (PIII002) for post-doctoral funding.

6. References

- [1] C. Schmidt, First deuterated drug approved, *Nat. Biotechnol.* 35 (2017) 493–494.
- [2] N. Miyaura, A. Suzuki, Palladium-Catalyzed Cross-Coupling Reactions of Organoboron Compounds, *Chem. Rev.* 95 (1995) 2457–2483.
- [3] N. Miyaura, K. Yamada, A. Suzuki, A new stereospecific cross-coupling by the palladium-catalyzed reaction of 1-alkenylboranes with 1-alkenyl or 1-alkynyl halides, *Tetrahedron Lett.* 20 (1979) 3437–3440.
- [4] J.W.B. Fyfe, A.J.B. Watson, Recent Developments in

Chem. 3 (2017) 31–55.

- [5] K. Shirakawa, A. Arase, M. Hoshi, Preparation of (*E*)-1-alkenylboronic acid pinacol esters via transfer of alkenyl group from boron to boron, *Synthesis* (Stuttg). (2004) 1814–1820.
- [6] N.W.J. Ang, C.S. Buettner, S. Docherty, A. Bismuto, J.R. Carney, J.H. Docherty, M.J. Cowley, S.P. Thomas, Borane-Catalysed Hydroboration of Alkynes and Alkenes, *Synth.* 50 (2018) 803–808.
- [7] D.G. Hall, *Boronic Acids: Preparation and Applications in Organic Synthesis, Medicine and Materials* (Volume 1 and 2), Second Edi, WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, 2011.
- [8] J. V Obligacion, P.J. Chirik, Bis(imino)pyridine Cobalt-Catalyzed Alkene Isomerization–Hydroboration: A Strategy for Remote Hydrofunctionalization with Terminal Selectivity, *J. Am. Chem. Soc.* 135 (2013) 19107–19110.
- [9] C.S. Wei, C.A. Jiménez-Hoyos, M.F. Videa, J.F. Hartwig, M.B. Hall, Origins of the Selectivity for Borylation of Primary over Secondary C–H Bonds Catalyzed by Cp*–Rhodium Complexes, *J. Am. Chem. Soc.* 132 (2010) 3078–3091.
- [10] J.Y. Wu, B. Moreau, T. Ritter, Iron-Catalyzed 1,4-Hydroboration of 1,3-Dienes, *J. Am. Chem. Soc.* 131 (2009) 12915–12917.
- [11] T. Xi, Z. Lu, Cobalt-Catalyzed Ligand-Controlled Regioselective Hydroboration/Cyclization of 1,6-Enynes, *ACS Catal.* 7 (2017) 1181–1185.
- [12] C. Chen, X. Shen, J. Chen, X. Hong, Z. Lu, Iron-Catalyzed Hydroboration of Vinylcyclopropanes, *Org. Lett.* 19 (2017) 5422–5425.
- [13] Q. Huang, M.-Y. Hu, S.-F. Zhu, Cobalt-Catalyzed Cyclization/Hydroboration of 1,6-Diynes with Pinacolborane, *Org. Lett.* 21 (2019) 7883–7887.
- [14] S.R. Tamang, D. Bedi, S. Shafiei-Haghighi, C.R. Smith, C. Crawford, M. Findlater, Cobalt-Catalyzed Hydroboration of Alkenes, Aldehydes, and Ketones, *Org. Lett.* 20 (2018) 6695–6700.
- [15] B. Sundararaju, A. Fürstner, A trans-Selective Hydroboration of Internal Alkynes, *Angew. Chemie Int. Ed.* 52 (2013) 14050–14054.
- [16] F. Labre, Y. Gimbert, P. Bannwarth, S. Olivero, E. Duñach, P.Y. Chavant, Application of Cooperative Iron/Copper Catalysis to a Palladium-Free Borylation of Aryl Bromides with Pinacolborane, *Org. Lett.* 16 (2014) 2366–2369.
- [17] C.E. Tucker, J. Davidson, P. Knochel, Mild and Stereoselective Hydroborations of Functionalized Alkynes and Alkenes Using Pinacolborane, *J. Org. Chem.* 57 (1992) 3482–3485.
- [18] Macgregor, R.L. Webster, Room Temperature Iron-Catalyzed Transfer Hydrogenation and Regioselective Deuteration of Carbon–Carbon Double Bonds, *J. Am. Chem. Soc.* 141 (2019) 572–582.
- [19] P.L. Callaghan, R. Fernández-Pacheco, N. Jasim, S. Lachaize, T.B. Marder, R.N. Perutz, E. Rivalta, S. Sabo-Etienne, Photochemical oxidative addition of B–H bonds at ruthenium and rhodium, *Chem. Commun.* (2004) 242–243.
- [20] S. Bontemps, L. Vendier, S. Sabo-Etienne, Borane-Mediated Carbon Dioxide Reduction at Ruthenium: Formation of C1 and C2 Compounds, *Angew. Chemie Int. Ed.* 51 (2012) 1671–1674.
- [21] H. Braunschweig, F. Guethlein, L. Mailänder, T.B. Marder, Synthesis of Catechol-, Pinacol-, and Neopentylglycolborane through the Heterogeneous Catalytic B–B Hydrogenolysis of Diboranes(4), *Chem. – A Eur. J.* 19 (2013) 14831–14835.
- [22] L. Qiao, L. Zhang, G. Liu, Z. Huang, A highly efficient cobalt-catalyzed deuterogenolysis of diboron: Synthesis of deuterated pinacolborane and vinylboronates, *Tetrahedron.* 75 (2019) 4138–4142.
- [23] J.H. Docherty, J. Peng, A.P. Dominey, S.P. Thomas, Activation and discovery of earth-abundant metal catalysts using sodium *tert*-butoxide, *Nat. Chem.* 9 (2017) 595–600.
- [24] D. Peng, Y. Zhang, X. Du, L. Zhang, X. Leng, M.D. Walter, Z. Huang, Phosphinite-Iminopyridine Iron Catalysts for Chemoselective Alkene Hydrosilylation, *J. Am. Chem. Soc.* 135 (2013) 19154–19166.
- [25] E. Balaraman, B. Gnanaprakasam, L.J.W. Shimon, D. Milstein, Direct Hydrogenation of Amides to Alcohols and Amines under Mild Conditions, *J. Am. Chem. Soc.* 132 (2010) 16756–16758.
- [26] L. Zhang, D. Peng, X. Leng, Z. Huang, Iron-Catalyzed, Atom-Economical, Chemo- and Regioselective Alkene Hydroboration with Pinacolborane, *Angew. Chemie Int. Ed.* 52 (2013) 3676–3680.
- [27] B. Moreau, J.Y. Wu, T. Ritter, Iron-Catalyzed 1,4-Addition of α -Olefins to Dienes, *Org. Lett.* 11 (2009) 337–339.
- [28] T.E. Elton, G.E. Ball, M. Bhadbhade, L.D. Field, S.B. Colbran, Evaluation of Organic Hydride Donors as Reagents for the Reduction of Carbon Dioxide and Metal-Bound Formates, *Organometallics.* 37 (2018) 3972–3982.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof