

# Extremely Stable Thorium(IV) Dialkyl Complexes Supported by Rigid Tridentate 4,5-Bis(anilido)xanthene and 2,6-Bis(anilidomethyl)pyridine Ligands

Carlos A. Cruz, David J. H. Emslie,\* Laura E. Harrington, James F. Britten, and Craig M. Robertson

Department of Chemistry, McMaster University, 1280 Main Street West, Hamilton, Ontario L8S 4M1, Canada

Received October 5, 2006

A new NON-donor ligand, 4,5-bis(2,6-diisopropylanilino)-2,7-di-*tert*-butyl-9,9-dimethylxanthene ( $H_2$ -[XA<sub>2</sub>], **1**), was prepared by palladium-catalyzed coupling of 2,6-diisopropylaniline with the appropriate dibromoxanthene precursor. Stable  $K_2(dme)_2[XA_2]$  (**2**) and  $Na_2[XA_2]$  (**3**) salts were accessible by deprotonation of  $H_2[XA_2]$  with KH in dme or NaH in toluene. The thermally unstable lithium salt of McConville's 2,6-bis(2,6-diisopropylanilidomethyl)pyridine ligand ( $Li_2[BDPP]$ , **4**) was isolated by deprotonation with *n*BuLi or  $LiCH_2SiMe_3$  in hexanes at low temperature. Reaction of  $[ThCl_4(dme)_2]$  with  $Li_2[BDPP]$  or  $M_2(dme)_n[XA_2]$  resulted in the formation of pentagonal bipyramidal  $[LThCl_2(dme)]$  complexes ( $L = BDPP$ , **5**;  $XA_2$ , **6**). Subsequent reaction of **5** or **6** with  $LiCH_2SiMe_3$  gave base- and salt-free dialkyl complexes,  $[LTh(CH_2SiMe_3)_2]$  ( $L = BDPP$ , **7**;  $XA_2$ , **8**), which are stable for days in solution at 90 and 70 °C, respectively. Complexes **5**, **7**, and **8** were also accessible by initial combination of 2 or 4 equiv of  $LiCH_2SiMe_3$  with  $[ThCl_4(dme)_2]$ , followed by addition of  $H_2L$ . These reactions likely proceed by alkane elimination, but dialkyl or tetraalkyl thorium intermediates were not identified. The X-ray crystal structure of **8** suggests the presence of  $\alpha$ -agostic C–H–Th interactions for both alkyl groups. In solution, **7** and **8** exhibit temperature-dependent  $^1J_{C,H}$  coupling constants for  $ThCH_2$ , demonstrating the presence of  $\alpha$ -agostic interactions which become increasingly favored at lower temperature. Reaction of **5** with  $Li_2[BDPP]$  at 0 °C or **7** with  $H_2[BDPP]$  at 100 °C resulted in the formation of extremely sterically encumbered  $[Th(BDPP)_2]$  (**9**), which adopts a highly distorted six-coordinate geometry with the four anilido groups arranged in an approximate tetrahedron around thorium. Bis-ligand complexes were not accessible with the  $XA_2$  platform, presumably due to increased ligand rigidity.

## Introduction

Complexes of the early actinide elements, of which only thorium and uranium are naturally abundant and readily available, are unique due to the potential for significant covalency and f-orbital involvement in bonding.<sup>1,2</sup> Actinides also tend to form complexes of high Lewis acidity and possess ionic radii that

are similar or larger than those of the lanthanide ions and significantly larger than those for transition metals in the same oxidation state.<sup>3</sup> These properties, in addition to the distinctly non-lanthanide-like variety of oxidation states accessible for several of the early actinides (e.g., U(III)–U(VI)),<sup>1,4</sup> lead to the potential for early actinide complexes to facilitate unusual reactivity<sup>5</sup> that may not be accessible elsewhere in the periodic table.

To date, organoactinide chemistry has been dominated by the use of carbocyclic ancillaries such as  $C_5R_5$  and related ligands,<sup>6,7</sup> the tetramethylphospholyl anion,<sup>8</sup> dianionic  $C_8R_8$  or pentalene<sup>9</sup> ligands, carboranes,<sup>10</sup> arenes,<sup>11</sup> and the cycloheptatrienyl trianion.<sup>12</sup> In contrast, the chemistry of non-carbocycle-supported organoactinide complexes is much less developed, but is of great interest due to the enormous structural and electronic versatility afforded by such ligands.

Non-cyclopentadienyl actinide(IV) bis-hydrocarbyl (alkyl, aryl, or allyl) complexes are particularly rare and generally rely on the coordination of monoanionic ancillaries such as alkoxy/aryloxy ( $Th/U$ ),<sup>13–15</sup> amidinate ( $U$ ),<sup>16</sup> and tris(pyrazolyl)borate ( $U$ )<sup>17</sup> ligands. Other ancillaries such as the popular  $\beta$ -diketiminate anion<sup>18</sup> have not been reported to be effective in the

\* To whom correspondence should be addressed. Phone: 905-525-9140. Fax: 905-522-2509. E-mail: emslied@mcmaster.ca.

(1) Burns, C. J.; Neu, M. P.; Boukhalfa, H.; Gutowski, K. E.; Bridges, N. J.; Rogers, R. D. Chapter 3.3: The Actinides. In *Comprehensive Coordination Chemistry II*; Parkin, G. F. R., Ed.; Elsevier Ltd.: San Diego, 2004; Vol. 3, p 189.

(2) For example: (a) Diaconescu, P. L.; Arnold, P. L.; Baker, T. A.; Mindiola, D. J.; Cummins, C. C. *J. Am. Chem. Soc.* **2000**, *122*, 6108. (b) Marks, T. J.; Kolb, J. R. *Chem. Rev.* **1977**, *77*, 263. (c) Kaltsoyannis, N.; Scott, P. *Chem. Commun.* **1998**, 1665. (d) Nakai, H.; Hu, X.; Zakharov, L. N.; Rheingold, A. L.; Meyer, K. *Inorg. Chem.* **2004**, *43*, 855. (e) Castro-Rodriguez, I.; Olsen, K.; Gantzel, P.; Meyer, K. *J. Am. Chem. Soc.* **2003**, *125*, 4565. (f) Mazzanti, M.; Wietzke, R.; Pécaut, J.; Latour, J.-M.; Maldivi, P.; Remy, M. *Inorg. Chem.* **2002**, *41*, 2389. (g) Brennan, J. G.; Stults, S. D.; Andersen, R. A.; Zalkin, A. *Organometallics* **1988**, *7*, 1329. (h) Sockwell, S. C.; Hanusa, T. P. *Inorg. Chem.* **1990**, *29*, 76. (i) Jensen, M. P.; Bond, A. H. *J. Am. Chem. Soc.* **2002**, *124*, 9870. (j) Guillaumont, D. J. *Phys. Chem. A* **2004**, *108*, 6893. (k) Petit, L.; Joubert, L.; Maldivi, P.; Adamo, C. *J. Am. Chem. Soc.* **2006**, *128*, 2190. (l) Cloke, F. G. N.; Green, J. C.; Kaltsoyannis, N. *Organometallics* **2004**, *23*, 832. (m) Straka, M.; Hrobárik, P.; Kaupp, M. *J. Am. Chem. Soc.* **2005**, *127*, 2591. (n) Streitwieser, A., Jr.; Dempf, D.; La, M.; Mar, G. N.; Karraker, D. G.; Edelstein, N. *J. Am. Chem. Soc.* **1971**, *93*, 7343. (o) Pepper, M.; Bursten, B. E. *Chem. Rev.* **1991**, *91*, 719.

(3) Revised effective ionic radii (C.N. 6) of Shannon and Prewitt:  $U^{3+}$  1.03,  $La^{3+}$  1.03,  $Lu^{3+}$  0.86,  $Th^{4+}$  0.94,  $U^{4+}$  0.89,  $Ce^{4+}$  0.87,  $Hf^{4+}$  0.71,  $Ti^{4+}$  0.61,  $U^{5+}$  0.76,  $Ta^{5+}$  0.64,  $V^{5+}$  0.54,  $U^{6+}$  0.73,  $W^{6+}$  0.60,  $Cr^{6+}$  0.44. A. Shannon, R. D. *Acta Crystallogr.* **1976**, *A32*, 751.

(4) Evans, W. J.; Kozimor, S. A. *Coord. Chem. Rev.* **2006**, *250*, 911.

preparation of actinide(IV) dialkyls,<sup>19–21</sup> perhaps due to provision of insufficient electronic saturation, a relatively small metal binding pocket, and a tendency toward ligand degradation reactions<sup>19</sup> under a variety of conditions.<sup>22</sup> The only dialkyl actinide(IV) complexes bearing a single dianionic, non-carbocyclic supporting ligand are [(<sup>DIPP</sup>NCOCN)U(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>2</sub>] and [(<sup>t</sup>BuNON)M(R)<sub>2</sub>] (M = Th or U; R = C<sub>3</sub>H<sub>5</sub> or CH<sub>2</sub>SiMe<sub>3</sub>)

(5) For example: (a) Moloy, K. G.; Marks, T. J. *J. Am. Chem. Soc.* **1984**, *106*, 7051. (b) Katahira, D. A.; Moloy, K. G.; Marks, T. J. *Organometallics* **1982**, *1*, 1723. (c) Edwards, P. G.; Parry, J. S.; Read, P. W. *Organometallics* **1995**, *14*, 3649. (d) Radu, N. S.; Engeler, M. P.; Gerlach, C. P.; Tilley, T. D.; Rheingold, A. L. *J. Am. Chem. Soc.* **1995**, *117*, 3621. (e) Summerscales, O. T.; Cloke, F. G. N.; Hitchcock, P. B.; Green, J. C.; Hazari, N. *Science* **2006**, *311*, 829. (f) Castro-Rodriguez, I.; Meyer, K. J. *Am. Chem. Soc.* **2005**, *127*, 11242. (g) Castro-Rodriguez, I.; Nakai, H.; Zakharov, L. N.; Rheingold, A. L.; Meyer, K. *Science* **2004**, *305*, 1757. (h) Peters, R. G.; Warner, B. P.; Burns, C. J. *J. Am. Chem. Soc.* **1999**, *121*, 5585. (i) Pool, J. A.; Scott, B. L.; Kiplinger, J. L. *Chem. Commun.* **2005**, 2591. (j) Parry, J.; Carmona, E.; Coles, S.; Hursthouse, M. J. *Am. Chem. Soc.* **1995**, *117*, 2649. (k) Diaconescu, P. L.; Odom, A. L.; Agapie, T.; Cummins, C. C. *Organometallics* **2001**, *20*, 4993. (l) Evans, W. J.; Kozimor, S. A.; Ziller, J. W. *J. Am. Chem. Soc.* **2003**, *125*, 14264. (m) Korobkov, I.; Gambarotta, S. *Organometallics* **2004**, *23*, 5379. (n) Castro-Rodriguez, I.; Nakai, H.; Gantzel, P.; Zakharov, L. N.; Rheingold, A. L.; Meyer, K. *J. Am. Chem. Soc.* **2003**, *125*, 15734. (o) Berthet, J.-C.; Ephritikhine, M. *New J. Chem.* **1992**, *16*, 767. (p) Lukens, W. W., Jr.; Beshouri, S. M.; Bloch, L. L.; Andersen, R. A. *J. Am. Chem. Soc.* **1996**, *118*, 901.

(6) (a) Marks, T. J.; Day, V. W. Actinide Hydrocarbyl and Hydride Chemistry. In *Fundamental and Technological Aspects of Organo-f-Element Chemistry*; Nato Science Series C: Vol. 155; Marks, T. J., Fraga, I. L., Eds.; D. Reidel Publishing Company: Dordrecht, 1985; p 115. (b) Takats, J. Organoactinide Complexes Containing Classical Ligands. In *Fundamental and Technological Aspects of Organo-f-Element Chemistry*; Nato Science Series C: Vol. 155; Marks, T. J., Fraga, I. L., Eds.; D. Reidel Publishing Company: Dordrecht, 1985; p 159. (c) Evans, W. J. *J. Organomet. Chem.* **2002**, *647*, 2. (d) Jia, L.; Yang, X.; Stern, C. L.; Marks, T. J. *Organometallics* **1997**, *16*, 842. (e) Golden, J. T.; Kazul'kin, D. N.; Scott, B. L.; Voskoboinikov, A. Z.; Burns, C. J. *Organometallics* **2003**, *22*, 3971. (f) Baudry, D.; Bulot, E.; Ephritikhine, M. *Chem. Commun.* **1989**, 1316. (g) Stubbart, B. D.; Stern, C. L.; Marks, T. J. *Organometallics* **2003**, *22*, 4836.

(7) Fendrick, C. M.; Schertz, L. D.; Day, V. W.; Marks, T. J. *Organometallics* **1988**, *7*, 1828.

(8) (a) Cendrowski-Guillaume, S. M.; Nierlich, M.; Ephritikhine, M. *J. Organomet. Chem.* **2002**, *643*, 209. (b) Gradoz, P.; Baudry, D.; Ephritikhine, M.; Lance, M.; Nierlich, M.; Vigner, J. *J. Organomet. Chem.* **1994**, *466*, 107. (c) Gradoz, P.; Boisson, C.; Baudry, D.; Lance, M.; Nierlich, M.; Vigner, J.; Ephritikhine, M. *Chem. Commun.* **1992**, 1720. (d) Gradoz, P.; Baudry, D.; Ephritikhine, M.; Nief, F.; Mathey, F. *Dalton Trans.* **1992**, 3047.

(9) Seyferth, D. *Organometallics* **2004**, *23*, 3562.

(10) Xie, Z. *Coord. Chem. Rev.* **2002**, *231*, 23.

(11) Baudry, D.; Bulot, E.; Charpin, P.; Ephritikhine, M.; Lance, M.; Nierlich, M.; Vigner, J. *J. Organomet. Chem.* **1989**, *371*, 155.

(12) (a) Arliguie, T.; Lance, M.; Nierlich, M.; Vigner, J.; Ephritikhine, M. *Chem. Commun.* **1995**, 183. (b) Arliguie, T.; Lance, M.; Nierlich, M.; Ephritikhine, M. *Dalton Trans.* **1997**, 2501.

(13) Clark, D. L.; Grumbine, S. K.; Scott, B. L.; Watkin, J. G. *Organometallics* **1996**, *15*, 949.

(14) (a) Korobkov, I.; Athimoolam, A.; Gambarotta, S. *Organometallics* **2004**, *23*, 6248. (b) Baudin, C.; Ephritikhine, M. *J. Organomet. Chem.* **1989**, *364*, C1. (c) Baudin, C.; Baudry, D.; Ephritikhine, M.; Lance, M.; Navaza, A.; Nierlich, M.; Vigner, J. *J. Organomet. Chem.* **1991**, *415*, 59. (d) Brunelli, M.; Perego, G.; Lugli, G.; Mazzei, A. *Dalton Trans.* **1979**, 861.

(15) Beshouri, S. M.; Fanwick, P. E.; Rothwell, I. P.; Huffman, J. C. *Organometallics* **1987**, *6*, 2498.

(16) Wedler, M.; Knösel, F.; Edelmann, F. T.; Behrens, U. *Chem. Ber.* **1992**, *125*, 1313.

(17) (a) Domingos, Â.; Marques, N.; de Matos, A. P.; Santos, I.; Silva, M. *Organometallics* **1994**, *13*, 654. (b) Campello, M. P. C.; Calhorda, M. J.; Domingos, Â.; Galvão, A.; Leal, J. P.; de Matos, A. P.; Santos, I. *J. Organomet. Chem.* **1997**, *538*, 223.


(18) Bourget-Merle, L.; Lappert, M. F.; Severn, J. R. *Chem. Rev.* **2002**, *102*, 3031.

(19) Hitchcock, P. B.; Lappert, M. F.; Liu, D.-S. *J. Organomet. Chem.* **1995**, *488*, 241.

(20) Hitchcock, P. B.; Hu, J.; Lappert, M. F.; Tian, S. *J. Organomet. Chem.* **1997**, *536–537*, 473.

(21) (a) Wright, R. J.; Power, P. P.; Scott, B. L.; Kiplinger, J. L. *Organometallics* **2004**, *23*, 4801. (b) Antunes, M. A.; Dias, M.; Monteiro, B.; Domingos, Â.; Santos, I. C.; Marques, N. *Dalton Trans.* **2006**, 3368.

## Scheme 1. Synthesis of Proligand H<sub>2</sub>XA<sub>2</sub> (1)



reported very recently by Leznoff [<sup>DIPP</sup>NCOCN = O(CH<sub>2</sub>CH<sub>2</sub>-NAr)<sub>2</sub>; <sup>t</sup>BuNON = O(SiMe<sub>2</sub>N<sup>t</sup>Bu)<sub>2</sub>; Ar = 2,6-diisopropylphenyl].<sup>23,24</sup>

We are interested in the preparation of non-carbocyclic organoactinide complexes as a result of their largely untapped potential in both traditional organometallic catalysis and the development of novel reactivity. Of particular interest are thorium(IV) dialkyl complexes of sufficient thermal stability to allow investigation of reactivity over a wide temperature range and to serve as precursors for the synthesis of new, highly reactive organometallic derivatives. With this goal in mind, we have focused our efforts on the use of *extremely rigid and planar* tridentate ancillary ligands, and in particular those that provide a binding pocket of a suitable size to accommodate a large actinide metal and contain only robust structural elements (e.g., avoidance of isolated imine groups).<sup>25,26</sup>

Our preference for rigid ligands stems from the expectation that they will (a) allow access to coordination environments that are dictated by design rather than the preferences of the central metal and/or coligands and (b) ensure that well-intentioned steric bulk is not positioned in such a way as to significantly limit its effectiveness. As a result, various modes of decomposition are expected to become less favorable, especially those involving sterically hindered transition states or the formation of dinuclear or bis-ligand complexes. Rigid ligands are also expected to be more amenable to steric tuning since the effects of steric bulk are not easily mitigated by alterations in the ligand geometry or hapticity and are, therefore, more predictable.

Herein we report the preparation of uniquely robust base- and salt-free thorium(IV) dialkyls, as well as dichloro and bis-ligand complexes, using the 2,6-bis(2,6-diisopropylanilido)methylpyridine dianion (BDPP, Scheme 1) and a new, extremely rigid and planar NON-donor, 4,5-bis(2,6-diisopropylanilido)-2,7-di-*tert*-butyl-9,9-dimethylxanthene (XA<sub>2</sub>, Scheme 2).

## Results and Discussion

**Ligand Synthesis.** The new NON-donor proligand H<sub>2</sub>[XA<sub>2</sub>] (1) was synthesized in 86% isolated yield by Hartwig–


(22) For example: (a) Bailey, B. C.; Basuli, F.; Huffman, J. C.; Mindiola, D. J. *Organometallics* **2006**, *25*, 3963. (b) Basuli, F.; Bailey, B. C.; Huffman, J. C.; Mindiola, D. J. *Organometallics* **2005**, *24*, 3321. (c) Basuli, F.; Bailey, B. C.; Watson, L. A.; Tomaszewski, J.; Huffman, J. C.; Mindiola, D. J. *Organometallics* **2005**, *24*, 1886. (d) Basuli, F.; Tomaszewski, J.; Huffman, J. C.; Mindiola, D. J. *J. Am. Chem. Soc.* **2003**, *125*, 10170. (e) Hamaki, H.; Takeda, N.; Tokitoh, N. *Organometallics* **2006**, *25*, 2457. (f) Gregory, E. A.; Lachicotte, R. J.; Holland, P. L. *Organometallics* **2005**, *24*, 1803. (g) Yokota, S.; Tachi, Y.; Itoh, S. *Inorg. Chem.* **2002**, *41*, 1342.

(23) Jantunen, K. C.; Haftbaradaran, F.; Katz, M. J.; Batchelor, R. J.; Schatte, G.; Leznoff, D. B. *Dalton Trans.* **2005**, 3083.

(24) Jantunen, K. C.; Batchelor, R. J.; Leznoff, D. B. *Organometallics* **2004**, *23*, 2186.

(25) (a) Black, D. G.; Swenson, D. C.; Jordan, R. F.; Rogers, R. D. *Organometallics* **1995**, *14*, 3539. (b) Woodman, P. R.; Alcock, N. W.; Munslow, I. J.; Sanders, C. J.; Scott, P. *Dalton Trans.* **2000**, 3340. (c) Emslie, D. J. H.; Piers, W. E.; Parvez, M. *Dalton Trans.* **2003**, 2615.

(26) Tsurugi, H.; Matsuo, Y.; Yamagata, T.; Mashima, K. *Organometallics* **2004**, *23*, 2797.

**Scheme 2. Preparation and Thermal Stability of 4**


Buchwald coupling of 4,5-dibromo-2,7-di-*tert*-butyl-9,9-dimethylxanthene with 2,6-diisopropylaniline (Scheme 1). Stirring  $H_2[XA_2]$  with excess KH in 1,2-dimethoxyethane (dme) at room temperature for 5 h gave base-stabilized  $K_2(dme)_2[XA_2]$  (**2**) in 81% yield. Alternatively, base-free  $Na_2[XA_2]$  (**3**) is accessible by refluxing  $H_2[XA_2]$  with excess NaH in toluene for several days.

The NNN-donor ligand  $H_2[BDPP]$  was prepared as reported by McConville et al. This ligand has previously been employed in titanium, zirconium, and tantalum chemistry, using amine ( $H_2[BDPP] + [M(NR_2)_n]$ ) or chlorotrimethylsilane ( $BDPP(SiMe_3)_2 + [MCl_n]$ ) elimination to effect ligand attachment.<sup>27–29</sup> An alkali or alkaline earth metal salt of the BDPP dianion, which would allow direct access to chloro complexes by salt metathesis, has not been reported. However, we found that reaction of  $H_2[BDPP]$  with 2 equiv of an alkyl lithium reagent ( $LiCH_2SiMe_3$  or  $nBuLi$ ) in hexanes at  $-78\text{ }^\circ\text{C}$  resulted in precipitation of base-free  $Li_2[BDPP]$  (**4**), which is isolated in 87% yield as a bright yellow solid (Scheme 2). This compound is unusually temperature sensitive, decomposing in minutes in benzene at room temperature.<sup>30</sup> Solid **4** is also significantly decomposed after several hours at room temperature, but can be stored for weeks without appreciable decomposition at  $-30\text{ }^\circ\text{C}$ .

Both the BDPP and  $XA_2$  ligands are structurally related to the diamidoamine ligand  $[(Me_3SiN(CH_2CH_2NSiMe_3)_2)]^{2-}$  previously employed by Cloke et al. for the synthesis of dichloro and bis(ligand) thorium(IV) complexes.<sup>31</sup> However, the rigidity of the ligand backbone increases dramatically in the order  $[(Me_3SiN(CH_2CH_2NSiMe_3)_2)]^{2-} < BDPP < XA_2$ .

**Dichloride Complexes.** Reaction of 1 equiv of  $Li_2[BDPP]$  (**4**) with  $[ThCl_4(dme)_2]$  in benzene<sup>30</sup> at  $0\text{ }^\circ\text{C}$  gave  $[(BDPP)ThCl_2(dme)]$  (**5**) in 51% yield. Similarly,  $[(XA_2)ThCl_2(dme)]$  (**6**) was isolated by reaction of  $Na_2[XA_2]$  (**3**) with  $[ThCl_4(dme)_2]$  in toluene at room temperature (69% yield; Scheme 3). In contrast, base-coordinated  $K_2(dme)_2[XA_2]$  (**2**) is considerably less reactive than **3**, despite similar solubilities (of **2** and **3** in  $C_6H_6$  or  $C_7H_8$ ), and the reaction is complete only after refluxing in toluene for 12 h.

Complex **5** could also be prepared by reaction of 2 equiv of  $LiCH_2SiMe_3$  with  $[ThCl_4(dme)_2]$  for 1 h at  $0\text{ }^\circ\text{C}$ , followed by cooling to  $-78\text{ }^\circ\text{C}$  and addition of  $H_2BDPP$ . In this case, ligand attachment likely proceeds by alkane elimination from a source of “ $ThCl_2(CH_2SiMe_3)_2$ ”. However, the reaction of  $LiCH_2SiMe_3$  with  $[ThCl_4(dme)_2]$  forms a mixture of products, the nature of

**Scheme 3. Synthesis of Dichloro Complexes 5 and 6**

which has not been determined (Scheme 3). Nevertheless, this is the preferred route for the synthesis of **5** due to greater simplicity and an improved yield (85%). By contrast, attempts to prepare **6** by this method gave solutions containing  $H_2[XA_2]$  and  $SiMe_4$  as the only soluble products, presumably due to thermal decomposition of the alkyl thorium precursor in preference to reaction with the rigid  $H_2[XA_2]$  proligand.

$^1H$  NMR spectra of **5** and **6** between  $20$  and  $-90\text{ }^\circ\text{C}$  show the presence of one molecule of symmetrically coordinated dme and a lone  $CHMe_2$  signal, consistent with  $C_{2v}$  symmetric pentagonal bipyramidal products. Single crystals of  $[(BDPP)ThCl_2(\kappa^2-dme)] \cdot 2\text{toluene}$  were grown by slow diffusion of hexanes into a toluene solution of **5** at  $-30\text{ }^\circ\text{C}$ . The X-ray crystal structure of **5** (Figure 1, Table 1) confirms a distorted pentagonal bipyramidal geometry with the two chloride anions occupying apical positions [ $Cl(1)-Th-Cl(2) = 156.38(10)^\circ$ ;  $Th-Cl = 2.698(3)$  and  $2.686(3)\text{ \AA}$ ]. As anticipated, the BDPP ligand is approximately planar and binds thorium via short  $Th-N_{anilido}$  contacts [ $2.305(9)$  and  $2.321(8)\text{ \AA}$ ] and a longer  $Th-N_{py}$  bond [ $2.568(9)\text{ \AA}$ ]. However, while  $N(1)$ ,  $N(3)$ ,  $O(1)$ ,  $O(2)$ , and  $Th$  lie in a plane, the metal is located  $0.33(1)\text{ \AA}$  above the  $N(1)-N(2)-N(3)$  plane of the BDPP ligand. In this way, only one chloro ligand is positioned directly between the bulky isopropyl groups, while the other is located in a more open region of the thorium coordination sphere. To accommodate the more sterically contentious chloride anion, both 2,6-diisopropylphenyl rings rotate to give  $C(14) \cdots C(26) = 6.35(2)\text{ \AA}$  and  $C(17) \cdots C(29) = 7.26(2)\text{ \AA}$ .

The dme molecule is  $\kappa^2$ -coordinated in the pentagonal plane and is bound via long and unequal  $Th-O$  contacts  $\{Th-O(1) = 2.674(8)\text{ \AA}$  and  $Th-O(2) = 2.724(8)\text{ \AA}\}$ , likely due to steric pressure at the metal; cf.  $2.564(8)-2.620(8)\text{ \AA}$  in  $[ThBr_4(\kappa^2-dme)_2]$ ,<sup>32</sup>  $2.620(5)\text{ \AA}$  in  $[LTh(NH_2)(\kappa^1-dme)]^-$ , and  $2.613(3)\text{ \AA}$  in  $[LTh(Cl)(\kappa^1-dme)]^-$   $\{L = 2,2'\text{-methylenebis}(6\text{-tert-butyl-4-methylphenolate})\}$ .<sup>33</sup> In contrast, the  $Th-N_{py}$  bond [ $2.568(9)\text{ \AA}$ ] is atypically short. For example,  $Th-N$  is  $2.72(1)-2.80(1)\text{ \AA}$  in  $[Th(\text{quiolinolate})_4(dmf)]$ ,<sup>34</sup>  $2.730(6)\text{ \AA}$  in  $\{[Th(OCHEt_2)_3(\mu-OCHEt_2)(Py)_2]\}_2$ ,  $2.752(7)\text{ \AA}$  in  $cis-[Th(OtBu)_4(Py)_2]$ ,<sup>35</sup> and  $2.662(8)$ ,  $2.696(8)\text{ \AA}$  in  $cis-[Th(OC_6H_3Me_2-2,6)_4(Py)_2]$ .<sup>36</sup> The only  $Th-NC_5R_5$  bonds of comparable length are found in  $[Th(OC_6H_3tBu-2,6)_2(C,N-2,6\text{-lutidinyl})_2]$  [ $2.61(1)\text{ \AA}$ ]<sup>15</sup> and  $[Th(BDPP)_2]$  (**9**) [ $2.615(6)\text{ \AA}$ , *vide infra*]. The former is a special case since significant delocalization of negative charge to the N-donor can occur. Therefore, the short  $Th-N_{py}$  bond lengths observed in this work are likely a result of incorporation of the pyridine unit into a rigid ligand framework and perhaps the

(27) Guérin, F.; McConville, D. H.; Vittal, J. J. *Organometallics* **1995**, *14*, 3154.

(28) (a) Guérin, F.; McConville, D. H.; Vittal, J. J.; Yap, G. A. P. *Organometallics* **1998**, *17*, 5172. (b) Guérin, F.; McConville, D. H.; Vittal, J. J.; Yap, G. A. P. *Organometallics* **1998**, *17*, 1290. (c) Guérin, F.; McConville, D. H.; Payne, N. C. *Organometallics* **1996**, *15*, 5085.

(29) Guérin, F.; McConville, D. H.; Vittal, J. J. *Organometallics* **1996**, *15*, 5586.

(30)  $Li_2[BDPP]$  (**4**) shows similar thermal stability in both benzene and THF (extensive decomposition in minutes at room temperature) and is considerably less stable in toluene (decomposition occurs rapidly even at  $-30\text{ }^\circ\text{C}$ ). The thermal decomposition of **4** in benzene is not clean, and a mixture of unidentified air-sensitive products is formed.

(31) Wilson, D. J.; Sebastian, A.; Cloke, F. G. N.; Avent, A. G.; Hitchcock, P. B. *Inorg. Chim. Acta* **2003**, *345*, 89.


(32) Rabinovich, D.; Scott, B. L.; Nielsen, J. B.; Abney, K. D. *J. Chem. Crystallogr.* **1999**, *29*, 243.

(33) Korobkov, I.; Gambarotta, S.; Yap, G. P. A. *Angew. Chem., Int. Ed.* **2003**, *42*, 4958.

(34) Barton, R. J.; Dabeka, R. W.; Shengzhi, H.; Mihichuk, L. M.; Pizzey, M.; Robertson, B. E.; Wallace, W. J. *Acta Crystallogr.* **1983**, *C39*, 714.

(35) Clark, D. L.; Watkin, J. G. *Inorg. Chem.* **1993**, *32*, 1766.

(36) Berg, J. M.; Clark, D. L.; Huffman, J. C.; Morris, D. E.; Sattelberger, A. P.; Streib, W. E.; van der Sluys, W. G.; Watkin, J. G. *J. Am. Chem. Soc.* **1992**, *114*, 10811.



**Figure 1.** Molecular structure of 5·2toluene with thermal ellipsoids at the 50% probability level. Hydrogen atoms and solvent omitted for clarity. In view b, only the N-C<sub>ippo</sub> carbon atoms of the N-Aryl rings are shown.

**Table 1. Crystallographic Data Collection and Refinement Parameters and Selected Bond Distances (Å) for Complexes 5, 8, and 9**


|                                                               | 5·2toluene                                                                                     | 8·toluene                                                          | 9                                                 |
|---------------------------------------------------------------|------------------------------------------------------------------------------------------------|--------------------------------------------------------------------|---------------------------------------------------|
| formula                                                       | C <sub>91</sub> H <sub>126</sub> Cl <sub>4</sub> N <sub>6</sub> O <sub>4</sub> Th <sub>2</sub> | C <sub>62</sub> H <sub>92</sub> N <sub>2</sub> OSi <sub>2</sub> Th | C <sub>62</sub> H <sub>82</sub> N <sub>6</sub> Th |
| fw                                                            | 1973.86                                                                                        | 1169.60                                                            | 1143.38                                           |
| cryst syst                                                    | monoclinic                                                                                     | monoclinic                                                         | monoclinic                                        |
| space group                                                   | C2/c                                                                                           | P2(1)/n                                                            | C2/c                                              |
| a (Å)                                                         | 28.7133(8)                                                                                     | 16.6445(5)                                                         | 14.901(15)                                        |
| b (Å)                                                         | 15.8253(5)                                                                                     | 22.5599(7)                                                         | 17.895(17)                                        |
| c (Å)                                                         | 21.3589(7)                                                                                     | 16.8509(5)                                                         | 21.354(16)                                        |
| α (deg)                                                       | 90                                                                                             | 90                                                                 | 90                                                |
| β (deg)                                                       | 107.398(2)                                                                                     | 107.928(2)                                                         | 106.72(3)                                         |
| γ (deg)                                                       | 90                                                                                             | 90                                                                 | 90                                                |
| volume (Å <sup>3</sup> )                                      | 9261.4(5)                                                                                      | 6020.2(3)                                                          | 5453(9)                                           |
| Z                                                             | 4                                                                                              | 4                                                                  | 4                                                 |
| density (calcd; mg/m <sup>3</sup> )                           | 1.449                                                                                          | 1.288                                                              | 1.393                                             |
| μ (mm <sup>-1</sup> )                                         | 3.374                                                                                          | 2.555                                                              | 2.779                                             |
| F(000)                                                        | 4074                                                                                           | 2408                                                               | 2344                                              |
| cryst size (mm <sup>3</sup> )                                 | 0.22 × 0.22 × 0.18                                                                             | 0.10 × 0.08 × 0.01                                                 | 0.12 × 0.10 × 0.03                                |
| θ range for collection (deg)                                  | 1.49 to 27.54                                                                                  | 1.50 to 27.49                                                      | 1.83 to 25.50                                     |
| no. of reflns collected                                       | 40 216                                                                                         | 32 615                                                             | 20 838                                            |
| no. of indep reflns                                           | 10 567                                                                                         | 13 480                                                             | 5071                                              |
| completeness to θ <sub>max</sub>                              | 98.9%                                                                                          | 97.5%                                                              | 100.0%                                            |
| max. and min. transmn                                         | 0.55 and 0.455                                                                                 | 0.98 and 0.8200                                                    | 0.920 and 0.711                                   |
| GOF on F <sup>2</sup>                                         | 1.003                                                                                          | 1.002                                                              | 1.100                                             |
| final R <sub>1</sub> [I > 2σ(I)] <sup>a</sup>                 | 0.0705                                                                                         | 0.0570                                                             | 0.0698                                            |
| Th–N <sub>amido</sub>                                         | 2.305(9), 2.321(8)                                                                             | 2.291(4), 2.312(4)                                                 | 2.373(7), 2.488(7)                                |
| Th–E (E = N <sub>py</sub> or O <sub>xant</sub> ) <sup>b</sup> | 2.568(9)                                                                                       | 2.535(4)                                                           | 2.614(7)                                          |
| Th–Cl or Th–C                                                 | 2.686(3), 2.698(3)                                                                             | 2.467(6), 2.484(6)                                                 |                                                   |
| Th–O <sub>dme</sub>                                           | 2.674(8), 2.724(8)                                                                             |                                                                    |                                                   |

<sup>a</sup> For **5**, **8**, and **9**: *T* = 173(2) K, wavelength = 0.71073 Å, absorption correction = semiempirical from equivalents, and refinement method = full-matrix least-squares on *F*<sup>2</sup>. <sup>b</sup>E = the central neutral donor of the BDPP and XA<sub>2</sub> ligands.

enhanced donor properties of N<sub>py</sub> (relative to an unsubstituted pyridine) as a result of 2,6-dialkyl substitution. All other metal–ligand bonds are in the usual range. For comparison, Th–N<sub>anilido</sub> is 2.327(6)–2.378(7) Å in [L<sub>2</sub>ThCl]<sup>−</sup> {L = 1,3-bis(2,6-diisopropylanilido)propane}<sup>37</sup> and 2.299(7), 2.304(6) Å in [Th{N(SiMe<sub>3</sub>)<sub>2</sub>(NMePh)<sub>2</sub>}],<sup>38</sup> while Th–Cl is 2.620(6)–2.697(1) Å in [L<sub>2</sub>ThCl<sub>2</sub>] {L = Tp, ArC(NSiMe<sub>3</sub>)<sub>2</sub>, and HC(CPhNSiMe<sub>3</sub>)<sub>2</sub>; Ar = C<sub>6</sub>H<sub>2</sub>(CF<sub>3</sub>)<sub>3</sub>-2,4,6}<sup>20,39</sup> and 2.673(1), 2.721(2) Å in [L'ThCl<sub>2</sub>(THF)] {L' = Me<sub>3</sub>SiN(C<sub>2</sub>H<sub>4</sub>NSiMe<sub>3</sub>)<sub>2</sub>}.<sup>31</sup>

**Dialkyl Complexes.** Reaction of the dichlorides, **5** and **6**, with 2 equiv of LiCH<sub>2</sub>SiMe<sub>3</sub> gave the base- and salt-free dialkyl complexes [LTh(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>2</sub>] {L = BDPP (**7**) and XA<sub>2</sub> (**8**)} in quantitative yield by <sup>1</sup>H NMR (68% and 63% isolated yields, respectively). However, a more straightforward route to these complexes involved reaction of [ThCl<sub>4</sub>(dme)<sub>2</sub>] with 4 equiv of LiCH<sub>2</sub>SiMe<sub>3</sub> at 0 °C, followed by addition of H<sub>2</sub>[BDPP] or H<sub>2</sub>–

#### Scheme 4. Synthesis of Dialkyl Complexes 7 and 8



[XA<sub>2</sub>]. Using this method, which likely proceeds by alkane elimination<sup>40</sup> from a source of “Th(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>4</sub>(dme)<sub>n</sub>”, tan **7** and white **8** may be isolated in 82% and 49% yield, respectively (Scheme 4).


The initial reaction between [ThCl<sub>4</sub>(dme)<sub>2</sub>] and LiCH<sub>2</sub>SiMe<sub>3</sub> proceeds cleanly and reproducibly to form a product or mixture of products giving rise to a single set of OMe, OCH<sub>2</sub>, SiMe<sub>3</sub>,

(37) Athimoolam, A.; Gambarotta, S.; Korobkov, I. *Organometallics* **2005**, *24*, 1996.

(38) Barnhart, D. M.; Clark, D. L.; Grumbine, S. K.; Watkin, J. G. *Inorg. Chem.* **1995**, *34*, 1695.

(39) (a) Domingos, Á.; Marçalo, J.; Santos, I.; de Matos, A. P. *Polyhedron* **1990**, *9*, 1645. (b) Wedler, M.; Knösel, F.; Noltemeyer, M.; Edelmann, F. T.; Behrens, U. *J. Organomet. Chem.* **1990**, *388*, 21.

(40) Propene elimination from the reaction of [U(allyl)<sub>3</sub>] with 2 ROH (R = Et, CHMe<sub>2</sub>, or CMe<sub>3</sub>) has been reported: see ref 14d.



**Figure 2.** Variable-temperature  $^1\text{H}$  NMR spectra of **7** in  $d_8$ -toluene (\* = toluene).

and  $\text{CH}_2\text{SiMe}_3$  signals in the  $^1\text{H}$  NMR spectrum at  $20^\circ\text{C}$ , with no change down to  $-90^\circ\text{C}$ . Unfortunately, the oily nature and thermal instability (decomposition is complete after 1.5 h at  $20^\circ\text{C}$ ) of this material precluded its isolation. However, the formation of a tetraalkyl derivative<sup>41–44</sup> under the conditions described above seems likely given that  $[\text{Th}(\text{CH}_3)_4(\text{dmpe})_2]$ ,<sup>41</sup>  $[\text{Th}(\text{CH}_2\text{Ph})_4]$ ,<sup>44</sup>  $[\text{Th}(\text{CH}_2\text{C}_6\text{H}_3\text{Me}_2-3,5)_4]$ ,<sup>43</sup> and  $[\text{Th}(\text{C}_3\text{H}_5)_4]$ <sup>43</sup> are formed in related reactions of  $[\text{ThCl}_4\text{L}_x]$   $\{\text{L}_x = (\text{dmpe})_2$  or  $(\text{THF})_3\}$  with 4 equiv of  $\text{LiR}$  at temperatures below  $0^\circ\text{C}$ . That said, the involvement of ate-complexes structurally related to  $[\text{MeTh}(\mu\text{-Me})_6\{\text{Li}(\text{TMEDA})\}_3]$ <sup>45</sup> and  $[\text{UR}_6\text{Li}_2\text{L}_x]$ <sup>46</sup> or *in-situ* ligand deprotonation by remaining  $\text{LiCH}_2\text{SiMe}_3$  cannot be ruled out.

At  $60^\circ\text{C}$  in  $d_8$ -toluene, the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of complexes **7** and **8** exhibit a single set of  $\text{ThCH}_2$  and  $\text{CHMe}_2$  resonances; the  $\text{ThCH}_2$  resonance is located at  $-0.32$  (**7**) or  $-0.17$  (**8**) ppm in the  $^1\text{H}$  NMR and 89 (**7**) or 97 (**8**) ppm in the  $^{13}\text{C}$  NMR. However, upon cooling to  $-80^\circ\text{C}$ , the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of both complexes show two distinct  $\text{CH}_2\text{SiMe}_3$  groups, two  $\text{CHMe}_2$  signals, and ligand backbone resonances consistent with loss of top-bottom symmetry to give a  $C_s$  symmetric product. The  $\text{ThCH}_2$  resonances are located at 0.31 and  $-0.70$  (**7**) or 0.52 and  $-0.68$  (**8**) ppm in the  $^1\text{H}$  NMR and 78 and 98 (**7**) or 90 and 103 (**8**) ppm in the  $^{13}\text{C}$  NMR (Figure 2).

X-ray-quality single crystals of  $[(\text{XA}_2)\text{Th}(\text{CH}_2\text{SiMe}_3)_2]\cdot\text{toluene}$  were grown by cooling a saturated toluene solution of **8** from room temperature to  $-30^\circ\text{C}$ , and to the best of our

knowledge, **8** represents the first structurally characterized thorium dialkyl complex supported by a multidentate non-carbocyclic ancillary. In the solid state (Figure 3, Table 1), **8** is pentacoordinate and adopts a square pyramidal geometry that is strongly distorted in the square plane as a result of  $\text{XA}_2$  ligand rigidity; the  $\text{XA}_2$  ligand backbone is approximately planar, but thorium is located  $0.475(6)$  Å above the NON ligand plane. Consistent with the low-temperature  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of **8**, the two alkyl groups are distinct; one is located approximately in the NON plane, while the other is located directly above the plane. This arrangement, in conjunction with rotation of the two 2,6-diisopropylphenyl groups to give  $\text{C}(34)\cdots\text{C}(44) = 7.514(9)$  Å and  $\text{C}(30)\cdots\text{C}(48) = 4.995(9)$  Å, is adopted in order to minimize unfavorable steric interactions between metal–alkyl and ligand–isopropyl groups. Similar behavior has been observed in other five-coordinate dialkyl complexes in which a metal is flanked by bulky 2,6-diisopropylphenyl substituents.<sup>26,27,47</sup>

As expected, the  $\text{Th}-\text{N}_{\text{anilido}}$   $\{2.292(4), 2.312(4)$  Å $\}$  bond lengths in **8** are very similar to those observed in complex **5**, and the  $\text{Th}-\text{C}$   $\{2.468(6), 2.485(6)$  Å $\}$  bond lengths fall within the range observed for the other crystallographically characterized thorium(IV) trimethylsilylmethyl complexes:  $2.48(2)$  and  $2.54(2)$  Å in  $[\{\text{Me}_2\text{Si}(\text{C}_5\text{Me}_4)_2\}\text{ThR}_2]$ ,<sup>48</sup>  $2.47(3)$  Å in  $[\text{Cp}^*\text{Th}(\text{CH}_2\text{tBu})\text{R}]$ ,<sup>49</sup>  $2.438(16)$  and  $2.485(18)$  Å in  $[(2,6\text{-tBu}_2\text{H}_3\text{C}_6\text{O})_2\text{-ThR}_2]$ ,<sup>13</sup> and  $2.488(2)$  and  $2.460(9)$  Å in  $[\text{Cp}^*(2,6\text{-tBu}_2\text{H}_3\text{C}_6\text{O})\text{-ThR}_2]$ <sup>50</sup> ( $\text{R} = \text{CH}_2\text{SiMe}_3$ ). Structurally characterized thorium diaryl ether or even aryl ether complexes have not previously been reported. However, the  $\text{Th}-\text{O}$  bond length in **8**  $\{2.534(3)$  Å $\}$  is similar to those observed for coordinated dme  $\{2.564(8)–2.620(8)$  Å $\}$ <sup>32,33</sup> or THF  $\{e.g., 2.53(1), 2.58(1)$  Å in  $[(\text{COT})\text{ThCl}_2(\text{THF})]\}$ <sup>51</sup> and  $2.520(7), 2.526(7)$  Å in  $[\text{Cp}^*\text{Th}(\text{NXyl})(\text{THF})]\}$ <sup>52</sup>. This is perhaps surprising since a diaryl ether should be a considerably less effective donor than dme or THF. However, a short bond between thorium and the neutral donor of the ligand backbone was also observed in **5**, and this feature is likely a consequence of tridentate binding and BDPP and  $\text{XA}_2$  ligand rigidity.

Of particular note are the  $\text{Th}-\text{C}-\text{Si}$  bond angles in **8**  $\{127.6(3)^\circ$  and  $126.8(3)^\circ\}$ , which are both significantly larger than is typically observed for an  $\text{sp}^3$ -hybridized carbon atom. Similar increases in  $\text{M}-\text{C}-\text{Si}$  or  $\text{M}-\text{C}-\text{C}$  ( $\text{M} = \text{Th}$  or  $\text{U}$ ) bond angles have been observed in other trimethylsilylmethyl or neopentyl complexes and are attributed to  $\alpha$ -agostic  $\text{C}-\text{H}-\text{Th}$  interactions.<sup>13,23,48–50</sup> Further evidence for the presence of agostic interactions is provided by a  $^1J_{\text{C,H}}$  of 102 Hz for the rapidly exchanging  $\text{CH}_2\text{SiMe}_3$  groups in the  $^1\text{H}$ -coupled  $^{13}\text{C}$  NMR spectrum of **8** in  $d_8$ -toluene at  $50^\circ\text{C}$ . This value is significantly lower than typically observed for an  $\text{sp}^3$ -hybridized carbon atom and compares well with literature values for related complexes: 104 Hz in Leznoff's  $[(^{\text{tBu}}\text{NON})\text{ThR}_2]$ ,<sup>24</sup> 100 Hz in  $[\text{Cp}^*(2,6\text{-tBu}_2\text{H}_3\text{C}_6\text{O})\text{ThR}_2]$ ,<sup>50</sup> 99 Hz in  $[\{\text{Me}_2\text{Si}(\text{C}_5\text{Me}_4)_2\}\text{ThR}_2]$ ,<sup>48</sup>

(41) Edwards, P. G.; Andersen, R. A.; Zalkin, A. *J. Am. Chem. Soc.* **1981**, *103*, 7792.

(42) (a) Edwards, P. G.; Andersen, R. A.; Zalkin, A. *Organometallics* **1984**, *3*, 293. (b) Carlson, C. N.; Hanusa, T. P.; Brennessel, W. W. *J. Am. Chem. Soc.* **2004**, *126*, 10550. (c) Lugli, G.; Marconi, W.; Mazzei, A.; Paladino, N.; Pedretti, U. *Inorg. Chim. Acta* **1969**, *3*, 253. (d) Brunelli, M.; Lugli, G.; Giacometti, G. *J. Magn. Reson.* **1973**, *9*, 247. (e) Thiele, K.-H.; Opitz, R.; Köhler, E. Z. *Anorg. Allg. Chem.* **1977**, *435*, 45.

(43) Eisen, M. S.; Marks, T. J. *J. Am. Chem. Soc.* **1992**, *114*, 10358.

(44) Köhler, E.; Brüser, W.; Thiele, K.-H. *J. Organomet. Chem.* **1974**, *76*, 235.

(45) Lauke, H.; Swepston, P. J.; Marks, T. J. *J. Am. Chem. Soc.* **1984**, *106*, 6841.

(46) Sigurdson, E. R.; Wilkinson, G. *Dalton Trans.* **1977**, 812.

(47) (a) Aizenberg, M.; Turculet, L.; Davis, W. M.; Schattenmann, F.; Schrock, R. R. *Organometallics* **1998**, *17*, 4795. (b) Cameron, T. M.; Gordon, J. C.; Michalczyk, R.; Scott, B. L. *Chem. Commun.* **2003**, 2282. (c) Reardon, D.; Conan, F.; Gambarotta, S.; Yap, G.; Wang, Q. *J. Am. Chem. Soc.* **1999**, *121*, 9318.


(48) Fendrick, C. M.; Mintz, E. A.; Schertz, L. D.; Marks, T. J.; Day, V. W. *Organometallics* **1984**, *3*, 819.

(49) Bruno, J. W.; Smith, G. M.; Marks, T. J.; Fair, C. K.; Schultz, A. J.; Williams, J. M. *J. Am. Chem. Soc.* **1986**, *108*, 40.

(50) Butcher, R. J.; Clark, D. L.; Grumbine, S. K.; Scott, B. L.; Watkin, J. G. *Organometallics* **1996**, *15*, 1488.

(51) Zalkin, A.; Templeton, D. H.; Le Vanda, C.; Streitwieser, A., Jr. *Inorg. Chem.* **1980**, *19*, 2560.

(52) Straub, T.; Haskel, A.; Neyroud, T. G.; Kapon, M.; Botoshansky, M.; Eisen, M. S. *Organometallics* **2001**, *20*, 5017.



**Figure 3.** Molecular structure of dialkyl complex **8**-toluene with thermal ellipsoids at the 50% probability level. Hydrogen atoms and solvent omitted for clarity. In view b,  $\text{CMe}_3$  groups have been removed and only the  $\text{N-C}_{\text{ipso}}$  carbon atoms of the  $\text{N-Aryl}$  rings are shown.

and 98 Hz in  $[(2,6\text{-tBu}_2\text{H}_3\text{C}_6\text{O})_2\text{ThR}_2]^{13}$  ( $\text{R} = \text{CH}_2\text{SiMe}_3$ ). A  $^1\text{J}_{\text{C,H}}$  value of 103 Hz is observed for the  $\text{CH}_2\text{SiMe}_3$  groups in **7** (60 °C,  $\text{C}_7\text{D}_8$ ).

Interestingly, upon lowering the temperature to  $-80$  °C, separate  $^1\text{J}_{\text{C,H}}$  coupling constants for each  $\text{CH}_2\text{SiMe}_3$  group were observed by fully coupled HSQC NMR. These values (88 and 91 Hz for **7** and 81 and 88 Hz for **8**) are considerably reduced relative to those obtained at 60 °C and suggest that for both complexes in solution an equilibrium exists between products participating in  $\alpha$ -agostic  $\text{C-H-Th}$  bonding to a greater or lesser extent (up to four  $\alpha$ -agostic  $\text{C-H-Th}$  interactions<sup>49</sup> could occur in either **7** or **8**) and that entropy favors less agostic products. Consequently, as the temperature is lowered, the equilibrium shifts toward more agostic products, resulting in a decrease in the average  $^1\text{J}_{\text{C,H}}$  coupling constant.<sup>53</sup>

The weak and nonstatic nature of  $\alpha$ -agostic interactions has previously been discussed,<sup>49,54</sup> and temperature-dependent  $^1\text{J}_{\text{E,H}}$  coupling constants as a result of an equilibrium between an agostic and a nonagostic isomer have been reported for several related complexes. For example,  $^1\text{J}_{\text{C,H}}$  for  $\alpha\text{-CHR}_2$  in  $[(\kappa^3\text{-Tp})\text{NbCl}(\text{cyclopropyl})(\text{C}_2\text{Me}_2)]$  varies from 102 Hz at 20 °C to 93 Hz at  $-80$  °C,<sup>55</sup> and  $^1\text{J}_{\text{Si,H}}$  in  $[\text{Cp}_2\text{Ti}(\text{tBuC}_2\text{SiHMe}_2)]$  varies from 123 Hz at 30 °C to 93 Hz at  $-80$  °C.<sup>56</sup> A temperature-dependent  $^1\text{J}_{\text{C,H}}$  constant was also observed for  $\alpha\text{-CH}_2\text{Me}$  in  $[(\text{P}_2\text{N}_2)\text{Ta}(\text{Et})(\text{C}_2\text{H}_4)]$  (123 Hz at 77 °C and 134 Hz at  $-40$  °C), but in this case it is due to a greater contribution from a  $\beta$ -agostic structure (versus an  $\alpha$ -agostic structure) at lower temperature.<sup>57</sup>

**Dialkyl Complex Stability.** Both dialkyl complexes exhibit remarkable thermal stability;  $[(\text{BDPP})\text{Th}(\text{CH}_2\text{SiMe}_3)_2]$  (**7**) is stable for days in toluene at 90 °C and is only fully decomposed after 3 days at 110 °C, while  $[(\text{XA}_2)\text{Th}(\text{CH}_2\text{SiMe}_3)_2]$  (**8**) is stable at 70 °C, but decomposes over several hours at 90 °C. Tetramethylsilane was the only soluble product observed when the thermal decomposition of **7** or **8** was monitored by  $^1\text{H}$  NMR

spectroscopy. The greater stability of **7** could be a result of decreased ligand rigidity, allowing the ligand to form a more optimal metal binding pocket. However,  $\text{Th-N}_{\text{amido}}$  bond lengths in **8** are very similar to those observed in the dichloro BDPP complex (**5**), and in both cases, the ligand is approximately planar. Therefore, the greater stability of **7** is more likely due to improved donor properties of a pyridine versus a diaryl ether neutral donor, leading to increased electronic saturation at the metal center.

The stability of **7** is particularly remarkable and highlights the suitability of rigid tridentate ligands such as BDPP for the stabilization of organoactinide complexes. The thermal stability of most other non-cyclopentadienyl thorium(IV) dialkyl complexes has not been reported. However, for comparison,  $[\text{Th}(\text{OC}_6\text{H}_3\text{tBu}_2\text{-2,6})_2(\text{CH}_2\text{SiMe}_3)_2]$  decomposes over 36 h at 60 °C,<sup>13</sup>  $[\text{Cp}^*\text{Th}(\text{OC}_6\text{H}_3\text{tBu}_2\text{-2,6})_2(\text{CH}_2\text{SiMe}_3)_2]$  decomposes over 12 h at 60 °C,<sup>50</sup>  $[\text{Cp}^*_2\text{Th}(\text{CH}_2\text{SiMe}_3)_2]$  decomposes over 36 h at 85 °C,<sup>49</sup> and  $[\text{Me}_2\text{Si}(\text{C}_5\text{Me}_4)_2]\text{Th}(\text{CH}_2\text{SiMe}_3)_2]$  decomposes at 60 °C.<sup>7</sup> In fact, only  $[\text{Cp}^*_2\text{ThMe}_2]$  is of similar thermal stability, being 50% decomposed after 1 week at 100 °C.<sup>58</sup> In addition, while  $[(\text{C}_5\text{R}_5)_2\text{ThR}_2]$  complexes are reported to be extremely air sensitive,<sup>48,59</sup> exposure of toluene solutions of **7** to air for 5 min resulted in only 20–30% decomposition.<sup>60</sup> By contrast, solid samples of **7** and **8** are more sensitive, being entirely decomposed after 5 min of exposure to air.

**Bis-ligand Complexes.** Addition of 1 equiv of  $\text{H}_2[\text{BDPP}]$  to a solution of  $[(\text{BDPP})\text{Th}(\text{CH}_2\text{SiMe}_3)_2]$  (**7**) followed by heating to 100 °C for 24 h in a sealed reaction vessel resulted in  $\text{SiMe}_4$  elimination to form the bis-ligand complex  $[\text{Th}(\text{BDPP})_2]$  (**9**) as a pale greenish-yellow solid in 37% isolated yield (Scheme 5). Alternatively, **9** may be synthesized by addition of benzene<sup>30</sup> to 2 equiv of  $\text{Li}_2[\text{BDPP}]$  (**4**) and  $[\text{ThCl}_4(\text{dme})_2]$  at  $-78$  °C followed by warming to room temperature.

Solution NMR spectra of **9** between 20 and  $-35$  °C ( $d_5$ -bromobenzene) are consistent with a highly symmetrical product containing a single  $\text{CHMe}_2$  environment. Crystals of **9** suitable for X-ray diffraction were grown by cooling a saturated toluene solution of **9** from room temperature to  $-30$  °C (Figure 4, Table 1). Complex **9** adopts an unusual six-coordinate geometry in which the four amido donors form a distorted tetrahedron around thorium while the pyridine units are directed

(53) An alternative explanation in which the strength of  $\text{E-H-M}$  interactions are strongly affected by temperature has been found not to be responsible for the observed temperature-dependent  $^1\text{J}_{\text{E,H}}$  coupling constants in related Ti and Nb complexes (see ref 56).

(54) (a) Brookhart, M.; Green, M. L. H.; Wong, L.-L. *Prog. Inorg. Chem.* **1988**, 36, 1. (b) Scherer, W.; McGrady, G. S. *Angew. Chem., Int. Ed.* **2004**, 43, 1782.

(55) Jaffart, J.; Cole, M. L.; Etienne, M.; Reinhold, M.; McGrady, J. E.; Maseras, F. *Dalton Trans.* **2003**, 4057.


(56) (a) Ohff, A.; Kosse, P.; Baumann, W.; Tillack, A.; Kempe, R.; Görls, H.; Burlakov, V. V.; Rosenthal, U. *J. Am. Chem. Soc.* **1995**, 117, 10399. (b) Peulecke, N.; Ohff, A.; Kosse, P.; Tillack, A.; Spannenberg, A.; Kempe, R.; Baumann, W.; Burlakov, V. V.; Rosenthal, U. *Chem.-Eur. J.* **1998**, 4, 1852.

(57) Fryzuk, M. D.; Johnson, S. A.; Rettig, S. J. *J. Am. Chem. Soc.* **2001**, 123, 1602.

(58) Manriquez, J. M.; Fagan, P. J.; Marks, T. J. *J. Am. Chem. Soc.* **1978**, 100, 3939.

(59) Fagan, P. J.; Manriquez, J. M.; Maatta, E. A.; Seyam, A. M.; Marks, T. J. *J. Am. Chem. Soc.* **1981**, 103, 6650.

(60) The rate at which benzene solutions of **8** decompose upon exposure to air was not reproducible (25–100% decomposition after 5 min), perhaps as a result of variations in atmospheric humidity.

**Scheme 5. Bis-ligand Complex 9 Formation by Salt Metathesis or Alkane Elimination**

toward two of the edges ( $\text{N}2\cdots\text{N}3$  and  $\text{N}2'\cdots\text{N}3'$ ) of the tetrahedron but offset toward the center of the two smaller faces ( $\text{N}1$  is offset toward the  $\text{N}3$ ,  $\text{N}3'$ ,  $\text{N}2$  face, and  $\text{N}1'$  is offset toward the,  $\text{N}3$ ,  $\text{N}3'$ ,  $\text{N}2'$  face). Above each of the two larger faces (defined by  $\text{N}2$ ,  $\text{N}2'$ ,  $\text{X}$ , where  $\text{X} = \text{N}3$  or  $\text{N}3'$ ) of the tetrahedron are located three of the eight isopropyl groups (Figure 4b). In order to adopt such a distorted geometry, the two identical BDPP ligands are considerably twisted away from planarity and the  $\text{C}(4)\text{--}\text{N}(1)\text{--}\text{Th}$  angle is far from linear ( $165^\circ$ ). This situation likely arises due to severe steric crowding at the metal, resulting in a geometry influenced more strongly by the anionic  $\text{N}_{\text{amido}}$  donors than the neutral pyridine units. Steric pressure at the metal center is also considered to be responsible for the significant difference in  $\text{Th}\text{--}\text{N}_{\text{amido}}$  bond lengths [ $\text{Th}\text{--}\text{N}3 = 2.368(7)$  Å and  $\text{Th}\text{--}\text{N}2 = 2.488(6)$  Å], as well as  $\text{Th}\text{--}\text{N}_{\text{amido}}$  and  $\text{Th}\text{--}\text{N}_{\text{py}}$  bonds [ $\text{Th}\text{--}\text{N}_{\text{py}} = 2.615(6)$  Å], which are considerably longer than those observed in the structures of the dichloro BDPP (5) and dialkyl  $\text{XA}_2$  (8) complexes.

Attempts to prepare the analogous  $[\text{Th}(\text{XA}_2)_2]$  complex by either alkane elimination or salt metathesis were unsuccessful:  $[(\text{XA}_2)\text{ThCl}_2(\text{dme})]$  (6) failed to react with  $\text{K}_2(\text{dme})_2[\text{XA}_2]$  (2) or  $\text{Na}_2[\text{XA}_2]$  (3) at temperatures up to  $110^\circ\text{C}$ , and  $[(\text{XA}_2)\text{Th}(\text{CH}_2\text{SiMe}_3)_2]$  (8) did not react with  $\text{H}_2[\text{XA}_2]$  (1) at temperatures ( $\leq 70^\circ\text{C}$ ) below the onset of thermal decomposition. A lack of reactivity was also observed in attempts to prepare the mixed ligand derivative  $[\text{Th}(\text{BDPP})(\text{XA}_2)]$  by salt metathesis ( $[\text{Th}(\text{BDPP})\text{Cl}_2(\text{dme})] + \text{Na}_2[\text{XA}_2]$ , up to  $110^\circ\text{C}$ ) or alkane elimination ( $[\text{Th}(\text{BDPP})(\text{CH}_2\text{SiMe}_3)_2] + \text{H}_2[\text{XA}_2]$ , up to  $110^\circ\text{C}$ ) in toluene. This marked difference in BDPP and  $\text{XA}_2$  ligand reactivity presumably stems from the enhanced rigidity of  $\kappa^3$ -coordinated  $\text{XA}_2$ , which does not permit the type of ligand twisting observed in  $[\text{Th}(\text{BDPP})_2]$  (9). As a result,  $\kappa^3\text{-N,O,N}$ - or even  $\kappa^2\text{-N,N}$ -donation, which would likely allow the xanthene backbone of  $\text{XA}_2$  to adopt a butterfly conformation,<sup>61</sup> must be rendered sterically inaccessible, precluding bis-ligand complex formation.

(61) Xantphos, the neutral 4,5-bis(diphenylphosphino) analogue of  $\text{XA}_2$ , is typically  $\kappa^2\text{-P,P}$ -coordinated with the xanthene backbone adopting a butterfly conformation: Kamer, P. C. J.; van Leeuwen, P. W. N. M.; Reek, J. N. H. *Acc. Chem. Res.* **2001**, 34, 895. However, a planar xanthene backbone has been observed in cationic complexes where  $\kappa^3\text{-P,O,P}$ -coordination is favored: (a) Zuideveld, M. A.; Swennenhuis, B. H. G.; Boele, M. D. K.; Guari, Y.; van Strijdonck, G. P. F.; Reek, J. N. H.; Kamer, P. C. J.; Goubitz, K.; Fraanje, J.; Lutz, M.; Spek, A. L.; van Leeuwen, P. W. N. M. *Dalton Trans.* **2002**, 2308. (b) Sandee, A. J.; van der Veen, L. A.; Reek, J. N. H.; Kamer, P. C. J.; Lutz, M.; Spek, A. L.; van Leeuwen, P. W. N. M. *Angew. Chem., Int. Ed.* **1999**, 38, 3231.

## Summary and Conclusions

A new, extremely rigid and planar NON-donor ligand,  $\text{H}_2\text{XA}_2$  (1), was prepared and is conveniently deprotonated to form stable disodium or dipotassium salts (2, 3). The thermally unstable dilithium salt of a structurally related 2,6-bis(2,6-diisopropylanilidomethyl)pyridine ligand ( $\text{Li}_2\text{BDPP}$ ; 4) was also isolated, and its availability is expected to extend the utility of this ligand, which until now could only be installed using amine or  $\text{Me}_3\text{SiCl}$  elimination.

For both the BDPP and  $\text{XA}_2$  ligands, pentagonal bipyramidal  $[\text{ThCl}_2(\text{dme})]$  (5, 6) and five-coordinate  $[\text{Th}(\text{CH}_2\text{SiMe}_3)_2]$  (7, 8) complexes were prepared by either salt metathesis or apparent alkane elimination. Alkane elimination has not previously been employed as a means of primary ancillary ligand attachment in actinide chemistry<sup>40</sup> but has the potential to serve as a valuable tool for the direct synthesis of thorium dichloro and dialkyl complexes, especially when salt metathesis proves ineffective.

Despite a coordination number of only five, complexes 7 and 8 are remarkably robust, showing no sign of decomposition after several days at 90 and  $70^\circ\text{C}$ , respectively. In fact, no other bis-trimethylsilylmethyl or bis-neopentyl thorium complexes, including  $[\text{Cp}^*_2\text{Th}(\text{CH}_2\text{EMe}_3)_2]$  ( $\text{E} = \text{Si}$  or  $\text{C}$ ), have been reported to be of comparable thermal stability to 7. The availability of 7 and 8 is expected to provide a unique opportunity to study the reactivity and catalytic potential of non-carbocyclic dialkyl and related thorium complexes over a significant temperature range.


The greater stability of 7 is suspected to result from reduced Lewis acidity as a result of the enhanced donor properties of the central pyridine unit in BDPP, relative to the diarylether group of  $\text{XA}_2$ . However, while the bis-ligand complex  $[\text{Th}(\text{BDPP})_2]$  (9) is readily obtained by either alkane elimination or salt metathesis,  $[\text{Th}(\text{XA}_2)_2]$  proved inaccessible, likely due to the greater rigidity of  $\text{XA}_2$ . Since ligand redistribution (to form a complex containing more than one ancillary ligand as well as a more highly alkylated product) is a common decomposition pathway for early transition metal and f-block dialkyl complexes at elevated temperatures,<sup>13,62</sup> the  $\text{XA}_2$  ligand represents an important step in the design of ligands suitable for the formation of uniquely robust thorium dialkyls.

Future work will investigate the potential of 7 and 8 and closely related thorium dialkyl complexes in catalysis and as starting materials for the preparation of dihydride complexes and cationic alkyls. Efforts toward a new ligand that combines the rigidity of the  $\text{XA}_2$  ligand with the more effective donor properties of BDPP are also underway.

## Experimental Section

**General Details.** An argon-filled MBraun UNIlab glove box was employed for the manipulation and storage of all oxygen- and moisture-sensitive compounds (2–9), and all thermally unstable compounds were stored in a  $-30^\circ\text{C}$  freezer within the glovebox. Reactions were performed on a double-manifold high-vacuum line using standard techniques,<sup>63</sup> and all reaction products were thor-

(62) For example: (a) Ernst, R. D.; Kennelly, W. J.; Day, C. S.; Day, V. W.; Marks, T. J. *J. Am. Chem. Soc.* **1979**, 101, 2656. (b) Woodman, T. J.; Schormann, M.; Hughes, D. L.; Bochmann, M. *Organometallics* **2004**, 23, 2972. (c) Bambirra, S.; Brandsma, M. J. R.; Brussee, E. A. C.; Meetsma, A.; Hessen, B.; Teuben, J. H. *Organometallics* **2000**, 19, 3197. (d) Emslie, D. J. H.; Piers, W. E.; Parvez, M.; McDonald, R. *Organometallics* **2002**, 21, 4226. (e) Trifonov, A. A.; Lyubov, D. M.; Fukin, G. K.; Baranov, E. V.; Kurskii, Y. A. *Organometallics* **2006**, 25, 3935. (f) Duncan, A. P.; Mullins, S. M.; Arnold, J.; Bergman, R. G. *Organometallics* **2001**, 20, 1808.



**Figure 4.** Molecular structure of bis-ligand complex **9** with hydrogen atoms omitted for clarity. (a and b) ORTEPs with thermal ellipsoids at the 50% probability level. (c) Space filling diagrams. Part b highlights the distorted tetrahedral arrangement of  $N_{\text{anilido}}$  donors around Th and the position of three isopropyl groups above each of the larger faces of the tetrahedron.

oughly dried *in vacuo*. Commonly utilized specialty glassware includes the swivel frit assembly, J-Young NMR tubes, and thick walled flasks equipped with Teflon stopcocks. A Fisher Scientific Ultrasonic FS-30 bath was used to sonicate reaction mixtures where indicated. Residual oxygen and moisture were removed from the argon stream by passage through an Oxisorb-W scrubber from Matheson Gas Products. Anhydrous 1,2-dme and diethyl ether were purchased from Aldrich. Hexanes, toluene, and THF were initially dried and distilled at atmospheric pressure from  $\text{CaH}_2$ , sodium, and sodium benzophenone ketal, respectively. Unless otherwise noted, all proteo solvents were stored over an appropriate drying agent (1,2-dme,  $\text{OEt}_2$ , THF, toluene,  $\text{C}_6\text{H}_6 = \text{Na/Ph}_2\text{CO}$ ; hexanes,  $\text{O}(\text{SiMe}_3)_2 = \text{Na/Ph}_2\text{CO/tetraglyme}$ ) and introduced to reactions via vacuum transfer with condensation at  $-78^\circ\text{C}$ . Deuterated solvents (ACP Chemicals) were dried over  $\text{CaH}_2$  ( $\text{CD}_2\text{Cl}_2$ ,  $\text{C}_6\text{D}_5\text{-Br}$ ) or  $\text{Na/Ph}_2\text{CO}$  ( $\text{C}_6\text{D}_6$ ,  $d_8$ -toluene,  $d_8$ -THF).

$\text{Th}(\text{NO}_3)_4(\text{H}_2\text{O})_4$  was purchased from Strem Chemicals.  $\text{SOCl}_2$ ,  $\text{O}(\text{SiMe}_3)_2$ , 2,6-bis(bromomethyl)pyridine, tetraglyme, xanthone,  $\text{AlMe}_3$  (2 M in toluene),  $\text{Pd}(\text{OAc})_2$ ,  $\text{NaOtBu}$ , DPEPhos [bis(2-(diphenylphosphino)phenyl) ether],  $\text{NaH}$ ,  $\text{KH}$  (30 wt % in mineral oil),  $\text{LiCH}_2\text{SiMe}_3$  (1.0 M in pentane),  $n\text{BuLi}$  (2.0 M in cyclohexane),<sup>64</sup> Fe powder, and  $\text{Br}_2$  were purchased from Sigma-Aldrich. 2,6-Diisopropylaniline was purchased from Lancaster. Prior to use, solid  $\text{LiCH}_2\text{SiMe}_3$  was obtained by removal of pentane *in vacuo*, solid  $\text{KH}$  was obtained by filtration and washing with hexanes, 2,6-diisopropylaniline was distilled from  $\text{CaH}_2$ , and tetraglyme was distilled from sodium/benzophenone ketal.  $\text{H}_2[\text{BDPP}]$ ,<sup>29</sup> 4,5-dibromo-2,7-di-*tert*-butyl-9,9-dimethylxanthene,<sup>65</sup> and  $\text{ThCl}_4(\text{dme})_2$ <sup>37</sup> were prepared by literature procedures.

NMR spectroscopy ( $^1\text{H}$ ,  $^{13}\text{C}\{^1\text{H}\}$ , DEPT-135, COSY, HSQC, HMBC) was performed on Bruker DRX-500 and AV-600 spectrometers. All  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra were referenced relative to  $\text{SiMe}_4$  through a resonance of the employed deuterated solvent or proteo impurity of the solvent:  $\text{C}_6\text{D}_6$  ( $\delta$  7.15 ppm),  $\text{C}_7\text{D}_8$  ( $\delta$  7.09, 7.00, 6.98, 2.09 ppm),  $\text{C}_6\text{D}_5\text{Br}$  ( $\delta$  7.30, 7.02, 6.94 ppm),  $d_8$ -THF (3.58, 1.73 ppm),  $\text{CD}_2\text{Cl}_2$  (5.32 ppm) for  $^1\text{H}$  NMR, and  $\text{C}_6\text{D}_6$  ( $\delta$  128.0 ppm),  $\text{C}_7\text{D}_8$  ( $\delta$  137.86, 129.24, 128.33, 125.49, 20.4 ppm),  $\text{C}_6\text{D}_5\text{Br}$  ( $\delta$  130.9, 129.3, 126.1, 122.3 ppm),  $d_8$ -THF (67.57, 25.37 ppm),  $\text{CD}_2\text{Cl}_2$  (54.0 ppm) for  $^{13}\text{C}$  NMR.

(63) Burger, B. J.; Bercaw, J. E. Vacuum Line Techniques for Handling Air-Sensitive Organometallic Compounds. In *Experimental Organometallic Chemistry—A Practicum in Synthesis and Characterization*; American Chemical Society: Washington, D.C., 1987; Vol. 357, p 79.

(64)  $n\text{BuLi}$  solutions were titrated using *N*-benzylbenzamide (Sigma-Aldrich) in THF at  $-45^\circ\text{C}$ : Burchat, A. F.; Chong, J. M.; Nielsen, N. J. *Organomet. Chem.* **1997**, 542, 281.

(65) Nowick, J. S.; Ballester, P.; Ebmeyer, F.; Rebek, J. J. *Am. Chem. Soc.* **1990**, 112, 8902.

Combustion elemental analyses were performed on a Thermo EA1112 CHNS/O analyzer by Dr. Steve Kornic of this department. X-ray crystallographic analyses were performed on suitable crystals coated in Paratone oil and mounted on a P4 diffractometer with a Bruker Mo rotating-anode generator and a SMART1K CCD area detector in the McMaster Analytical X-Ray (MAX) Diffraction Facility. Herein, Q = quaternary, Ar = 2,6-diisopropylphenyl, the *ipso* carbon refers to the carbon attached to nitrogen, Py and Xanth refer to the pyridine and xanthene backbones of the BDPP and  $\text{XA}_2$  ligands, respectively, and  $\text{CH}^1$  and  $\text{CH}^3$  refer to the 1- and 3-positions of the xanthene backbone (*para* and *ortho*, respectively, to the amido group in  $\text{XA}_2$ ).

**$\text{H}_2(\text{XA}_2)$  (1).** 4,5-Dibromo-2,7-di-*tert*-butyl-9,9-dimethylxanthene (10.00 g, 20.82 mmol), 2,6-diisopropylaniline (7.85 mL, 41.64 mmol),  $\text{NaOtBu}$  (5.60 g, 58.30 mmol),  $\text{Pd}(\text{OAc})_2$  (40 mg, 0.21 mmol), and DPEPhos (167 mg, 0.31 mmol) in toluene (150 mL) were heated to  $100^\circ\text{C}$  for 16 h. The reaction mixture was then quenched with water, extracted into toluene ( $3 \times 50$  mL), dried over  $\text{MgSO}_4$ , and concentrated to approximately 30 mL. Recrystallization from a hot ethanol/toluene (10:1) solution gave **1** as a white solid (12.02 g, 17.86 mmol) in 86% yield.  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ , 600 MHz):  $\delta$  7.26–7.21 (m, 6H, Ar-*H*), 6.99 (d, 2H,  $J$  1.94 Hz,  $\text{CH}^3$ ), 6.51 (d, 2H,  $J$  1.94 Hz,  $\text{CH}^1$ ), 5.93 (s, 2H, NH), 3.47 (sept, 4H,  $J$  6.88 Hz,  $\text{CHMe}_2$ ), 1.98 (s, 18H,  $\text{CMe}_3$ ), 1.68 (s, 6H,  $\text{CMe}_2$ ), 1.18 (d, 12H,  $J$  6.88 Hz,  $\text{CHMe}_2$ ), 1.14 (d, 12H,  $J$  6.88 Hz,  $\text{CHMe}_2$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{C}_6\text{D}_6$ , 150 MHz):  $\delta$  147.8 (Ar- $\text{C}_{\text{ipso}}$ ), 146.1, 136.4, 135.9 (Xanth-Q), 136.2 (Ar- $\text{C}_{\text{ortho}}$ ), 129.4 (Ar- $\text{CH}_{\text{para}}$ ), 124.3 (Ar- $\text{CH}_{\text{meta}}$ ), 111.9 ( $\text{CH}^3$ ), 108.1 ( $\text{CH}^1$ ), 35.1 ( $\text{CMe}_2$ ), 34.8 ( $\text{CMe}_3$ ), 32.9 ( $\text{CMe}_2$ ), 31.7 ( $\text{CMe}_3$ ), 28.7 ( $\text{CHMe}_2$ ), 24.7, 23.5 ( $\text{CHMe}_2$ ). Anal. Calcd for  $\text{C}_{47}\text{H}_{64}\text{N}_2\text{O}$ : C, 83.88; H, 9.58; N, 4.16. Found: C, 83.91; H, 9.64; N, 4.00.

**$\text{K}_2(\text{dme})_2[\text{XA}_2]$  (2).**  $\text{KH}$  (0.120 g, 3.00 mmol) and  $\text{H}_2[\text{XA}_2]$  (0.750 g, 1.11 mmol) in dme (60 mL) were stirred at room temperature for 5 h. The solution was filtered to remove excess  $\text{KH}$ . Solvent was removed *in vacuo*, and hexamethyldisiloxane (30 mL) was added, followed by sonication. The solution was cooled to  $-78^\circ\text{C}$  and filtered on a precooled frit to obtain 0.756 g (0.81 mmol, 73%) of **2** as a white solid.  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ , 600 MHz):  $\delta$  7.29 (d, 4H,  $J$  7.53 Hz, Ar-*H*), 7.14 (t, 2H,  $J$  7.53 Hz, Ar-*H*), 6.57 (d, 2H,  $J$  1.95 Hz,  $\text{CH}^3$ ), 6.18 (d, 2H,  $J$  1.95 Hz,  $\text{CH}^1$ ), 3.00 (sept, 4H,  $J$  6.90 Hz,  $\text{CHMe}_2$ ), 2.97 (s, 8H,  $\text{OCH}_2$ ), 2.83 (s, 12H,  $\text{OCH}_3$ ), 1.90 (s, 6H,  $\text{CMe}_2$ ), 1.40 (s, 18H,  $\text{CMe}_3$ ), 1.29 (d, 12H,  $J$  6.77 Hz,  $\text{CHMe}$ ), 1.16 (d, 12H,  $J$  7.04 Hz,  $\text{CHMe}_2$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{C}_6\text{D}_6$ , 150 MHz):  $\delta$  154.1 (Ar- $\text{C}_{\text{ipso}}$ ), 149.2 (Ar- $\text{C}_{\text{ortho}}$ ), 147.7, 143.0, 138.6, 132.9 (Xanth-Q), 124.3 (Ar- $\text{CH}_{\text{para}}$ ), 120.6 (Ar- $\text{CH}_{\text{meta}}$ ), 109.1 ( $\text{CH}^1$ ), 100.8 ( $\text{CH}^3$ ), 71.7 ( $\text{OCH}_2$ ), 58.7 ( $\text{OMe}$ ), 36.4 ( $\text{CMe}_2$ ), 35.6 ( $\text{CMe}_3$ ), 32.8 ( $\text{CMe}_3$ ), 31.2 ( $\text{CMe}_2$ ), 28.7 ( $\text{CHMe}_2$ ), 25.5, 24.8

(CHMe<sub>2</sub>). Anal. Calcd for C<sub>55</sub>H<sub>82</sub>N<sub>2</sub>O<sub>5</sub>K<sub>2</sub>: C, 71.07; H, 8.89; N, 3.01. Found: C, 71.16; H, 8.86; N, 3.10.

**Na<sub>2</sub>[XA<sub>2</sub>] (3).** NaH (0.100 g, 4.17 mmol) and H<sub>2</sub>[XA<sub>2</sub>] (0.700 g, 1.04 mmol) were refluxed in toluene (60 mL) for 2–12 days.<sup>66</sup> The solution was then filtered to remove excess NaH, solvent was removed *in vacuo*, and hexamethyldisiloxane (30 mL) was added. Sonication and filtration gave 0.555 g (0.866 mmol, 84%) of **3** as a white solid. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 600 MHz): δ 7.28 (d, 4H, *J* 7.65 Hz, Ar-*H*), 7.19 (t, 2H, *J* 7.6 Hz, Ar-*H*), 6.67 (d, 2H, *J* 2.0 Hz, CH<sup>3</sup>), 6.34 (d, 2H, *J* 2.0 Hz, CH<sup>1</sup>), 2.98 (sept, 4H, *J* 6.9 Hz, CHMe<sub>2</sub>), 1.81 (s, 6H, CMe<sub>2</sub>), 1.35 (s, 18H, CMe<sub>3</sub>), 1.29 (d, 12H, *J* 6.8 Hz, CHMe<sub>2</sub>), 0.99 (d, 12H, *J* 7.1 Hz, CHMe<sub>2</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>, 150 MHz): δ 151.8, 148.8, 146.2, 133.0 (Xanth-Q), 137.4 (Ar-C<sub>ipso</sub>), 128.6 (Ar-C<sub>ortho</sub>), 125.0 (Ar-CH<sub>meta</sub>), 121.2 (Ar-CH<sub>para</sub>), 109.4 (CH<sup>3</sup>), 101.2 (CH<sup>1</sup>), 35.8 (CMe<sub>2</sub>), 34.4 (CMe<sub>3</sub>), 31.4 (CMe<sub>3</sub>), 28.3 (CMe<sub>2</sub>), 27.7 (CHMe<sub>2</sub>), 24.0, 23.4 (CHMe<sub>2</sub>). Anal. Calcd for C<sub>47</sub>H<sub>62</sub>N<sub>2</sub>Na<sub>2</sub>O: C, 78.73; H, 8.72; N, 3.91. Found: C, 79.09; H, 8.65; N, 3.96.

**Li<sub>2</sub>[BDPP] (4).** *n*BuLi (2.0 M) in cyclohexane (2.20 mL, 4.37 mmol) was added dropwise to 2,6-bis(2,6-diisopropylanilinoethyl)pyridine (1.00 g, 2.19 mmol) in hexanes (30 mL) at –78 °C. After stirring at –78 °C for 5 min the solution was warmed to –45 °C for 5 min to give a yellow-brown solution with large amounts of yellow precipitate. The mixture was then recooled to –78 °C and filtered quickly on a precooled frit to provide **4** as a yellow solid (0.898 g, 1.91 mmol) in 87% yield. <sup>1</sup>H NMR (d<sub>8</sub>-THF, –30 °C, 500 MHz): δ 7.58 (t, 1H, *J* 7.3 Hz, py-CH), 7.44 (d, 2H, *J* 7.3 Hz, py-CH), 6.71 (d, 4H, *J* 7.3 Hz, Ar-*H*), 6.23 (t, 2H, *J* 7.3 Hz, Ar-*H*), 4.78 (s, 4H, NCH<sub>2</sub>), 3.73 (sept, *J* 6.9 Hz, CHMe<sub>2</sub>), 1.11 (d, 24H, *J* 6.7 Hz, CHMe<sub>2</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (d<sub>8</sub>-THF, –30 °C, 125 MHz): δ 166.4 (Py-C<sub>ortho</sub>), 160.1 (Ar-C<sub>ipso</sub>), 138.5 (Ar-C<sub>ortho</sub>), 133.0 (Py-CH<sub>para</sub>), 120.8 (Ar-CH<sub>meta</sub>), 115.9 (Py-CH<sub>meta</sub>), 110.2 (Ar-CH<sub>para</sub>), 62.4 (NCH<sub>2</sub>), 25.5 (CHMe<sub>2</sub>), 21.0 (CHMe<sub>2</sub>). Anal. Calcd for C<sub>31</sub>H<sub>41</sub>N<sub>3</sub>Li<sub>2</sub>: C, 79.29; H, 8.80; N 8.95. Found: C, 79.15; H, 9.21; N, 8.49.

**[(BDPP)ThCl<sub>2</sub>(dme)] (5): Method A.** ThCl<sub>4</sub>(dme)<sub>2</sub> (0.600 g, 1.08 mmol) and LiCH<sub>2</sub>SiMe<sub>3</sub> (0.204 g, 2.17 mmol) in toluene (60 mL) were stirred for 1 h at –78 °C followed by 1 h at 0 °C. A solution of 2,6-bis(2,6-diisopropylanilinoethyl)pyridine (0.620 g, 1.35 mmol) in toluene (10 mL) was then added dropwise. The solution was allowed to warm to room temperature over ca. 2 h, stirred for an additional 12 h, and then filtered to remove lithium salts. Solvent was removed *in vacuo*, and hexanes (30 mL) were added, followed by sonication and filtration to give **5** as a white solid (0.781 g, 0.92 mmol) in 85% yield. **Method B.** ThCl<sub>4</sub>(dme)<sub>2</sub> (0.140 g, 0.25 mmol) in benzene (10 mL) was placed in an ice–water bath (the majority of the solution does not freeze, despite the 6 °C freezing point of benzene). A solution of Li<sub>2</sub>[BDPP] (0.119 g, 0.25 mmol) in benzene<sup>30</sup> (5 mL) was added dropwise (over 1–2 min). The solution was stirred for 1 h, followed by filtration and removal of solvent *in vacuo*. Hexanes (10 mL) were added, followed by sonication and filtration to give **5** as a white solid (0.109 g, 0.13 mmol) in 51% yield. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>5</sub>Br, 600 MHz): δ 7.24 (t, 1H, *J* 7.6 Hz, py-CH), 7.17 (m, 6H, Ar-*H*), 6.85 (d, 2H, *J* 7.6 Hz, py-CH), 5.27 (s, 4H, NCH<sub>2</sub>), 4.19 (sept, *J* 6.7 Hz, CHMe<sub>2</sub>), 3.42 (s, 2H, OCH<sub>2</sub>), 2.38 (s, 3H, OCH<sub>3</sub>), 1.42, 1.15 (d, 24H, *J* 6.7 Hz, CHMe<sub>2</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>, 150 MHz): δ 165.0 (Py-C<sub>ortho</sub>), 148.1 (Ar-C<sub>ipso</sub>), 147.6 (Ar-C<sub>ortho</sub>), 137.4 (Py-CH<sub>para</sub>), 125.3 (Ar-CH<sub>meta</sub>), 116.8 (Py-CH<sub>meta</sub>), 124.2 (Ar-CH<sub>para</sub>), 70.6 (NCH<sub>2</sub>), 27.8 (CHMe<sub>2</sub>), 27.0, 24.3 (CHMe<sub>2</sub>). Anal. Calcd for C<sub>35</sub>H<sub>51</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub>Th: C, 49.53; H, 6.06; N 4.95. Found: C, 49.85; H, 6.35; N, 4.37.

**[(XA<sub>2</sub>)ThCl<sub>2</sub>(dme)] (6).** ThCl<sub>4</sub>(dme)<sub>2</sub> (0.416 g, 0.75 mmol) and K<sub>2</sub>(dme)<sub>2</sub>[XA<sub>2</sub>] (0.700 g, 0.75 mmol) in toluene (60 mL) were stirred for 16 h at 100 °C. The solution was cooled to room

temperature and filtered, and the solvent was removed *in vacuo*. Hexanes (30 mL) were added, followed by sonication and filtration to give **6** as a white solid (0.378 g, 0.36 mmol) in 59% yield. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 600 MHz): δ 7.31 (m, 4H, *J* 7.5 Hz, Ar-*H*), 7.25 (m, 2H, *J* 7.5 Hz, Ar-*H*), 6.85 (d, 2H, *J* 1.8 Hz, CH<sup>3</sup>), 5.89 (d, 2H, *J* 1.8 Hz, CH<sup>1</sup>), 4.06 (sept, 4H, *J* 6.8 Hz, CHMe<sub>2</sub>), 3.04 (s, 4H, OCH<sub>2</sub>), 2.25 (s, 6H, OCH<sub>3</sub>), 1.70 (s, 6H, CMe<sub>2</sub>), 1.47 (d, 12H, *J* 6.7 Hz, CHMe<sub>2</sub>), 1.25 (s, 18H, CMe<sub>3</sub>), 1.09 (d, 12H, *J* 6.9 Hz, CHMe<sub>2</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>, 150 MHz): δ 149.4 (Ar-C<sub>ortho</sub>), 147.1, 145.9, 141.3 (Xanth-Q), 142.0 (Ar-C<sub>ipso</sub>), 127.3 (Ar-CH<sub>para</sub>), 125.4 (Ar-CH<sub>meta</sub>), 111.7 (CH<sup>1</sup>), 111.4 (CH<sup>3</sup>), 71.0 (OCH<sub>2</sub>), 61.3 (OMe), 34.9 (CMe<sub>3</sub>), 34.1 (CMe<sub>2</sub>), 33.8 (CMe<sub>2</sub>), 31.7 (CMe<sub>3</sub>), 28.1 (CHMe<sub>2</sub>), 27.2, 24.8 (CHMe<sub>2</sub>). Anal. Calcd for C<sub>51</sub>H<sub>72</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>3</sub>Th: C, 57.57; H, 6.82; N, 2.63. Found: C, 57.76; H, 6.89; N, 2.49.

**[(BDPP)Th(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>2</sub>] (7): Method A.** ThCl<sub>4</sub>(dme)<sub>2</sub> (0.750 g, 1.35 mmol) and LiCH<sub>2</sub>SiMe<sub>3</sub> (0.510 g, 5.41 mmol) in toluene (60 mL) were stirred for 1 h at –78 °C followed by 1 h at 0 °C. The cloudy, colorless solution was then recooled to –78 °C, and a solution of 2,6-bis(2,6-diisopropylanilinoethyl)pyridine (0.620 g, 1.35 mmol) in toluene (10 mL) was added dropwise. The solution was allowed to warm to room temperature over ca. 2 h, stirred for an additional 12 h, and then filtered to remove lithium salts. Solvent was removed *in vacuo* and hexamethyldisiloxane (30 mL) was added, followed by sonication and filtration to afford **7** as an off-white solid (0.949 g, 1.10 mmol) in 82% yield. **Method B.** [(BDPP)ThCl<sub>2</sub>(dme)] (0.150 g, 0.18 mmol) and LiCH<sub>2</sub>SiMe<sub>3</sub> (0.033 g, 0.35 mmol) in toluene (15 mL) were stirred for 30 min at room temperature. The solution was filtered, and solvent was removed *in vacuo*. Hexamethyldisiloxane (10 mL) was added, followed by sonication and filtration to afford **7** as an off-white solid (0.106 g, 0.12 mmol) in 68% yield. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>): δ 7.24 (m, 6H, Ar-*H*), 6.90 (t, 1H, *J* 7.7 Hz, py-CH), 6.49 (d, 2H, *J* 7.7 Hz, py-CH), 5.24 (s, 4H, NCH<sub>2</sub>), 3.75 (sept, 4H, *J* 6.8 Hz, CHMe<sub>2</sub>), 1.52 (d, 12H, *J* 6.8 Hz, CHMe<sub>2</sub>), 1.26 (d, 12H, *J* 6.8 Hz, CHMe<sub>2</sub>), –0.02 (s, 18H, SiMe<sub>3</sub>), –0.32 (s, 4H, ThCH<sub>2</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>, 150 MHz): δ 164.8 (Py-C<sub>ortho</sub>), 148.1 (Ar-C<sub>ipso</sub>), 141.9 (Ar-C<sub>ortho</sub>), 138.4 (Py-CH<sub>para</sub>), 126.8 (Ar-CH<sub>meta</sub>), 125.0 (Ar-CH<sub>para</sub>), 117.8 (Py-CH<sub>meta</sub>), 89.9 (ThCH<sub>2</sub>), 68.7 (NCH<sub>2</sub>), 28.7 (CHMe<sub>2</sub>), 27.3, 24.9 (CHMe<sub>2</sub>), 3.8 (SiMe<sub>3</sub>). Anal. Calcd for C<sub>39</sub>H<sub>63</sub>N<sub>3</sub>Si<sub>2</sub>Th: C, 54.33; H, 7.37; N, 4.87. Found: C, 54.10; H, 7.35; N, 4.69.

**[(XA<sub>2</sub>)Th(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>2</sub>] (8): Method A.** Complex **8** was prepared in a similar fashion to **7** (method A) using 0.750 g (1.35 mmol) of ThCl<sub>4</sub>(dme)<sub>2</sub>, 0.510 g (5.41 mmol) of LiCH<sub>2</sub>SiMe<sub>3</sub>, and 0.909 g (1.35 mmol) of H<sub>2</sub>[XA<sub>2</sub>]. However, the crude filtered product was sonicated in hexanes (20 mL) before filtration to give 0.710 g of **8** as a white solid (0.711 g, 0.66 mmol) in 49% yield. **Method B.** Complex **8** was prepared in a similar fashion to **7** (method B) using [(XA<sub>2</sub>)ThCl<sub>2</sub>(dme)] (0.250 g, 0.25 mmol) and LiCH<sub>2</sub>SiMe<sub>3</sub> (0.048 g, 0.51 mmol). However, the crude filtered product was sonicated in hexanes (10 mL) before filtration to afford **8** as a white solid (0.175 g, 0.16 mmol) in 65% yield. <sup>1</sup>H NMR (d<sub>8</sub>-toluene, 600 MHz): δ 7.26 (m, 6H, Ar-*H*), 6.77 (d, 2H, *J* 1.5 Hz, CH<sup>3</sup>), 6.00 (d, 2H, *J* 1.5 Hz, CH<sup>1</sup>), 3.54 (sept, 4H, *J* 7.0 Hz, CHMe<sub>2</sub>), 1.66 (s, 6H, CMe<sub>2</sub>), 1.40 (d, 12H, *J* 7.0 Hz, CHMe<sub>2</sub>), 1.16 (d, 12H, *J* 7.0 Hz, CHMe<sub>2</sub>), 0.03 (s, 18H, SiMe<sub>3</sub>), –0.17 (s, 4H, ThCH<sub>2</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR (d<sub>8</sub>-toluene, 150 MHz): δ 146.2 (Ar-C<sub>ortho</sub>), 148.5 (Xanth-Q), 142.6 (Ar-C<sub>ipso</sub>), 130.3 (Ar-CH<sub>para</sub>), 129.7 (Ar-CH<sub>meta</sub>), 110.7 (CH<sup>1</sup>), 110.9 (CH<sup>3</sup>), 35.6 (CMe<sub>3</sub>), 35.5 (CMe<sub>2</sub>), 31.4 (CMe<sub>2</sub>), 32.1 (CMe<sub>3</sub>), 29.5 (CHMe<sub>2</sub>), 26.8, 25.5 (CHMe<sub>2</sub>), 3.4 (SiMe<sub>3</sub>). Note: ThCH<sub>2</sub> was not observed at room temperature but was observed at +50 or –80 °C (see main text). Anal. Calcd for C<sub>55</sub>H<sub>84</sub>N<sub>2</sub>OSi<sub>2</sub>Th: C, 61.31; H, 7.86; N, 2.60. Found: C, 61.41; H, 8.06; N, 2.37.

**[Th(BDPP)] (9): Method A.** [(BDPP)Th(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>2</sub>] (**7**) (0.200 g, 0.23 mmol) and 2,6-bis(2,6-diisopropylanilinoethyl)pyridine (0.106 g, 0.23 mmol) in benzene (15 mL) were heated at 100 °C for 24 h in a sealed flask. Upon cooling, the reaction mixture

(66) The reaction time was not reproducible. Typically, no reaction is observed for days, but once the reaction begins, it takes less than 24 h to reach completion.

was filtered and the solvent was reduced to ca. 1 mL *in vacuo*. Addition of hexamethyldisiloxane (10 mL) and sonication afforded **9** as a pale greenish-yellow solid (0.097 g, 0.085 mmol) in 37% isolated yield. **Method B.**  $\text{ThCl}_4(\text{dme})_2$  (0.100 g, 0.18 mmol) and  $\text{Li}_2\text{BDPP}$  (0.170 g, 0.36 mmol) in benzene<sup>30</sup> (15 mL) were stirred at 0 °C for 1 h (the majority of the solution does not freeze, despite the 6 °C freezing point of pure benzene). The reaction mixture was then warmed to room temperature and filtered, and the solvent was removed *in vacuo*. Addition of hexanes (5 mL) and sonication gave a brown solid, which was collected by filtration and washed with hexanes several times. Recrystallization by cooling a hot, concentrated toluene solution of **9** to −30 °C afforded the product as a pale greenish-yellow solid (0.109 g, 0.095 mmol) in 26% isolated yield.  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ , 600 MHz):  $\delta$  7.10 (m, 12H, Ar-*H*), 6.81 (t, 2H, *J* 7.7 Hz, py-*CH*), 6.48 (d, 4H, *J* 7.7 Hz, py-*CH*), 4.87 (s, 8H,  $\text{NCH}_2$ ), 3.00 (sept, 8H, *J* 6.5 Hz,  $\text{CHMe}_2$ ), 1.01 (d, 24H, *J* 6.5 Hz,  $\text{CHMe}_2$ ), 0.97 (d, 24H, *J* 6.6 Hz,  $\text{CHMe}_2$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CD}_2\text{-Cl}_2$ , 150 MHz):  $\delta$  167.5 (Py- $\text{C}_{\text{ortho}}$ ), 156.1 (Ar- $\text{C}_{\text{ortho}}$ ), 146.2 (Ar-

$\text{C}_{\text{ipso}}$ ), 138.0 (Py- $\text{CH}_{\text{meta}}$ ), 123.9 (Ar- $\text{CH}_{\text{meta}}$ ), 123.4 (Ar- $\text{CH}_{\text{para}}$ ), 117.5 (Py- $\text{CH}_{\text{para}}$ ), 68.0 ( $\text{NCH}_2$ ), 29.5 ( $\text{CHMe}_2$ ), 27.7, 24.1 ( $\text{CHMe}_2$ ). Anal. Calcd for  $\text{C}_{62}\text{H}_{82}\text{N}_6\text{Th}$ : C, 65.13; H, 7.23; N, 7.35. Found: C, 65.22; H, 7.36; N, 7.02.

**Acknowledgment.** D.J.H.E. thanks NSERC of Canada for a Discovery Grant and Canada Foundation for Innovation (CFI) and Ontario Innovation Trust (OIT) for New Opportunities Grants. D.J.H.E. also thanks McMaster University for support in the form of a start-up grant, and C.A.C. thanks the Government of Ontario for an Ontario Graduate Scholarship (OGS).

**Supporting Information Available:** X-ray crystallographic data in PDF format and CIF files are available free of charge via the Internet at <http://pubs.acs.org>.

OM060914F