

## C–C Bond Elimination from High-Valent Mn Aryl Complexes

Yu-Tao He, Ayumu Karimata, Olga Gladkovskaya, Eugene Khaskin, Robert R. Fayzullin, Abir Sarbajna, and Julia R. Khusnutdinova\*

Cite This: <https://doi.org/10.1021/acs.organomet.1c00047>

Read Online

ACCESS |

Metrics &amp; More

Article Recommendations

Supporting Information

**ABSTRACT:** Manganese complexes have been considered as a cheap and readily available alternative to commonly used precious metal catalysts in C–C bond coupling reactions. Although high-valent Mn aryl intermediates have been proposed in such reactions, a mechanistic understanding of possible organometallic intermediates in Mn-mediated C–C coupling is still lacking due to their high reactivity. We report the synthesis of stable, isolable Mn(III) aryl complexes obtained by oxidative addition of aryl bromide or aryl chloride. These complexes react with a range of organometallic alkylating or arylating reagents (alkyl and aryl Grignard reagents, MeLi, ZnMe<sub>2</sub>) to undergo C(sp<sup>2</sup>)–C(sp<sup>3</sup>) or C(sp<sup>2</sup>)–C(sp<sup>2</sup>) bond coupling, and a preliminary catalytic system could be demonstrated. The reagent scope and yield of the C(sp<sup>2</sup>)–C(sp<sup>3</sup>) coupled product can further be increased by addition of TEMPO as an oxidant, generating alkyl radicals.



## INTRODUCTION

Transition-metal-catalyzed C–C bond coupling is one of the most powerful and appealing methods for the construction of C–C bonds in synthetic chemistry.<sup>1,2</sup> For many decades, precious metals, in particular palladium complexes, have been predominant as highly efficient homogeneous catalysts in this transformation.<sup>3</sup> The commonly proposed mechanism of palladium-catalyzed cross-coupling typically involves two-electron oxidative addition, transmetalation, and reductive elimination as crucial steps in the catalytic cycle.<sup>4</sup>

The replacement of precious metals with abundant, cheap, and less toxic first-row transition metals has recently become a perennial goal of increasing urgency in applied synthetic chemistry.<sup>5–12</sup> Despite this, Mn-catalyzed cross-coupling of organic halides with organometallic reagents remains relatively underdeveloped. There are notable examples of Mn-catalyzed cross-coupling and homocoupling reactions reported by Cahiez,<sup>13–20</sup> Madsen,<sup>21,22</sup> Rueping,<sup>23</sup> Falck,<sup>24</sup> and others.<sup>25</sup> However, a mechanistic understanding of Mn-catalyzed C–C bond formation reactions is complicated by the paramagnetism of the vast majority of Mn complexes and intermediates due to the large variety of available oxidation states to this metal, and the frequently encountered low stability of organomanganese compounds that leads to a larger number of paramagnetic decomposition products.<sup>12</sup> In particular, the C–C bond elimination from Mn has sometimes been proposed to occur via either Mn<sup>III</sup> or Mn<sup>IV</sup> organometallic intermediates. For example, in the homocoupling of alkenyl lithium derivatives reported by Cahiez, Normant et al., the C–C elimination step was proposed to occur via an initial formation of a Mn<sup>IV</sup> species, followed by bond homolysis and reductive elimination from Mn<sup>III</sup> organometallic intermediates.<sup>26</sup> Intermediacy of high-valent Mn species in C(sp<sup>3</sup>)–C(sp<sup>2</sup>) bond elimination


has also been proposed by Falck and co-workers as one of the steps in Mn- or Mn/Cu(OTf)<sub>2</sub>-co-catalyzed Stille cross-coupling.<sup>24</sup> In more recent examples of homo- and cross-coupling of Grignard reagents in the presence of oxygen as an oxidant, C–C reductive elimination from Mn<sup>IV</sup> bis-organyl species has been proposed based on experimental and theoretical studies. Another important class of reactions in which intermediacy of either Mn<sup>III</sup> or Mn<sup>IV</sup> has been implicated is the directed C–H bond functionalization that leads to C–C bond formation. Nakamura and Ilies reported Mn-catalyzed alkylation of C(sp<sup>2</sup>)–H bonds using MeMgBr, in which a (cyclometalated aryl)(alkyl)Mn<sup>III</sup> intermediate was proposed to undergo C–C bond reductive elimination.<sup>27</sup> Ackerman and co-workers proposed that a C–C bond formation step occurs from a cyclometalated aryl Mn<sup>III</sup> intermediate in Mn-catalyzed C–H bond arylation or alkylation.<sup>28,29</sup> Interestingly, Ar–Ar elimination from Mn<sup>IV</sup> has been proposed by Uzelac et al. in Mn-catalyzed homocoupling, and Mn<sup>III</sup> species were isolated from the reaction mixtures containing Mn<sup>II</sup> aryl complexes under O<sub>2</sub>.<sup>30</sup> However, to the best of our knowledge, direct observation of the C–C bond coupling step from isolated Mn<sup>III</sup> or Mn<sup>IV</sup> aryl complexes has not been reported, although the intermediacy of Mn<sup>IV</sup> diaryl complexes in C–C bond coupling has been demonstrated by Taillefer and co-workers by electrospray ionization mass spectrometry.<sup>31</sup>

**Special Issue:** Organometallic Solutions to Challenges in Cross-Coupling

**Received:** January 26, 2021

Recently, we reported the isolation and characterization of (aryl)Mn<sup>III</sup> complexes stabilized by chelation to the macrocyclic pyridinophane <sup>t</sup>Bu<sub>3</sub>N3C<sup>-</sup> ligand (Scheme 1).<sup>32</sup> These

### Scheme 1. Oxidative Addition and Reductive Elimination of Ar-X Using (<sup>t</sup>Bu<sub>3</sub>N3C)Mn Aryl Complexes




complexes enabled us to study direct Ar–Br oxidative addition to a Mn<sup>I</sup> complex in the presence of light, to give stable and fully characterized (<sup>t</sup>Bu<sub>3</sub>N3C)Mn<sup>III</sup> monoaryl complexes. We also observed the subsequent oxidatively induced two-electron nonradical reductive elimination. Notably, Ar–X (X = Br, I, CN) bond elimination did not occur directly from Mn<sup>III</sup> complexes, presumably due to the lack of a driving force to prevent the reverse Ar–X bond oxidative addition and high stability. Thus, these stable aryl Mn<sup>III</sup> complexes served as convenient model compounds to study C–heteroatom bond oxidative addition and demonstrated the necessity of one-electron oxidation to induce aryl–X bond elimination.

In this work, we decided to study the possibility of C–C bond coupling from stable isolated Mn<sup>III</sup> aryl complexes in the absence and presence of oxidants. Notably, unlike in the case of Ar–X elimination which did not occur from Mn<sup>III</sup>, we observed C–C bond formation even from Mn<sup>III</sup> aryl complexes in the presence of methylating or arylating reagents even in the absence of oxidants, albeit with moderate or low efficiency and limited by the alkylating reagent scope. Interestingly, we found that addition of TEMPO or other one-electron oxidants to the aryl Mn<sup>III</sup> complex in the presence of alkylating agents promotes more selective and efficient aryl–C(sp<sup>3</sup>) bond elimination, presumably via an alternative pathway involving high-valent Mn<sup>IV</sup> intermediates. This study confirms that (aryl)Mn<sup>III</sup> complexes can be considered as possible intermediates in C–C bond coupling. However, there is an alternative pathway for Ar–C(sp<sup>3</sup>) bond coupling that is strongly affected by the presence of oxidative additives.

## RESULTS AND DISCUSSION

**Synthesis of (Aryl)Mn<sup>III</sup> Complexes.** We have previously reported the synthesis of complex **1** via oxidative addition of an aryl bromide <sup>t</sup>Bu<sub>3</sub>N3CBr to MnBr(CO)<sub>5</sub> at room temperature (RT) under mercury lamp irradiation in 1,2-dichloroethane (DCE) as a solvent (Scheme 1).<sup>32</sup> Screening of solvents showed that comparable yields can be obtained when the reaction is performed in THF to afford **1** in 51% isolated yield (Scheme 2). We hypothesized that the role of UV light was to remove CO ligands from the Mn center, and that the oxidative addition occurs by a nonradical pathway as confirmed by the


### Scheme 2. Synthesis of (<sup>t</sup>Bu<sub>3</sub>N3C)Mn<sup>III</sup>(X)(Y) Complexes



previous study.<sup>32</sup> Therefore, we decided to test whether the formation of **1** can be achieved under thermal conditions in the absence of UV light. Indeed, heating a mixture of <sup>t</sup>Bu<sub>3</sub>N3CBr and MnBr(CO)<sub>5</sub> in refluxing dichloroethane or toluene solution at 110 °C for 16 h also produced **1**, albeit in lower isolated yields, 32% and 16%, respectively.

We further examined the reactivity of aryl chloride derivative and reacted a chloro-substituted pyridinophane ligand <sup>t</sup>Bu<sub>3</sub>N3CCl with MnCl(CO)<sub>5</sub> in THF under UV light irradiation, as photochemical conditions gave the optimal yield of the desired product in the case of aryl bromide. Aryl chloride <sup>t</sup>Bu<sub>3</sub>N3CCl was synthesized by the reaction of 2-chloro-1,3-bis(bromomethyl)benzene with 2,6-bis(*tert*-butylamino-methyl)pyridine in THF. The expected dichloro complex **2**, was obtained in 22% isolated yield after complexation with the Mn precursor under standard conditions and recrystallization (Scheme 2). This indicates that, although aryl chloride undergoes oxidative addition, it is significantly less reactive compared to the bromide analogue. The structure of the dichloride complex **2** was confirmed by single-crystal X-ray diffraction (Figure 1a). The octahedral complex **2** is characterized by the positional C1/N1 disordering in the crystal.

In a similar way, we attempted the synthesis of mixed chloro bromo complexes by the oxidative addition of <sup>t</sup>Bu<sub>3</sub>N3CBr to MnCl(CO)<sub>5</sub> and the oxidative addition of <sup>t</sup>Bu<sub>3</sub>N3CCl to



**Figure 1.** ORTEP of complexes **2** (a) and **3** (b) at 70% probability level according to single-crystal X-ray diffraction data. Hydrogen atoms and minor disorder components are omitted for clarity. Selected interatomic distances [Å]: Mn1–Cl1 2.3087(8), Mn1–Cl2 2.3266(8), Mn1–C1 2.02(3), Mn1–N1 2.07(2), Mn1–N2 2.442(3), Mn1–N3 2.449(2) for **2**; Mn1–Br1 2.4576(14), Mn1–Cl2 2.321(2), Mn1–C1 1.98(3), Mn1–N1 2.10(2), Mn1–N2 2.442(3), Mn1–N3 2.465(2) for **3**.


MnBr(CO)<sub>5</sub>. As expected, oxidative addition of aryl bromide afforded the product **3** in a good yield of 74%, while the reaction with aryl chloride afforded the analogous mixed chloro bromo complex **3** in a much lower yield of 23%. The X-ray structure of the complex obtained by the first method is shown in Figure 1b. The complex appeared to be almost equally disordered with respect to Cl/Br and C1/N1 sites in the crystal studied. Molecule **3** shows distorted octahedral geometry at the Mn center with elongated Mn–N<sub>amine</sub> distances (2.442(3)–2.465(2) Å) as compared to Mn–N<sub>pyridine</sub> distances (2.10(2) Å). The Mn–C<sub>ipso</sub> bond distances in complexes **2** (2.02(3) Å) and **3** (1.98(3) Å) are comparable to that previously reported for complex **1** (2.027(14) Å). The analogous structure was established for complex **3** obtained by the oxidative addition of <sup>t</sup>BuN<sub>3</sub>CCl to MnBr(CO)<sub>5</sub>, and its X-ray structure is shown in Figure S60 (see the Supporting Information (SI)). Besides single-crystal X-ray diffraction, complexes **2** and **3** were characterized by UV–vis, IR spectroscopy, high-resolution mass spectrometry (HRMS), and elemental analysis. Complex **3** shows two peaks at 440.1641 and 484.1138 (*z* = 1) assigned to [M – Br]<sup>+</sup> and [M – Cl]<sup>+</sup>, respectively, displaying the expected isotopic patterns. Similarly, two peaks at 440.1637 and 484.1119 were also observed in complex **3** obtained by the inverted halogen precursor method.

The effective magnetic moment of new complexes **2** and **3** in solution measured by the Evans method in CD<sub>2</sub>Cl<sub>2</sub> solution was determined as 4.71 and 4.78 μ<sub>B</sub>, consistent with the presence of a high-spin d<sup>4</sup> Mn<sup>III</sup> center (*S* = 2), similar to previously reported complex **1**. The magnetic susceptibility measurements in the solid state for **2** and **3** also have χ<sub>M</sub>*T* values of 3.1–3.2 T·cm<sup>3</sup>·mol<sup>−1</sup> and 2.9–3.1 T·cm<sup>3</sup>·mol<sup>−1</sup>, respectively, in the temperature range from 10 to 300 K, corresponding to a high-spin d<sup>4</sup> center (μ<sub>eff</sub> of 5.0–5.1 and 4.8–4.9 μ<sub>B</sub>, respectively).

**C–C Elimination from Mn<sup>III</sup> Complexes without Oxidants.** Our previous study of Ar–X (*X* = Br, I, or CN) reductive elimination from Mn<sup>III</sup> showed that Ar–X bond

formation must be preceded by oxidation with a strong one-electron oxidant.<sup>32</sup> As C–C bond elimination is a different type of reaction, where the metal center in the intermediate has stronger σ-donor ligands, we set out to investigate whether Mn<sup>III</sup> complexes are capable of C–C bond elimination directly, in the absence of any oxidants. We hypothesized that the reaction of complex **1** with common organometallic reagents might lead to the initial transmetalation to form Mn<sup>III</sup> aryl alkyl or diaryl complexes, followed by the C–C reductive elimination (Scheme 3).

**Scheme 3.** Proposed C–C Elimination from (<sup>t</sup>BuN<sub>3</sub>)Mn<sup>III</sup> Complexes




We reacted complex **1** with 3 equiv of MeMgBr in THF solution in the absence of any additives at room temperature (RT). Attempts to isolate or spectroscopically detect transmetalation intermediates either at RT or low temperature were not successful, and the reaction proceeded immediately to undergo C–C bond coupling and give the expected C–C coupling product, <sup>t</sup>BuN<sub>3</sub>CMe, which was obtained after quenching the excess MeMgBr with aqueous ammonium chloride and basic workup to remove Mn residues (58% crude NMR yield, Table 1). The C–C coupling product was further isolated and characterized by NMR and HRMS. The low yield of the desired C–C coupling product is likely due to the formation of the protonated ligand <sup>t</sup>BuN<sub>3</sub>CH as a side-product, obtained in 28% NMR yield after workup of the reaction mixture. Reaction selectivity was decreased when only 1 equiv of MeMgBr was used, giving <sup>t</sup>BuN<sub>3</sub>CMe and <sup>t</sup>BuN<sub>3</sub>CH in 48% and 46% yields, respectively (entry 2).

Under analogous conditions, with 3 equiv of MeMgBr, dichloro complex **2** gave a 56% yield of <sup>t</sup>BuN<sub>3</sub>CMe and a 39% yield of <sup>t</sup>BuN<sub>3</sub>CH (entry 3).

The formation of <sup>t</sup>BuN<sub>3</sub>CH likely results from quenching starting material, aryl complex **1**, with aqueous ammonia solution used to remove an excess of methylating reagent. Thus, if a typical workup procedure involving treatment with aqueous NH<sub>4</sub>Cl solution and extraction is applied to complex **1** in the absence of methylating agents, formation of <sup>t</sup>BuN<sub>3</sub>CH is also detected. Although the alternative route to the formation of <sup>t</sup>BuN<sub>3</sub>CH could involve formation of aryl radical, the reaction with MeMgBr under typical conditions (RT, 16 h) in THF-*d*<sub>8</sub> showed that the <sup>t</sup>BuN<sub>3</sub>CH byproduct is formed exclusively and no deuterated product <sup>t</sup>BuN<sub>3</sub>CD was detected by GC-MS, ESI-MS, or NMR.

**Table 1. C–C Coupling from Complexes 1, 2, and 3 in the Presence of Organomagnesium, Zinc, and Lithium Reagents without Oxidants<sup>a</sup>**



Entry	Complex	Reagent (3 equiv)	<sup>t</sup> BuN <sub>3</sub> CR, % yield <sup>b</sup>	<sup>t</sup> BuN <sub>3</sub> CH, % yield <sup>b</sup>
1	<b>1</b>	MeMgBr	R = Me 58	28
2 <sup>c</sup>	<b>1</b>	MeMgBr (1 equiv)	R = Me 48	46
3	<b>2</b>	MeMgBr	R = Me 56	39
4	<b>3</b>	MeMgBr	R = Me 65	27
5 <sup>d</sup>	<b>1</b>	MeMgBr (-80° to RT)	R = Me 75	1
6	<b>1</b>	ZnMe <sub>2</sub>	R = Me 41	21
7 <sup>d</sup>	<b>1</b>	ZnMe <sub>2</sub> (-80° to RT)	R = Me 35	22
8	<b>1</b>	MeLi	R = Me 22	70
9 <sup>d</sup>	<b>1</b>	MeLi (-80° to RT)	R = Me 53	11
10	<b>1</b>	<sup>i</sup> PrMgCl	R = <i>i</i> Pr n.d. <sup>e</sup>	88
11	<b>1</b>	MgBr	R = CH <sub>2</sub> =CH-CH <sub>2</sub> - n.d. <sup>e</sup>	90
12	<b>1</b>	MgBr	R = Ph 67	16
13 <sup>f</sup>	<b>1</b>	MgBr	R = C <sub>6</sub> F <sub>5</sub> 49 (isolated)	n.d.

<sup>a</sup>Reaction was performed in THF for 16 h at RT in the presence of 3 equiv of alkylating or arylating reagent. <sup>b</sup>Yields were determined after quenching excess reagent by NMR integration against 1,3,5-trimethoxybenzene as an internal standard. <sup>c</sup>1 equiv of MeMgBr was added at RT. <sup>d</sup>Methylating reagent was added at -80 °C, and then the reaction mixture was slowly warmed up to RT. <sup>e</sup>Not detected. <sup>f</sup>Heated at 80 °C in benzene solution for 40 h in the presence of 6 equiv of C<sub>6</sub>F<sub>5</sub>MgBr.

Although the standard reaction time of 16 h was used in all experiments to ensure completion, the reaction in the case of MeMgBr occurs within several minutes at RT. Therefore, we also attempted methylation by slow addition of MeMgBr at


-80 °C, followed by slow warming up to RT. Under these conditions, a higher yield of methylation product was obtained, while formation of <sup>t</sup>BuN<sub>3</sub>CH was significantly suppressed, leading to more selective reaction (entry 5).

We have also tested the reactivity with other methylating reagents, ZnMe<sub>2</sub> and MeLi. In these cases, lower yields of the C–C coupling product <sup>t</sup>BuN<sub>3</sub>CMe were obtained under analogous conditions. In the case of MeLi, low temperature also had a beneficial effect, leading to higher yield of <sup>t</sup>BuN<sub>3</sub>CMe and suppressed formation of <sup>t</sup>BuN<sub>3</sub>CH as compared to the analogous reaction at RT (entries 8 and 9), whereas no significant effect of temperature was observed for ZnMe<sub>2</sub> (entries 6 and 7), probably due to its lower reactivity compared to MeLi and MeMgBr. Notably, in the presence of <sup>i</sup>PrMgBr and (allyl)MgBr, no desired C–C bond coupling product was obtained, with the major product after workup being <sup>t</sup>BuN<sub>3</sub>CH (Table 1).

We have further tested the possibility of aryl–aryl coupling and reacted complex **1** with 3 equiv of PhMgBr in THF at RT. This resulted in the formation of 67% of phenylated product, <sup>t</sup>BuN<sub>3</sub>CPh, after 16 h. The reaction with less nucleophilic pentafluorophenyl Grignard reagent, C<sub>6</sub>F<sub>5</sub>MgBr, gave no product at RT. However, when complex **1** was heated in the presence of 6 equiv of C<sub>6</sub>F<sub>5</sub>MgBr in benzene at 80 °C, the pentafluorophenyl derivative, <sup>t</sup>BuN<sub>3</sub>CC<sub>6</sub>F<sub>5</sub>, was obtained in 49% yield.

These results show that, unlike Ar–X reductive elimination, aryl–C(sp<sup>3</sup>) and aryl–C(sp<sup>2</sup>) bond elimination in (<sup>t</sup>BuN<sub>3</sub>C)–Mn<sup>III</sup> complexes does not necessarily require an oxidant and likely proceeds directly via the Mn<sup>III</sup> oxidation state, although the proposed methyl–aryl or bis(aryl) intermediates could not be isolated or observed, presumably due to fast C–C coupling following transmetalation (Scheme 3). We attempted a different approach to obtain such intermediates using MeMn(CO)<sub>5</sub> as a precursor and reacted it with <sup>t</sup>BuN<sub>3</sub>CBr under UV irradiation (Scheme 4). If the reactivity of MeMn(CO)<sub>5</sub> is

**Scheme 4. Reaction of MeMn(CO)<sub>5</sub> with <sup>t</sup>BuN<sub>3</sub>CBr**



similar to that of MnBr(CO)<sub>5</sub>, we could expect formation of the Ar–Br oxidative addition product where a methyl group is present instead of one bromo ligand. As anticipated, when MeMn(CO)<sub>5</sub> was reacted in the presence of 1 equiv of <sup>t</sup>BuN<sub>3</sub>CBr under UV light irradiation in THF or toluene solution, the reaction proceeded to form the C–C coupled product, <sup>t</sup>BuN<sub>3</sub>CMe, in 7–13% yield. No intermediates could be isolated, suggesting that, once the expected aryl methyl complex is formed, it immediately undergoes C–C elimination.

The low yield of the coupling product could be due to only one Me group being present in the oxidative addition product, while the reaction of **1** with excess MeMgBr could lead to replacement of both bromo ligands with the Me groups. However, formation of a small amount of (<sup>t</sup>BuN<sub>3</sub>C)Mn<sup>III</sup>Me<sub>2</sub> in this reaction cannot be excluded if Me group exchange

between two monomethyl intermediates is involved, similar to the Schlenk equilibrium<sup>33</sup> present in solutions of Grignard reagents. Alternatively, the low yield in the case of the  $\text{MeMn}(\text{CO})_5$  reaction could be due to the amount of the isomer in which Ar and Me groups are present in mutual *cis*-positions.<sup>32</sup>


Unfortunately, the  $\text{Mn}^{\text{I}}$  product of C–C bond coupling could not be isolated due to the absence of stabilizing ligands and basic workup of the reaction mixture leading to precipitation of all Mn residues or their removal to the aqueous phase.

Overall, these experiments indicate that C–C coupling proceeds upon the reaction of common organometallic reagents with isolated  $\text{Mn}^{\text{III}}$  complexes in the absence of oxidative additives or chlorinated solvents that could act as oxidants, suggesting that the C–C coupling step likely occurs at the  $\text{Mn}^{\text{III}}$  aryl center, although the selectivity of this reaction remains limited.


To demonstrate that the proposed sequence of oxidative addition/transmetalation/C–C elimination steps may give rise to a catalytic turnover, we performed a catalytic experiment using a mixture of only 10 mol % of  $\text{MnBr}(\text{CO})_5$ ,  ${}^t\text{Bu}_3\text{N3CBr}$ , and 3 equiv of Grignard reagent in THF solution (Scheme 5,

#### Scheme 5. Mn-Catalyzed ${}^t\text{Bu}_3\text{N3CBr}$ Methylation with MeMgBr

Catalytic experiment:




Stoichiometric experiment:



top). Interestingly, we observed catalytic turnover (TON 8.5) under these conditions, leading to the formation of  ${}^t\text{Bu}_3\text{N3CMe}$  in 85% yield after 20 h. This result confirms that  $({}^t\text{Bu}_3\text{N3C})\text{Mn}$  complexes not only are suitable model compounds to study Ar–X and Ar–C bond formation but also mimic Mn-catalyzed Ar–C bond coupling. Notably, ICP-MS analysis of the catalyst showed that no platinum group metals were present in the sample at the detectable level (see the SI).

The control experiment showed that no reaction occurs in the absence of Mn complexes when  ${}^t\text{Bu}_3\text{N3CBr}$  is treated with an excess of  $\text{MeMgBr}$ ,  $\text{ZnMe}_2$ , or  $\text{MeLi}$  in THF under analogous conditions (Scheme 6), confirming that manganese complexes are indeed required to mediate the cross-coupling reaction. To further confirm that the C–C coupling is mediated by the formation of  $\text{Mn}^{\text{III}}$  aryl species, we performed the reaction in the presence of equivalent amounts of  $\text{Mn}^{\text{I}}\text{Br}(\text{CO})_5$ ,  ${}^t\text{Bu}_3\text{N3CBr}$ , and 3 equiv of  $\text{MeMgBr}$  (Scheme

#### Scheme 6. Control Experiment in the Absence of Mn Complexes



5, bottom). The ESI-MS analysis of the reaction mixture confirmed the presence of the characteristic peak with  $m/z$  486.1114 corresponding to the  $[\text{M} - \text{Br}]^+$  fragment of complex **1**, along with peaks of  ${}^t\text{Bu}_3\text{N3CMe}$ ,  ${}^t\text{Bu}_3\text{N3CBr}$ , and  ${}^t\text{Bu}_3\text{N3CH}$  (see the SI). This experiment also shows that the reaction of  $\text{Mn}^{\text{I}}\text{Br}(\text{CO})_5$  with  ${}^t\text{Bu}_3\text{N3CBr}$  occurs prior to C–C elimination. When  $\text{Mn}^{\text{I}}\text{Br}(\text{CO})_5$  was reacted with  $\text{MeMgBr}$  in the absence of  ${}^t\text{Bu}_3\text{N3CBr}$ , no  $\text{MeMn}^{\text{I}}(\text{CO})_5$  was observed.

**C–C Elimination in the Presence of Oxidants.** To further test the mechanism of C–C coupling from complex **1** in the reaction with  $\text{MeMgBr}$ , we attempted to perform the reaction in the presence of TEMPO as a radical trap. Surprisingly, when C–C coupling from **1** using  $\text{MeMgBr}$  (3 equiv) as a methylating reagent was performed in the presence of 1.1 equiv of TEMPO, we observed almost clean formation of the C–C coupling product in 92% yield, while no protonation product,  ${}^t\text{Bu}_3\text{N3CH}$ , was detected under these conditions (Table 2). Notably, the yield was high even in the

Table 2. C–C Coupling from Complex **1** in the Presence of TEMPO<sup>a</sup>

entry	equiv of MeMgBr	equiv of TEMPO	${}^t\text{Bu}_3\text{N3CMe}$ , % yield <sup>b</sup>	${}^t\text{Bu}_3\text{N3CH}$ , % yield <sup>b</sup>
1	3.0	none	58	28
2	3.0	0.1	72	15
3	3.0	1.1	92	n.d. <sup>c</sup>
4	3.0	2.0	93	n.d.
5	3.0	3.0	97	n.d.
6	2.0	2.0	90	n.d.
7	1.0	1.0	59	13


presence of 1.1 equiv of TEMPO and only slightly increased when the amount of TEMPO was changed to 2 and 3 equiv, showing that even 1.1 equiv was sufficient to promote selective C–C bond formation (Table 2). A more selective C–C coupling was also observed in the reaction where only 1 equiv of  $\text{MeMgBr}$  and 1 equiv of TEMPO were used, giving 59% of  ${}^t\text{Bu}_3\text{N3CMe}$  and 13% of protonated product  ${}^t\text{Bu}_3\text{N3CH}$ , as




presence of 1.1 equiv of TEMPO and only slightly increased when the amount of TEMPO was changed to 2 and 3 equiv, showing that even 1.1 equiv was sufficient to promote selective C–C bond formation (Table 2). A more selective C–C coupling was also observed in the reaction where only 1 equiv of  $\text{MeMgBr}$  and 1 equiv of TEMPO were used, giving 59% of  ${}^t\text{Bu}_3\text{N3CMe}$  and 13% of protonated product  ${}^t\text{Bu}_3\text{N3CH}$ , as

compared to the reaction with 1 equiv of MeMgBr in the absence of TEMPO, which gave only 43% of  ${}^t\text{BuN3CMe}$  (vide supra).

Encouraged by these results, we also examined if the scope of alkylating reagents could be improved using these conditions (Table 3). Interestingly, in the presence of 2

**Table 3. C–C Coupling from Complexes 1, 2, and 3 with Other Alkylating and Arylating Reagents in the Presence of TEMPO as an Oxidant<sup>a</sup>**



Entry	Complex	RM	${}^t\text{BuN3CR}$ , % yield <sup>b</sup>
1	1	ZnMe <sub>2</sub>	R = Me 95
2	1	MeLi	R = Me 40
3	1	 MgCl	R = <i>i</i> Pr 47
4	1	 MgBr	R = allyl 72
5	1	 MgBr	R = Ph 71
6	2	MeMgBr	R = Me 87
7	3	MeMgBr	R = Me 92

<sup>a</sup>Reaction was performed in THF for 16 h at RT. <sup>b</sup>Yields were determined after quenching excess reagent by NMR integration against 1,3,5-trimethoxybenzene as an internal standard.


equiv of TEMPO, the C–C coupling product was also obtained using *i*PrMgBr and (allyl)MgBr as coupling reagents in 47% and 72% yields, respectively. High yield of methylation product was also observed using ZnMe<sub>2</sub> (95%). The reaction with MeLi in the presence of TEMPO produced  ${}^t\text{BuN3CMe}$  in a moderate 40% yield, as compared to only 22% obtained in the absence of additives. High yields of  ${}^t\text{BuN3CMe}$  were also obtained using the similar dichloro and chloro bromo complexes 2 and 3 as coupling partners.

Incongruently, the presence of TEMPO did not play a significant role in the promotion of PhMgBr reaction, and  ${}^t\text{BuN3CPh}$  was obtained in almost the same yield with and without TEMPO.

To clarify the role of TEMPO, we first tested the reactivity of complex 1 with 2 equiv of TEMPO; however, no reaction was observed, and the starting material remained unchanged.


When MeMgBr was allowed to react with 1 equiv of TEMPO in the absence of complex 1, we observed the formation of Me-TEMPO in 52% yield, which indicates that the Me radical is formed under these conditions trapped by TEMPO. This observation is consistent with the reported detailed studies of the reactivity of TEMPO with alkyl organometallic compounds.<sup>34–37</sup> TEMPO is known to react with alkyl organometallic species first via a one-electron oxidation to produce an alkyl radical, which is then trapped by another equivalent of TEMPO (Scheme 7) to produce a Me-TEMPO adduct.

**Scheme 7. Reactivity of TEMPO with Grignard Reagents in the Absence of Mn Complexes (This Work)**




Independently prepared Me-TEMPO did not react with complex 1 under typical reaction conditions. Interestingly, aryl Grignard reagents react by a different pathway which does involve the formation of high energy free aryl radicals.<sup>34</sup> Indeed, the reaction of PhMgBr with 1 equiv of TEMPO did not produce Ph-TEMPO, but led to homocoupling to give biphenyl, as is consistent with literature precedents (Scheme 8).

**Scheme 8. Literature-Reported Reactivity of TEMPO with Alkylmetal Reagents**



On the basis of these observations, we propose that the increased yield of C–C coupling product from complex 1 in the presence of TEMPO could involve a different pathway via the formation of an alkyl radical, which could then either react with a Mn<sup>III</sup> center to give a  $({}^t\text{BuN3C})\text{Mn}^{\text{IV}}(\text{Me})\text{X}_2$  (X = halogen or Me) intermediate (Path 1a, Scheme 9) or directly attack a Mn-bound aryl ligand without changes in Mn oxidation state (Path 1b, Scheme 9). The absence of the boosting TEMPO effect in the reaction with aryl Grignard reagents could be due to the absence of free radicals that would

### Scheme 9. Proposed Mechanism of C–C Coupling in the Presence of TEMPO or Other Oxidants



react with a Mn<sup>III</sup> complex, further suggesting that the formation of alkyl radicals may be a prerequisite for the observed more efficient C(sp<sup>2</sup>)–C(sp<sup>3</sup>) coupling reactivity.


The fast reactions of Me or other alkyl radicals with transition metals, leading to their formal one-electron oxidation and M–C bond formation, have been studied extensively for some first-row transition metals,<sup>38,39</sup> and a similar mechanism was proposed for C(sp<sup>3</sup>)–C(sp<sup>3</sup>) bond elimination from a monomethyl Pd<sup>III</sup> complex via oxidation to Pd<sup>IV</sup> by a Me radical.<sup>40</sup> Although the Mn<sup>III</sup> complex **1** is six-coordinate, with a κ<sup>4</sup>-bound tBu<sub>3</sub>N<sub>3</sub>C<sup>−</sup> ligand, a vacant coordination site in Path 1b could form via dissociation of a bulky tBu-substituted amine group. The hemilability of tBu-amine groups in structurally similar tBu<sub>3</sub>N<sub>4</sub>-pyridinophane ligands has been studied extensively, showing facile inter-conversion between the κ<sup>4</sup>-bound and κ<sup>3</sup>-bound ligand coordination modes.<sup>41–43</sup>


Interestingly, higher yields of C–C coupling product were also observed in the presence of some other one-electron oxidants including Cu<sup>II</sup>(OTf)<sub>2</sub>, ferrocenium hexafluorophosphate, and thianthrenyl hexachloroantimonate, whereas the yield remained unchanged or decreased in the presence of silver tetrafluoroborate and tris(4-bromophenyl)aminium hexachloroantimonate (Magic Blue), respectively, compared to the reaction in the absence of any additives (Table 4). The increased yield of tBu<sub>3</sub>N<sub>3</sub>CMe coupling product in the presence of other one-electron oxidants could similarly be explained by the generation of a Me radical. However, we cannot exclude an alternative mechanism (Path 2, Scheme 9) via initial transmetalation, followed by oxidation to a Mn<sup>IV</sup> species and oxidatively induced reductive elimination similar to the mechanism proposed by us for Ar–X (X = Br, I, CN) elimination in the presence of strong oxidants (Scheme 1).

## SUMMARY AND CONCLUSION

Mn-mediated C–C bond coupling reactions attract attention as an inexpensive alternative to the use of precious metal in C–C cross-coupling reactions. The intermediacy of Mn<sup>III</sup> or Mn<sup>IV</sup> species in C–C bond coupling has been a subject of debate, and such complexes have been proposed as intermediates in

### Table 4. C–C Coupling from Complex **1** in the Presence of MeMgBr and One-Electron Oxidants<sup>a</sup>



Entry	Oxidant	tBu <sub>3</sub> N <sub>3</sub> CMe, % yield <sup>b</sup>	tBu <sub>3</sub> N <sub>3</sub> CH, % yield <sup>b</sup>
1	Cu(OTf) <sub>2</sub>	88	n.d. <sup>c</sup>
2	 SbF <sub>6</sub> <sup>−</sup>	90	n.d.
3	[(p-BrC <sub>6</sub> H <sub>4</sub> ) <sub>3</sub> N]SbCl <sub>6</sub>	32	n.d.
4	AgBF <sub>4</sub>	68	n.d.
5	Fc <sup>+</sup> PF <sub>6</sub> <sup>−</sup>	37	n.d.

<sup>a</sup>Reaction was performed in THF for 16 h at RT in the presence of 3 equiv of MeMgBr and 2 equiv of oxidant. <sup>b</sup>Yields were determined after quenching excess reagent by NMR integration against 1,3,5-trimethoxybenzene as an internal standard. <sup>c</sup>Not detected.

several examples of Mn-catalyzed C–C bond coupling reactions. In this work, we demonstrate that Mn<sup>III</sup> aryl species are indeed viable intermediates in C–C bond elimination and they undergo C–C bond coupling with a range of alkylating and arylating organometallic reagents under mild conditions. Although the couplings occur in the absence of any oxidants, the presence of oxidative additives, in particular TEMPO, significantly improves the reaction selectivity and the scope of alkylating agents, presumably via oxidation to Mn<sup>IV</sup> and/or alkyl radical formation. Our results suggest that two pathways, via either a Mn<sup>III</sup> or a Mn<sup>IV</sup> center, may be operative in C–C coupling bond elimination reactions. While we were able to develop an initial catalytic protocol via the former pathway, the latter leads to greater yield and selectivity despite requiring a sacrificial oxidant. We are currently focusing on trying to develop a more practical catalytic system via both the Mn<sup>III</sup> and Mn<sup>IV</sup> oxidation states for a greater variety of substrates.

## EXPERIMENTAL SECTION

All manipulations, unless stated otherwise, were performed using Schlenk or glovebox techniques under a dry argon atmosphere. Anhydrous solvents were dispensed from an MBRAUN solvent purification system and degassed before use. Anhydrous deuterated solvents were purchased from Eurisotop and stored over 4 Å molecular sieves. All chemicals unless noted otherwise were purchased from major commercial suppliers (TCI, Sigma-Aldrich, and Nacalai Tesque) and used without purification. tBu<sub>3</sub>N<sub>3</sub>CBr was synthesized as described previously.<sup>32,44</sup> 2-Chloro-1,3-bis-(bromomethyl)benzene was synthesized according to the literature procedure.<sup>45</sup> 2,6-Bis(tert-butylaminomethyl)pyridine was synthesized according to the modified literature method.<sup>46</sup> MnCl(CO)<sub>5</sub><sup>47</sup> and MeMn(CO)<sub>5</sub><sup>48</sup> were obtained according to literature procedures.

**Instrumentation.** NMR spectra were measured on JEOL ECZ600R 600 MHz, JEOL ECZ400S 400 MHz, Bruker Avance II 400 MHz, and Bruker Avance III Neo 500 MHz (CryoProbe) spectrometers. ESI-MS measurements were performed on a Thermo Scientific ETD apparatus. Elemental analyses were performed using an Exeter Analytical CE440 instrument. FT-IR spectra were measured

using an Agilent Cary 630 with an ATR module in an argon-filled glovebox. The following abbreviations are used for describing FT-IR spectra: s (strong), m (medium), w (weak), br (broad). UV–vis absorbance spectra were collected using an Agilent Cary 60 instrument. The magnetic properties were measured using a 9T physical properties measurement system PPMS Dynacool from Quantum Design, equipped with the vibrating sample magnetometer (VSM) option, in a 2–300 K temperature range under a magnetic field of 10 000 Oe. For these measurements, samples were ground into powder and placed in plastic capsules. The Evans method measurements were performed in the coaxial NMR tube at 298 K; diamagnetic correction was applied.<sup>49</sup>

**X-ray Structure Determination Details.** The X-ray diffraction data were collected on a Rigaku XtaLab PRO instrument in an  $\omega$ -scan mode with a PILATUS3 R 200 K hybrid pixel array detector and MicroMax-003 microfocuss X-ray tubes using Cu  $K\alpha$  (1.54184 Å) or Mo  $K\alpha$  (0.71073 Å) radiation at low temperature. Images were indexed and integrated using the CrysAlis<sup>Pro</sup> (version 1.171.41.93a) data reduction package. Data were corrected for systematic errors and absorption using the ABSPACK module: Numerical absorption correction based on Gaussian integration over a multifaceted crystal model and empirical absorption correction based on spherical harmonics according to the point group symmetry using equivalent reflections. The GRAL module was used for the analysis of systematic absences and space-group determination. All structures were solved by the direct methods using SHELXT-2018/2<sup>50</sup> and refined by the full-matrix least-squares on  $F^2$  using SHELXL-2018/3.<sup>51</sup> Non-hydrogen atoms were refined anisotropically. The hydrogen atoms were inserted at the calculated positions and refined as riding atoms. The positions of the hydrogen atoms of methyl groups were found using rotating group refinement with idealized tetrahedral angles. The disorder was resolved using free variables and reasonable restraints on geometry and anisotropic displacement parameters. The crystal structures of **3** obtained by different methods were refined as two-component inversion twins.

**Crystallographic Data for 2.** C<sub>23</sub>H<sub>32</sub>Cl<sub>2</sub>MnN<sub>3</sub>, brown prism (0.111 × 0.061 × 0.046 mm<sup>3</sup>), formula weight 476.35; orthorhombic, *Pbca* (No. 61), *a* = 14.38378(7) Å, *b* = 11.29423(6) Å, *c* = 27.85495(19) Å, *V* = 4525.14(5) Å<sup>3</sup>, *Z* = 8, *Z'* = 1, *T* = 95(2) K, *d*<sub>calc</sub> = 1.398 g cm<sup>-3</sup>,  $\mu$ (Cu  $K\alpha$ ) = 7.016 mm<sup>-1</sup>, *F*(000) = 2000; *T*<sub>max/min</sub> = 1.000/0.636; 44 277 reflections were collected (4.419° ≤  $\theta$  ≤ 78.086°, index ranges: -18 ≤ *h* ≤ 18, -14 ≤ *k* ≤ 13, -27 ≤ *l* ≤ 34), 4819 of which were unique, *R*<sub>int</sub> = 0.0277, *R* <sub>$\sigma$</sub>  = 0.0158; completeness to  $\theta$ <sub>max</sub> 99.4%. The refinement of 275 parameters with 84 restraints converged to *R*<sub>1</sub> = 0.0466 and *wR*<sub>2</sub> = 0.1086 for 4789 reflections with *I* > 2 $\sigma$ (*I*) and *R*<sub>1</sub> = 0.0468 and *wR*<sub>2</sub> = 0.1087 for all data with *S* = 1.159 and residual electron density,  $\rho$ <sub>max/min</sub> = 0.544 and -0.394 e Å<sup>-3</sup>. The crystals were grown by diethyl ether vapor diffusion to a dichloromethane solution.

**Crystallographic Data for 3.** C<sub>23</sub>H<sub>32</sub>Br<sub>0.88</sub>Cl<sub>1.12</sub>MnN<sub>3</sub>, red prism (0.194 × 0.170 × 0.094 mm<sup>3</sup>), formula weight 515.62; orthorhombic, *Pca*<sub>21</sub> (No. 29), *a* = 14.6467(3) Å, *b* = 13.8349(3) Å, *c* = 11.2247(3) Å, *V* = 2274.53(9) Å<sup>3</sup>, *Z* = 4, *Z'* = 1, *T* = 95(2) K, *d*<sub>calc</sub> = 1.506 g cm<sup>-3</sup>,  $\mu$ (Mo  $K\alpha$ ) = 2.278 mm<sup>-1</sup>, *F*(000) = 1064; *T*<sub>max/min</sub> = 1.000/0.629; 26 895 reflections were collected (2.719° ≤  $\theta$  ≤ 32.230°, index ranges: -19 ≤ *h* ≤ 21, -19 ≤ *k* ≤ 20, -15 ≤ *l* ≤ 16), 7151 of which were unique, *R*<sub>int</sub> = 0.0339, *R* <sub>$\sigma$</sub>  = 0.0344; completeness to  $\theta$ <sub>max</sub> 93.3%. The refinement of 296 parameters with 113 restraints converged to *R*<sub>1</sub> = 0.0321 and *wR*<sub>2</sub> = 0.0737 for 6532 reflections with *I* > 2 $\sigma$ (*I*) and *R*<sub>1</sub> = 0.0366 and *wR*<sub>2</sub> = 0.0747 for all data with *S* = 1.055 and residual electron density,  $\rho$ <sub>max/min</sub> = 0.423 and -0.415 e Å<sup>-3</sup>. The crystals were grown by diethyl ether vapor diffusion to a dichloromethane solution.

**Crystallographic Data for 3 Obtained by the Oxidative Addition of <sup>t</sup>BuN<sub>3</sub>CCl to MnBr(CO)<sub>5</sub>.** C<sub>23</sub>H<sub>32</sub>Br<sub>0.84</sub>Cl<sub>1.16</sub>MnN<sub>3</sub>, red prism (0.223 × 0.126 × 0.103 mm<sup>3</sup>), formula weight 513.86; orthorhombic, *Pca*<sub>21</sub> (No. 29), *a* = 14.6407(4) Å, *b* = 13.8107(4) Å, *c* = 11.2124(4) Å, *V* = 2267.12(12) Å<sup>3</sup>, *Z* = 4, *Z'* = 1, *T* = 95(2) K, *d*<sub>calc</sub> = 1.505 g cm<sup>-3</sup>,  $\mu$ (Cu  $K\alpha$ ) = 7.830 mm<sup>-1</sup>, *F*(000) = 1061; *T*<sub>max/min</sub> = 1.000/0.368; 14 985 reflections were collected (3.200° ≤  $\theta$  ≤ 77.818°, index ranges: -18 ≤ *h* ≤ 18, -15 ≤ *k* ≤ 17, -13 ≤ *l* ≤ 13), 4393 of which

were unique, *R*<sub>int</sub> = 0.0746, *R* <sub>$\sigma$</sub>  = 0.0526; completeness to  $\theta$ <sub>max</sub> 97.6%. The refinement of 296 parameters with 113 restraints converged to *R*<sub>1</sub> = 0.0626 and *wR*<sub>2</sub> = 0.1667 for 4193 reflections with *I* > 2 $\sigma$ (*I*) and *R*<sub>1</sub> = 0.0643 and *wR*<sub>2</sub> = 0.1679 for all data with *S* = 1.075 and residual electron density,  $\rho$ <sub>max/min</sub> = 1.191 and -0.862 e Å<sup>-3</sup>. The crystals were grown by diethyl ether vapor diffusion to a dichloromethane solution.

**Synthesis of <sup>t</sup>BuN<sub>3</sub>CCl.** *Synthesis of 2,6-Bis(tert-butylaminomethyl)pyridine.* A 500 mL round-bottom flask equipped with a magnetic stirring bar and a reflux condenser was charged with chloroform (100 mL); *tert*-butylamine (100 mL, 950 mmol, 50 equiv) was added to stirred chloroform slowly, in several portions. A solution of 2,6-bis(bromomethyl)pyridine (5 g, 19 mmol, 1 equiv) in CHCl<sub>3</sub> (150 mL) was slowly added to the stirred solution (**Caution! exothermic**) at room temperature. The resulting solution was heated at 50 °C for 5 h under a N<sub>2</sub> atmosphere. The sample of the reaction mixture after heating for 5 h was evaporated to dryness and redissolved in CDCl<sub>3</sub> to check the completeness of the reaction; according to <sup>1</sup>H NMR, no starting material was present after 5 h and the diamine was the only pyridine-containing product present in solution. The reaction mixture was cooled down to RT; the chloroform solution was washed with saturated aqueous NaHCO<sub>3</sub> solution (2 × 150 mL) and water (150 mL). The organic layer was separated and dried over MgSO<sub>4</sub>. The solvents were removed by rotary evaporation at RT to give a colorless or pale yellow oil, which was further dried under high vacuum for 1 h at RT and stored under N<sub>2</sub> at -20 °C. Yield 4.58 g, 97%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz),  $\delta$  (ppm): 7.56 (t, *J* = 7.5 Hz, 1H, Py CH<sub>para</sub>), 7.17 (d, *J* = 7.5 Hz, 2H, Py CH<sub>meta</sub>), 3.86 (s, 4H, CH<sub>2</sub>), 1.67 (br s, 2H, NH), 1.19 (s, 18H, *t*Bu).

*Synthesis of <sup>t</sup>BuN<sub>3</sub>CCl.* A round-bottom flask was charged with 50 mL of toluene, 30 mL of 10 wt % Na<sub>2</sub>CO<sub>3</sub> solution, and 2,6-bis(*tert*-butylaminomethyl)pyridine (1.10 g, 4.4 mmol, 1 equiv). The flask was supplemented with a two-neck extension; one neck was connected to a reflux condenser, and the other one with an addition funnel. The reaction mixture was heated up to 80 °C under a nitrogen atmosphere. A solution of 2-chloro-1,3-bis(bromomethyl)benzene (1.18 g, 4.0 mmol, 0.9 equiv) in 50 mL of toluene was added dropwise through the addition funnel to the reaction mixture over 3 h. After complete addition, the reaction mixture was heated with stirring at 90 °C for 36 h and then cooled down. The aqueous layer was discarded, and the pale yellow toluene layer was washed three times with saturated Na<sub>2</sub>CO<sub>3</sub> aqueous solution and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The solvent was removed, and the solid residue was washed with a minimal amount of ethanol and filtered off. The solid product was recrystallized from warm Et<sub>2</sub>O solution and cooled down in a freezer overnight to give a pure white crystalline product, which was filtered off and dried under vacuum. Yield 450 mg (1.17 mmol, 29%). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.13 (t, *J* = 7.6 Hz, 1H), 6.82 (d, *J* = 7.4 Hz, 2H), 6.75 (d, *J* = 7.6 Hz, 2H), 6.59 (t, *J* = 7.4 Hz, 1H), 4.15 (dd, *J* = 16.7, 13.3 Hz, 4H), 3.96 (d, *J* = 12.9 Hz, 2H), 3.52 (d, *J* = 13.4 Hz, 2H), 1.32 (s, 18H). <sup>13</sup>C NMR (126 MHz, CDCl<sub>3</sub>)  $\delta$  159.86, 137.26, 136.84, 135.00, 131.13, 124.99, 121.38, 56.77, 55.79, 51.38, 27.71. HRMS (ESI) (*m/z*): calculated for C<sub>23</sub>H<sub>32</sub>ClN<sub>3</sub> [M + H]<sup>+</sup> 386.2358; found 386.2335.

**Synthesis of (<sup>t</sup>BuN<sub>3</sub>C)Mn<sup>III</sup>Cl<sub>2</sub> (2).** 38.5 mg of <sup>t</sup>BuN<sub>3</sub>CCl (0.1 mmol) and 23 mg of MnCl(CO)<sub>5</sub> (0.1 mmol) were combined in a flame-dried Schlenk flask inside a glovebox, and 2 mL of THF was added to give a yellow suspension. The flask was taken outside the glovebox and stirred in a water bath in front of a mercury lamp. The reaction vessel was subjected to vacuum for 1 s every hour, then stirred under static vacuum, and after 3 h, the reaction was then stirred under a static vacuum overnight. After 13 h, the solution color changed to wine-red and the entire solvent was evaporated under reduced pressure. The solid obtained was redissolved in a minimal amount of dichloromethane and filtered through Celite. The filtrate was evaporated under reduced pressure to yield a red solid which was washed three times with copious amounts of ether (ca. 3 mL) and then dried under vacuum. Deep red crystals were grown by vapor diffusion of ether into a dichloromethane solution of the complex (about 2 mL of DCM); yield of isolated product (<sup>t</sup>BuN<sub>3</sub>C)Mn<sup>III</sup>Cl<sub>2</sub>,

(2) 10.4 mg, 22% yield. Evans method ( $\text{CD}_2\text{Cl}_2$ , 298 K):  $\mu_{\text{eff}} = 4.71 \mu_{\text{B}}$ . ESI-HRMS of  $\text{C}_{23}\text{H}_{32}\text{ClN}_3\text{Mn}$  in MeOH ( $m/z$ ): calculated for  $[\text{C}_{23}\text{H}_{32}\text{ClN}_3\text{Mn}]^+$ ,  $([\text{M} - \text{Cl}]^+)$  440.1660; found 440.1632 ( $z = 1$ ). Anal. Calcd. for  $\text{MnC}_{23}\text{H}_{32}\text{ClN}_3$ : C, 57.99; H, 6.77; N, 8.82. Found: C, 57.74; H, 6.65; N, 8.73. UV-vis,  $\lambda$ , nm ( $\epsilon$ ,  $\text{M}^{-1} \text{cm}^{-1}$ ),  $\text{CH}_2\text{Cl}_2$ : 534 (328), 301 (3412), 235 (8934). FT-IR (ATR, solid,  $\text{cm}^{-1}$ ):  $\nu$  2966 (w), 2893 (w), 1601 (w), 1576 (w), 1428 (w), 1376 (s), 1189 (s), 909 (w), 849 (w), 772 (s).

**Synthesis of  $(^{\text{tBu}}\text{N}_3\text{C})\text{Mn}^{\text{III}}\text{BrCl}$  (3) by Oxidative Addition of  $^{\text{tBu}}\text{N}_3\text{CBr}$ .** 42.9 mg of  $^{\text{tBu}}\text{N}_3\text{CBr}$  (0.1 mmol) and 23 mg of  $\text{MnCl}(\text{CO})_5$  (0.1 mmol) were combined in a flame-dried Schlenk flask inside a glovebox, and 2 mL of THF was added to give a yellow suspension. The flask was taken outside the glovebox and stirred in a water bath in front of a mercury lamp. The reaction vessel was subjected to vacuum for 1 s every hour, then stirred under static vacuum, and after 3 h, the reaction was then stirred under a static vacuum overnight. After 13 h, the solution appeared wine-red and the entire solvent was evaporated under reduced pressure. The solid obtained was redissolved in a minimal amount of dichloromethane and filtered through Celite. The filtrate was evaporated under reduced pressure to yield a red solid which was washed three times with copious amounts of ether (ca. 3 mL) and then dried under vacuum. Deep red crystals were grown by vapor diffusion of ether into a dichloromethane solution of the complex (about 2 mL of DCM); yield of isolated product  $(^{\text{tBu}}\text{N}_3\text{C})\text{Mn}^{\text{III}}\text{BrCl}$  (3) 38.2 mg, 74% yield. HRMS (ESI) in MeOH for  $\text{C}_{23}\text{H}_{32}\text{BrClN}_3\text{Mn}$  ( $m/z$ ): calculated for  $[\text{C}_{23}\text{H}_{32}\text{ClN}_3\text{Mn}]^+$  ( $[\text{M} - \text{Br}]^+$ ) 440.1660; found 440.1641 ( $z = 1$ ) and calculated for  $[\text{C}_{23}\text{H}_{32}\text{BrN}_3\text{Mn}]^+$  ( $[\text{M} - \text{Cl}]^+$ ): 484.1155; found: 484.1138 ( $z = 1$ ), both peaks showing expected isotopic patterns. Evans method ( $\text{CD}_2\text{Cl}_2$ , 298 K):  $\mu_{\text{eff}} = 4.78 \mu_{\text{B}}$ . Anal. Calcd. for  $\text{Mn}_1\text{C}_{23}\text{H}_{32}\text{BrClN}_3$ : C, 53.04; H, 6.19; N, 8.07. Found: C, 53.45; H, 6.05; N, 8.07. UV-vis,  $\lambda$ , nm ( $\epsilon$ ,  $\text{M}^{-1} \text{cm}^{-1}$ ),  $\text{CH}_2\text{Cl}_2$ : 539 (854), 304 (6637), 260 (10817). FT-IR (ATR, solid,  $\text{cm}^{-1}$ ):  $\nu$  2968 (w), 2898 (w), 1601 (w), 1462 (w), 1429 (w), 1376 (s), 1188 (s), 910 (w), 849 (w), 770 (s).

**Synthesis of  $(^{\text{tBu}}\text{N}_3\text{C})\text{Mn}^{\text{III}}\text{BrCl}$  (3) by Oxidative Addition of  $^{\text{tBu}}\text{N}_3\text{CCl}$ .** 19.3 mg of  $^{\text{tBu}}\text{N}_3\text{CCl}$  (0.05 mmol) and 13.7 mg of  $\text{MnBr}(\text{CO})_5$  (0.05 mmol) were combined in a flame-dried Schlenk flask inside a glovebox, and 2 mL of THF was added to give a yellow suspension. The flask was taken outside the glovebox and stirred in a water bath in front of a mercury lamp. The reaction vessel was subjected to vacuum for 1 s every hour, then stirred under static vacuum, and after 3 h, the reaction was then stirred under a static vacuum overnight. After 13 h, the solution turned wine-red and the entire solvent was evaporated under reduced pressure. The solid obtained was redissolved in a minimal amount of dichloromethane and filtered through Celite. The filtrate obtained was evaporated under reduced pressure to yield a red solid which was washed three times with copious amounts of ether (ca. 3 mL) and then dried under vacuum. Deep red crystals were grown by vapor diffusion of ether into a dichloromethane solution of the complex (about 2 mL of DCM); yield of isolated product  $(^{\text{tBu}}\text{N}_3\text{C})\text{Mn}^{\text{III}}\text{BrCl}$  (3) 6.0 mg, 23% yield. HRMS (ESI) in MeOH for  $\text{C}_{23}\text{H}_{32}\text{BrClN}_3\text{Mn}$  ( $m/z$ ): calculated for  $[\text{C}_{23}\text{H}_{32}\text{ClN}_3\text{Mn}]^+$  ( $[\text{M} - \text{Br}]^+$ ) 440.1660; found: 440.1637 ( $z = 1$ ) and calculated for  $[\text{C}_{23}\text{H}_{32}\text{BrN}_3\text{Mn}]^+$  ( $[\text{M} - \text{Cl}]^+$ ): 484.1155; found: 484.1119 ( $z = 1$ ), both peaks showing expected isotopic patterns.

**General Procedure for C–C Coupling without Oxidant.** 0.05 mmol of a manganese complex (**1**, **2**, or **3**) was dissolved in 2 mL of THF. Then, 3.0 equiv of an organometal reagent were added and the reaction mixture was stirred at RT in a glovebox for 16 h. The vial was then taken out of the glovebox and quenched by addition of a saturated aqueous solution of ammonium chloride (0.2 mL); then a 2 mL saturated solution of  $\text{K}_2\text{CO}_3$  was added to it and vigorously stirred for 30 min. Workup using basic aqueous solutions was needed to remove all unreacted Mn residues after reaction and avoid the presence of unreacted paramagnetic species in solution that can potentially lead to broadening of NMR spectra. The mixture was extracted with ethyl acetate ( $3 \times 10$  mL). The combined organic extracts were washed with saturated brine (15 mL) and dried over  $\text{Na}_2\text{SO}_4$ . The solvent was removed by rotary evaporation. 1,3,5-

Trimethoxybenzene (0.05 mmol, 1.0 equiv) was then added to this mixture as an internal standard, and NMR yields of the products were determined in  $\text{C}_6\text{D}_6$  (see the SI). Then the above crude product was purified by flash column chromatography on a silica gel ( $\text{CH}_2\text{Cl}_2/\text{MeOH} = 20/1-15/1$ ) give the desired product.

For low temperature experiments, the solution of **1** (0.05 mmol) in THF (8 mL) was placed in a Schlenk tube, cooled down to  $-80^\circ\text{C}$  using a methanol-dry ice cooling bath. The solution of  $\text{MeMgBr}$  (3.0 M in diethyl ether) (0.05 mL, 0.150 mmol, 3 equiv) was added slowly to the reaction mixture under argon gas flow. The reaction mixture was warmed up to RT slowly over the course of 4 h, then stirred at RT for another 16 h. The workup procedure was carried out as described above. The same procedure was followed for the reactions with  $\text{MeLi}$  and  $\text{ZnMe}_2$ .

**General Procedure for C–C Coupling with Oxidant.** 0.05 mmol of a manganese complex (**1**, **2**, or **3**) was dissolved in 2 mL of THF. Then, 15.6 mg of TEMPO (2 equiv) and 3.0 equiv of an organometal reagent were added and stirred at room temperature in a glovebox for 16 h. The vial was then taken out of the glovebox and quenched by addition of a saturated aqueous solution of ammonium chloride (0.2 mL); then a 2 mL saturated solution of  $\text{K}_2\text{CO}_3$  was added to it and vigorously stirred for 30 min. The mixture was extracted with ethyl acetate ( $3 \times 10$  mL). The combined organic extracts were washed with saturated brine (15 mL) and dried over  $\text{Na}_2\text{SO}_4$ . The solvent was removed by rotary evaporation. 1,3,5-Trimethoxybenzene (0.05 mmol, 1.0 equiv) was then added to this mixture as an internal standard, and NMR yields of the products were determined in  $\text{C}_6\text{D}_6$  (see the SI). Then the above crude product was purified by flash column chromatography on a silica gel ( $\text{CH}_2\text{Cl}_2/\text{MeOH} = 20/1-15/1$ ) to give the desired product.

**Isolation and Characterization of C–C Coupling Products.**  
 **$^{\text{tBu}}\text{N}_3\text{CMe}$ .** Following the general procedure for C–C coupling with oxidant, the crude product of the reaction was collected and purified by flash column chromatography on silica gel ( $\text{CH}_2\text{Cl}_2/\text{MeOH}$ , 20:1) to provide  $^{\text{tBu}}\text{N}_3\text{CMe}$  as a white solid in 93% isolated yield. The desired product was confirmed by ESI-(HR)MS and NMR.  $^1\text{H}$  NMR (500 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  6.98–6.95 (m, 3H), 6.84 (t,  $J = 7.4$  Hz, 1H), 6.61 (d,  $J = 7.6$  Hz, 2H), 4.14 (d,  $J = 12.0$  Hz, 2H), 3.73 (d,  $J = 13.2$  Hz, 2H), 3.65 (d,  $J = 11.8$  Hz, 2H), 3.59 (d,  $J = 14.3$  Hz, 2H), 2.84 (s, 3H), 1.15 (s, 18H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  161.15, 138.55, 135.08, 130.98, 128.68, 124.74, 120.32, 56.10, 53.73, 27.73, 18.17. ESI-(HR)MS of the product from methanol solution ( $m/z$ ): calculated for  $\text{M}^+\text{H}^+$ ,  $\text{C}_{24}\text{H}_{35}\text{N}_3$ , 366.2904; found 366.2884.

**$^{\text{tBu}}\text{N}_3\text{C}i\text{Pr}$ .** 28.3 mg (0.05 mmol) of  $(^{\text{tBu}}\text{N}_3\text{C})\text{Mn}^{\text{III}}\text{Br}_2$  (**1**) was dissolved in 2 mL of THF. Then, 15.6 mg of TEMPO (2 equiv) and 75  $\mu\text{L}$  of isopropylmagnesium chloride (2.0 M in THF, 3 equiv) were added to the mixture and stirred at room temperature in a glovebox for 16 h. The mixture was quenched by addition of a saturated aqueous solution of ammonium chloride (0.2 mL); then a 2 mL saturated solution of  $\text{K}_2\text{CO}_3$  was added to it and vigorously stirred for 30 min. The mixture was extracted with ethyl acetate ( $3 \times 10$  mL). The combined organic extracts were washed with saturated brine (15 mL) and dried over  $\text{Na}_2\text{SO}_4$ . The ethyl acetate was completely evaporated by a rotavapor, and the solid left was isolated by a short flash column chromatography on silica gel with  $\text{DCM}/\text{MeOH}$  (15/1) to provide  $^{\text{tBu}}\text{N}_3\text{C}i\text{Pr}$  (9.2 mg, 47%).  $^1\text{H}$  NMR (500 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  7.00–6.97 (m, 3H), 6.66–6.63 (m, 3H), 4.13–4.09 (m, 2H), 3.92–3.88 (m, 4H), 3.77–3.71 (m, 2H), 3.59–3.55 (m, 1H), 1.83 (s, 6H), 1.19 (s, 18H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  160.99, 138.61, 134.65, 131.85, 125.12, 120.54, 57.72, 56.01, 31.26, 28.00, 24.55, 23.39. ESI-(HR)MS of the product from methanol solution ( $m/z$ ): calculated for  $[\text{M} + \text{H}]^+$ ,  $\text{C}_{26}\text{H}_{39}\text{N}_3$ , 394.3217; found  $m/z$  394.3215.

**$^{\text{tBu}}\text{N}_3\text{CC}i\text{H}_5$ .** 28.3 mg (0.05 mmol) of  $(^{\text{tBu}}\text{N}_3\text{C})\text{Mn}^{\text{III}}\text{Br}_2$  (**1**) was dissolved in 2 mL of THF. Then, 15.6 mg of TEMPO (2 equiv) and 150  $\mu\text{L}$  of allylmagnesium bromide (1 M in diethyl ether, 3 equiv) were added to the mixture and stirred at room temperature for 16 h. The vial was then taken out of the glovebox and quenched by addition of a saturated aqueous solution of ammonium chloride (0.2 mL); then a 2 mL saturated solution of  $\text{K}_2\text{CO}_3$  was added to it and vigorously stirred for 30 min. The mixture was extracted with EtOAc ( $3 \times 10$

mL). The combined organic extracts were washed with brine (15 mL) and dried over  $\text{Na}_2\text{SO}_4$ . The solvent was removed by rotary evaporation, and a 72% yield of product  $^{18}\text{BuN3CC}_3\text{H}_5$  was obtained and confirmed by the NMR spectrum. This product was not stable and decomposed after isolation from column chromatography, but it was stable in solution for a few days.  $^1\text{H}$  NMR (400 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  7.01–6.97 (m, 3H), 6.92–6.88 (m, 1H), 6.60 (d,  $J = 7.6$  Hz, 2H), 6.20–6.11 (m, 1H), 5.06–5.05 (m, 2H), 4.58 (d,  $J = 5.5$  Hz, 2H), 4.22 (d,  $J = 12.0$  Hz, 2H), 3.71 (d,  $J = 14.5$  Hz, 2H), 3.63–3.59 (m, 4H), 1.14 (s, 18H).  $^{13}\text{C}$  NMR (101 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  161.15, 140.06, 138.32, 135.18, 131.80, 125.42, 120.34, 114.28, 56.09, 55.70, 53.66, 33.98, 27.65. ESI-(HR)MS of the product from methanol solution ( $m/z$ ): calculated for  $[\text{M} + \text{H}]^+$ ,  $\text{C}_{26}\text{H}_{37}\text{N}_3$ ,  $m/z$  392.3060; found 392.3058.

$^{18}\text{BuN3CPh}$ . 28.3 mg (0.05 mmol) of  $(^{18}\text{BuN3C})\text{Mn}^{\text{III}}\text{Br}_2$  (**1**) was dissolved in 2 mL of tetrahydrofuran. Then, 15.6 mg of TEMPO (2 equiv) and 50  $\mu\text{L}$  of phenylmagnesium bromide (3 M in diethyl ether, 3 equiv) were added and stirred at room temperature in a glovebox for 16 h. The vial was then taken out of the glovebox and quenched by addition of a saturated aqueous solution of ammonium chloride (0.2 mL); then a 2 mL saturated solution of  $\text{K}_2\text{CO}_3$  was added to it and vigorously stirred for 30 min. The mixture was extracted with ethyl acetate ( $3 \times 10$  mL). The combined organic extracts were washed with saturated brine (15 mL) and dried over  $\text{Na}_2\text{SO}_4$ . The solvent was removed by rotary evaporation, and a 71% yield was shown from crude NMR using 1,3,5-trimethoxybenzene as an internal standard. The crude product was purified by flash column chromatography on silica gel (DCM/MeOH, 20:1) to provide  $^{18}\text{BuN3CC}_6\text{H}_5$  (12.8 mg, 60%) as a white solid.  $^1\text{H}$  NMR (500 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  9.81 (d,  $J = 7.1$  Hz, 1H), 7.57 (t,  $J = 7.5$  Hz, 1H), 7.25–7.22 (m, 2H), 7.01–6.96 (m, 4H), 6.81 (t,  $J = 7.5$  Hz, 1H), 6.66 (s, 2H), 3.85 (s, 4H), 3.76 (s, 4H), 1.00 (s, 18H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  161.54, 142.04, 137.93, 134.84, 132.04, 130.97, 128.89, 127.57, 127.22, 126.64, 120.30, 56.75, 55.92, 53.29, 27.72. ESI-(HR)MS of the product from methanol solution ( $m/z$ ): calculated for  $[\text{M} + \text{H}]^+$ ,  $\text{C}_{29}\text{H}_{38}\text{N}_3$ ,  $m/z$  428.3060; found 428.3049.

$^{18}\text{BuN3CC}_6\text{F}_5$ . 28.3 mg (0.05 mmol) of  $(^{18}\text{BuN3C})\text{Mn}^{\text{III}}\text{Br}_2$  (**1**) was weighed out in a scintillation vial inside a glovebox, and 2 mL of benzene was added and stirred for 20 min at RT. To the red solution was added 600  $\mu\text{L}$  (0.3 mmol) of pentafluorophenylmagnesium bromide solution (0.5 M in diethyl ether, 6 equiv) in one portion, and the mixture was allowed to stir for 40 h at 80  $^\circ\text{C}$  over which time period the color of the solution gradually changed to brown and the reaction was stopped. The vial was then taken out of the glovebox, and then 2 mL of a saturated aqueous solution of  $\text{K}_2\text{CO}_3$  was added to it and vigorously stirred for 30 min. The aqueous solution was extracted with ethyl acetate, filtered, and then dried over anhydrous  $\text{Na}_2\text{SO}_4$ . The ethyl acetate was completely evaporated by a rotavapor, and the solid left was isolated by a short flash column chromatography on silica gel with hexane/ethyl acetate (2/1–1.5/1) to provide  $^{18}\text{BuN3CC}_6\text{F}_5$  (12.6 mg, 49%).  $^1\text{H}$  NMR (400 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  7.10–7.04 (m, 3H, ArH), 6.95 (t,  $J = 7.6$  Hz, 1H, ArH), 6.49 (d,  $J = 7.6$  Hz, 2H, ArH), 3.70–3.65 (m, 2H,  $\text{CH}_2$ ), 3.54–3.46 (m, 4H,  $\text{CH}_2$ ), 3.27–3.21 (m, 2H,  $\text{CH}_2$ ), 0.85 (s, 18H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  161.06, 146.14, 141.11, 135.29, 131.90, 128.68, 119.95, 56.21, 54.70, 53.52, 27.09.  $^{19}\text{F}$  NMR (376 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  –106.51, –117.34, –138.00 (dt,  $J = 25.4$ , 6.1 Hz), –158.34, –164.83 (d,  $J = 111.4$  Hz).  $^{19}\text{F}$  NMR (376 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  –106.5, –117.3, –138.0 (dt,  $J = 25.4$ , 6.1 Hz), –158.3, –164.8 (d,  $J = 111.4$  Hz). ESI-(HR)MS of the product from methanol solution ( $m/z$ ): calculated for  $[\text{M} + \text{H}]^+$ ,  $\text{C}_{29}\text{H}_{33}\text{F}_5\text{N}_3$ ,  $m/z$  518.2589; found 518.2601.

## ■ ASSOCIATED CONTENT

### SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.organomet.1c00047>.

Experimental details and characterization data (PDF)

## Accession Codes

CCDC 2056056, 2056057, and 2056899 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

## ■ AUTHOR INFORMATION

### Corresponding Author

Julia R. Khusnutdinova – *Coordination Chemistry and Catalysis Unit, Okinawa Institute of Science and Technology Graduate University, Onna-son, Okinawa 904-0495, Japan*;  
[orcid.org/0000-0002-5911-4382](https://orcid.org/0000-0002-5911-4382); Email: [juliak@oist.jp](mailto:juliak@oist.jp)

### Authors

Yu-Tao He – *Coordination Chemistry and Catalysis Unit, Okinawa Institute of Science and Technology Graduate University, Onna-son, Okinawa 904-0495, Japan*  
 Ayumu Karimata – *Coordination Chemistry and Catalysis Unit, Okinawa Institute of Science and Technology Graduate University, Onna-son, Okinawa 904-0495, Japan*  
 Olga Gladkovskaya – *Coordination Chemistry and Catalysis Unit, Okinawa Institute of Science and Technology Graduate University, Onna-son, Okinawa 904-0495, Japan*  
 Eugene Khaskin – *Coordination Chemistry and Catalysis Unit, Okinawa Institute of Science and Technology Graduate University, Onna-son, Okinawa 904-0495, Japan*;  
[orcid.org/0000-0003-1790-704X](https://orcid.org/0000-0003-1790-704X)  
 Robert R. Fayzullin – *Arbuzov Institute of Organic and Physical Chemistry, FRC Kazan Scientific Center, Russian Academy of Sciences, Kazan 420088, Russian Federation*;  
[orcid.org/0000-0002-3740-9833](https://orcid.org/0000-0002-3740-9833)  
 Abir Sarbajna – *Coordination Chemistry and Catalysis Unit, Okinawa Institute of Science and Technology Graduate University, Onna-son, Okinawa 904-0495, Japan*;  
[orcid.org/0000-0003-2478-5044](https://orcid.org/0000-0003-2478-5044)

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.organomet.1c00047>

## Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This work was supported by JSPS Kakenhi Grant 20K15305. We thank the Instrumental Analysis Section (Dr. Michael Roy and Dr. Yoshiteru Iinuma) and Engineering Support Section (Dr. Hyung Been Kang) for technical support. The authors acknowledge the Okinawa Institute of Science and Technology Graduate University for start-up funding.

## ■ REFERENCES

- (1) Beller, M.; Varela-Fernandez, A.; Vries, J. G.; Man Wong, S.; Ming So, C.; Kwong, F. Y. *Cross Coupling Reactions. Appl. Homogeneous Catal. Organomet. Compd. (3rd Ed.)* **2017**, *2*, 411–464.
- (2) de Meijere, A.; Diederich, F. *Metal-Catalyzed Cross-Coupling Reactions*; Wiley-VCH, 2004.
- (3) Negishi, E.; de Meijere, A. *Handbook of Organopalladium Chemistry for Organic Synthesis*; Wiley, 2003.
- (4) Johansson Seechurn, C. C. C.; Kitching, M. O.; Colacot, T. J.; Snieckus, V. *Palladium-Catalyzed Cross-Coupling: A Historical Contextual Perspective to the 2010 Nobel Prize. Angew. Chem., Int. Ed.* **2012**, *51*, 5062–5085.

- (5) Carney, J. R.; Dillon, B. R.; Thomas, S. P. Recent Advances of Manganese Catalysis for Organic Synthesis. *Eur. J. Org. Chem.* **2016**, *2016*, 3912–3929.
- (6) Bauer, I.; Knölker, H.-J. Iron Catalysis in Organic Synthesis. *Chem. Rev.* **2015**, *115*, 3170–3387.
- (7) Bolm, C.; Legros, J.; Le Pailh, J.; Zani, L. Iron-Catalyzed Reactions in Organic Synthesis. *Chem. Rev.* **2004**, *104*, 6217–6254.
- (8) Liu, W.; Ackermann, L. Manganese-Catalyzed C–H Activation. *ACS Catal.* **2016**, *6*, 3743–3752.
- (9) Hu, Y.; Zhou, B.; Wang, C. Inert C–H Bond Transformations Enabled by Organometallic Manganese Catalysis. *Acc. Chem. Res.* **2018**, *51*, 816–827.
- (10) Liu, W.; Groves, J. T. Manganese Catalyzed C–H Halogenation. *Acc. Chem. Res.* **2015**, *48*, 1727–1735.
- (11) Bedford, R. B. How Low Does Iron Go? Chasing the Active Species in Fe-Catalyzed Cross-Coupling Reactions. *Acc. Chem. Res.* **2015**, *48*, 1485–1493.
- (12) Valyaev, D. A.; Lavigne, G.; Lugan, N. Manganese organometallic compounds in homogeneous catalysis: Past, present, and prospects. *Coord. Chem. Rev.* **2016**, *308*, 191–235.
- (13) Bottoni, A.; Cahiez, G.; Calvaresi, M.; Moyeux, A.; Giacinto, P.; Miscione, G. P. A mechanistic insights into manganese-catalyzed oxidative homocoupling reactions of Grignard reagents: A computational DFT investigation. *J. Organomet. Chem.* **2016**, *814*, 25–34.
- (14) Cahiez, G.; Duplais, C.; Buendia, J. Manganese-Catalyzed Oxidative Cross-Coupling of Grignard Reagents with Oxygen as an Oxidant. *Angew. Chem., Int. Ed.* **2009**, *48*, 6731–6734.
- (15) Cahiez, G.; Duplais, C.; Buendia, J. Chemistry of Organomanganese(II) Compounds. *Chem. Rev.* **2009**, *109*, 1434–1476.
- (16) Cahiez, G.; Gager, O.; Lecomte, F. Manganese-Catalyzed Cross-Coupling Reaction between Aryl Grignard Reagents and Alkenyl Halides. *Org. Lett.* **2008**, *10*, 5255–5256.
- (17) Cahiez, G.; Moyeux, A.; Buendia, J.; Duplais, C. Manganese- or Iron-Catalyzed Homocoupling of Grignard Reagents Using Atmospheric Oxygen as an Oxidant. *J. Am. Chem. Soc.* **2007**, *129*, 13788–13789.
- (18) Cahiez, G.; Luart, D.; Lecomte, F. Unexpected Cross-Coupling Reaction between o-Chloroaryl Ketones and Organomanganese Reagents. *Org. Lett.* **2004**, *6*, 4395–4398.
- (19) Cahiez, G.; Lepifre, F.; Ramiandrasoa, P. Manganese-catalyzed substitution of activated aryl halides (X = Cl, Br, and F) and aryl ethers by organomagnesium reagents. *Synthesis* **1999**, *1999*, 2138–2144.
- (20) Alami, M.; Ramiandrasoa, P.; Cahiez, G. Stereoselective Manganese-Catalyzed Coupling Reaction of Conjugated Chloroenynes and Chlorodienes with Grignard Reagents. *Synlett* **1998**, *09*, 325–327.
- (21) Santilli, C.; Beigbaghlou, S. S.; Ahlburg, A.; Antonacci, G.; Fristrup, P.; Norrby, P.-O.; Madsen, R. The Manganese-Catalyzed Cross-Coupling Reaction and the Influence of Trace Metals. *Eur. J. Org. Chem.* **2017**, *2017*, 5269–5274.
- (22) Antonacci, G.; Ahlburg, A.; Fristrup, P.; Norrby, P.-O.; Madsen, R. Manganese-Catalyzed Cross-Coupling of Aryl Halides and Grignard Reagents by a Radical Mechanism. *Eur. J. Org. Chem.* **2017**, *2017*, 4758–4764.
- (23) Rueping, M.; Ieawsuwan, W. A manganese-catalyzed cross-coupling reaction. *Synlett* **2007**, *2007*, 247–250.
- (24) Dakarapu, R.; Falck, J. R. Stereospecific Stille Cross-Couplings Using Mn(II)Cl<sub>2</sub>. *J. Org. Chem.* **2018**, *83*, 1241–1251.
- (25) Kang, S.-K.; Kim, J.-S.; Choi, S.-C. Copper- and Manganese-Catalyzed Cross-Coupling of Organostannanes with Organic Iodides in the Presence of Sodium Chloride. *J. Org. Chem.* **1997**, *62*, 4208–4209.
- (26) Cahiez, G.; Bernard, D.; Normant, J. F. Catalysis by manganese(II). I. Coupling reactions of iodoalkenes to dienes initiated by organolithium reagents in the presence of manganese(II). *J. Organomet. Chem.* **1976**, *113*, 99–106.
- (27) Sato, T.; Yoshida, T.; Al Mamari, H. H.; Ilies, L.; Nakamura, E. Manganese-Catalyzed Directed Methylation of C(sp<sup>2</sup>)-H Bonds at 25 °C with High Catalytic Turnover. *Org. Lett.* **2017**, *19*, 5458–5461.
- (28) Zhu, C.; Oliveira, J. C. A.; Shen, Z.; Huang, H.; Ackermann, L. Manganese(II/III/I)-Catalyzed C–H Arylations in Continuous Flow. *ACS Catal.* **2018**, *8*, 4402–4407.
- (29) Liu, W.; Cera, G.; Oliveira, J. C. A.; Shen, Z.; Ackermann, L. MnCl<sub>2</sub>-Catalyzed C–H Alkylations with Alkyl Halides. *Chem. - Eur. J.* **2017**, *23*, 11524–11528.
- (30) Uzelac, M.; Mastropierro, P.; de Tullio, M.; Borilovic, I.; Tarrés, M.; Kennedy, A. R.; Aromí, G.; Hevia, E. Tandem Mn–I Exchange and Homocoupling Processes Mediated by a Synergistically Operative Lithium Manganate. *Angew. Chem., Int. Ed.* **2021**, *60*, 3247–3253.
- (31) Liu, Y.; Berges, J.; Zaid, Y.; Chahdi, F. O.; Van Der Lee, A.; Harakat, D.; Clot, E.; Jaroschik, F.; Taillefer, M. Aerobic and Ligand-Free Manganese-Catalyzed Homocoupling of Arenes or Aryl Halides via in Situ Formation of Aryllithiums. *J. Org. Chem.* **2019**, *84*, 4413–4420.
- (32) Sarbajna, A.; He, Y.-T.; Dinh, M. H.; Gladkovskaya, O.; Rahaman, S. M. W.; Karimata, A.; Khaskin, E.; Lapointe, S.; Fayzullin, R. R.; Khusnutdinova, J. R. Aryl–X Bond-Forming Reductive Elimination from High-Valent Mn–Aryl Complexes. *Organometallics* **2019**, *38*, 4409–4419.
- (33) Smith, M. B. *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*; Wiley, 2020.
- (34) Murarka, S.; Möbus, J.; Erker, G.; Mück-Lichtenfeld, C.; Studer, A. TEMPO-mediated homocoupling of aryl Grignard reagents: mechanistic studies. *Org. Biomol. Chem.* **2015**, *13*, 2762–2767.
- (35) Whitesides, G. M.; Newirth, T. L. Reaction of n-butyllithium and 2,2,6,6-tetramethylpiperidine nitroxyl. *J. Org. Chem.* **1975**, *40*, 3448–3450.
- (36) Nagashima, T.; Curran, D. P. Reactions of Tempo with Alkylsamarium and Other Organometallic Reagents. *Synlett* **1996**, *1996*, 330–332.
- (37) Budny-Godlewski, K.; Kubicki, D.; Justyniak, I.; Lewiński, J. A New Look at the Reactivity of TEMPO toward Diethylzinc. *Organometallics* **2014**, *33*, 5093–5096.
- (38) Espenson, J. H. Chemistry of organochromium(III) complexes. *Acc. Chem. Res.* **1992**, *25*, 222–227.
- (39) Kochi, J. K. Electron-transfer mechanisms for organometallic intermediates in catalytic reactions. *Acc. Chem. Res.* **1974**, *7*, 351–360.
- (40) Khusnutdinova, J. R.; Rath, N. P.; Mirica, L. M. Stable Mononuclear Organometallic Pd(III) Complexes and Their C–C Bond Formation Reactivity. *J. Am. Chem. Soc.* **2010**, *132*, 7303–7305.
- (41) Meneghetti, S. P.; Lutz, P. J.; Kress, J. Neutral and Cationic Palladium(II) Complexes of a Diazapyridinophane. Structure, Fluxionality, and Reactivity toward Ethylene. *Organometallics* **2001**, *20*, 5050–5055.
- (42) Khusnutdinova, J. R.; Rath, N. P.; Mirica, L. M. The Conformational Flexibility of the Tetradentate Ligand tBuN4 is Essential for the Stabilization of (tBuN4)Pd<sup>III</sup> Complexes. *Inorg. Chem.* **2014**, *53*, 13112–13129.
- (43) Patil, P. H.; Filonenko, G. A.; Lapointe, S.; Fayzullin, R. R.; Khusnutdinova, J. R. Interplay between the Conformational Flexibility and Photoluminescent Properties of Mononuclear Pyridinophane-copper(I) Complexes. *Inorg. Chem.* **2018**, *57*, 10009–10027.
- (44) Zhou, W.; Schultz, J. W.; Rath, N. P.; Mirica, L. M. Aromatic Methoxylation and Hydroxylation by Organometallic High-Valent Nickel Complexes. *J. Am. Chem. Soc.* **2015**, *137*, 7604–7607.
- (45) Casitas, A.; Canta, M.; Solà, M.; Costas, M.; Ribas, X. Nucleophilic Aryl Fluorination and Aryl Halide Exchange Mediated by a Cu<sup>I</sup>/Cu<sup>III</sup> Catalytic Cycle. *J. Am. Chem. Soc.* **2011**, *133*, 19386–19392.
- (46) Gemel, C.; Folting, K.; Caulton, K. G. New Approach to Ru(II) Pincer Ligand Chemistry. Bis(tert-butylaminomethyl)pyridine Coordinated to Ruthenium(II). *Inorg. Chem.* **2000**, *39*, 1593–1597.

- (47) Motterlini, R. A.; Mann, B. E.; Scapens, D. A. WO 2008003953 A3, 2008.
- (48) Kadassery, K. J.; Lacy, D. C. Pentacarbonylmethylmanganese(I) as a synthon for Mn(I) pincer catalysts. *Dalton Trans.* **2019**, 48, 4467–4470.
- (49) McQuarrie, D. A.; Cox, H.; Simon, J. D.; Choi, J. *Physical Chemistry: A Molecular Approach*; University Science Books, 1997.
- (50) Sheldrick, G. M. SHELXT - Integrated space-group and crystal-structure determination. *Acta Crystallogr., Sect. A: Found. Adv.* **2015**, 71, 3–8.
- (51) Sheldrick, G. M. Crystal structure refinement with SHELXL. *Acta Crystallogr., Sect. C: Struct. Chem.* **2015**, 71, 3–8.