

Article

Triflic acid-promoted adamantylation and tert-butylation of pyrene: fluorescent properties of pyrene-decorated adamantanes and a channeled crystal structure of 1,3,5-tris(pyren-2-yl)adamantane.

Anna Wrona-Piotrowicz, Anna Makal, and Janusz Zakrzewski

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.0c01060 • Publication Date (Web): 07 Aug 2020

Downloaded from pubs.acs.org on August 7, 2020

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.


Triflic acid-promoted adamantylation and tert-butylation of pyrene: fluorescent properties of pyrene-decorated adamantanes and a channeled crystal structure of 1,3,5-tris(pyren-2-yl)adamantane.

Anna Wrona-Piotrowicz,^{*,†} Anna Makal[‡] and Janusz Zakrzewski[†]

[†]Department of Organic Chemistry, Faculty of Chemistry, University of Lodz, Tamka 12, 91-403 Łódź, Poland

[‡]Biological and Chemical Research Center, Faculty of Chemistry, University of Warsaw, Żwirki i Wigury 101, 02-089 Warszawa, Poland

Supporting Information Placeholder



ABSTRACT: Triflic acid-promoted 1-adamantylation and *tert*-butylation of pyrene at positions 2 and 7 along with the synthesis of compounds having one-, two- and three pyrenyl groups attached to the adamantane scaffold is disclosed. Fluorescent properties of these compounds and channeled crystal structure of the 1,3,5-tris(pyren-2-yl)adamantane containing chloroform as a guest are also presented.

INTRODUCTION

Pyrene **1** (Figure 1) is a fascinating fluorophore¹ and one of the most promising starting materials for preparing a plethora of organic optoelectronic materials.² Two alkyl-substituted pyrene derivatives, 2-*tert*-butylpyrene **2** and 2,7-di-*tert*-butylpyrene **3** have been widely used as useful starting materials in numerous syntheses of dyes that feature special molecular shapes and suppressed aggregation properties.³ Moreover, the latter compound proved to be an efficient acceptor for triplet-triplet annihilation⁴ and a π -donor for semiconductive charge transfer cocrystals.⁵ To date, both compounds could be obtained via Friedel-Crafts alkylation of pyrene with *tert*-butyl chloride in the presence of a Lewis acid (aluminum chloride or bromide). This reaction is unusual because electrophiles usually attack pyrene at positions 1, 3, 6 or 8.^{2a} Moreover, there

have been no reports on the extension of its scope to other alkyl halides.





Figure 1. Formulae of compounds 1-3.

RESULTS AND DISCUSSION

In continuation of our earlier work on triflic acid (TfOH)-promoted Friedel-Crafts pyrene chemistry,⁶ we became interested in elaborating a synthetic route to pyrenes bearing other bulky tertiary alkyl groups at C(2) and C(7) using this superacid as a promoter. The adamantyl group seemed especially interesting because physicochemical properties of adamantyl-substituted compounds may be different from those of their *tert*-butyl-substituted counterparts.⁷ Moreover, in contrast to the *tert*-butyl group, the rigid adamantane scaffold offers a possibility of attachment, via the Friedel-Crafts chemistry, of two, three or four aryl group to the tertiary carbons. It has been demonstrated that di-, tri- and tetraaryl adamantanes have a great tendency to form various types of supramolecular

Table 1. Optimized conditions for the adamantylation of pyrene.



Entry	AdaX	Pyrene: AdaX: TfOH (mmole)	Reaction time	Product (Isolated yield, %) ^a
1	1-AdaBr	1 : 1 : 4	5 min	4 (93)
2	1-AdaOH	1 : 1 : 4	5 min	4 (84)
3	1-AdaBr	1 : 2.2 : 4	30 min	5 (91)
4	1-AdaCl	1 : 2.2 : 4	30 min	5 (87)
5	1-AdaOH	1 : 2.2 : 4	1 h	5 (95)
6	2-AdaBr	1 : 2.2 : 4	30 min	5 (83)
7	2-AdaOH	1 : 2.2 : 4	1 h	5 (87)
8	1,3-AdaBr ₂	3 : 1 : 12	4 h	6 (62)
9	1,3-Ada(OH) ₂	3 : 1 : 12	4 h	6 (49)
10	1,3,5-Ada(OH) ₃	4.5 : 1 : 18	6 h	7 (63)

^aReactions were carried out in dichloromethane (10 mL) at r.t.; Ada = adamantyl.

We found that pyrene **1** smoothly reacts with 1-adamantanol or 1-bromoadamantane in the presence of excess of TfOH in dichloromethane at room temperature to afford, depending on the reaction conditions compound **4** or **5**. To obtain pure **4**, not contaminated with **5**, pyrene and adamantyl compound were used at a molar ratio 1:1 and the reaction was stopped after 5 min (Entry 1, 2). On the other hand, at the pyrene to

systems in the solid state (e.g. encapsulating crystals⁸ and were used in syntheses of porous aromatic frameworks (PAFs)).⁹ The adamantane skeleton was also used as a rigid linker between chromophoric groups.¹⁰ Therefore, molecules bearing several pyrenyl groups in a well-defined spatial arrangement might be of interest for developing solid-state emitters, sensors, or self-assembled nanomaterials.

In this study, we report the results of this research along with the basic fluorescent properties of the synthesized pyrene derivatives **4-7** (Table 1) and the crystal structure of tripyrenyladamantane **7** encapsulating molecule of the solvent (chloroform).

adamantyl compound molar ratio of 1:2.2 and prolonged reaction times (30min - 1h) diadamantylpyrene **5** was formed in high yields (Entries 3-7). It is also worthy to note that compound **4** can also be obtained in practically the same yield using 2-bromoadamantane or 2-adamantanol as alkylating agents (Table 1, Entries 6-7). This suggests the intermediacy

of 2-adamantyl cation, which is known to rapidly isomerize to its 1-isomer.^{11, 12}

Having in hands an efficient protocol for the synthesis of **4** and **5** we became interested in its application for the synthesis of compounds having two and three pyrenyl units linked to the adamantane skeleton. To our pleasure, we found that the reaction of 3 equiv. of pyrene with 1,3-dibromoadamantane or adamantane-1,3-diol in the presence of TfOH yielded, after 4 h, 1,3-bis(pyren-2-yl)adamantane **6** in 62% and 49% yield, respectively (Entries 8, 9). Furthermore, the reaction of 4.5 equiv. of pyrene with adamantane-1,3,5-triol gave after 6 h, 1,3,5-tris(pyren-2-yl)adamantane (**7**) in 63% yield (Table 1, Entry 10).

We also found out that our protocol can be used for alternative synthesis of *tert*-butylpyrenes **2** and **3** (see the Supporting Information for details). Both *tert*-butyl halides and *tert*-butanol afford the expected products in high yields (86-96%). In our opinion, this procedure is more convenient than the procedures described in literature, which use aluminum halides as promoter.

Formation of the 2- and 2,7- substituted products in the reaction of pyrene with *tert*-butyl halides is usually explained as due to bulkiness of the *tert*-butyl group (these positions are relatively less sterically hindered)^{2a}. In our opinion the same explanation may be valid for reactions with adamantyl compounds.

We studied electronic absorption and emission spectra of compounds **2-7**. The spectra obtained for diluted solutions in dichloromethane are shown in Figure 2 (Table contains the photophysical data in Supporting Information).




Figure 2. Electronic absorption (a) and emission (b) spectra of compounds **2-7** in argon-saturated dichloromethane solutions. Absorption spectra were run at $c = 5 \times 10^{-6}$ M. Emission spectra were obtained with an excitation at 340 nm for solution having the same absorbance (0.10) at this wavelength.

All investigated compounds showed absorption and emission bands assignable to the isolated, monomeric pyrene fluorophore. There was no evidence for the formation of intramolecular excimers by **6** and **7** (negligible emission at ~450 nm). This means that the rigid adamantane skeleton efficiently keeps pyrenyl moieties electronically isolated from each other. However, there was a significant difference in the solid-state emission of compounds **3** and **5** (normalized emission spectra in the solid state in Supporting Information). While the former compound showed the structured emission characteristic of isolated pyrene fluorophore, the latter displayed excimer emission ($\lambda_{\text{max}} \sim 450$ nm). Therefore the 1-adamantyl group, unlike the *tert*-butyl group, does not protect 2,7-disubstituted pyrene against the aggregation.

Unexpectedly, we found that solution of **7** in CDCl_3 in a NMR tube slowly deposited crystals in a form of large, yellowish elongated blocs. They tended to lose transparency and transform into a powder within seconds after extracting from mother solution. Nevertheless, we succeeded in X-ray diffraction structure determination of a single crystal suspended in a drop of the solvent. The molecular structure of **7** is shown in Figure 3. The three pyrene units are differently inclined with respect to the idealized plane containing the main adamantane axis and C10, H10 atoms. The inclination angles are $67.4(5)^\circ$, $15.6(6)^\circ$ and $21.9(5)^\circ$ for pyrene moieties I, II and III respectively. They also show some deviation from planarity and are to some extent bent roughly along the short pyrene axis. The angles between the average planes of the terminal rings within the pyrene fragments are $4.2(3)^\circ$, $2.9(3)^\circ$ and $6.9(3)^\circ$ for pyrene I, II and III, respectively.




Figure 3. Molecular structure of **7**. Atomic displacement parameters represented at 50% probability level. C atoms numbered 11 – 26 constitute pyrene I, C atoms 31 – 46 constitute pyrene II and C atoms 51 – 66 constitute pyrene III moieties.

The crystal contains infinite channels (~7.5% of the unit cell volume) spreading in [100] direction filled with highly disordered chloroform molecules (Figure 4).




Figure 4. Crystal packing of **7** viewed in [100] direction. Pyrene I, II and III are highlighted in red, blue and green. The disordered molecules of chloroform occupy channels highlighted in yellow.

Analysis of intermolecular interactions in the crystal of **7** shows that pyrene II and III moieties are involved in a partial π - π stacking with their inversion-related counterparts (inter-ring distances are equal to 3.402(4) Å for pyrenes II and 3.489(5) Å for pyrenes III). On the other hand pyrene I, is involved in a C10 \cdots H10 \cdots π stacking interaction with the adamantane core of the inversion-related molecule. The distance between the planes of the symmetry-related pyrenes in that case is as long as 5.318(4) Å (Figure 5).




Figure 5. The most important intermolecular interactions formed by **7** in the crystal structure viewed along [100]. Interatomic distances shorter than the sum of the Van der Waals radii presented as red dashed lines.

The transformation of the crystals of **7** into a powder is caused by the loss of the chloroform guest, which apparently stabilize the crystal structure. Therefore, this compound is able to bind chloroform in its crystal structure in chloroform solution and

release it when chloroform is absent in the surrounding medium. Such a behavior promises possible applications of **7** and its derivatives as crystal containers for various substances. In addition, it is worthy to note that all of the synthesized pyrenyladamantanes are expected to be easily functionalized by electrophilic substitution, making possible a tuning of their photophysical and supramolecular properties.

In conclusion, we have elaborated simple and efficient synthetic route for introduction of *tert*-butyl and 1-adamantyl groups into positions 2- and 2,7- of pyrene. It allowed synthesis of compounds having two- and three pyrenyl groups attached to the adamantane scaffold. They show a solvent-dependent pyrene-like fluorescence which may be of interest for various applications. Furthermore, a synthesized tripyrenyladamantane is able to reversibly bind and release chloroform in the solid state. Future work will consist of derivatization of the synthesized pyrenyladamantanes and testing their ability to bind various guests in the solid state.

EXPERIMENTAL SECTION

General Remarks. Solvents were purified prior to use by reported methods. All reagents were purchased from Sigma-Aldrich and used without further purification. The reaction progress was monitored by means of thin-layer chromatography (TLC) which was performed on aluminum foil plates covered with silica gel 60 F₂₅₄. Column chromatography was performed on silica gel 60 (0.040–0.063 mm, 230–400 mesh, Fluka). ¹H and ¹³C NMR spectra were recorded in CDCl₃ on a Bruker ARX 600 MHz (600 MHz for ¹H and 151 MHz for ¹³C). Chemical shifts were referenced relative to solvent signals: δ = 7.26 ppm for ¹H and δ = 77.00 ppm for ¹³C. Spectra were recorded at room temperature (291 K), chemical shifts are presented in ppm, and coupling constants are presented in Hz. Electronic absorption spectra were run on a Perkin Elmer Lambda 45 UV/Vis spectrometer. Corrected emission spectra were obtained on a Perkin Elmer LS55 Fluorescence spectrometer. The emission quantum yields were determined for argon-purged solutions using quinine sulfate in 0.5 M sulfuric acid (Φ_F = 0.546) as a reference.

General procedure for synthesis of 2-7. Alkylating agent RX (for amounts of reactants, see Table 1 (S1) or S2) and TfOH were added to a solution of pyrene in 10 mL CH₂Cl₂ at room temperature. After stirring for 5 min to 6 h (Table 1 (S1), S2), the reaction mixture was poured into water (100 mL) and extracted several times with dichloromethane. The combined extracts were dried over anhydrous Na₂SO₄ and evaporated to dryness. The crude products were purified by column chromatography on silica gel (hexane or petroleum ether/ ethyl acetate as eluent) to give the desired products.

2-*tert*-butylpyrene (2). Purification by chromatography eluting with petroleum ether/ethyl acetate (95/5, v/v) gave **2** (248 mg, 96% (Table S2, Entry 2)) as a white powder. mp 110–111 °C (lit.¹³ mp 110–112 °C). ¹H NMR (600 MHz, CDCl₃) δ 8.27 (s, 2H), 8.18 (d, J = 7.8 Hz, 2H), 8.10 (d, J = 8.7 Hz, 2H), 8.08 (d, J = 8.7 Hz, 2H), 8.00 (t, J = 7.8 Hz, 1H), 1.64 (s, 9H); ¹³C{¹H} NMR (151 MHz, CDCl₃) δ 148.9, 130.9, 130.9, 127.5, 127.2, 125.5, 124.7, 124.6, 122.9, 122.1, 35.2, 31.9. MS (EI): m/z : 259 [M]⁺, 281 [M + Na]⁺; Anal. calcd. for C₂₀H₁₈: C, 92.98; H, 7.02; found: C, 92.77; H, 6.91.

2,7-di-*tert*-butylpyrene (3). Purification by chromatography eluting with petroleum ether/ethyl acetate (95/5, v/v) gave **3** (289 mg, 92% (Table S2, Entry 5)) as a white powder. mp 210–212 °C (lit.¹⁴ mp 210–212 °C). ¹H NMR (600 MHz, CDCl₃) δ 8.19 (s, 4H), 8.03 (s, 4H), 1.59 (s, 18H); ¹³C{¹H} NMR (151 MHz, CDCl₃) δ 148.5, 130.8, 127.4, 122.8, 122.0, 35.2, 31.9; MS (EI): *m/z*: 315 [M]⁺, 337 [M + Na]⁺; Anal. calcd. for C₂₄H₂₆: C, 91.67; H, 8.33; found: C, 91.76; H, 8.20.

2-(adamant-1-yl)pyrene (4). Purification by chromatography eluting with hexane/ethyl acetate (95/5, v/v) gave **4** (312 mg, 93% (Table 1, Entry 1)) as a white powder. mp 218–219 °C. ¹H NMR (600 MHz, CDCl₃) δ 8.21 (s, 2H), 8.14 (d, *J* = 7.2 Hz, 2H), 8.07 (d, *J* = 9.0 Hz, 2H), 8.04 (d, *J* = 9.0 Hz, 2H), 7.96 (t, *J* = 7.2 Hz, 1H), 2.21 (s, 9H), 1.89 (s, 6H); ¹³C{¹H} NMR (151 MHz, CDCl₃) δ 149.2, 131.0, 131.0, 127.6, 127.1, 125.4, 124.6, 123.0, 121.9; MS (EI): *m/z*: 337 [M]⁺, 359 [M + Na]⁺; Anal. calcd. for C₂₆H₂₄: C, 92.81; H, 7.19; found: C, 92.67; H, 7.02.

2,7-di(adamant-1-yl)pyrene (5). Purification by chromatography eluting with hexane/ethyl acetate (95/5, v/v) gave **5** (447 mg, 95% (Table 1, Entry 5)) as a white powder. mp >300 °C. ¹H NMR (600 MHz, CDCl₃) δ 8.17 (s, 4H), 8.03 (s, 4H), 2.21 (s, 18H), 1.89 (s, 12H); ¹³C{¹H} NMR (151 MHz, CDCl₃) δ 148.8, 130.8, 127.4, 123.0, 121.6, 43.8, 36.9, 29.1; MS (EI): *m/z*: 471 [M]⁺, 493 [M + Na]⁺; Anal. calcd. for C₃₆H₃₈: C, 91.86; H, 8.14; found: C, 91.96; H, 8.01.

1,3-bis(pyren-2-yl)adamantane (6). Purification by chromatography eluting with hexane/ethyl acetate (90/10, v/v) gave **6** (332 mg, 62% (Table 1, Entry 8)) as a white powder. mp 274–275 °C. ¹H NMR (600 MHz, CDCl₃) δ 8.31 (s, 4H), 8.15 (d, *J* = 7.8 Hz, 4H), 8.09 (d, *J* = 9.0 Hz, 4H), 8.06 (d, *J* = 9.0 Hz, 4H), 7.96 (t, *J* = 7.8 Hz, 2H), 2.11 (s, 2H), 2.57 (s, 2H), 2.36 (q, *J* = 12.0 Hz, 8H), 2.02 (s, 2H); ¹³C{¹H} NMR (151 MHz, CDCl₃) δ 148.5, 131.1, 130.9, 127.6, 127.3, 125.5, 124.7, 124.6, 123.1, 121.9, 50.2, 43.0, 38.0, 36.1, 29.9; MS (EI): *m/z*: 537 [M]⁺, 559 [M + Na]⁺; Anal. calcd. for C₄₂H₃₂: C, 93.89; H, 6.01; found: C, 94.02; H, 5.88.

1,3,5-tris(pyren-2-yl)adamantane (7). Purification by chromatography eluting with hexane/ethyl acetate (85/15, v/v) gave **7** (464 mg, 63% (Table 1, Entry 10)) as a white powder. mp 221–222 °C. ¹H NMR (600 MHz, CDCl₃) δ 8.42 (s, 6H), 8.17 (d, *J* = 7.8 Hz, 6H), 8.12 (d, *J* = 9.0 Hz, 6H), 8.08 (d, *J* = 9.0 Hz, 6H), 7.99 (t, *J* = 7.8 Hz, 3H), 2.86 (m, 1H), 2.82 (d, *J* = 12.0 Hz, 3H), 2.75 (d, *J* = 12.6 Hz, 3H), 2.48 (d, *J* = 2.4 Hz, 6H); ¹³C{¹H} NMR (151 MHz, CDCl₃) δ 147.8, 131.2–, 131.0, 127.6, 127.4, 125.6, 124.8, 124.5, 123.–3, 121.9, 49.4, 42.2, 39.2, 30.7; MS (EI): *m/z*: 737 [M]⁺, 759 [M + Na]⁺; Anal. calcd. for C₅₈H₄₀: C, 94.53; H, 5.47; found: C, 94.41; H, 5.35.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

¹H and ¹³C NMR spectra, absorption and emission spectra and photophysical data for compounds **2–7** (PDF)

Crystal structure data of compound **7** (CIF)

AUTHOR INFORMATION

Corresponding Author

*E-mail: anna.wrona@chemia.uni.lodz.pl

ORCID

Anna Wrona-Piotrowicz: 0000-0001-6930-6989

Anna Makal: 0000-0001-9357-2670

Janusz Zakrzewski: 0000-0002-3403-1126

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

Financial support from the Polish National Science Centre (NCN, Grant Miniatura 2018/02/X/ST5/01472) is gratefully acknowledged.

REFERENCES

- (a) Figueira-Duarte, T. M.; Müllen, K. Pyrene-based materials for organic electronics. *Chem. Rev.* **2011**, *111* (11), 7260–7314; (b) Islam, M. M.; Hu, Z.; Wang, Q.; Redshaw, C.; Feng, X. Pyrene-based aggregation-induced emission luminogens and their applications. *Mater. Chem. Front.* **2019**, *3*, 762–781.
- (a) Feng, X.; Hu, J.-Y.; Redshaw, C.; Yamato, T. Functionalization of pyrene to prepare luminescent materials—typical examples of synthetic methodology. *Chem. Eur. J.* **2016**, *22* (34), 11898–11916; (b) Casas-Solvas, J. M.; Howgego, J. D.; Davis, A. P. Synthesis of substituted pyrenes by indirect methods. *Org. Biomol. Chem.* **2014**, *12* (2), 212–232; (c) Mochida, K.; Kawasumi, K.; Segawa, Y.; Itami, K. Direct Arylation of Polycyclic Aromatic Hydrocarbons through Palladium Catalysis. *J. Am. Chem. Soc.* **2011**, *133* (28), 10716–10719; (d) Ji, L.; Lorbach, A.; Edkins, R. M.; Marder, T. B. Synthesis and Photophysics of a 2,7-Disubstituted Donor–Acceptor Pyrene Derivative: An Example of the Application of Sequential Ir-Catalyzed C–H Borylation and Substitution Chemistry. *J. Org. Chem.* **2015**, *80* (11), 5658–5665;
- (a) Feng, X.; Hu, J.-Y.; Iwanaga, F.; Seto, N.; Redshaw, C.; Elsegood, M. R. J.; Yamato, T. Blue-emitting butterfly-shaped 1,3,5,9-tetraarylpyrenes: synthesis, crystal structures, and photophysical properties. *Org. Lett.* **2013**, *15* (6), 1318–1321; (b) Feng, X.; Hu, J.-Y.; Tomiyasu, H.; Seto, N.; Redshaw, C.; Elsegood, M. R. J.; Yamato, T. Synthesis and photophysical properties of novel butterfly-shaped blue emitters based on pyrene. *Org. Biomol. Chem.* **2013**, *11* (48), 8366–8374; (c) Liu, R.; Ran, H.; Zhao, Z.; Yang, X.; Zhang, J.; Chen, L.; Sun, H.; Hu, J.-Y. Synthesis and Optical Properties of Donor–Acceptor-Type 1,3,5,9-Tetraarylpyrenes: Controlling Intramolecular Charge-Transfer Pathways by the Change of π -Conjugation Directions for Emission Color Modulations. *ACS Omega* **2018**, *3* (5), 5866–5875; (d) Maeda, H.; Hironishi, M.; Ishibashi, R.; Mizuno, K.; Segi, M. Synthesis and fluorescence properties of dioxo-, dithia-, and diselena-[3.3](1,3)pyrenophanes. *Photochem. Photobiol. Sci.* **2017**, *16* (2), 228–237.
- (a) Peng, J.; Guo, X.; Jiang, X.; Zhao, D.; Ma, Y. Developing efficient heavy-atom-free photosensitizers applicable to TTA upconversion in polymer films. *Chem. Sci.* **2016**, *7* (2), 1233–1237; (b) Peng, J.; Jiang, X.; Guo, X.; Zhao, D.; Ma, Y. Sensitizer design for efficient triplet–triplet annihilation upconversion: annihilator-appended tris-cyclometalated Ir(III) complexes. *Chem. Comm.* **2014**, *50* (58), 7828–7830; (c) Duan, P.; Yanai, N.; Kimizuka, N. A bis-cyclometalated iridium complex as a benchmark sensitizer for efficient visible-to-UV photon upconversion. *Chem. Comm.* **2014**, *50* (86), 13111–13113; (d) Castellano, F. N.; McCusker, C. E. MLCT sensitizers in photochemical upconversion: past, present, and potential future directions. *Dalton Trans.* **2015**, *44* (41), 17906–17910.

5. Mandal, A.; Swain, P.; Nath, B.; Sau, S.; Mal, P. Unipolar to ambipolar semiconductivity switching in charge transfer cocrystals of 2,7-di-*tert*-butylpyrene. *CrystEngComm* **2019**, *21* (6), 981-989.
6. (a) Wrona-Piotrowicz, A.; Witalewska, M.; Zakrzewski, J.; Makal, A. Regioselective (thio)carbamoylation of 2,7-di-*tert*-butylpyrene at the 1-position with iso(thio)cyanates. *Beilstein J. Org. Chem.* **2017**, *13*, 1032-1038; (b) Wrona-Piotrowicz, A.; Zakrzewski, J.; Metivier, R.; Brosseau, A.; Makal, A.; Wozniak, K. Efficient synthesis of pyrene-1-carbothioamides and carboxamides. Tunable solid-state fluorescence of pyrene-1-carboxamides. *RSC Adv.* **2014**, *4* (99), 56003-56012.
7. (a) Chen, L.; Ren, P.; Carrow, B. P. Tri(1-adamantyl)phosphine: Expanding the Boundary of Electron-Releasing Character Available to Organophosphorus Compounds. *J. Am. Chem. Soc.* **2016**, *138* (20), 6392-6395; (b) Carrow, B. P.; Chen, L. Tri (1-adamantyl) phosphine: exceptional catalytic effects enabled by the synergy of chemical stability, donicity, and polarizability. *Synlett* **2017**, *28* (03), 280-288.
8. (a) Schwenger, A.; Frey, W.; Richert, C. Reagents with a Crystalline Coat. *Angew. Chem. Int. Ed.* **2016**, *55* (44), 13706-13709; (b) Tominaga, M.; Ukai, H.; Katagiri, K.; Ohara, K.; Yamaguchi, K.; Azumaya, I. Tubular structures bearing channels in organic crystals composed of adamantane-based macrocycles. *Tetrahedron* **2014**, *70* (15), 2576-2581; (c) Kawahata, M.; Hyodo, T.; Tominaga, M.; Yamaguchi, K. Separation of *p*-xylene from aromatic compounds through specific inclusion by acyclic host molecule. *CrystEngComm* **2018**, *20* (38), 5667-5671; (d) Tominaga, M.; Iekushi, A.; Katagiri, K.; Ohara, K.; Yamaguchi, K.; Azumaya, I. Organic crystals bearing both channels and cavities formed from tripodal adamantane molecules. *J. Mol. Struct.* **2013**, *1046*, 52-56; (e) Richert, C.; Krupp, F. Encapsulating Organic Crystals (EnOCs). *Synlett* **2017**, *28* (14), 1763-1766.
9. Tian, Y.; Zhu, G. Porous Aromatic Frameworks (PAFs). *Chem. Rev.* **2020** (DOI 10.1021/acs.chemrev.9b00687)
10. (a) Tan, Z.; Kote, R.; Samaniego, W. N.; Weininger, S. J.; McGimpsey, W. G. Intramolecular Singlet-Singlet and Triplet-Triplet Energy Transfer in Adamantyl-Linked Trichromophores. *J. Phys. Chem. A* **1999**, *103* (38), 7612-7620; (b) Meineke, D. N. H.; Bossi, M. L.; Ta, H.; Belov, V. N.; Hell, S. W. Bichromophoric Compounds with Orthogonally and Parallely Arranged Chromophores Separated by Rigid Spacers. *Chem. Europ. J.* **2017**, *23* (10), 2469-2475.
11. Olah, G. A.; Liang, G.; Mateescu, G. D. Stable carbocations. CLXXII. 2-Adamantyl cations. *J. Org. Chem.* **1974**, *39* (25), 3750-3754.
12. Laali, K. K.; Sarca, V. D.; Okazaki, T.; Brock, A.; Der, P. Triflic acid-catalyzed adamantylation of aromatics in [BMIM][OTf] ionic liquid; synthetic scope and mechanistic insight. *Org. Biomol. Chem.* **2005**, *3* (6), 1034-1042.
13. Feng, X.; Hu, J.-Y.; Seto, N.; Tao, Z.; Redshaw, C.; Elsegood, M. R. J.; Yamato, T. Pyrene-Based Y-shaped Solid-State Blue Emitters: Synthesis, Characterization, and Photoluminescence. *Chem.-Asian J.* **2012**, *7* (12), 2854-2863.
14. Tashiro, M.; Yamato, T. Halogenation of 2,7-di-*tert*-butyl-trans-10b,10c-dialkyl-10b,10c-dihydropyrenes. *Chem. Lett.* **1980**, *9* (9), 1127-1130.