

Article

Switchable Palladium-Catalyst Reaction of Bromomethyl Sulfoxides, CO and N-Nucleophiles: Aminocarbonylation at Csp³ versus Oxidative Carbonylation of Amines

Cristian Mollar, Carmen Ramirez de Arellano, Mercedes Medio-Simon, and Gregorio Asensio

J. Org. Chem., **Just Accepted Manuscript** • Publication Date (Web): 05 Oct 2012

Downloaded from <http://pubs.acs.org> on October 6, 2012

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications
High quality. High impact.


The Journal of Organic Chemistry is published by the American Chemical Society.
1155 Sixteenth Street N.W., Washington, DC 20036
Published by American Chemical Society. Copyright © American Chemical Society.
However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Switchable Palladium-Catalyst Reaction of Bromomethyl Sulfoxides, CO and N-Nucleophiles: Aminocarbonylation at Csp³ versus Oxidative Carbonylation of Amines

Cristian Mollar, Carmen Ramirez de Arellano, Mercedes Medio-Simón* and Gregorio Asensio

Departamento de Química Orgánica Universidad de Valencia 46100-Burjassot-Valencia-Spain

email: Mercedes.Medio@uv.es



Abstract: The palladium-catalyzed reaction of α -bromomethyl sulfoxides, carbon monoxide and N-nucleophiles follows different reaction pathways according to the catalytic system and the reaction conditions. The Pd-xantphos catalyst affords high yields of α -sulfinyl amides by an aminocarbonylation process and is the first example of this type of transformation for a non benzylic sp³ hybridized carbon. On the other hand, the oxidative carbonylation of amines occurs with α -bromomethyl sulfoxides, carbon monoxide and catalytic $\text{Pd(PPh}_3)_4$ under aerobic conditions, yielding ureas and oxalamides from either primary or secondary amines. The reaction with ambident nucleophiles such as aminoalcohols was highly selective and took place exclusively at the amino group despite the presence of the alcohol functionality. In parallel to the reaction paths for simple amines, aminoalcohols were converted into hydroxy sulfinyl amides when the reaction was catalyzed by Pd-xantphos, while $\text{Pd(PPh}_3)_4$ catalyst afforded cyclic carbamates. The alkoxycarbonylation reaction of bromomethyl sulfoxides with simple alcohols and CO leading to the corresponding sulfinyl esters is also described.

Introduction

Palladium-catalyzed cross-coupling reactions are now a mainstay of fine chemical synthesis. The good compatibility of palladium-catalyzed synthetic procedures with a wide range of subsidiary

functional groups on the reactive partners justifies the continuous efforts being made to extend the scope of the organic transformations promoted by this metal.¹ In this area, carbonylation reactions² occupy a prominent place since they provide direct, convenient access to the omnipresent carbonyl building blocks. In general, the three-component reaction is carried out with carbon, oxygen or nitrogen nucleophiles, carbon monoxide, and aryl or alkyl halides as electrophiles. The alkoxycarbonylation reaction is perhaps the most robust among palladium-catalyzed carbonylation processes since alternative side reactions are limited and poorly favored.³ The opposite can be said of the carbonylative reaction of aryl halides with aryl metal reagents⁴ or amines.⁵ Nevertheless, the use of the carbonylation reaction has been progressively extended by challenging contributions. In this way, the development of new and effective palladium ligands allowed to overcome the drawbacks deriving from competing side reactions; moreover, the array of suitable substrates has been extended to include reluctant electrophiles such as aryl chlorides or aryl bromides in the case of carbonylative Sonogashira reactions or nucleophiles, like ammonia, phenols or anilines.⁶ In general, alkyl halides prove to be worse substrates than aryl halides, and only a few examples of the carbo-⁷ and alkoxycarbonylation⁸ of alkyl derivatives have been described to date. In addition, the intermolecular aminocarbonylation reaction with alkyl halides, benzyl halides⁹ or α -halomethyl ketones is a difficult transformation since direct nucleophilic substitution is highly favored. In sharp contrast, α -bromomethyl sulfoxides do not react with amines at the electrophilic α carbon¹⁰ and, for this reason, they are singular substrates for the aminocarbonylation at sp^3 -hybridized carbon atoms, as shown in this paper. This transformation is also interesting from a synthetic point of view since α -sulfinyl carbonyl derivatives are reliable synthetic intermediates¹¹ and metal ligands.¹² These compounds are readily available in the homochiral pure form following the procedure reported herein.


Furthermore, interesting singular structural properties of these compounds, such as the S--O interaction between sulfur and the carbonyl oxygen atom, have been described.¹³ This intramolecular interaction seems to play a critical role in, for instance, the stereochemical control of the Pummerer rearrangement. Previously, we showed that bromomethyl sulfoxides give ketosulfoxides with moderate to good yields in carbonylative Suzuki reactions with boronic acids and CO at atmospheric pressure at 60°C. The high chemoselectivity of the three-component reaction *versus* direct coupling is observed under these conditions.¹⁴ The new carbon-carbon bond formation, which occurs at the α position of a sulfinyl group from a halosulfoxide, overcomes the limitation of uncatalyzed reactions where nucleophiles such as Grignard reagents react at the sulfur atom with displacement of the halomethyl group.¹⁵ Alternatively, other α -functionalizations of sulfoxides described to date usually occur through the corresponding anions,¹¹ and require strong bases that often favor secondary competing reactions.¹⁶ This scenario prompted us to further explore the palladium-catalyzed carbonylation of halosulfoxides by testing their reaction with amines, alcohols and aminoalcohols.

Results and discussion

The palladium-catalyzed aminocarbonylation reaction in primary alkyl halides is a difficult transformation because oxidative addition in this kind of substrates is slow,¹⁷ while the competitive direct nucleophilic substitution with amines proves favorable. This trend is observed even with benzyl halides,⁹ which are activated toward the palladium oxidative addition. In contrast bromomethyl sulfoxides are good substrates in palladium-catalyzed cross-coupling reactions,¹⁸ but are poor substrates in S_N2 reactions with usual nucleophiles.^{10, 15} By bearing this background in mind, we first explored the possibility of carrying out the aminocarbonylation of the sp³-hybridized carbon of bromomethyl sulfoxides. The aminocarbonylation of sulfoxides **1a**

or (*S*)-**1b** took place adequately with aliphatic **2a-f** and aromatic amines **2g-i** with CO at atmospheric pressure and catalytic Pd₂(dba)₃-xantphos affording the corresponding racemic (**3aa-3ah**) or chiral sulfinyl amides (**3ba**, **3be**, and **3bi**) with high yields (93-99%) (Table 1). Palladium-catalyzed aminocarbonylation requires the presence of the base in stoichiometric amount or an excess of base to neutralize the acid formed during the reaction. Common organic trialkyl amines and inorganic salts (K₃PO₄, K₂CO₃, CsF) were tested to promote the reaction. CsF was found to be the most effective, which is in agreement with previous results in several carbonylative Suzuki reactions.¹⁴

Table 1. Palladium-Catalyzed Synthesis of Chiral and Racemic Sulfinyl Amides.



Sulfinyl amides prove to be attractive substrates as potential chiral ligands. The scarce references available on the synthesis of this class of compounds involve the use of dicyclohexylcarbodiimide to activate the corresponding acid¹³ or the use of amide anions previously generated under

strongly basic conditions. Therefore, the aminocarbonylation described herein is a competitive alternative for these compounds in simplicity and atom economy terms. The absolute configuration of the starting chiral sulfoxides is retained in the formation of the corresponding sulfinyl amides. The palladium-catalyzed cross-coupling reactions of chiral bromomethyl sulfoxide (*S*)-**1b** and boronic acids, previously described by us,¹⁸ also takes place with retention of the configuration. As expected, the chiral HPLC analysis of sulfinyl amides **3ba**, **3be** and **3bi** proved that chiral purity is preserved during transformation. The absolute configuration of sulfinyl amides **3be** and **3bi** has been established by single-crystal X-ray diffraction (for ORTEP diagrams, see Supporting Information).

The crystal structure of sulfinyl amide (*R*)-**3bi** is shown in Figure 1.





Figure 1. X-ray thermal ellipsoid plot of compound (*R*)-**3bi** (50% probability level).

The palladium-catalyzed reaction of amines, carbon monoxide and bromomethyl sulfoxides depends on the ligand, and also on the presence or absence of oxygen. Sulfinyl amides **3** were prepared in high yields with Pd₂(dba)₃-xantphos as the catalyst under aerobic or anaerobic conditions. Conversely, the preparation of compounds **3** using Pd(PPh₃)₄ requires the rigorous exclusion of oxygen. Under aerobic conditions (see Experimental Section¹⁹) oxidative carbonylation of the



Table 2. Palladium-Catalyzed Oxidative Carbonylation of Amines mediated by α -Bromomethyl Sulfoxide **1a**.

Conditions

Products



A

B

**3aa** (5%)**3aa** (52%)**4a** (95%)**4a** (24%)**6** (18%)



A

B

**3ab** (3%)**3ab** (53%)**4b** (93%)**4b** (20%)**6** (19%)



A

B

**3ac** (5%)**3ac** (51%)**4c** (93%)**4c** (20%)**6** (19%)



A

B

**3ad** (9%)**3ad** (89%)**5a** (60%)**5a** (5%)**6** (9%)**6** (3%)



A

B

**3ae** (5%)**3ae** (90%)**5b** (48%)**5b** (4%)**6** (8%)**6** (4%)

A

B

**3af** (5%)**3af** (89%)**5c** (55%)**5c** (5%)**6** (8%)**6** (6%)

amine²⁰ occurs, while ureas **4** or oxalamides **5** were the main products from the primary or secondary amines (Table 2). Ureas **4a-c**, slightly contaminated by sulfinyl amides **3aa-ac**, were obtained with very high yields from amines **2a-c**. Oxalamides **5**, the common by-products in the carbonylation of amines, were not detected from amines **2a-c** under our reaction conditions. In contrast, oxalamides **5a-c** were the major products of secondary aliphatic amines **2d-2f**, which were contaminated only by sulfinyl amides **3ad-af**, the three-component cross-coupling products (Table 2). Primary amines gave symmetrical ureas **4**, most probably through an isocyanate intermediate (*vide infra*), which is an unfeasible reaction pathway in the case of secondary amines. Detection of isocyanates in the oxidative carbonylation of amines has been previously reported.²⁰


The formation of ureas and oxalamides described herein occurred under very mild conditions if compared with the precedents in this area. In this sense, it is noteworthy that: i) carbonylation takes place with CO at an atmospheric pressure. Previous procedures require up to 50 atm of CO; ii) ureas are the only product from primary amines that are free of contamination by oxalamides, the usual by-products in these amine carbonylations.

Next, we turned our attention to explore selective aminocarbonylation reactions using dinucleophiles, such as aminoalcohols, where amino- and/or alkoxycarbonylation could take place alternatively. The alkoxycarbonylation of bromomethyl sulfoxides has not yet been described; hence, the reaction of 1-bromomethyl sulfoxides, CO and simple alcohols was first evaluated (Table 3). To optimize the reaction conditions, sulfoxide **1a**, ethanol and Pd(PPh₃) were selected. The initial screening was conducted in alcohol with an excess or stoichiometric hydroxylic base, which is usual in palladium-catalyzed alkoxycarbonylation reactions.^{2b}

Reactions took place in aprotic-coordinating solvents, such as THF, containing two equivalents of alcohol, but did not proceed in ethanol. A moderate base like CsF proved more effective than

other usual bases employed to promote the reaction. The yields obtained using $\text{Pd}_2(\text{dba})_3$ -xantphos or $\text{Pd}(\text{PPh}_3)_4$ in the alkoxy carbonylation of bromomethyl sulfoxide were similar.

Table 3. Palladium-Catalyzed Alkoxy carbonylation of α -Bromomethyl Sulfoxides **1**.




The carbonylation of bromomethyl sulfoxide with aliphatic low-boiling point primary alcohols and catalytic $\text{Pd}(\text{PPh}_3)_4$ gave the corresponding sulfinyl esters with excellent yields (Table 3). Reactions were conducted below the alcohol boiling temperature and did not require special equipment. On the other hand, benzyl alcohol, which is easily oxidized under palladium catalysis, gave only a moderate yield (Table 3). Substituted alcohols containing additional functional coordinating groups also reacted efficiently in the three-component process (Table 3).

Bromomethyl sulfoxide behaves better than halomethyl ketones^{8b-d} in the alkoxy carbonylation reaction since the competitive dehalogenation reaction is more difficult in the former case.

A three-component reaction between bromomethyl sulfoxide **1**, CO and aminoalcohols occurred and exclusively gave aminoacylation products **10** when $\text{Pd}(\text{dba})_2$ -xantphos was used as the catalyst. However, selectivity dramatically changed when $\text{Pd}(\text{PPh}_3)_4$ was used to catalyze the reaction; cyclic carbamates **11** were obtained in the latter case (Scheme 1). The different

coordination possibilities of compounds **10** confer them interesting features for the facultative use as ligands.



Scheme 1. Palladium-catalyzed carbonylation of α -bromomethyl sulfoxides **1** and aminoalcohols **9**.

Mechanisms

The most frequently postulated catalytic cycle^{2a,2b} to account for amino- and alkoxy carbonylation involves the following steps: oxidative addition of the electrophile to Pd(0) (step 1), migratory insertion of carbon monoxide (step 2) and attack of the nucleophile (alcohol or amine) to the acyl complex (step 3). Nevertheless, based on our previous findings in the palladium-catalyzed carbonylative Suzuki reaction with these compounds,¹⁴ the above proposal hardly explains the results obtained in the case of bromomethyl sulfoxides. In fact, the experimental data from this work suggest an atypical sequence in the catalytic cycle, with the insertion of carbon monoxide taking place after the transmetallation step instead of the usual CO insertion into the oxidative addition complex. Moreover, we proved the reluctance of the oxidative addition complex **I** to insert CO by monitoring this process by NMR.

Therefore, the amino and alkoxy carbonylation of bromomethyl sulfoxides could be explained better as follows²¹: first, oxidative addition of sulfoxide **1** to palladium (0), followed by the attack

of the nucleophile (amine or alcohol) to the coordinated CO leading to complexes **II** or **III**, and then, the reductive elimination step (Scheme 2). Formation of intermediate $\text{ArS(O)CH}_2\text{PdOR}$ via metathesis from the oxidative addition complex, followed by insertion of carbon monoxide into the Pd-OR bond and, finally, reductive elimination, cannot be ruled out to explain the alkoxycarbonylation reaction. In fact, this alternative involving Pd-OR species could account for the competitive oxidation of alcohols observed in some cases. A similar catalytic cycle based on kinetic studies to account for the alkoxycarbonylation of aryl iodides in the presence of sodium alkoxides has been previously postulated; however, this sequence is less likely in absence of strong bases.²²


The $\text{Pd}_2(\text{dba})_3$ -xantphos catalyzed aminocarbonylation reaction of α -bromomethyl sulfoxides **1** occurs in the absence of competitive side reactions (Scheme 2). Conversely, dehalogenation of the sulfoxide and oxidative carbonylations of the amine²⁰ leading to ureas or oxalamides resulted in detectable side pathways with $\text{Pd}(\text{PPh}_3)_4$ under anaerobic conditions. In this case, bromomethyl sulfoxide behaves as an oxidant²³ (Scheme 3). An schematic catalytic cycle 2 (Scheme 3) to account for the oxidative carbonylation of amines is proposed to involve a complex **Ia** formed in the oxidative addition of bromomethyl sulfoxide to Pd(0). Accordingly, **Ia** would react with CO and amine to give **IIa**, a carbamoyl dioorganopalladium complex. In the case of primary amines, the subsequent hydrogen transfer to the metal leads to palladium hydride complex **IV** and an isocyanate (*via a*). Ureas **4** are formed by the reaction of the isocyanate with a second equivalent of amine, while methyl phenyl sulfoxide **6** results by reductive elimination from complex **IV**. Alternatively, oxalamides **5** and methyl phenyl sulfoxide **6** would be produced from the reaction of complex **IIa** and a second equivalent of the corresponding secondary amine and carbon monoxide (*via b*).

Conversely under aerobic conditions, the oxidative carbonylation of amines became the main process in the reaction of bromomethyl sulfoxide **1a** with amines and Pd(PPh₃)₄. Ureas **4** or oxalamides **5** were formed by a catalytic cycle in which molecular oxygen acted as the oxidant (Scheme 4). Oxidative carbonylation could be explained by an alternative catalytic cycle involving a peroxo complex²⁴ of palladium without the participation of bromomethyl sulfoxide. However, the catalyst displays longer activity and proves highly efficient in the presence of **1**, which is not consumed during the process, but is recovered (about 80%) after completing the reaction. Thus, a catalytic cycle 3 is proposed to account for the oxidative carbonylation of amines (Scheme 4). Intermediate complex **IV** participates in cycles 2 and 3 (Schemes 3 and 4), but its evolution differs in the presence or absence of O₂. Insertion of molecular oxygen into the palladium-hydrogen bond in complex **IV** and loss of hydrogen peroxide by ligand exchange with CO and amine under aerobic conditions close the catalytic cycle, with the regeneration of complex **IIa**. The generated hydrogen peroxide could play the role of a further oxidant, but palladium has been reported to accelerate its decomposition in O₂ and water.²⁵ Hydrogen peroxide is a stronger oxidant than molecular oxygen. Nevertheless, the data from other palladium-catalyzed oxidation reactions suggest that it is not a kinetically competent oxidant.²⁵ The formation of cyclic carbamates **11** from aminoalcohols **9** is explained by following an equivalent reaction pathway.


Formation of oxalamides **5** from secondary amines cannot be explained in the same way as ureas **4** given the lack of a β-hydrogen to be transferred from the corresponding carbamoyl intermediate **IIa**. The oxidative carbonylation of secondary amines could be explained by an alternative catalytic cycle, where molecular oxygen (*via* a peroxo complex of palladium)²⁴ acts as the oxidant. In addition, the formation of dehalogenated sulfoxide **6** under aerobic conditions

1
2
3 suggests that bromomethyl sulfoxide acts concomitantly with O₂ as oxidant. In contrast for
4
5 primary amines, bromomethyl sulfoxide seems to act as an oxidant, but only under anaerobic
6
7 conditions since dehalogenated sulfoxide is only detected in this way.
8
9


10
11
12 The ligand effect encountered in the palladium-catalyzed carbonylation reaction of sulfoxides **1**
13
14 with amines could be attributed to differences in the ligand-dependent *trans-cis* isomerization in
15
16 complex **II**, the common intermediate in both cycles. The *trans-cis* isomerization process in
17
18 complex **II** with monophosphine ligands opens temporary coordination sites in palladium, which
19
20 allows either the transfer of hydrogen in the case of a primary amine or the coordination of CO
21
22 followed by amine attack in the case of secondary amines, thus favoring the occurrence of the
23
24 oxidative carbonylation of amines instead of the aminocarbonylation reaction. These events may
25
26 be minimized in the case of the diphosphine xantphos which promotes a faster isomerization.²⁶
27
28 Besides, the wide bite angle of xantphos favors the reductive elimination process and,
29
30 consequently, the formation of α -sulfinyl amides results in the prevailing pathway. The slower
31
32 reductive elimination promoted by monophosphines enhances the competitive oxidative
33
34 carbonylation of amines as an alternative reaction pathway.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Scheme 2. Cycle 1. Proposed catalytic cycle for the palladium-catalyzed alkoxy and aminocarbonylation of sulfoxides **1**.



Scheme 3. Cycle 2. Proposed catalytic cycles for the palladium-catalyzed oxidative carbonylation of amines with sulfoxide **1a** as oxidant.



Scheme 4. Cycle 3. Proposed catalytic cycles for the palladium-catalyzed oxidative carbonylation of amines with O_2 as oxidant.

Conclusion

In conclusion, we describe palladium-catalyzed carbonylation involving a sp^3 -hybridized carbon and N- and N,O-nucleophiles for the first time. The reaction takes advantage of the reluctance of the electrophilic partner, that is, α -bromomethyl sulfoxides, to participate in nucleophilic substitution reactions. Consequently the aminocarbonylation reaction can be carried out with high selectivity for a wide range of amines and aminoalcohols. The developed method requires neither overpressure of carbon monoxide nor sophisticated palladium ligands. The mild reaction conditions allow us to obtain chiral and racemic α -sulfinyl amides with very high efficiency. The effect of the ligand and reaction conditions driving the outcome of the process to the aminocarbonylation reaction or the oxidative carbonylation of amines in almost a completely

selective manner shows the high potential of the three-component reactions presented herein, which allows access to a wide range of products with minimal changes in simple and easily available starting materials and catalyst systems.

Experimental Section

General methods. Solvents were degassed by the Freeze-Pump-Thaw method (five cycles) to ensure the absolute strict absence of oxygen. Proton magnetic resonance and carbon magnetic resonance were recorded at 300 MHz and 75 Hz, respectively. Chemical shifts are reported in δ ppm in relation to the CHCl_3 peak at 7.20 ppm (^1H) or 77.0 ppm (^{13}C). Multiplicities are reported as follows: s= singlet, d= doublet, t= triplet, q= quartet, m= multiplet. The ee values of the products were determined by HPLC (Chiralcel IA). Optical rotation measurements were determined at room temperature. All the melting points were uncorrected. Reactions were monitored by analytical thin layer chromatography using commercial aluminium sheets pre-coated (0.2-mm layer thickness) with silica gel 60 F₂₅₄. Product purification was performed by flash chromatography with silica gel (230-400 mesh).

Starting materials

Bromomethyl phenyl sulfoxide (1a); ²⁷ (*S*)-1-(bromomethylsulfinyl)-4-methylbenzene (**1b**),^{18a} and *trans*-[Pd(CH₂SOPh)Br(PPh₃)₂]^{18a} were prepared according to the previously described procedures.

General procedure for the carbonylation of amines, aminoalcohols and alcohols

Aerobic conditions. A mixture of α -bromomethyl sulfoxide **1** (0.3 mmol), N- or O-nucleophile (0.6 mmol), CsF (1.2 mmol) and the appropriate catalyst (0.03 mmol) was added to a flask fitted with a reflux condenser and a septum inlet. The flask was fluxed with carbon monoxide and then charged with THF (10 mL). The mixture was stirred at 65°C under an atmospheric pressure of carbon monoxide (balloon).¹⁹ After the appropriate time, the mixture was cooled at room

temperature, quenched with water (10mL) and extracted with diethyl ether (2x 15mL) and dichloromethane (3 x15 mL). The combined organic extracts were dried with Na₂SO₄ and evaporated under reduced pressure.

Anaerobic conditions. A mixture of α -bromomethyl sulfoxide **1** (0.3 mmol), N-nucleophile (0.6 mmol), CsF (1.2 mmol) and Pd(PPh₃)₄ (0.03 mmol) was added to a flask fitted with a reflux condenser and a septum inlet. The flask was fluxed several times with carbon monoxide and then charged with degassed THF (10 mL). The mixture was stirred at 65°C under an atmospheric pressure of carbon monoxide (the balloon was immediately inflated before starting the reaction to minimize air diffusion).¹⁹ After the appropriate time, the mixture was cooled at room temperature, quenched with water (10mL) and extracted with diethyl ether (2x 15mL) and dichloromethane (3 x15 mL). The combined organic extracts were dried with Na₂SO₄ and evaporated under reduced pressure

N-Octyl-2-(phenylsulfinyl)acetamide (3aa). Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 84 mg, 95% yield; mp 40–44 °C; ¹H NMR (CDCl₃) δ 7.70-7.67 (m, 2H), 7.46-7.44 (m, 3H), 3.62 (d, *J*= 14.1 Hz, 1H), 3.41 (d, *J*= 14.1 Hz, 1H), 3.18-3.05 (m, 2H), 1.19 (m, 12H), 0.81 (t, *J*= 6.6 Hz, 3H); ¹³CNMR (CDCl₃) δ 163.7, 143.8, 131.3, 129.1, 124.4, 58.8, 39.7, 31.7, 29.3, 29.2, 29.1, 26.8, 22.6, 14.0; HRMS (EI) *m/z* (*M*⁺) calcd for C₁₆H₂₅NO₂S 295.1606, found 295.1600.

N-Benzyl-2-(phenylsulfinyl)acetamide (3ab).²⁸ Chromatographic purification on a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). Pale yellow liquid: 79 mg, 96% yield; ¹H-NMR (CDCl₃) δ 7.49-7.40 (m, 5H), 7.25-7.13 (m, 5H), 4.35 (dd, *J*=14.7 and 6 Hz, 1H), 4.27 (dd, *J*=14.7 and 6 Hz, 1H), 3.69 (d, *J*= 14.4 Hz, 1H), 3.42 (d, *J*= 14.4 Hz, 1H); ¹³C-NMR (CDCl₃) δ 163.5, 141.3, 137.5, 129.4, 128.7, 131.5, 127.9, 127.5, 123.9, 58.3, 43.6; HRMS (E.I.) *m/z* (*M*⁺) calcd. for C₁₅H₁₅NO₂S 273.0823, found 273.0811.

N-(4-phenylbutyl)-2-(phenylsulfinyl)acetamide (3ac). Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 88 mg, 93% yield; mp 55–59 °C; ¹H-NMR (CDCl₃) δ 7.53-7.50 (m, 2H), 7.44-7.42 (m, 3H), 7.14-7.08 (m, 5H), 6.90-6.80 (m, 1H), 3.61 (d, *J*= 14.3 Hz, 1H), 3.38 (d, *J*= 14.3 Hz, 1H), 3.24-3.06 (m, 2H), 2.54 (t, *J*= 7.2 Hz, 2H), 1.59-1.36 (m, 4H); ¹³C-NMR (CDCl₃) δ 163.9, 142.4, 141.9, 132.0, 129.8, 128.8, 128.7, 126.2, 124.3, 59.0, 39.9, 35.8, 29.4, 29.0; HRMS (E.I.) *m/z* (*M*⁺) calcd. for C₁₈H₂₁NO₂S 315.1293, found 315.1283.

***N,N*-Dibutyl-2-(phenylsulfinyl)acetamide (3ad).**^{12a} Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 83 mg, 94% yield; mp 41–45 °C; ¹H-NMR (CDCl₃) δ 7.76–7.73 (m, 2H), 7.52–7.50 (m, 3H), 3.98 (d, *J* = 13.9 Hz, 1H), 3.64 (d, *J* = 13.9 Hz, 1H), 3.29–3.00 (m, 4H), 1.46–1.15 (m, 8H), 0.87 (t, *J* = 7.2 Hz, 6H); ¹³C-NMR (CDCl₃): δ = 163.7, 143.8, 131.3, 129.1, 124.5, 62.2, 48.2, 46.0, 31.0, 29.6, 20.0, 19.8, 13.7, 13.6; HRMS (E.I.) *m/z* (*M*⁺) calcd. for C₁₆H₂₅NO₂S 295.1606, found 295.1593.

2-(Phenylsulfinyl)-1-(pyrrolidin-1-yl)ethanone (3ae). Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 69 mg, 97% yield; mp 104–108 °C; ¹H-NMR (CDCl₃) δ 7.69–7.66 (m, 2H), 7.47–7.41 (m, 3H), 3.93 (d, *J* = 13.2 Hz, 1H), 3.60 (d, *J* = 13.2 Hz, 1H), 3.45–3.32 (m, 3H), 3.09–3.01 (m, 1H), 1.86–1.63 (m, 4H); ¹³C-NMR (CDCl₃) δ 162.3, 143.7, 129.2, 131.4, 124.2, 63.2, 47.3, 46.0, 25.9, 24.3; HRMS (E.I.) *m/z* (*M*⁺) calcd. for C₁₂H₁₅NO₂S 237.0823, found 237.0814.

1-Morpholino-2-(phenylsulfinyl)ethanone (3af). Chromatographic purification on a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 74 mg, 98% yield; mp 146–150 °C; ¹H-NMR (CDCl₃) δ 7.67–7.64 (m, 2H), 7.49–7.47 (m, 3H), 3.92 (d, *J* = 13.2 Hz, 1H), 3.66 (d, *J* = 13.2 Hz, 1H), 3.41–3.63 (m, 8H); ¹³C-NMR (CDCl₃) δ 162.7, 143.3, 131.6, 129.3, 124.1, 66.6, 66.5, 60.9, 46.8, 42.3; HRMS (E.I.) *m/z* (*M*⁺) calcd. for C₁₂H₁₅NO₃S 253.0773, found 253.0761

***N*-(4-Methoxyphenyl)-2-(phenylsulfinyl)acetamide (3ag).** Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 86 mg, 99% yield; mp 124–126 °C; ¹H-NMR (CDCl₃) δ 7.58–7.54 (m, 2H), 7.46–7.44 (m, 3H), 7.28 (d, *J* = 9.0 Hz, 2H), 6.75 (d, *J* = 9.0 Hz, 2H), 3.84 (d, *J* = 14.4 Hz, 1H), 3.49 (d, *J* = 14.4 Hz, 1H), 3.71 (s, 3H); ¹³C-NMR (CDCl₃) δ 161.4, 156.6, 140.8, 131.8, 130.4, 129.5, 123.9, 122.0, 114.1, 58.5, 55.4; HRMS (E.I.) *m/z* (*M*⁺) calcd. for C₁₅H₁₅NO₃S 289.0773, found 289.0779.

***N*-(4-chlorophenyl)-2-(phenylsulfinyl)acetamide (3ah).** Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 84 mg, 95% yield; mp 162–166 °C; ¹H-NMR (CDCl₃) δ 7.56–7.52 (m, 2H), 7.46–7.44 (m, 3H), 7.34 (d, *J* = 9.0 Hz, 2H), 7.17 (d, *J* = 9.0 Hz, 2H), 3.86 (d, *J* = 14.1 Hz, 1H), 3.52 (d, *J* = 14.1 Hz, 1H); ¹³C-NMR (CDCl₃) δ 161.6, 140.6, 135.9, 131.9, 129.6, 129.0, 123.8, 121.4, 58.5; HRMS (E.I.) *m/z* (*M*⁺): calcd. for C₁₄H₁₂ClNO₂S 293.0277, found 293.0269

(*R*)-2-(*p*-tolylsulfinyl)-*N*-octylacetamide (3ba). Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 92 mg, 99% yield; mp 170–174

°C; ¹H-NMR (CDCl₃) δ 7.26 (d, *J* = 8.1 Hz, 2H), 7.22 (d, *J* = 8.1 Hz, 2H), 3.60 (d, *J* = 14.1 Hz, 1H), 3.35 (d, *J* = 14.1 Hz, 1H), 3.24-3.03 (m, 2H), 2.34 (s, 3H), 1.20 (m, 12H), 0.81 (t, *J* = 6.9 Hz, 3H); ¹³C-NMR (CDCl₃) δ 163.6, 142.1, 138.2, 130.1, 123.9, 58.5, 39.7, 31.8, 29.2, 29.1, 26.9, 22.6, 21.4, 14.1; HRMS (EI) *m/z* (*M*⁺) calcd. for C₁₇H₂₇NO₂S 309.1762, found 309.1755; [α]_D²⁵ +208.3 (c:1, acetone), ee > 99% (Chiralcel IA, Hexane:2-Propanol 95:5, flow 1 mL/min).

(*R*)-2-(*p*-tolylsulfinyl)-1-(pyrrolidin-1-yl)ethanone (3be). Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 70 mg, 93% yield; mp 121–125 °C; ¹H-NMR (CDCl₃) δ 7.55 (d, *J* = 8.1 Hz, 2H), 7.26 (d, *J* = 8.1 Hz, 2H), 3.91 (d, *J* = 12.9 Hz, 1H), 3.57 (d, *J* = 12.9 Hz, 1H), 3.49-3.05 (m, 4H), 2.35 (s, 3H), 1.86-1.68 (m, 4H); ¹³C-NMR (CDCl₃) δ 162.4, 141.9, 140.4, 129.9, 124.2, 63.2, 47.3, 46.0, 25.9, 24.3, 21.4; HRMS (E.I.) *m/z* (*M*⁺) calcd. for C₁₃H₁₇NO₂S 251.0980, found 251.0970; [α]_D²⁵ +140.2 (c: 1, acetone), ee > 99% (Chiralcel IA, Hexane: 2-Propanol 96:4, flow 4 mL/min).

(*R*)-2-(*p*-tolylsulfinyl)-*N*-*p*-tolylacetamide (3bi). Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 82 mg, 96% yield; mp 170–172 °C. ¹H-NMR (CDCl₃) δ 7.43 (d, *J* = 8.1 Hz, 2H), 7.27 (d, *J* = 8.1 Hz, 2H), 7.22 (d, *J* = 8.1 Hz, 2H), 7.00 (d, *J* = 8.1 Hz, 2H), 3.80 (d, *J* = 14.1 Hz, 1H), 3.49 (d, *J* = 14.1 Hz, 1H), 2.35 (s, 3H), 2.33 (s, 3H); ¹³C-NMR (CDCl₃) δ 161.6, 142.3, 137.6, 134.8, 134.2, 130.2, 129.4, 120.3, 60.3, 21.4, 21.0; HRMS (E.I.) *m/z* (*M*⁺) calcd. for C₁₆H₁₇NO₂S 287.0980, found 287.0985; [α]_D²⁵ +211.4 (c:1, acetone), ee > 99% (Chiralcel IA, Hexane: 2-Propanol 85:15, flow 3 mL/min).

1,3-dioctylurea (4a).²⁹ Chromatographic purification on a silica gel column with *n*-hexane–ethyl acetate (1 : 1). White solid: 81 mg, 95% yield; mp 89–90 °C; ¹H-NMR (CDCl₃) δ 5.00-5.01 (bb, 2H), 3.03-3.09 (m, 4H), 1.42-1.37 (m, 4H), 1.30-1.10 (m, 20H), 0.80 (t, *J* = 6.0 Hz, 6H); ¹³C-NMR (CDCl₃) δ 158.9, 40.2, 31.8, 30.4, 29.4, 29.3, 27.0, 22.6, 14.0; HRMS (E.I.) *m/z* (*M*⁺) calcd. for C₁₇H₃₆N₂O 284.2827, found 284.2812.

1,3-Dibenzylurea (4b).³⁰ Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1). White solid: 67 mg, 93% yield; mp 170–171 °C; ¹H-NMR (CDCl₃) δ 7.25-7.15 (m, 10H), 4.80 (bs, 2H), 4.26 (s, 2H), 4.24 (s, 2H); ¹³C-NMR (CDCl₃) δ 158.0, 139.1, 128.6, 127.4, 127.3, 44.5; HRMS (E.I.) *m/z* (*M*⁺) calcd. for C₁₅H₁₆N₂O 240.1263, found 240.1255.

1,3-Bis(4-phenylbutyl)urea (4c). Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1). White solid: 90 mg, 93% yield; mp 76–80 °C; ¹H-NMR (CDCl₃) δ 7.22 -7.12 (m,

4H), 7.10-7.07 (m, 6H), 4.33-4.30 (bb, 2H), 3.11-3.04 (m, 4H), 2.57-2.51 (m, 4H), 1.61-1.54 (m, 4H), 1.51-1.37 (m, 4H); ^{13}C -NMR (CDCl_3) δ 158.2, 142.2, 128.4, 128.3, 125.8, 40.3, 35.5, 29.8, 28.6; HRMS (E.I.) m/z (M^+) calcd. for $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}$ 324.2216, found 324.2217.

***N*¹,*N*¹,*N*²,*N*²-tetrabutylloxalamide (5a).**³¹ Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1). Colorless oil: 56 mg, 60% yield; ^1H -NMR (CDCl_3) δ 3.38-3.13 (m, 8H), 1.63-1.50 (m, 8H), 1.39-1.21 (m, 8H), 0.95-0.88 (m, 12H); ^{13}C -NMR (CDCl_3): δ = 165.0, 47.7, 43.5, 30.5, 29.2, 20.1, 20.0, 13.7, 13.6; HRMS (E.I.) m/z (M^+) calcd. for $\text{C}_{18}\text{H}_{36}\text{N}_2\text{O}_2$ 312.2777, found 312.2766.

1,2-Di(pyrrolidin-1-yl)ethane-1,2-dione (5b).³² Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1). White solid: 28 mg, 48% yield; mp 73–75 °C; ^1H -NMR (CDCl_3) δ 3.49-3.42- (m, 8H), 1.90-1.83 (m, 8H); ^{13}C -NMR (CDCl_3) δ 163.1, 46.8, 45.0, 25.8, 24.1; HRMS (E.I.) m/z (M^+) calcd. for $\text{C}_{10}\text{H}_{16}\text{N}_2\text{O}_2$ 196.1212, found 196.1199.

1,2-Dimorpholinoethane-1,2-dione (5c).³³ Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1). White solid: 38 mg, 55% yield; mp 181–185 °C; ^1H -NMR (CDCl_3) δ 3.68-3.57 (m, 12H), 3.40-3.37 (m, 4H); ^{13}C -NMR (CDCl_3): δ = 162.7, 66.7, 66.4, 46.4, 41.4; HRMS (E.I.) m/z (M^+) calcd. for $\text{C}_{10}\text{H}_{16}\text{N}_2\text{O}_4$ 228.1110, found 228.1103.

Ethyl 2-(phenylsulfinyl)acetate (8aa).³⁴ Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (5 : 1). Colorless oil: 53 mg, 83% yield; ^1H NMR (CDCl_3) δ 7.63-7.61 (m, 2H), 7.48-7.46 (m, 3H), 4.08 (q, J = 6.9 Hz, 2H), 3.78 (d, J = 13.5 Hz, 1H), 3.60 (d, J = 13.5 Hz, 1H), 1.14 (t, J = 7.2 Hz, 3H); ^{13}C NMR (CDCl_3) δ 164.0, 143.0, 131.7, 129.3, 124.1, 61.9, 61.6, 13.9; HRMS (EI) m/z (M^+) calcd for $\text{C}_{10}\text{H}_{12}\text{O}_3\text{S}$ 212.0507, found 212.0500.

Butyl 2-(phenylsulfinyl)acetate (8ab). Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (5 : 1). Colorless oil: 56 mg, 78% yield; ^1H NMR (CDCl_3) δ 7.64-7.62 (m, 2H), 7.48-7.46 (m, 3H), 4.03 (t, J = 6.6 Hz, 2H), 3.76 (d, J =13.5 Hz, 1H), 3.60 (d, J =13.5 Hz, 1H), 1.28-1.18 (m, 4H), 0.83 (t, J = 7.2 Hz, 3H); ^{13}C NMR (CDCl_3) δ 164.7, 143.1, 131.7, 129.3, 124.1, 65.8, 61.7, 30.3, 18.9, 13.6; HRMS (EI) m/z (M^+) calcd for $\text{C}_{12}\text{H}_{16}\text{O}_3\text{S}$ 240.08202, found 240.0806.

2-Chloroethyl 2-(phenylsulfinyl)acetate (8ac). Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (5 : 1). Colorless oil: 46 mg, 63% yield; ^1H NMR (CDCl_3) δ 7.65-7.62 (m, 2H), 7.50-7.47 (m, 3H), 4.27 (t, J = 5.7 Hz, 2H), 3.82 (d, J = 13.8 Hz, 1H), 3.67 (d,

$J=13.8$ Hz, 1H), 3.54 (t, $J=5.7$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 164.3, 142.8, 131.9, 129.4, 124.1, 65.2, 61.3, 40.9; HRMS (EI) m/z (M^+) calcd for $\text{C}_{10}\text{H}_{11}\text{ClO}_3\text{S}$ 246.0117, found 246.0110.

2-Methoxyethyl 2-(phenylsulfinyl)acetate (8ad). Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (5 : 1). Colorless oil: 44 mg, 61% yield; ^1H NMR (CDCl_3) δ 7.63–7.61 (m, 2H), 7.48–7.46 (m, 3H), 4.18 (t, $J=4.8$ Hz, 2H), 3.83 (d, $J=13.8$ Hz, 1H), 3.65 (d, $J=13.8$ Hz, 1H), 3.47 (t, $J=4.8$ Hz, 2H), 3.29 (s, 3H); ^{13}C NMR (CDCl_3) δ 164.7, 143.0, 131.8, 129.4, 124.2, 69.9, 64.8, 61.5, 58.9; HRMS (EI) m/z (M^+) calcd for $\text{C}_{11}\text{H}_{14}\text{O}_4\text{S}$ 242.0613, found 242.0612.

Phenylsulfinyl-2-ethoxy-2-oxoethyl ester (8ae). Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (5 : 1). Colorless oil: 48 mg, 59% yield; ^1H NMR (CDCl_3) δ 7.63–7.61 (m, 2H), 7.48–7.47 (m, 3H), 4.57 (s, 2H), 4.17 (q, $J=7.2$ Hz, 2H), 3.84 (d, $J=13.8$ Hz, 1H), 3.72 (d, $J=13.8$ Hz, 1H), 1.22 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (CDCl_3) δ 167.0, 164.2, 142.9, 133.1, 129.4, 124.1, 61.7, 61.6, 61.5, 14.0; HRMS (EI) m/z (M^+) calcd for $\text{C}_{12}\text{H}_{14}\text{O}_5\text{S}$ 270.0562, found 270.0554.

Benzyl 2-(phenylsulfinyl)acetate (8af). Chromatographic purification in a silica gel column with *n*-hexane–AcOEt (5 : 1). Colorless oil: 21 mg, 25% yield; ^1H NMR (CDCl_3) δ 7.59–7.56 (m, 3H), 7.30–7.20 (m, 5H), 7.44–7.42 (m, 2H), 5.05 (s, 2H), 3.84 (d, $J=13.5$ Hz, 1H), 3.64 (d, $J=13.5$ Hz, 1H); ^{13}C NMR (CDCl_3) δ 164.5, 143.0, 134.7, 131.8, 129.4, 128.7, 128.6, 128.5, 124.2, 67.7, 61.6; HRMS (EI) m/z (M^+) calcd for $\text{C}_{15}\text{H}_{14}\text{O}_3\text{S}$ 274.0664, found 274.0657.

(*R*)-Ethyl 2-(*p*-tolylsulfinyl)acetate (8ba). ^{13}C Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (5 : 1). Colorless oil: 55 mg, 81% yield; ^1H NMR (CDCl_3) δ 7.51 (d, $J=8.1$ Hz, 2H), 7.27 (d, $J=8.1$ Hz, 2H), 4.09 (q, $J=6.9$ Hz, 2H), 3.80 (d, $J=13.5$ Hz, 1H), 3.57 (d, $J=13.5$ Hz, 1H), 2.36 (s, 3H), 1.15 (t, $J=7.2$ Hz, 3H); ^{13}C -NMR (CDCl_3): $\delta=14.0, 21.5, 61.7, 62.0, 124.2, 130.0, 139.8, 142.4, 164.8$; HRMS (EI) m/z (M^+) calcd for $\text{C}_{11}\text{H}_{14}\text{O}_3\text{S}$ 226.0664, found 226.0660; ee > 99%, $[\alpha]_{\text{D}}^{25} +155.3$ (c: 1, acetone). (Chiralcel OD-H, Hexane: 2-Propanol 99:1, flow 2 mL/min).

***N*-(2-Hydroxyethyl)-2-(phenylsulfinyl)acetamide (10aa).** Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 65 mg, 96% yield; mp 101–104 °C; ^1H -NMR (CDCl_3) δ 7.59–7.55 (m, 2H), 7.49–7.47 (m, 3H), 3.67 (d, $J=13.8$ Hz, 1H), 3.60 (q, $J=4.8$ Hz, 2H), 3.50 (d, $J=13.8$ Hz, 1H), 3.33 (q, $J=5.4$ Hz, 2H); ^{13}C -NMR (CDCl_3) δ

164.2, 141.4, 131.7, 129.5, 124.0, 61.4, 59.9, 42.6; HRMS (E.I.) m/z (M^+) calcd. for $C_{10}H_{13}NO_3S$ 227.0616, found 227.0626.

***N,N*-Bis(2-hydroxyethyl)-2-(phenylsulfinyl)acetamide (10ab).** Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 77 mg, 95% yield; mp 101–104 °C; 1H -NMR ($CDCl_3$) δ 7.65–7.63 (m, 2H), 7.47–7.45 (m, 3H), 4.20 (d, J = 14.1 Hz, 1H), 3.81 (d, J = 14.1 Hz, 1H), 3.71–3.47 (m, 10H); ^{13}C -NMR ($CDCl_3$) δ 166.1, 142.4, 131.6, 129.4, 124.4, 72.2, 61.6, 60.0, 52.6, 50.4; HRMS (E.I.) m/z (M^+) calcd. for $C_{12}H_{17}NO_4S$ 271.0878, found 271.0885.

(*R*)-2-(*p*-tolylsulfinyl)-*N*-(2-hydroxyethyl)acetamide (10ba). Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1 to 1 : 2). White solid: 73 mg, 99% yield; mp 123–125 °C; 1H -NMR ($CDCl_3$) δ 7.44 (d, J =8.1Hz, 2H), 7.24 (d, J =8.1Hz, 2H), 3.65–3.27 (m, 7H), 2.33 (s, 3H); ^{13}C -NMR ($CDCl_3$) δ 164.3, 142.3, 138.3, 130.1, 124.1, 61.1, 60.8, 42.5, 21.4; HRMS (E.I.) m/z (M^+): calcd. for $C_{11}H_{15}NO_3S$ 241.0773, found 241.0771; $[\alpha]_D^{25}$ +198.1 (c: 1, acetone), ee > 99% (Chiralcel IA, Hexane: 2-Propanol 90:10, flow 2 mL/min).

Oxazolidin-2-one (11aa). Obtained under aerobic conditions. Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1). White solid: 41 mg, 78% yield; mp 86–91 °C; 1H -NMR ($CDCl_3$) δ 6.40–6.10 (br s, 1H), 4.37 (t, J = 7.9 Hz, 2H), 3.55 (t, J = 7.9 Hz, 2H); ^{13}C -NMR ($CDCl_3$) δ 160.7, 64.9, 40.6; HRMS (E.I.) m/z (M^+) calcd. for $C_3H_5NO_2$ 87.0320, found 87.0299

3-(2-Hydroxyethyl)oxazolidin-2-one (11ab). Obtained under aerobic conditions. Chromatographic purification in a silica gel column with *n*-hexane–ethyl acetate (1 : 1). Colorless oil: 61 mg, 77% yield; 1H -NMR ($CDCl_3$) δ 4.32 (t, J = 7.9 Hz, 2H), 3.75–3.65 (m, 4H), 3.60–3.50 (brs, 1H), 3.36 (t, J = 5.6 Hz, 2H); ^{13}C -NMR ($CDCl_3$) δ 159.3, 62.1, 59.9, 46.6, 45.4; HRMS (E.I.) m/z (M^+) calcd. for $C_5H_9NO_3$ 131.0582, found 131.0611.

Acknowledgment This work was supported by the Spanish Ministerio de Ciencia e Innovación and European Community Funds (FEDER) Grants (CTQ 2010-19999) and Consolider-Ingenio2010 (CSD2007-00006). We acknowledge the SCSIE (Universidad de Valencia) for access to the instrumental facilities.

Supporting information

Copies of ^1H and ^{13}C NMR spectra for all compounds and HPLC chromatograms for chiral compounds. The crystallographic information file (CIF) contains the supplementary crystallographic data for this paper. This material is available free of charge over the Internet at <http://pubs.acs.org>.

References

- ¹ *Transition Metals for Organic Synthesis* M. Beller, C. Bolm, Eds.; Wiley-VCH: Weinheim, 2004; vol. 2, 437.
- ² Reviews: (a) Grigg, R.; Mutton, S. P. *Tetrahedron* **2010**, *66*, 5515. (b) Brennführer, A.; Neumann, H.; Beller, M. *Angew. Chem., Int. Ed.* **2009**, *48*, 4114; (c) Brennführer, A.; Neumann, H.; Beller, M. *ChemCatChem* **2009**, *1*, 28. (d) Banard, C.F.J. *Organometallics* **2008**, *27*, 5402.
- ³ (a) Cho, C. H.; Larock, R. C. *Tetrahedron Lett.* **2010**, *51*, 3417. (b) Rami, S.; Bassam, E. A. *Tetrahedron Lett.* **2010**, *51*, 3211. (c) Schareina, T.; Zapf, A.; Cotte, A.; Gotta, M.; Beller, M. *Adv. Synt & Cat.* **2010**, *352*, 1205. (d) Gabriele, B.; Veltri, L.; Mancuso, R.; Plastina, P.; Salerno, G.; Costa, M. *Tetrahedron Lett.* **2010**, *51*, 1663. (e) Salvadori, J.; Balducci, E.; Zaza, S.; Petricci, E.; Taddei, M. *J. Org. Chem.* **2010**, *75*, 1841. (f) Yamamoto, Y. *Adv. Synt & Cat.* **2010**, *352*, 478. (g) Munday, R. H.; Martinelli, J. R.; Buchwald, S. L. *J. Am. Chem. Soc.* **2008**, *130*, 2754.
- ⁴ (a) Wakita, Y.; Yasunaga, T.; Akita, M.; Kojima, M. *J. Organomet. Chem.* **1986**, *301*, C17. (b) Kondo, T.; Tsuji, Y.; Watanabe, Y. *J. Organomet. Chem.* **1988**, *345*, 397. (c) Suzuki, A. *Pure Appl. Chem.* **1994**, *66*, 213. (d) Ishikura, M.; Terashima, M. *J. Org. Chem.* **1994**, *59*, 2634. (e) Ishiyama, T.; Kizaki, H.; Hayashi, T.; Suzuki, A.; Miyaura, N. *J. Org. Chem.* **1998**, *63*, 4726. (f) Andrus, M. B.; Ma, Y.; Zang, Y.; Song, C. *Tetrahedron Lett.* **2002**, *43*, 9137. (g) Mingji, D.; Liang, B.; Wang, C.; You, Z.; Xiang, J.; Dong, G.; Chen, J.; Yang, Z. *Adv. Synth. Catal.* **2004**, *346*, 1669. (h) Petz, A.; Péczely, G.; Pintér, Z.; Kollár, L. *J. Mol. Catal. A: Chem.* **2006**, *255*, 97. (i) Larini, P.; Guarna, A.; Occhiato, E. G. *Org. Lett.* **2006**, *8*, 781. (j) Maerten, E.; Sauthier, M.; Mortreux, A.; Castanet, Y. *Tetrahedron* **2007**, *63*, 682. (k) Maerten, E.; Hassouna, F.; Couve-Bonnaire, S.; Montreux, A.; Carpentier, J.-F.; Castanet, Y. *Synlett* **2003**, 1874.
- ⁵ (a) Ozawa, F.; Soyama, H.; Yanagihara, H.; Aoyama, I.; Takino, H.; Izawa, K.; Yamamoto, T.; Yamamoto, A. *J. Am. Chem. Soc.* **1985**, *107*, 3235. (b) Yamamoto, A. *Bull. Soc. Chem. Jpn.* **1985**, *68*, 433.
- ⁶ (a) Wu, X.-F.; Neumann, H.; Beller, M. *Chem. Eur. J.* **2012**, *18*, 419. (b) Wu, X.-F.; Neumann, H.; Beller, M. *Chem. Eur. J.* **2010**, *16*, 12104. (c) Wu, X.-F.; Neumann, H.; Beller, M. *ChemCatChem* **2010**, *2*, 509. (d) Wu, X.-F.; Schranck, J.; Neumann, H.; Beller, M. *Tetrahedron Lett.* **2011**, *52*, 3702. (e) Martinelli, J. R.; Clark, T. P.; Watson, D. A.; Munday, R. H.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2007**, *46*, 8460.
- ⁷ Ishiyama, T.; Miyaura, N.; Suzuki, A. *Tetrahedron Lett.* **1991**, *47*, 6923
- ⁸ (a) Lindsell, W. E.; Palmer, D. D. D.; Preston, P. N.; Rosair, G. M.; Jones, R. V. H.; Whitton, A. J. *Organometallics* **2005**, *24*, 1119. (b) Lapidus, A. L.; Eliseev, O. L.; Bondarenko, T. N.; Sizan, O. E.; Ostapenko, E. G.; Beletskaya I. P. *Kinet. Catal.* **2004**, *45*, 234. (c) Cavinato, G.; Pasqualetto, M.; Ronchin, L.; Tonilo, L. *J. Mol. Catal. A: Chem.* **1997**, *125*, 15. (d) Cavinato, G.; Tonilo, L. *J. Mol. Catal. A: Chem.* **1996**, *104*, 221.
- ⁹ (a) Troisi, L.; Granito, C.; Rosato, F.; Videtta, V. *Tetrahedron Lett* **2010**, *51*, 371. (b) For intramolecular versions see: Wu, X.-F.; Schranck, J.; Neumann, H.; Beller, M. *ChemCatChem* **2012**, *4*, 69. (c) Marosvölgyi-Hask, D.; Attila Takács, A.; Riedl, S.; Kollar, L. *Tetrahedron* **2011**, *67*, 1036.
- ¹⁰ (a) Cinquini, M.; Landini, D.; Maia, A. *J. Chem. Soc. Chem. Commun.* **1972**, 734. (b) Numata, T.; Oae, S. *Bull. Soc. Chem. Jpn.* **1972**, *45*, 2794.
- ¹¹ (a) García-Ruano, J. L.; Alvarado, C.; Díaz-Tendero, S.; Alemán, J. *Chem.-Eur. J.* **2011**, *17*, 4030. (b) Obringer, M.; Barbarotto, M.; Choppin, M. S.; Colobert, F. *Org. Lett.* **2009**, *11*, 3542. (c) Colobert, F.; Choppin, S.; Ferreiro-Mederos, L.; Obringer, M.; Luengo-Arratta, S.; Urbano, A.; Carreño, M. C. *Org. Lett.* **2007**, *9*, 4451. (d)

- Ishigami, K.; Katsuta, R.; Watanabe, H. *Tetrahedron* **2006**, *62*, 2224. (e) Carreño, M. C.; Hernández-Torres, G.; Urbano, A.; Colobert, F. *Org. Lett.* **2005**, *7*, 5517. (f) Lanners, S.; Khiri, N.; Solladié, G.; Hanquet, G. *Tetrahedron Lett.* **2005**, *46*, 619. (g) García-Ruano, J. L.; Rodríguez-Fernández, M. M.; Maestro, M. C. *Tetrahedron* **2004**, *60*, 5701. (h) Silveira, C. C.; Braga, A. L.; Kaufman, T. S.; Lenardao, E. J. *Tetrahedron* **2004**, *60*, 8295. (i) Carreño, M. C.; Des Mazery, R.; Urbano, A.; Colobert, F.; Solladié, G. *J. Org. Chem.* **2003**, *68*, 7779. (j) Nakamura, S.; Nakayama, J.; Toru, T. *J. Org. Chem.* **2003**, *68*, 5766. (k) Ghosh, A. K.; Lei, H. *Tetrahedron: Asymmetry* **2003**, *14*, 629. (l) Colobert, F.; Des Mazery, R.; Solladié, G.; Carreño, M. C. *Org. Lett.* **2002**, *4*, 1723. (m) Nokami, J.; Kataoka, K.; Shiraishi, K.; Osafune, M.; Hussain, I.; Sumida, S. *J. Org. Chem.* **2001**, *66*, 1228. (n) Toró, A.; L'Heureux, A.; Deslongchamps, P. *Org. Lett.* **2000**, *2*, 2737. (o) García-Ruano, J. L. *Top. Curr. Chem.* **1999**, *204*, 1. (p) Carreño, M. C. *Chem. Rev.* **1995**, *95*, 1717.
- ¹² (a) Kankan, S.; Fun, H. K. *J. Chem. Res.* **2007**, 668. (b) García-Ruano, J. L.; Gonzalez, A. M.; Lopez-Solera, I.; Masaguer, J. R.; Navarro-Ranninger, C.; Raithby, P. R.; Rodríguez, J. H. *Angew. Chem., Int. Ed.* **1995**, *34*, 1351. For further examples of complexes with sulfoxide ligands: (c) Chen, M. S.; Prabakaran, N.; Labenz, N. A.; White, M. C. *J. Am. Chem. Soc.* **2005**, *127*, 6970. (d) Pettinari, C.; Pellei, M.; Cavicchio, G.; Crucianelli, M.; Panzeri, W.; Colapietro, M.; Cassetta, A. *Organometallics* **1999**, *18*, 555. (e) Ogura, K.; Aizawa, T.; Uguiyama, K.; Iida, H. *Bull. Soc. Chem. Jpn.* **1983**, *56*, 953. (f) Price, J. H.; Williamson, A. N.; Schramm, R. F.; Wayland, B. B. *Inorg. Chem.* **1972**, *11*, 1280.
- ¹³ Nagao, Y.; Miyamoto, S.; Miyamoto, M.; Takeshige, H.; Hayashi, K.; Sano, S.; Shiro, M.; Yamaguchi, K.; Sei, Y. *J. Am. Chem. Soc.* **2006**, *128*, 9722.
- ¹⁴ Medio-Simon, M.; Mollar, C.; Rodriguez, N.; Asensio, G. *Org. Lett.* **2005**, *7*, 4669.
- ¹⁵ (a) Hojo, M.; Masuda, R.; Saeki, T.; Fujimori, K.; Tsutsumi, S. *Synthesis* **1977**, 789. (b) Cardellicchio, C.; Fiandanese, V.; Naso, F.; Scilimati, A. *Tetrahedron. Lett.* **1992**, *33*, 5121.
- ¹⁶ Corey, E. J.; Chavkovsky, M. *J. Am. Chem. Soc.* **1965**, *87*, 1345.
- ¹⁷ (a) Kirchhoff, J. H.; Netherton, M. R.; Hill, I. D.; Fu, G. C. *J. Am. Chem. Soc.* **2002**, *124*, 13662. (b) Hill, I. D.; Netherton, M. R.; Fu, G. C. *Angew. Chem., Int. Ed.* **2003**, *42*, 5749. (c) Zhou, J.; Fu, G. C. *J. Am. Chem. Soc.* **2003**, *125*, 12527. (d) Zhou, J.; Fu, G. C. *J. Am. Chem. Soc.* **2003**, *125*, 14726. (e) Netherton, M. R.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, *41*, 3910. (f) Kirchhoff, J. H.; Dai, C.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, *41*, 1945. (g) Netherton, M. R.; Dai, C.; Neuschütz, K.; Fu, G. C. *J. Am. Chem. Soc.* **2001**, *123*, 10099. (h) Frisch, A. C.; Beller, M. *Angew. Chem., Int. Ed.* **2005**, *44*, 674.
- ¹⁸ (a) Rodríguez, N.; Cuenca, A.; Ramírez de Arellano, C.; Medio-Simón, M.; Peine, D.; Asensio, G. *J. Org. Chem.* **2004**, *69*, 8070. (b) Rodríguez, N.; Cuenca, A.; Ramírez de Arellano, C.; Medio-Simón, M.; Asensio, G. *Org. Lett.* **2003**, *5*, 1705. (c) Rodríguez, N.; Ramírez de Arellano, C.; Asensio, G.; Medio-Simón, M. *Chem. Eur. J.* **2007**, *13*, 4223. (d) Gurlaouen, C.; Ujaque, G.; Lledós, A.; Medio-Simón, M.; Asensio, G.; Maseras, F. *J. Org. Chem.* **2009**, *74*, 4049.
- ¹⁹ For demonstration of diffusion of air into the balloon see Lee, J.-Y.; Yoo, H.-S.; Park, J. S.; Hwang, K.-J.; Kim, J. S. *J. Chem. Educ.* **2005**, *82*, 288.
- ²⁰ (a) Ragaini, F. *Dalton Trans.* **2009**, 6251. (b) Gabriele, B.; Salerno, G.; Mancuso, R.; Costa, M. *J. Org. Chem.* **2004**, *69*, 4741. (c) Gabriele, B.; Salerno, G.; Mancuso, R.; Costa, M. *Chem. Commun.* **2003**, 486.
- ²¹ Cowell, A.; Stille, J. F. *J. Am. Chem. Soc.* **1980**, *102*, 4193.
- ²² Hu, Y.; Liu, J.; Lu, Z.; Luo, X.; Zhang, H.; Lan, Y.; Lei, A. *J. Am. Chem. Soc.* **2010**, *132*, 3153.
- ²³ Rodriguez, N.; Medio-Simón, M.; Asensio, G. *Adv. Synth. Cat.* **2007**, *349*, 987.
- ²⁴ Steinhoff, B. A.; Fix, S. R.; Stahl, S. S. *J. Am. Chem. Soc.* **2002**, *124*, 766.
- ²⁵ (a) Roussel, M.; Mimoun, H. *J. Org. Chem.* **1980**, *45*, 5387. (b) Nashimura, T.; Onoue, T.; Ohe, K.; Uemura, S. *J. Org. Chem.* **1999**, *64*, 6750. (c) Remias, J. E.; Sen, A. *J. Mol. Catal. A: Chem.* **2002**, *189*, 33.
- ²⁶ For examples of *trans*-chelating xantphos-Pd(II) complexes: (a) Yin, J.; Buchwald, S. L. *J. Am. Chem. Soc.* **2002**, *124*, 6043. (b) Grushin, V. V.; Marshall, W. J. *J. Am. Chem. Soc.* **2006**, *128*, 12644.
- ²⁷ Cinquini, M.; Colonna, S. *J. Chem. Soc. Perkin 1* **1972**, 1883.
- ²⁸ Huang, C. G.; Chang, B. R.; Chang, N. C. *Tetrahedron Lett.* **2002**, *43*, 2721.
- ²⁹ Bigi, F.; Frullanti, B.; Maggi, R.; Sartori, G.; Zamboni, E. *J. Org. Chem.* **1999**, *64*, 1004.
- ³⁰ Pasha, M. A.; Jayashankara, V. P. *Synth. Comm.* **2006**, *36*, 1787.
- ³¹ Tsuda, T.; Isegawa, Y.; Saegusa, T. *J. Org. Chem.* **1972**, *37*, 2670.
- ³² Crumrine, D. S.; Dieschbourg, T. A.; O'Toole, J. G.; Tassone, B. A.; Vandeburg, S. C. *J. Org. Chem.* **1987**, *52*, 3699.
- ³³ Lemoucheux, L.; Seitz, T.; Rouden, J.; Lasne, M. C. *Org. Lett.* **2004**, *6*, 3703.
- ³⁴ Zhang, Q.; Wu, Y. *Tetrahedron*, **2007**, *63*, 10189.