View Article Online View Journal



Organic & Biomolecular Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: M. März, J. Chudoba, M. Kohout and R. Cibulka, *Org. Biomol. Chem.*, 2017, DOI: 10.1039/C6OB02770A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the **author guidelines**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the ethical guidelines, outlined in our <u>author and reviewer resource centre</u>, still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



rsc.li/obc

Journal Name



Photocatalytic Esterification under Mitsunobu Reaction Conditions Mediated by Flavin and Visible Light †

Received 00th January 20xx, Accepted 00th January 20xx

M. März,^a J. Chudoba,^b M. Kohout,^a and R. Cibulka*^a

DOI: 10.1039/x0xx00000x

www.rsc.org/

Published on 03 February 2017. Downloaded by University of Newcastle on 04/02/2017 16:22:39

The usefulness of flavin-based aerial photooxidation in Mitsunobu esterification under reaction conditions was demonstrated, providing aerial dialkyl azodicarboxylate recycling/generation from the corresponding dialkyl hydrazine dicarboxylate. Simultaneously, activation of triphenylphosphine (Ph₃P) by photoinduced electron transfer from flavin allows azoreagent-free esterification. An optimized system with 3methylriboflavin tetraacetate (10%), oxygen (terminal oxidant), visible light (450 nm), Ph_3P , and dialkyl hydrazine dicarboxylate (10%) has been shown to provide efficient and stereoselective coupling of various alcohols and acids to esters with retention of configuration.

A systems of triphenylphosphine (Ph₃P) or a related phosphine activated by an oxidant allow efficient esterification under mild conditions.¹ In the Mitsunobu reaction,^{2,3} PPh₃ is oxidized by a dialkyl azodicarboxylate (usually DEAD [**1a**] or DIAD [**1b**]) to form a betain species, which facilitates transformation of an alcohol to a reactive alkoxyphosphonium intermediate (Scheme 1). The latter undergoes $S_N 2$ substitution with carboxylate giving an ester with inversion of configuration. There are a few cases in which an acyloxyphosphonium intermediate predominates under Mitsunobu reaction conditions, subsequent $S_N Ac$ substitution of which gives the product with retention of configuration.

The Mitsunobu reaction has become an extremely useful tool in organic synthesis.^{3b,7} However, its further expansion, in particular towards large-scale applications, is limited by the need for a stoichiometric amount of oxidant, azodicarboxylate **1**, which is toxic and unstable and the use of which generates dialkyl hydrazine dicarboxylate **2** as a waste by-product. An attempt to solve this problem led to recent pioneering studies on catalytic Mitsunobu reaction.^{1b} Toy and co-workers applied a stoichiometric amount of the oxidant PhI(OAc)₂ to convert the generated hydrazine **2a** back to azo compound **1a**,

This journal is C The Royal Society of Chemistry 20xx

allowing only 10 mol% 1a to be used.8 Taniguchi, et al developed and optimized catalytic Mitsunobu reaction using ethyl N-aryl-azocarboxylates 1e (10 mol%) instead of 1a or 1b, in conjunction with a catalytic amount (10 mol%) of Fe(II)phthalocyanine (Fe[Pc]), which re-oxidized arylhydrazine, being simultaneously re-generated by air oxygen.⁹ Notably, Fe[Pc] was not able to recycle original Mitsunobu reagents, DEAD or DIAD, due to high oxidation potential of corresponding hydrazines 2. Another approach is to use a procedure free from azo reagent 1 in which PPh₃ is activated by aerial oxidation catalyzed by Fe[Pc]. This esterification occurs via an acylphosphonium intermediate giving an ester with retention of configuration.¹⁰ Another problem of the Mitsunobu type reactions is bulk production of phosphine oxide. O'Brien¹¹ and later Aldrich and Buonomo¹² showed that phosphine oxide can be reduced in situ with phenylsilane, thus allowing use of the phosphine reagent in a catalytic amount.¹³



^{a.} Department of Organic Chemistry, University of Chemistry and Technology,

Prague, Technická 5, 166 28 Prague, Czech Republic.

^{b.} Central Laboratories, University of Chemistry and Technology, Prague, Technická 5, 166 28 Prague, Czech Republic.

⁺Electronic Supplementary Information (ESI) available: [experimental section and copies of NMR spectra and HPLC chromatograms]. See DOI: 10.1039/x0xx00000x

Scheme 1. Catalytic esterification mediated by activated Ph₃P under Mitsunobu (A) and azo-reagent-free (B) conditions.

COMMUNICATION

Published on 03 February 2017. Downloaded by University of Newcastle on 04/02/2017 16:22:39

We surmise that, due to their redox character, Mitsunobu-type reactions represent a typical challenge for photoredox catalysis,¹⁴ which have undergone rapid development in recent years. Upon excitation, molecules become stronger oxidizing agents compared to their ground-state forms, which considerably extends the possibilities for regeneration of Mitsunobu reagents or alternative activation of phosphines. However, to the best of our knowledge, no application of photoredox catalysis in catalytic Mitsunobu reactions or phosphine-mediated esterifications has hitherto been reported. Herein, we present a photocatalytic system (Scheme 1) based on flavin 3d (a derivative of vitamin B2 [3a]), oxygen, and visible light, which is able to regenerate commercial dialkyl azodicarboxylates 1. The system was found to simultaneously activate Ph₃P by photoinduced electrontransfer, thus providing azo-reagent-free esterification.

Our original idea to regenerate azodicarboxylates 1 from corresponding hydrazides 2 led us to flavin the photocatalysts.^{15,16} Upon excitation with blue light (450 nm), flavins become oxidizing agents¹⁷ (E_{red} *=1.67 V vs. SCE for riboflavin tetraacetate (3b)), strong enough to oxidize 2 (a value of E_{ox} *=1.62 V vs. SCE has been reported for BocNHNHBoc¹⁸), but not so strong as to mediate undesired oxidation of alcohols, even when they are activated, e.g. benzyl alcohols, which are often substrates of Mitsunobu reactions. The only exceptions are electron-rich benzyl alcohols such as 4-methoxybenzyl alcohol (E_{ox} *=1.43 V vs. SCE¹⁹), which is a traditional substrate for testing flavins in photooxidations.^{16e,17,20} Importantly, flavins can be converted back to their oxidized forms by oxygen, and can thus be applied in catalytic amounts (cf. Scheme 2).¹⁵

Preliminary experiments confirmed that 3b oxidizes dialkyl hydrazine-dicarboxylates 2a-d to the corresponding azo compounds 1 when irradiated by light of wavelength 450 nm (see ESI). The oxidation proceeded remarkably in acetonitrile, and thus we used this solvent in model photocatalytic esterification with 3-nitrobenzoic acid 4a and benzyl alcohol 5a using 10 mol% of **3b**, 10 mol% of **1b**, and 2 equivalents of PPh₃. The reaction was performed at 25 °C under oxygen and irradiation with a blue diode (450 nm), in the presence of molecular sieves (4 Å) to remove hydrogen peroxide formed during flavin regeneration.9c To our delight, we observed formation of ester 6a in moderate conversion after 24 h (entry 1), and a positive result was found even when 2b (10 mol%) was used instead of 1b (entry 2). Only a small amount of aldehyde was formed with 3b, confirming the suitability of flavins in this reaction. Use of stronger oxidants, such as 9mesityl-10-methylacridinium perchlorate (7; E_{red} *=2.08 V vs. SCE^{14a}) or triphenylpyrylium tetrafluoroborate (**8**; E_{red} *=2.45 V vs. SCE^{14a}) preferentially led to benzylic oxidation (entries 3 and 4). Notably, the N-H bond in flavins is sufficiently acidic for 3b to be a substrate of Mitsunobu reaction. We observed formation of N-benzyl derivative 3c, which was still active in esterification, but less than 3b (cf. entries 2 and 5). Thus, Nmethylflavin 3d seemed to be a good choice of photocatalyst (entries 6 and 7).

Journal Name

Page 2 of 5

Monitoring the reaction course showed undesired direct oxidation of Ph₃P to Ph₃P=O, which retarded ester formation. We assume that excited flavin participates in this oxidation by electron transfer from Ph_3P ($E_{ox} = 1.06$ vs. SCE^{21}) (for quenching experiment, see ESI). The formed Ph₃P^{·+} reacts with oxygen to form phosphine oxide, which is known to occur by several mechanisms.²² This side reaction can be suppressed by subsequent addition of PPh₃, which increases the yield significantly (entry 8). A similar effect was achieved by adding $PhSiH_3^{12}$ for in situ reduction of $Ph_3P=O$ back to Ph_3P (entry 9). Notably, positive effects of elevated temperature on the reaction in both the absence and presence of PhSiH₃ were observed, achieving yields of up to 92% (entries 10 and 11). It should be noted that formation of **6a** was not observed in blank experiments (see ESI) in the absence of the flavin photocatalyst, light, Ph₃P or molecular sieves.[‡] On the other hand, a little 6a was formed with omission of the azo compound or its precursor, irrespective of the presence of PhSiH₃ (entries 12 and 13). The amount of ester formed by the azo-free process was increased at elevated temperature (entries 14 and 15).

Taking into account conversions of ester **6a** achieved in the presence and absence of **2b** (cf. entries 7 vs. 12, 9 vs. 13, 10 vs. 14, and 11 vs. 15), the initial results gave evidence that: i) after irradiation, flavin **3d** can mediate a Mitsunobu reaction that is catalytic in azo reagent by virtue of its generation/recycling from the corresponding hydrazine; ii) another mechanistic pathway not requiring the azo component is involved, especially at elevated temperature. Such a "background" reaction has been analogously observed using recycling systems with PhI(OAc)₂⁸ and Fe(II)-phthalocyanine.^{9c}

Table 1. Optimization of protocol for photocatalytic Mitsunobu reaction with recycling/generation of ${\bf 1}.$

⁰ 2 ^N CO 4a	он + ОН 5а	$\begin{tabular}{lllllllllllllllllllllllllllllllllll$	O₂N _	6a	
Entry	Catalytic	Temp	Conversion after 24h [%]		
	system	[°C]	Ester	Aldehyde	
1	3b/1b	25	58	2	
2	3b/2b	25	48	2	
3	7/2b	25	46	31	
4	8/2b	25	8	20	
5	3c/2b	25	41	7	
6	3d/1b	25	64	2	
7	3d/2b	25	60	2	
8 ^{<i>a</i>}	3d/2b	25	79	trace	
9 ^b	3d/2b	25	85	trace	
10	3d/2b	50	66	trace	
11 ^b	3d/2b	50	92	trace	
12	3d/-	25	17	2	
13 ^b	3d/-	25	26	-	
14	3d/-	50	47	2	
15 ^b	3d/-	50	69	-	

 a PPh_3 added in three portions at the time 0, 4 and 8 hours. b PhSiH_3 (2 equiv.) added.

Journal Name

Next, we examined the substrate scope of photocatalytic esterification alternating various alcohols **5** (Table 2) and acids **4** (Table 3) under selected conditions: i) without PhSiH₃ with 10% of **1b** at 25 °C, characterized by a major contribution from the azo-compound-mediated reaction (Method I, analogous to entry 6 in Table 1), and ii) with PhSiH₃ and 10% of **2b** at 50 °C, whereupon the non-azo-reagent-free pathway predominated (Method II, analogous to entry 11 in Table 1). In the Tables 2 and 3, the efficacy of both methods is characterized by conversions and preparative yields of esters after 24 h. Conversions of photocatalytic esterifications in the absence of azo reagent **1b** or its precursor **2b** are given for comparison to estimate the contribution of azo-reagent-free pathway (blank).

Method I (Table 2, odd entries) provided moderate to good conversions and yields of esters **6a–h** by esterification of 3-nitrobenzoic acid **(4a)** with substituted benzyl alcohols regardless of the character and position of the substituent.

Table 2. Substrate scope of photocatalytic esterification (various alcohols).							
Entry	Product	Method ^a	Conv. [%]	Yield ^b [%]	Conv. (blank) ^c [%]		
1		I	64 ^d	53	17		
2	6a 6a	П	92	82	69		
3		I	73	65	21		
4	6b Cl	П	quant.	85	80		
5		I	67	60	25		
6	6c NO ₂	П	quant.	79	80		
7	O ₂ N	I	64	54	60		
8	6d CF ₃	П	82	72	75		
9	O ₂ N	I	62 ^e	54	28		
10	6e Me	П	76 ^{<i>f</i>}	65	40		
11	O ₂ N	I	50 ^{<i>g</i>}	39	39		
12	6f OMe	П	85 ^{<i>h</i>}	71	68		
13		I	52	46	46		
14	6g	П	78	60	59		
15		I	53	44	43		
16	6h	П	78	71	68		
17	O ₂ N	I	18	n.d.	10		
18	6i	П	quant.	80	20		
19	02N C6H17	I	51	39	22		
20	6k	П	72	61	63		

^{*a*} Conditions for method I: $n(\mathbf{5}) = 0.15 \text{ mmol}$, $n(\mathbf{4}) = 0.18 \text{ mmol}$, $n(\mathbf{3d}) = 0.015 \text{ mmol}$, $n(\mathbf{1b}) = 0.015 \text{ mmol}$, $n(\mathbf{PPh}_3) = 0.3 \text{ mmol}$, MS 4 Å (150 mg), 2 ml CH₃CN, 455 nm, 25 °C, 24h; for method II: **2b** instead of **1b**; additionally 0.3 mmol of PhSiH₃; 50 °C. ^{*b*} Preparative yield. ^{*c*} Conversion of esterification in the absence of **1b** (method I) or **2b** (method II) from ¹H NMR data. ^{*d*} 80% after 72h. ^{*e*} 25% of aldehyde. ^{*f*} 18% of aldehyde. ^{*h*} 15% of aldehyde.

COMMUNICATION

Octyl ester **6k** (representative of esters with aliphatic alcohols) was also obtained in good yield by method? While ester **6** With 2-phenylethanol (**5**i) was formed in poor conversion only (entries 17 and 19). On the other hand, at elevated temperature (method II), high conversions and good to high yields of esters **6a**–**k** were achieved with all alcohols investigated (Table 2, even entries). As could be expected, significant amount of benzaldehyde (15–28%) was formed during esterification of 4-methyl- (**5e**) and 4-methoxybenzyl (**5f**) alcohols because of their low oxidation potentials (entries 9–12).¹⁹ Both methods proved to be effective for secondary alcohols, albeit requiring longer reaction times (Table 4).

Besides **4a**, other aromatic acids (4-nitrobenzoic **[4b]** and benzoic acid **[4c]**) as well as phenylalkanoic acids (phenylacetic **[4d]** and 3-phenylpropanoic acid **[4f]**) gave good to high yields of esters **6l–n** and **6p** with 4-chlorobenzyl alcohol (**5b**) by method II (Table 3, even entries). For these acids, method I seemed to be less efficient giving lower yields of **6n** and **6p** (entries 5 and 9) and almost no ester **6m** (entry 3). Interestingly, both methods afforded ester **6o** with sterically hindered α, α -disubstituted acid **4e** albeit in low yields and after prolonged reaction time (entries 7 and 8). Low efficacy of of our protocols was shown for esterification of hexanoic acid (**4g**) as a representative of non-substituted aliphatic acids (entries **11** and **12**). Method II was proved to be usefull also for phthalimide as a representative of N- nucleophile (entry **14**).

Table 3. Su	bstrate scope of photocatalytic e	esterification	(various ac	ids).	
Entry	Product	Method ^a	Conv. [%]	Yield ^b [%]	Conv. (blank) ^c [%]
1		I	30	19	27
2	O ₂ N 6I CI	Ш	quant.	79	80
3		I	5	n.d.	n.d.
4	6m CI	Ш	58	43	5
5		I	68	45	12
6	6n Cl	Ш	quant.	75	60
7		ľ	33	25	7
8	60 GO	Π^d	40	32	16
9		I	44	32	31
10	6p 6p	Ш	65	58	44
11	C-Hu O	I	8	n.d.	n.d.
12	6q CI	Ш	16	n.d.	11
13		I	6	n.d.	n.d.
14		П	66	57	20

^{*a*} Conditions for method I: n(5) = 0.15 mmol, n(4) = 0.18 mmol, n(3d) = 0.015 mmol, n(1b) = 0.015 mmol, $n(PPh_3) = 0.3$ mmol, MS 4 Å (150 mg), 2 ml CH₃CN, 455 nm, 25 °C, 24h; for method II: **2b** instead of **1b**; additionally 0.3 mmol of PhSiH₃; 50 °C. ^{*b*} Preparative yield. ^{*c*} Conversion of esterification in the absence of **1b** (method I) or **2b** (method II) from ¹H NMR data. ^{*d*} 72h.

COMMUNICATION

Published on 03 February 2017. Downloaded by University of Newcastle on 04/02/2017 16:22:39

Notably, a significant contribution of azodicarboxylate 1b or its precursor 2b was observed with most substrates (Tables 2 and 3, cf. conversions of methods I or II with azo-free pathway [blank]). This contribution became even more pronounced for esterification of some benzylic esters, e.g. 6a-6c and 6e (Table 2, entries 1, 3, 5 and 9), provided by method I or for some esterifications by method II, e.g. towards 6i (Table 2, entry 18), 6m and 9 (Table 3, entries 4 and 14). On the other hand, in some cases, especially for method II, azo-free reaction pathway predominates as indicated by high conversions of blank. The conversions of esters are indeed high under optimal conditions (method II) but not quantitative in many cases. The reason is probably bleaching of the flavin photocatalyst which was observed during esterifications. Similar photodecomposition of flavins was reported to occur also during other photocatalytic processes.^{15,16,20}

Irrespective of the method, highly stereoselective production of esters (Table 4) with retention of configuration was observed for two types of secondary alcohols, 1phenylethanol (5l, entries 1-4) and ethyl 2-lactate (5m, entries 5-8), indicating that our photocatalytic esterification occurs through alkoxyphosphonium **10** but through not acylphosphonium species 11 (see Scheme 2). This intermediate must also predominate in the photocatalytic "Mitsunobu reaction pathway" occurring through betain intermediate 12 (Scheme 2A), as indicated by the very high stereoselectivities observed with 5I at 25 °C (Method I), i.e. at the conditions where a significant contribution of the catalytic amount of 1b was detected (entries 1 and 3, cf. conversions in the presence and the absence of 1b). Most probably, our photocatalytic the conditions esterification just met favoring acyloxyphosphonium species 11 in the delicate equilibrium.⁵ acyloxy/alkoxyphosphonium Notably, the inversion product predominated (e.r. = 85:15[§]) in standard

Table 4. Stereoselectivity of photocatalytic esterification. Conv Yield er^{d} Entry Alcohol Product Method [%] [%] 1 А 46/7 30 98:2 CH 2 (R)**-5** В 76/61^e 65 99:1 3 A 44 /7 25 99:1 4 В 70/61 59 99:1 5 48/44 39 99:1 Δ (S)-6s 6 В 98:2 58/49 50 7 А 45/44 38 97:3 8 R 60/36 48 98:2

^a For conditions, see Table 2; reaction time: 72h. ^b Conversion of esterification in the presence/in the absence of 1b (method I) or 2b (method II) from ¹H NMR data. ^c Preparative yields. ^d From HPLC, see ESI. ^e e.r. 98:2 was found in the absence of 2b

Based on literature data and our own experimental results, we propose a mechanism for the pathway not involving the azo compound or its hydrazine precursor (Scheme 2B). Excited flavin **FI**^{*} oxidizes Ph_3P to Ph_3P^{+} , as it is evident from the redox potentials (see above) and confirmed by the observation of efficient emission quenching ($K_s = 22 \text{ Lmol}^{-1}$; for the Stern– Volmer plot, see ESI). The flavin radical anion (pK_a of conjugated acid is 8.4, ref.²³) is immediately protonated by a carboxylic acid to form radical **FIH**. Ph₃P⁺⁺ then reacts with an alcohol to form (after subsequent oxidation and deprotonation) an alkoxyphosphonium species 10, which is in equilibrium with the corresponding acyloxyphosphonium species 11. Nevertheless, 11 may also be formed directly from Ph₃P⁺⁺ and carboxylate.^{§§} Finally, **11** undergoes substitution with alkoxide to form ester and the reduced flavin FIH₂ is reoxidized by oxygen in a dark procedure.^{17,20a,24} Hydrogen peroxide, formed as a by-product, is decomposed by molecular sieves.⁹⁰

Quantum yields of ester 6a production by methods I and II was found to be 0.04 and 0.07, respectively, thus supporting closed catalytic cycle and indicating that an open radical propagation mechanism is not involved. We also not observed corresponding anhydride when monitoring the reaction mixture with 4b and 5b by ¹H NMR. Nevertheless presence of by-product/intermediate^{3b,5} this alternative in low concentration cannot be excluded.





Conclusions

In conclusion, we have shown for the first time that the original Mitsunobu reagents, DIAD (1b) and DEAD (1a), can be regenerated from the corresponding hydrazines bv photocatalytic and/or organocatalytic system,²⁵ thus allowing

Journal Name

⁽stoichiometric) Mitsunobu esterification of 51, but Ait is not formed exclusively indicating not only 5,12.1979 Charris and starting formed exclusively indicating not only 5,12.1979 Charrier and the second starting indicating indicating the second starting indicating indic involved.

COMMUNICATION

Journal Name

esterifications that are catalytic in **1**. The method is based on visible light and readily available riboflavin derivative **3d**, which is used in a catalytic amount, being recycled with molecular oxygen, an inexpensive and green terminal oxidant. The only drawback is side photooxidation of Ph_3P to $Ph_3P=O$ under aerial conditions, which can be eliminated by in-situ back-reduction. We have observed a completely new esterification pathway being involved in our photocatalytic esterification that occurs without any contribution from the azo compound **1** or hydrazine **2**. It proceeds through Ph_3P^{-+} generated by photoinduced electron transfer to flavin. To the best of our knowledge, no similar synthetic application of photochemically generated Ph_3P^{++} has hitherto been reported. As there is still room for improvement, optimization of both concepts is currently being pursued in our laboratories.

Acknowledgement

This project was supported by the Czech Science Foundation (Grant No. 16-09436S). R.C. thanks the German National Science Foundation (GRK 1626 "Chemical Photocatalysis").

Notes and references

 \ddagger Notably only trace of ester was formed in stoichiometric Mitsunobu reaction in the presence of hydrogen peroxide (1 equiv.) in acetonitrile while 90% of ester was formed after addition of MS 4A thus demonstrating H₂O₂-quenching of Mitsunobu reaction. Hydrogen peroxide was not detected by iodometry after photocatalytic esterifications in the presence of MS 4A.

§ This value was not affected by either light or the flavin (see ESI).

\$\$ Notably analogous formation of alkoxyphosphonium and acyloxyphosphonium species was observed when Ph_3P^+ was generated electrochemically, see ref.26.

- a) A. Voituriez, A., N. Saleh *Tetrahedron Lett.* 2016, **57**, 4443;
 b) J. An, R. M. Denton, T. H. Lambert, E. D. Nacsa *Org. Biomol. Chem.* 2014, **12**, 2993;
 c) J. Otera, J. Nishikido, In *Esterification*; Wiley-VCH Verlag GmbH & Co. KGaA: 2010.
- 2 O. Mitsunobu, Y. Masaaki Bull. Chem. Soc. Jpn. 1967, 40, 2380.
- 3 a) S. Fletcher Org. Chem. Front. 2015, 2, 739; b) K. C. K. Swamy, N. N. B. Kumar, E. Balaraman, K. V. P. P. Kumar Chem. Rev. 2009, 109, 2551; c) T. Y. S. But, P. H. Toy Chem. Asian J. 2007, 2, 1340.
- 4 a) J. McNulty, A. Capretta, V. Laritchev, J. Dyck, A. J. Robertson Angew. Chem. Int. Ed. 2003, 42, 4051; b) G. Wang, J. –R. Ella-Menye, M. St. Martin, H. Yang, K. Williams Org. Lett. 2008, 10, 4203; c) A. B. Hughes, M. M. Sleebs J. Org. Chem. 2005, 70, 3079; d) A. B. Smith, I. G. Safonov, R. M. Corbett J. Am. Chem. Soc. 2002, 124, 11102; e) C. Ahn, P. DeShong J. Org. Chem. 2002, 67, 1754.
- 5 S. Schenk, J. Weston, E. Anders J. Am. Chem. Soc. 2005, **127**, 12566.
- 6 T. Mukaiyama Angew. Chem. Int. Ed. 1976, 15, 94.
- 7 A. J. Reynolds, M. Kassiou Curr. Org. Chem. 2009, 13, 1610.
- 8 T. Y. S. But, P. H. Toy J. Am. Chem. Soc. 2006, 128, 9636.
- 9 a) D. Hirose, M. Gazvoda, J. Kosmrlj, T. Taniguchi Chem. Sci. 2016, 7, 5148; b) T. Hashimoto, D. Hirose, T. Taniguchi Adv. Synth. Catal. 2015, 357, 3346; c) D. Hirose, T. Taniguchi, H. Ishibashi Angew. Chem. Int. Ed. 2013, 52, 4613.

- 10 T. Taniguchi, D. Hirose, H. Ishibashi ACS Catalysis 2011, 1. Wew Article Online DOI: 10.1039/C6OB02770A
- 11 O'Brien, C. J. PCT Int. Appl. WO2010/118042A2.2010, 2010.
- 12 J. A. Buonomo, C. C. Aldrich Angew. Chem. Int. Ed. 2015, 54, 13041.
- 13 Very recently, idea on system catalytic both from the point of view of azo-compound and phosphine based on the above mentioned recycling agents was introduced (ref.¹²) but realization is still subject of discussion; see D. Hirose, M. Gazvoda, J. Kosmrlj, T. Taniguchi *Org Lett* 2016, **18**, 4036.
- 14 a) N. A. Romero, D. A. Nicewicz Chem. Rev. 2016, 116, 10075;
 b) D. M. Schultz, T. P. Yoon Science 2014, 343, 985; c) S. Fukuzumi, K. Ohkubo Org. Biomol. Chem. 2014, 12, 6059; d) D. Ravelli, M. Fagnoni, A. Albini Chem. Soc. Rev. 2013, 42, 97.
- 15 S. Kümmel, R. Cibulka, B. König, B. In *Chemical Photocatalysis*, B. König, Ed.; de Gruyter Berlin: 2013, p 45.
- 16 a) T. Neveselý, E. Svobodová, J. Chudoba, M. Sikorski, R. Cibulka Adv. Synth. Catal. 2016, **358**, 1654; b) B. Mühldorf, R. Wolf Angew. Chem. Int. Ed. Engl. 2016, **55**, 427; c) J. B. Metternich, R. Gilmour J. Am. Chem. Soc. 2016, **138**, 1040; d) T. Hartman, R. Cibulka Org. Lett. 2016, **18**, 3710.
- S. Fukuzumi, K. Yasui, T. Suenobu, K. Ohkubo, M. Fujitsuka, O. Ito J. Phys. Chem. A 2001, 105, 10501.
- 18 G. Jürmann, O. Tšubrik, K. Tammeveski, U. Mäeorg J. Chem. Res. 2005, 661.
- 19 M. Yasuda, M.; Nakai, Y. Kawahito, T. Shiragami Bull. Chem. Soc. Jpn. 2003, 76, 601.
- 20 a) J. Daďová, S. Kümmel, C. Feldmeier, J. Cibulková, R. Pažout, J. Maixner, R. M. Gschwind, B. König, R. Cibulka *Chem. Eur. J.* 2013, **19**, 1066; b) U. Megerle, R. Lechner, B. Konig, E. Riedle, E. *Photochem. Photobiol. Sci.* 2010, **9**, 1400; c) J. Svoboda, H. Schmaderer, B. König, B. *Chem. Eur. J.* 2008, **14**, 1854.
- 21 S. Fukuzumi, K. Shimoosako, T. Suenobu, Y. Watanabe J. Am. Chem. Soc. 2003, 125, 9074.
- 22 a) S. Yasui, S. Kobayashi, M. Mishima J. Phys. Org. Chem. 2016, 29, 443; b) S. Tojo, S. Yasui, M. Fujitsuka, T. Majima J. Org. Chem. 2006, 71, 8227; c) O. Kei, N. Takashi, F. Shunichi, Bull. Chem. Soc. Jpn. 2006, 79, 1489; d) S. Yasui, S. Tojo, T. Majima J. Org. Chem. 2005, 70, 1276.
- 23 A. Ehrenberg, F. Müller, P. Hemmerich *Eur. J. Biochem.* 1967, 2, 286.
- 24 C. Feldmeier, H. Bartling, K. Magerl, R. M. Gschwind Angew. Chem. Int. Ed. 2015, 54, 1347.
- 25 Oxidation of di-*tert*-butyl hydrazinedicarboxylate to azo compound with Cul/O_2 system was reported during preparation of the manuscript: D. Jung, M. H. Kim, J. Kim *Org. Lett.* **2017**, DOI: 10.1021/acs.orglett.6b03166.
- 26 a) H. Maeda, T. Koide, T. Maki, H. Ohmori *Chem. Pharm. Bull.* 1995, 43, 1076; b) H. Ohmori, M. Maeda, M. Kikuoka, T. Maki, M. Masui, M. *Tetrahedron* 1991, 47, 767.